Check list of parasites and bacteria recorded from pangolins (*Manis* sp.)

Rajesh Kumar Mohapatra · Sudarsan Panda ·
Manoj V. Nair · Lakshmi Narayan Acharjyo

Received: 25 February 2014 / Accepted: 24 January 2015 / Published online: 14 February 2015
© Indian Society for Parasitology 2015

Abstract Sound knowledge on parasite fauna of pangolins is crucial for evaluation of their health status. In the present review, a checklist of 34 genera of parasites and bacteria, including 4 genera of protozoan, 13 genera of helminthes, 8 genera of ticks, 2 genera of mites and 7 genera of bacteria reported from pangolins was compiled and their zoonotic potential were discussed. The aim of this checklist is to underline the information gap and to provide a reference list of parasites and bacteria known for pangolins to assist in their further investigation.

Keywords Pangolin · *Manis* · Bacteria · Ecto-parasite · Endo-parasite, Zoonoses

Introduction

Pangolins are toothless placental mammals represented by eight species belonging to a single family Manidae of order

Pholidota. Four pangolin species viz. Chinese pangolin (*Manis pentadactyla*), Indian pangolin (*Manis crassicaudata*), Philippine pangolin (*Manis culionensis*) and Malayan pangolin (*Manis javanica*) are native to Asia, whereas, other four pangolin species viz. Temminck's ground pangolin (*Manis temminckii*), giant ground pangolin (*Manis gigantea*), long-tailed pangolin (*Manis tetradactyla*) and tree pangolin (*Manis tricuspis*) occur in Africa. The Asiatic pangolins such as *M. javanica* and *M. pentadactyla* are classified as critically endangered, while *M. culionensis* and *M. crassicaudata* are classified as endangered and all African pangolin are classified as vulnerable by the International Union for Conservation of Nature (Challender et al. 2014). Limited information are available about the biology, ecology, behaviour, diseases and parasites of pangolins both in captivity as well as in wild (Narayanan et al. 1977; Acharjyo 2000; Lim and Ng 2008; Challender 2009; Clark et al. 2009; Bao et al. 2013; Mohapatra and Panda 2014). Therefore, it is necessary to collect more data and carry out more field research on these animals, to promote their conservation. Considering the above necessities, the present communication is intended to compile all the available published literature on parasites and bacteria recorded from pangolins. The aim of this review is to prepare a single source to provide informations to researchers and veterinarians concerned with the parasitological and bacteriological aspects of pangolins as the pertinent sources reporting parasites and bacteria from pangolins were scattered over a number of publications covering about nine decades.

Materials and methods

Available published literatures were reviewed for compilation of informations on parasites (ecto and endo) and

R. K. Mohapatra (✉)
Pangolin Conservation Breeding Center, Nandankanan
Zoological Park, Baranga, Khurda 754005, Odisha, India
e-mail: rajesh.wildlife@gmail.com

S. Panda
Nandankanan Biological Park, Mayur Bhawan, Saheed Nagar,
Bhubaneswar 751007, Odisha, India
e-mail: sudarsanpanda61@gmail.com

M. V. Nair
Nandankanan Zoological Park,
Baranga, Khurda 754005, Odisha, India
e-mail: manojnair74@gmail.com

L. N. Acharjyo
House No.-M 71, Housing Board Colony, Bermunda,
Bhubaneswar 751003, Odisha, India
e-mail: lnacharjyo@gmail.com

bacteria reported from pangolins. Many of the report on these parasites was found incidental or constitute a part of the other studies. Inspection of wild, confiscated and captive pangolins, their routine health examinations and post mortem findings contributed to the parasitological profile of pangolins. A checklist was prepared basing on these published literatures. The major listing is grouped into endo-parasites (i.e., protozoa and helminthes), ecto-parasites (i.e., ticks and mites) and bacteria. Within these groups the parasites and bacteria species were arranged in order of their host pangolin species. Parasite species identified up to genus level (undetermined species) were also included in the checklist. In addition, the parasites and bacteria common to both pangolins and human, and their zoonotic significance were also reviewed and discussed. A draft checklist was circulated among IUCN-SSC Pangolin Specialist Group to give their comments and further information on this subject, if any. The new informations received were included in the final checklist.

Results and discussion

A total of 34 genera of parasites and bacteria, including 4 genera of protozoan, 13 genera of helminthes, 8 genera of ticks, 2 genera of mites and 7 genera of bacteria were check listed in the present review (Table 1). Out of eight pangolin species, parasites had been reported from seven species and no parasites were reported from one pangolin species (*Manis culionensis*). Even though pangolins had been on exhibits since 1892 (Sanyal 1892), the first report on parasites on pangolins dated back only to 1927 (Neveu-Lemaire 1927) and the first report on bacteria dated back to 1977 (Narayanan et al. 1977). Parasites reported from pangolins were identified to their species level in most of the cases, except eight cases where they were identified to their genus level. There were reports of six new parasite species described from pangolins, including two protozoan parasites (*Eimeria tenggilingi* and *Eimeria nkaka*), three helminth parasites (*Dipetalonema fausti*, *Leiperinema leiperi*, *Prolospirura hamospiculala*) and one mite species (*Manisicola africana*) (Else and colley 1976; Kvicerova and Hypsa 2013; Esslinger 1966; Singh 1976; Neveu-Lemaire 1927; Lawrence 1939). It appears that the present checklist is the first comprehensive review of parasites and bacteria of pangolins.

Out of four genera of protozoan parasites reported from pangolins, *Eimeria* was intestinal parasite and *Trypanosoma* and *Toxoplasma* were haemo-protozoan parasite and the site of parasitism for *Piroplasma* in pangolins could not be found from available literature (Rewell 1950; Kageruka and Willaert 1971; Else and Colley 1976; Njiokou et al. 2004, 2006; Jirku et al. 2013). Clark et al. (2009) reported

death of pangolin (*M. javanica*) secondary to gastrointestinal coccidiosis. Clinical coccidiosis is most commonly seen when animals are kept in overcrowded conditions with poor sanitation and poor nutrition (Kahn 2008). The conditions in which pangolins transported in the illegal wildlife trade are highly suited to the development of clinical coccidiosis (Clark et al. 2009). Njiokou et al. (2004) reported the occurrence of *Trypanosoma brucei* and *T. vivax* in different wild animals of Cameron including primates, artiodactyls, rodents, carnivores and pangolins. Such observations strengthen the hypothesis that wild animals including pangolins may serve as reservoirs for Trypanosomiasis infections (Njiokou et al. 2004). Toxoplasmosis caused by *Toxoplasma gondii* is a potential zoonotic disease reported from animal hosts including human and pangolin (Kageruka and Willaert 1971; Tenter et al. 2000; Hill and Dubey 2002).

Out of 13 reported helminth parasites from pangolins, eight genera of helminths including, *Cylicospirura*, *Leiperinema*, *Manistrongylus*, *Necator*, *Strongyloides*, *Trichocheenia*, *Ancylostoma*, *Gendrespirura* were intestinal parasites, occurring in stomach, small intestine and large intestine (Naidu and Naidu 1981; Sharma 1955; Heath and Vanderlip 1988; Baylis 1933; Hanna 1933; Esslinger 1966) and three genera including *Dipetalonema*, *Brugia* and *Microfilaria* were blood parasites (Pais Caeiro 1959; Laing et al. 1960). The helminth *Brugia malayi* was reported in a wide range of hosts including long tailed macaque, leaf-monkey, cat, civet cat, pangolin (*M. javanica*) and human in Southeast Asia (Wilson 1961; Kazura 2002). Similarly *Brugia pahangi* was reported from cat, civet cats, dog, tiger, slow loris, moonrat, giant squirrel and pangolin (*M. javanica*) in Malaysia (Wilson 1961). The helminth *Necator americanus* was reported both from pangolin (*M. crassicaudata* and *M. javanica*) and human (Baylis 1933; Sharma 1955; Ashford and Crewe 2003; Hasegawa et al. 2014). Besides, there were records of *N. americanus* as an occasional parasite of the chimpanzee, gorilla, monkey, dog, pig and rhinoceros (Hanna 1933). Hydatid cysts by the *Echinococcus* sp. also reported from lungs of an Indian pangolin (Rao et al. 1972; Acharjyo 2013).

Ticks are well represented in pangolins comprising 8 genera and 20 species reported from six pangolin species (*M. crassicaudata*, *M. pentadactyla*, *M. javanica*, *M. temminckii*, *M. tricuspis*, *M. gigantea*). Genus *Amblyomma* was reported from all the above six pangolin species. It is interesting to know that unidentified ticks had been reported from *M. tetradactyla* (Robinson 1983). Three genera of ticks (*Amblyomma*, *Rhipicephalus* and *Aponomma*) reported from both Asian and African pangolin species. *Amblyomma javanense* found to occur in three Asian pangolins (*M. crassicaudata*, *M. pentadactyla*, *M. javanica*). Similarly *Amblyomma compressum* recorded

Table 1 Parasites and bacteria reported from pangolins (*Manis* sp.)

Pangolin host (scientific name)	Parasite species	Reference (s)
Endo-parasites		
Protozoa		
Indian pangolin (<i>Manis crassicaudata</i>)	<i>Toxoplasma gondii</i>	Kageruka and Willaert (1971)
Long-tailed pangolin (<i>Manis tetradactyla</i>)	<i>Trypanosoma brucei</i>	Njiokou et al. (2004, 2006)
	<i>Trypanosoma vivax</i>	Njiokou et al. (2004)
Malayan pangolin (<i>Manis javanica</i>)	<i>Eimeria tenggilingi</i> (n. sp.)	Else and Colley (1976)
Temminck's ground pangolin (<i>Manis temminckii</i>)	<i>Piroplasma</i> sp.	Rewell (1950)
Tree pangolin (<i>Manis tricuspis</i>)	<i>Eimeria</i> sp.	Kvicerova and Hypsa (2013)
	<i>Eimeria nkaka</i> (n. sp.)	Jirku et al. (2013)
	<i>Trypanosoma brucei</i>	Njiokou et al. (2004, 2006)
	<i>Trypanosoma vivax</i>	Njiokou et al. (2004)
Helminths		
Chinese pangolin (<i>Manis pentadactyla</i>)	<i>Cylicospirura</i> sp.	Heath and Vanderlip (1988)
	<i>Dipetalonema fausti</i> (n. sp.)	Esslinger (1966)
	<i>Leiperinema leiperi</i> (n. sp.)	Singh (1976)
	<i>Manistrongylus meyeri</i> (Travassos 1937)	Cameron and Myers (1960)
	<i>Necator americanus</i> (Stiles 1902)	Cameron and Myers (1960)
	<i>Strongyloides</i> sp.	Singh (1976), Heath and Vanderlip (1988)
Indian pangolin (<i>Manis crassicaudata</i>)	<i>Echinococcus</i> sp. (Hydatid cysts)	Rao et al. (1972), Acharjyo (2013)
	<i>Necator americanus</i> (Stiles 1902)	Sharma (1955)
	<i>Trichocheenia meyeri</i> (Travassos 1937)	Naidu and Naidu (1981)
Giant ground pangolin (<i>Manis gigantea</i>)	<i>Ancylostoma</i> sp. (ova)	T-W-Fiennes (1968)
Malayan pangolin (<i>Manis javanica</i>)	<i>Brugia malayi</i>	Laing et al. (1960)
	<i>Brugia pahangi</i>	Laing, et al. (1960)
	<i>Necator americanus</i> (Stiles 1902)	Baylis (1933)
	<i>Gendrespirura</i> sp.	Gardiner and Werner (1984)
Temminck's ground pangolin (<i>Manis temminckii</i>)	<i>Protospirura</i> (<i>Habronema</i>) <i>hamospiculata</i> (n. sp.)	Neveu-Lemaire (1927), Baylis (1931)
Tree pangolin (<i>Manis tricuspis</i>)	<i>Habronema hamospiculatum</i> (Neveu-Lemaire 1927)	Baylis (1931, 1936)
	<i>Microfilaria lukakae</i>	Pais Caeiro (1959)
	<i>Microfilaria lundae</i>	Pais Caeiro (1959)
	<i>Microfilaria nobrei</i>	Pais Caeiro (1959)
	<i>Microfilaria vilhenae</i>	Pais Caeiro (1959)
Ecto-parasites		
Ticks		
Chinese pangolin (<i>Manis pentadactyla</i>)	<i>Amblyomma javanense</i> (Supino 1897)	Li et al. (2010), Kollars and Sithiprasasna (2000)
Giant ground pangolin (<i>Manis gigantea</i>)	<i>Amblyomma compressum</i> (Macalister 1872)	Uilenberg et al. (2013)
Indian pangolin (<i>Manis crassicaudata</i>)	<i>Amblyomma javanense</i> (Supino 1897)	Sanyal et al. (1987), Nandi (1981), Acharjyo (2013)
	<i>Aponomma gerviasi</i> (Lucas 1847)	Seniviratna (1965), Pillai and George (1997)
	<i>Rhipicephalus</i> sp.	Anonymous (2010)
Malayan pangolin (<i>Manis javanica</i>)	<i>Amblyomma cordiferum</i> (Neumann 1899)	Hafiz et al. (2012)
	<i>Amblyomma javanense</i> (Supino 1897)	Kollars and Sithiprasasna (2000), Parola et al. (2003), Hafiz et al. (2012), Hassan et al. (2013)

Table 1 continued

Pangolin host (scientific name)	Parasite species	Reference (s)
Temminck's ground pangolin (<i>Manis temminckii</i>)	<i>Aponomma varanensis</i> (Supino 1897)	Burridge (2001)
	<i>Amblyomma compressum</i> (Macalister 1872)	Uilenberg et al. (2013)
	<i>Ixodes rarus</i> (Neumann 1899)	Uilenberg et al. (2013)
	<i>Ornithodoros moubata</i> (Murray 1877)	Jacobsen et al. (1991)
	<i>Rhipicentor longus</i> Neumann 1907	Uilenberg et al. (2013)
	<i>Rhipicentor muhsamae</i> (Morel and Vassiliades 1964)	Uilenberg et al. (2013)
Tree pangolin (<i>Manis tricuspis</i>)	<i>Rhipicephalus simus</i> (Koch 1844)	Tandon (1991)
	<i>Amblyomma compressum</i> (Macalister 1872)	Ntiamoa-Baidu et al. (2005), Uilenberg et al. (2013)
	<i>Aponomma exornatum</i> (Koch 1844)	Burridge (2001)
	<i>Aponomma flavomaculatum</i> (Lucas 1846)	Burridge (2001)
	<i>Aponomma latum</i> (Koch 1844)	Burridge (2001)
	<i>Haemaphysalis parvata</i> (Neumann 1905)	Ntiamoa-Baidu et al. (2005)
	<i>Rhipicentor muhsamae</i> (Morel and Vassiliades 1964)	Uilenberg et al. (2013)
Unidentified pangolin species (<i>Manis</i> sp.)	<i>Amblyomma arcanum</i> (Karsch 1879)	Guglielmone et al. (2014)
	<i>Amblyomma cordiferum</i> (Neumann 1899)	Guglielmone et al. (2014)
	<i>Amblyomma geoemydae</i> (Cantor 1847)	Guglielmone et al. (2014)
	<i>Amblyomma testudinarium</i> (Koch 1844)	Durden et al. (2008)
	<i>Dermacentor (Indocentor) atrosignatus</i> (Neumann 1906)	Durden et al. (2008)
	<i>Dermacentor (Indocentor) steini</i> (Schulze 1933)	Durden et al. (2008)
	<i>Ixodes oldi</i> (Nuttall 1913)	Guglielmone et al. (2014)
Mite		
Temminck's ground pangolin (<i>Manis temminckii</i>)	<i>Manicola africana</i> (n. sp.)	Lawrence (1939)
	<i>Manitherionyssus heterotarsus</i> (Vitzthum 1925)	Jacobsen et al. (1991)
Bacteria		
Chinese pangolin (<i>Manis pentadactyla</i>)	<i>Escherichia coli</i>	Heath and Vanderlip (1988)
	<i>Klebsiella aerogenes</i>	Heath and Vanderlip (1988)
	<i>Klebsiella pneumoniae</i>	Heath and Vanderlip (1988)
	<i>Proteus vulgaris</i>	Heath and Vanderlip (1988)
	<i>Pseudomonas fluorescens</i>	Heath and Vanderlip (1988)
Indian pangolin (<i>Manis crassicaudata</i>)	<i>Escherichia coli</i>	Narayanan et al. (1977)
	<i>Klebsiella aerogenes</i>	Narayanan et al. (1977)
	<i>Proteus vulgaris</i>	Narayanan et al. (1977)
	<i>Pseudomonas aeruginosa</i>	Narayanan et al. (1977)
	<i>Staphylococcus</i> sp.	Narayanan et al. (1977)
	<i>Streptococcus faecalis</i>	Narayanan et al. (1977)
Malayan pangolin (<i>Manis javanica</i>)	<i>Mycoplasma</i> sp.	Jamnah et al. (2014)

from three African pangolin species (*M. temminckii*, *M. tricuspis*, *M. gigantea*). Ticks were reported to occur underneath the scales and ventral portions of the body in pangolins (Seniviranta 1965; Sanyal et al. 1987; Pillai and George 1997; Parola et al. 2003; Anonymous 2010; Hafiz et al. 2012). Apart from pangolins *Amblyomma javanense* ticks had been reported from water monitor lizard, python, skink, hill turtle, bat, wild boar, hyena, bear and sambar deer (Kollars and Sithiprasasna 2000; Hassan 2013). Similarly there were reports of *Amblyomma latum* ticks from Python; *Amblyomma exornatum* ticks from monitor lizards; *Rhipicephalus simus* ticks from dogs (Navajas et al. 2010; Reeves et al. 2006; Parola et al. 2013). Guglielmone et al. (2014) reported ten species of ticks found in pangolins also occur in different animal hosts (classified up to family level) including mammals, birds and reptiles. Some of the ticks reported from pangolins were also reported to feed on humans and may serve as potential vectors for transmission of disease. The tick *Ornithodoros moubata*, transmits *Borrelia duttoni*, the agent of African relapsing fever in human (Estrada-Pena and Jongejan 1999). In Sub-Saharan Africa, *Amblyomma compressum* ticks transmit African tick bite fever caused by *Rickettsia africae*; *Rhipicephalus simus* ticks transmit Mediterranean spotted fever caused by *Rickettsia conorii* subsp. *conorii*; *Rhipicephalus muhsamae* ticks transmit spotted fever caused by *Rickettsia massiliae* in human (Parola et al. 2013). The bacteria *Anaplasma* sp. strain AnAj360 associated with the ehrlichiosis disease was also detected from the tick *Amblyomma javanense* collected from *Manis javanica* (Parola et al. 2003). Besides, *Aponomma latum*, *Amblyomma cordiferum*, *Amblyomma geoemydae*, *Amblyomma javanense*, *Amblyomma testudinarium*, *Dermacentor atrosignatus* and *Rhipicephalus longus*, recorded from pangolins, have also been reported to feed on human (Estrada-Pena and Jongejan 1999; Durden et al. 2008; Burridge 2001; Audy et al. 1960; Gould et al. 1970). Mites recorded from pangolins were reported from the body surface (Jacobsen et al. 1991).

Some bacteria were isolated and identified from faeces, blood and urine samples of the diseased Indian pangolin (Narayanan et al. 1977) and Chinese pangolin (Heath and Vanderlip 1988). The bacteria species *Escherichia coli*, *Klebsiella aerogenes*, *Proteus vulgaris* were recorded from both Indian pangolin (Narayanan et al. 1977) and Chinese pangolin (Heath and Vanderlip 1988). Bacteria *Escherichia coli* were recorded from faeces, blood and urine of diseased Indian pangolin (Narayanan et al. 1977). *Klebsiella aerogenes* and *Klebsiella pneumonia*, *Proteus vulgaris*, *Pseudomonas fluorescens*, *Pseudomonas aeruginosa* were isolated from faecal samples of pangolins (Narayanan et al. 1977; Heath and Vanderlip 1988). *Sterptococcus* was isolated from faeces and blood of Indian pangolins

(Narayanan et al. 1977). *Mycoplasma* was isolated from blood of Malayan pangolin (Jamnah et al. 2014). Bacteria are ubiquitous in nature and some bacteria reported from pangolin also found as commensal organisms of human gastrointestinal tract (e.g. *Proteus vulgaris*, *Klebsiella pneumonia*, *Escherichia coli*) (Muller 1986; Lightfoot 2003). *Pseudomonas aeruginosa* and *Klebsiella pneumonia* are opportunistic pathogen in human that causes a variety of bronchopulmonary infections including pneumonia (Domingo et al. 1998; Lightfoot 2003). Though most *Escherichia coli* are not pathogenic, but some serotypes possess virulence factors and cause gastroenteritis in humans (Lightfoot 2003).

Acknowledgments Authors are thankful to Professor Wu Shibao, School of Life Science, South China Normal University, China; Dr Ranjana Jaiswara, Post Doctoral Fellow, Museum national d'Histoire naturelle, France; Gopinath Panda, Research Fellow, Institute of Physics, Bhubaneswar, India; Suraj Rajan Prasanjit, BS-MS Student, Indian Institute of Science Education and Research, Kolkata, India; Helen Nash, PhD candidate, National University of Singapore, Singapore; Louise Fletchere, Field Researcher, Carnivore and Pangolin Conservation Program, Vietnam for making available some relevant literatures for the present review. We are also thankful to members of IUCN-SSC Pangolin Specialist Group for their valuable comments on earlier draft of the checklist.

References

- Acharjyo LN (2000) Management of Indian pangolin in captivity. In: The managing committee (eds) Souvenir, 125 years of Calcutta zoo (1875-2000). Zoological Garden, Alipore, Calcutta. pp 27–30
- Acharjyo LN (2013) Check list of parasites recorded from wild animals and birds of Nandankanan Zoological Park, Odisha. In: Parida GS, Panda N, Mohanty B, Das D, Behura K, Dehuri M (eds) Souvenir-National symposium 'Newer approaches in welfare and health management of captive and free ranging wild animals', 19-12 April, 2013. Orissa University of Agriculture Technology and Association of Indian Zoo and Wildlife Veterinarians, Bhubaneswar, pp 15–29
- Anonymous (2010) Project report. Ecto and endo parasites of captive animals and birds of Nandankanan zoo. Nandankanan Zoological Park and Orissa Veterinary College, Bhubaneswar (Supported by Central Zoo Authority, New Delhi). pp 1–79
- Ashford RW, Crewe W (2003) The parasites of Homo sapiens. Taylor & Francis, London, pp 1–142. doi:10.1007/springerreference_227283
- Audy JR, Nadchatram M, Lim B-L (1960) Malaysian parasites. XLIX. Host distribution of Malayan ticks (Ixodoidea). Stud Inst Med Res Malaya 29:225–246
- Bao F, Wu S, Su C, Yang L, Zhang F, Ma G (2013) Air temperature changes in a burrow of Chinese pangolin, *Manis pentadactyla*, in winter. Folia Zool 62(1):42–47
- Baylis HA (1931) On a nematode parasite of pangolin. Ann Mag Nat Hist 10(8):191–194. doi:10.1080/00222933108673379
- Baylis HA (1933) On some parasitic worms from Java, with remarks on the Acanthocephalan genus *Pallisentis*. Ann Mag Nat Hist 12(70):443–449. doi:10.1080/00222933308673708

- Baylis HA (1936) Some parasitic worms from the British Cameroons. *J Nat Hist Ser* 17(10):257–272. doi:[10.1080/00222933608655118](https://doi.org/10.1080/00222933608655118)
- Burridge MJ (2001) Ticks (Acari: Ixodidae) spread by the international trade in reptiles and their potential roles in dissemination of diseases. *Bull Entomol Res* 91:3–23
- Cameron TWM, Myers BJ (1960) *Manistrongylus meyeri* (Travassos, 1937) Gen. Nov., and *Necator americanus* from the pangolin. *Can J Zool* 38(4):781–786. doi:[10.1139/z60-082](https://doi.org/10.1139/z60-082)
- Challender DWS (2009) Asian Pangolins: how behavioural research can contribute to their conservation. In: Pantel S, Chin SY (eds) Proceedings of the workshop on trade and conservation of pangolins native to south and southeast Asia. Singapore Zoo, Singapore. TRAFFIC southeast Asia, Petaling Jaya, Selangor, Malaysia, 30 Jun–2 Jul 2008, pp 95–102
- Challender DWS, Waterman C, Baillie JEM (2014) Scaling up pangolin conservation. IUCN SSC Pangolin specialist group conservation action plan. Zoological Society of London, London
- Clark L, Thai NV, Phuong TQ (2009) A long way from home: the health status of Asian pangolins confiscated from the illegal wildlife trade in Viet Nam. In: Pantel S, Yun CS (eds) Proceedings of the workshop on trade and conservation of Pangolins native to south and southeast Asia, Singapore Zoo, Singapore. TRAFFIC southeast Asia, Petaling Jaya, Selangor, Malaysia, 30 Jun–2 Jul 2008, pp 111–118
- Domingo P, Ferre A, Baraldes MA, Ris J, Sanchez F (1998) *Pseudomonas aeruginosa* bronchopulmonary infection in patients with AIDS, with emphasis on relapsing infection. *Eur Respir J* 12:107–112. doi:[10.1183/09031936.98.12010107](https://doi.org/10.1183/09031936.98.12010107)
- Durden LA, Merker S, Beati L (2008) The tick fauna of Sulawesi, Indonesia (Acari: Ixodoidea: Argasidae and Ixodidae). *Exp Appl Acarol* 45:85–110. doi:[10.1007/s10493-008-9144-z](https://doi.org/10.1007/s10493-008-9144-z)
- Else JG, Colley FC (1976) *Eimeria tenggilingi* sp. n. from the scaly anteater *Manis javanica* Desmarest in Malaysia. *J Eukaryot Microbiol* 23(4):487–488. doi:[10.1111/j.1550-7408.1976.tb03820.x](https://doi.org/10.1111/j.1550-7408.1976.tb03820.x)
- Esslinger JH (1966) *Dipetalonema Fausti* sp. n. (Filarioidea: Onchocercidae), a filarial parasite of the scaly anteater, *Manis pentadactyla* L. (Pholidota), from China. *J Parasitol* 54(3):494–497. doi:[10.2307/3276315](https://doi.org/10.2307/3276315)
- Estrada-Pena A, Jongejan F (1999) Ticks feeding on humans: a review of records on human-biting Ixodoidea with special reference to pathogen transmission. *Exp Appl Acarol* 23:685–715
- Gardiner CH, Werner RM (1984) Polypoid gastritis in a scaly anteater caused by larvae of *Gendrespirura* sp. *J Wildl Dis* 20(1):68–70. doi:[10.7589/0090-3558-20.1.68](https://doi.org/10.7589/0090-3558-20.1.68)
- Gould DJ, Clifford CM, Lakshana P et al. (1970) Distribution and ecology of ectoparasites of vertebrates in Southeast Asia. Annual Program Report SEATO Medical Research Laboratory, United States Army Medical Component, Bangkok, Apr 1969–Mar 1970, pp 97–105
- Guglielmone AA, Robbins RG, Apanaskevich DA, Petney TN, Estrada-Pena A, Horak IG (2014) The hard ticks of the world (Acari: Ixodida: Ixodidae). Springer, Dordrecht Heidelberg, New York. doi:[10.1007/978-94-007-7497-1_22](https://doi.org/10.1007/978-94-007-7497-1_22)
- Hafiz MS, Marina H, Afzan AW, Chong JL (2012) Ectoparasite from confiscated Malayan pangolin (*Manis javanica* Desmarest) in peninsular Malaysia. UMT 11th International annual symposium on sustainability science and management, Terengganu, Malaysia, 09–11 Jul 2012, pp. 1225–1229
- Hanna GD (1933) The dates of publication of ‘Synopsis des Diatomées de Belgique,’ by Henri Van Heurck. *Ann Mag Nat Hist* 12(70): 442–443 doi:[10.1080/00222933308673707](https://doi.org/10.1080/00222933308673707)
- Hasegawa H, Modry D, Kitagawa M, Shutt KA, Todd A et al (2014) Humans and great apes cohabiting the forest ecosystem in Central African Republic Harbour the same hookworms. *PLoS*. doi:[10.1371/journal.pntd.0002715](https://doi.org/10.1371/journal.pntd.0002715)
- Hassan M, Hafiz MS, Chong JL (2013) The prevalence and intensity of *Amblyomma javanense* infestation on Malayan pangolins (*Manis javanica* Desmarest) from Peninsular Malaysia. *Acta Trop* 126:142–145. doi:[10.1016/j.actatropica.2013.02.001](https://doi.org/10.1016/j.actatropica.2013.02.001)
- Heath ME, Vanderlip SL (1988) Biology, husbandry, and veterinary care of captive Chinese pangolins (*Manis pentadactyla*). *Zoo Biol* 7:293–312. doi:[10.1002/zoo.1430070402](https://doi.org/10.1002/zoo.1430070402)
- Hill D, Dubey JP (2002) *Toxoplasma gondii*: transmission, diagnosis and prevention. *Clin Microbiol Infect* 8(10):634. doi:[10.1046/j.1469-0691.2002.00485.x](https://doi.org/10.1046/j.1469-0691.2002.00485.x)
- Jacobsen NHG, Newbery RE, Dewet MH, Viljoen PC, Pietersen E (1991) A contribution to the ecology of the steppe pangolin *Manis temminckii* in the transvaal. *Ziet-schrift fur Saugetierkunde* 56:94–100
- Jamnah O, Faizal H, Chandrawathani P, Premaalatha B, Erwanas AI, Lily R, Ramlan M (2014) Eperythrozoonosis (*Mycoplasma* sp.) in Malaysian pangolin. *Malays J Vet Res* 5(1):65–69
- Jirku M, Kvcierova J, Modry D, Hypsa V (2013) Evolutionary plasticity in coccidia—striking morphological similarity of unrelated coccidia (Apicomplexa) from related hosts: *Eimeria* spp. from African and Asian pangolins (Mammalia:Pholidota). *Protist* 164:470–481. doi:[10.1016/j.protis.2013.04.001](https://doi.org/10.1016/j.protis.2013.04.001)
- Kageruka P, Willaert E (1971) *Toxoplasma gondii* (Nicolle et Manceaux 1908) isole chez Goura cristata Pallas et *Manis crassicaudata* Geoffroy. *Acta Zool Pathol Antverp* 52:3–10
- Kahn CM (ed) (2008) Merck veterinary manual, 9th edn. Merck and Co. Ltd, Whitehouse Station, NJ
- Kazura JW (2002) Lymphatic filarial infections: an introduction to the filariae. In: Klei TR, Rajan TV (eds) The filaria. Kluwer, New York, pp 1–8. doi:[10.1007/0-306-47661-4_1](https://doi.org/10.1007/0-306-47661-4_1)
- Kollars TM, Sithiprasasna R (2000) New host and distribution record of *Amblyomma javanense* (Acari: Ixodidae) in Thailand. *J Med Entomol* 37(4):640. doi:[10.1603/0022-2585-37.4.640](https://doi.org/10.1603/0022-2585-37.4.640)
- Kvcierova J, Hypsa V (2013) Host-parasite incongruences in rodent *Eimeria* suggest significant role of adaptation rather than copyphylogeny in maintenance of host specificity. *PLoS One* 8(7):e63601. doi:[10.1371/journal.pone.0063601](https://doi.org/10.1371/journal.pone.0063601)
- Laing ABG, Edeson JFB, Wharton RH (1960) Studies on filariasis in Malaya: the vertebrate hosts of *Brugia malayi* and *Brugia pahangi*. *Ann Trop Med Parasit* 53(4):92–99
- Lawrence RF (1939) A new mite from the South African Pangolin. *Parasitology* 31(04):451–457. doi:[10.1017/s0031182000012968](https://doi.org/10.1017/s0031182000012968)
- Li Y, SU C, Fu-hua Z, Shi-bao W, Guang-zhi M (2010) Age structure and parasites of Malayan Pangolin (*Manis javanica*). *J Econ Anim* 14(1):22–25
- Lightfoot NF (2003) Bacteria of potential health concern. In: Bartram J, Cotruvo J, Exner M, Fricker C, Glasmacher A (eds) Heterotrophic plate counts and drinking-water safety. pp 119–137
- Lim NTL, Ng PKL (2008) Home range, activity cycle and natal den usage of a female Sunda pangolin *Manis javanica* (Mammalia: Pholidota) in Singapore. *Endanger Species Res* 4:233–240. doi:[10.3354/esr00032](https://doi.org/10.3354/esr00032)
- Mohapatra RK, Panda S (2014) Husbandry, behaviour and conservation breeding of Indian pangolin. *Folia Zool* 63(2):73–80
- Muller HE (1986) Occurrence and pathogenic role of Morganella-Proteus-Providencia group bacteria in human feces. *J Clin Microbiol* 23(2):404–405
- Naidu KV, Naidu KA (1981) *Trichocheenia meyeri* (Travassos 1937) Naidu K. V. and Naidu K. A. Comb. Nov. (Nematoda: Trichostrongylidae) from pangolin in south India. *Proc Indian Acad Sci (Anim Sci)* 90(6):615–618
- Nandi NC (1981) *Amblyomma javanense* (Supino, 1897) from a new host, *Manis crassicaudata* Gray together with a description of its nymph from Goa, India. *Bull Zool Surv India* 4(1):23–26

- Narayanan SE, Kirchheimer WF, Bedi BMS (1977) Some bacteria isolated from the Indian pangolin (*Manis crassicaudata*), Geoffroy. Indian Vet J 54(9):692–988
- Navajas M, Migeon A, Estrada-Pena A, Mailleux AC, Servigne P, Petanovic R (2010) Mites and ticks (Acari). In: Roques A et al. (eds) Arthropod invasions in Europe. Bio risk, vol 4, pp 149–192. doi: [10.3897/biorisk.4.58](https://doi.org/10.3897/biorisk.4.58)
- Neveu-Lemaire M (1927) *Protophysa hamospiculata* n. sp. nematode parasite d'un pangolin africain *Manis (Pholodatus) temminckii*. Ann Parasit 5:107–113
- Njiokou F, Simo G, Nkinin SW, Laveissiere C, Herder S (2004) Infection rate of *Trypanosoma brucei* s.l., *T. vivax*, *T. congolense* “forest type”, and *T. simiae* in small wild vertebrates in south Cameroon. Acta Trop 92:139–146. doi: [10.1016/j.actatropica.2004.04.011](https://doi.org/10.1016/j.actatropica.2004.04.011)
- Njiokou F, Laveissiere C, Simo G, Nkinin S, Grebaut P, Cuny G, Herder S (2006) Wild fauna as a probable animal reservoir for *Trypanosoma brucei gambiense* in Cameroon. Infect Genet and Evol 6:147–153. doi: [10.1016/j.meegid.2005.04.003](https://doi.org/10.1016/j.meegid.2005.04.003)
- Ntiemoa-Baidu Y, Carr-Saunders C, Matthews BE, Preston PM, Walker AR (2005) Ticks associated with wild mammals in Ghana. Bull Entomol Res 95:205–219. doi: [10.1079/ber.2004.352](https://doi.org/10.1079/ber.2004.352)
- Pais Caeiro VM (1959) Quatro especies de microfilarias do *Phataginus manis tricusps* (Rafinesque). An Escola Superior Med Vet Lisbon 2:83–94
- Parola P, Cornet J, Sanogo Y et al (2003) Detection of *Ehrlichia* spp, *Anaplasma* spp, *Rickettsia* spp, and other eubacteria in ticks from the Thai-Myanmar border and Vietnam. J Clin Microbiol 41:1600–1608. doi: [10.1128/jcm.41.4.1600-1608.2003](https://doi.org/10.1128/jcm.41.4.1600-1608.2003)
- Parola P, Paddock CD, Socolovschi C, Labruna MB, Mediannikov O, Kernif T, Abdad MY, Stenos J, Bitam I, Fournier P-E, Raoult D (2013) Update on tick-borne rickettsioses around the world: a geographic approach. Clin Microbiol Rev 26(4):657–702. doi: [10.1128/CMR.00032-13](https://doi.org/10.1128/CMR.00032-13)
- Pillai KM, George PO (1997) Tick infestation in an Indian pangolin (*Manis crassicaudata*). Indian Vet J 74(1):71–72
- Rao AT, Misra SC, Acharjyo LN (1972) Pulmonary hydatidosis in captive animals at Nandankanan Zoo. Indian Vet J 49(8): 842–843
- Reeves WK, Durden LA, Dasch GA (2006) A spotted fever group Rickettsia from an exotic tick species, *Amblyomma exornatum* (Acari: Ixodidae), in a reptile breeding facility in the United States. J Med Entomol 43(5):1099–1101
- Rewell RE (1950) Report of the Society's pathologist for the year 1949. Proc Zool Soc London 120(3):485–495. doi: [10.1111/j.1096-3642.1950.tb00658.x](https://doi.org/10.1111/j.1096-3642.1950.tb00658.x)
- Robinson PT (1983) The use of ketamine in restraint of a black-bellied pangolin (*Manis tetradactyla*). J Zoo Anim Med 14:19–23. doi: [10.2307/20094623](https://doi.org/10.2307/20094623)
- Sanyal RB (1892) A hand book of the management of animals in captivity in lower Bengal. Secretariat Press, Calcutta
- Sanyal AK, De SK, Rao AT, Acharjyo LN (1987) Metastigmata (Ticks), Fauna of Orissa. Publ. Zool. Surv., India. State Fauna Series No. 1, pp 273–287
- Seneviratna P (1965) The Ixodoidea (ticks) of Ceylon. Ceylon Vet J 13:28–54
- Sharma SN (1955) *Necater americanus* sp. in an Indian pangolin. Curr Sci 24(8):287
- Singh SN (1976) On a new nematode *Leiperinema leiperi* n.g., n.sp. (Strongyloididae), parasitic in the pangolin *Manis pentadactyla* from Hyderabad, India. J Helminthol 50:267–274. doi: [10.1017/s0022149x00026705](https://doi.org/10.1017/s0022149x00026705)
- Tandon SK (1991) The Ixodid tick of Zambia (Acarina: Ixodidae) (A study on distribution, ecology and host relationship). Director, Zoological Survey of India (eds) Records of Zoological Society of India. Occasional paper No. 134. Zoological Survey of India Calcutta, pp 1–175
- Tenter AM, Heckeroth AR, Weiss LM (2000) *Toxoplasma gondii*: from animals to humans. Int J Parasitol 30(12–13):1217–1258. doi: [10.1016/0264-410x\(85\)90041-6](https://doi.org/10.1016/0264-410x(85)90041-6)
- T-W-Fiennes RN (1968) Report of the Society's Pathologist for the year 1965. Proc Zool Soc Lond 154:173–191. doi: [10.1111/j.1469-7998.1968.tb01658.x](https://doi.org/10.1111/j.1469-7998.1968.tb01658.x)
- Uilenberg G, Estrada-Pena A, Thal J (2013) Ticks of the Central African Republic. Exp Appl Acarol 60(1):1–40. doi: [10.1007/s10493-012-9605-2](https://doi.org/10.1007/s10493-012-9605-2)
- Wilson T (1961) Filariasis in Malaya—a general review. Trans R Soc Trop Med Hyg 55(2):107–129