ABSTRACTS

Neuro-Ophthalmic Literature Review

Carmen K. M. Chan, Evan Price, John H. Pula, Michael Vaphiades,
An-Guor Wang, Sui Wong

CLINICAL CHARACTERISTICS OF PARAINFECTIOUS OPTIC NEURITIS IN CHILDREN AND ADULTS

Rappoport D, Goldenberg-Cohen N, Luckman J, Leiba H. Parainfectious optic neuritis: manifestations in children vs adults. *J Neuroophthalmol* 2014;34:122–129.

In this retrospective case series, the authors described the clinical features of 18 patients with parainfectious optic neuritis who presented to two neuro-ophthalmology practices in Israel between 2005 and 2012. In this study, parainfectious optic neuritis was defined as acute optic neuritis within 1 month after an infectious disease. Patients with multiple sclerosis and other causes of optic neuropathy were excluded. Of the 18 patients, 10 were children (mean age 12.7 years, 50% bilateral) and 8 adults (mean age 31 years, 38% bilateral). The male:female ratios in both groups were 1:1. All the patients had disc swelling. The most common pathogen for the preceding infection was *Mycoplasma pneumoniae* in the paediatric group, whereas no organism predominated in the adult group. Acute disseminated encephalomyelitis was diagnosed in 4/10 children but none in adults. The final visual outcome was good (20/30 or better) in all patients and was not affected by whether the patient was treated with corticosteroids or not. None of the patients had a recurrence within the follow-up period. This study demonstrated that the clinical course of parainfectious optic neuritis is quite different from that of demyelinating optic neuritis.

Carmen Chan

CHILDREN WITH OCULAR TICS—WHAT HAPPENS TO THEM IN THE LONG TERM?

Bisker ER, McClelland CM, Brown LW, Liu GT. The long-term outcomes of ocular tics in a pediatric neuro-ophthalmology practice. *J AAPOS* 2014;18:31–35.

Tics are involuntary, rapid, repetitive, purposeless, and stereotyped intermittent muscle movements or vocalizations. They are not associated with alterations in consciousness and can partially be suppressed voluntarily. Although transient tic disorder has been described in up to 3% of school-aged children, the symptoms can still be alarming for the parents. Therefore, it would be very helpful to let them know the long-term clinical outcome of these children. In this retrospective case series, the authors described the clinical features of 43 children (63% boys) who presented to a paediatric neuro-ophthalmology practice with idiopathic ocular tics, and provided follow-up information for 32 of them (mean follow-up period 6.1 years). The mean age of presentation was 7.8 years, and the most common ocular tics were eye blinking and eye rolling. Amongst the group with follow-up information, 14/32 (44%) had persistent ocular tics during the follow-up period, 10 of which had improvement. Out of the 32 patients, 1 was diagnosed with Tourette Syndrome and 3 diagnosed with attention deficit hyperactive disorder; all

4 were children whose ocular tics resolved. None developed obsessive compulsive disorder. Three children (9%) reported new non-ocular motor tics and 5 (16%) developed new vocal tics.

Carmen Chan

PATIENTS WITH OPTIC PATHWAY GLIOMAS NEED LONG-TERM FOLLOW-UP

Shofty B, Constantini S, Bokstein F, Ram Z, Ben-Sira L, Freedman S, Vainer G, Kesler A. Optic pathway gliomas in adults. *Neurosurgery* 2014;74:273–279; discussion 279–280.

Optic pathway gliomas (OPG) are generally considered benign when diagnosed in childhood, especially when associated with neurofibromatosis type 1 (NF-1), with only about 30% of patients experiencing visual impairment. The onset of OPG classically follows a bimodal distribution, with one peak below 7 years of age, and another in the 50s. In this retrospective case series, the authors described the clinical characteristics of 22 adults who were followed up in a tertiary referral centre for neurosurgery in Israel between 1990 and 2012. The patients were divided into 2 groups: group A (12 patients, mean follow-up 9.5 years) who were diagnosed in childhood; and group B (10 patients, mean follow-up 13.8 years) who were diagnosed in adulthood. 7/12 patients and 5/10 patients had NF-1 in groups A and B, respectively. In group A, 6 patients who had tumours that were stable or regressed in childhood also remained stable in adulthood. Of the remaining 6 patients whose tumour progressed in childhood, 3 suffered additional tumour progression or visual decline in adulthood despite earlier treatment (one with NF-1). Of the 10 patients in group B, 8 had low-grade tumours, 4 (1 with NF-1) of which progressed. Two patients in group B had high-grade gliomas and had rapid progression. The authors recommended prolonged monitoring for paediatric OPG patients who had progression in childhood, and for adult OPG patients, even if they had NF-1. The authors also admitted that the sample could be biased towards the more severe cases, since these patients were seen in a tertiary neurosurgical centre.

Carmen Chan

REDUCING B CELLS WITH RITUXIMAB IN ORBITAL INFLAMMATORY SYNDROME

Suhler EB, Lim LL, Beardsley RM, Giles TR, Pasadhika S, Lee ST, de Saint Sardos A, Butler NJ, Smith JR, Rosenbaum JT. Rituximab therapy for refractory orbital inflammation: results of a phase 1/2, dose-ranging, randomized clinical trial. *JAMA Ophthalmol* 2014;132:572–578.

The consequences of long-term corticosteroid use have prompted investigations into alternative therapies for steroid-refractory orbital inflammatory disease (OID). Rituximab, a monoclonal antibody to the CD20 antigen on mature B cells, has been identified as one such therapy. In this phase 1/2 clinical trial, 10 patients with non-infectious OID were given either 500 or 1000 mg rituximab infusions on study days 1 and 15, with the first 5 patients receiving open-label infusions of 1000 mg and the second 5 randomized to either 500 or 1000 mg in a double-masked fashion. Patients who showed an initial treatment response at week 24 but later had waning of this effect were given 2 additional doses of 1000 mg. All patients had been on a stable dose of prednisone and at least one steroid-sparing agent in the 30 days before enrolment. Of the 10 patients in the study, 2 carried a diagnosis of granulomatosis with polyangiitis (GPA), 3 had Graves disease, and 5 had been diagnosed with idiopathic OID. Seven patients had decreased inflammation on an OID grading scale composed of clinical indicators such as pain, proptosis, extraocular motility, corneal disease, visual acuity, visual field, and optic nerve head appearance. Four of these 7 patients were on corticosteroids at the time of enrolment and all of them had their dose reduced by at least 50%. Visual acuity remained stable in all patients, and 7 had at least a 25% reduction in pain on a visual analogue scale at 24 weeks. Although the authors note that only a large disparity would be detectable in a study of this size, no difference was noted between the dosing arms in terms of efficacy, toxicity, or B-cell depletion. Of the 7 patients who responded to treatment, 3 had idiopathic OID, 2 had GPA, and 2 had Graves disease. Four patients received another 2-dose infusion after week 24, half from each dosing arm. One of these patients had idiopathic

OID, 2 had GPA, and 1 had Graves disease. As such, no clear difference in need for retreatment by dosing arm or effectiveness of treatment by diagnosis was identified either.

Evan Price

A PROMISING SEMI-AUTOMATED METHOD FOR DIAGNOSIS AND MANAGEMENT OF PAPILLOEDEMA

Moss HE, Treadwell G, Wanek J, DeLeon S, Shahidi M. Retinal vessel diameter assessment in papilledema by semi-automated analysis of SLO images: feasibility and reliability. *Invest Ophthalmol Vis Sci* 2014;55:2049–2054.

As the authors point out in this study, there is a need for identifiable markers of disease status in idiopathic intracranial hypertension (IIH). Currently, notable clinical findings such as optic nerve head appearance and loss of visual field serve as driving factors in the management of IIH patients, but these are effects that are downstream and often delayed from the onset of increased intracranial pressure. Building upon prior findings seen in optic nerve sheath fenestration patients and animals that had undergone experimental elevation of ICP, the authors sought to determine the feasibility and reliability of using vessel diameter as a suitable marker of IIH disease status. Scanning laser ophthalmoscopy (SLO) images of the optic nerve head and vessels were analysed by a semi-automated processing algorithm to assess for possible differences in vessel diameter between papilloedema patients, pseudopapilloedema patients, and normal controls. The difference in vessel size was also compared in papilloedema patients before and after treatment. Artery diameters were not statistically different in any of the above comparisons. However, vein diameters were significantly larger in papilloedema as compared with both pseudopapilloedema and normal subjects. After treatment, vein diameter decreased in papilloedema patients. In addition, whereas manual analysis of images yielded an interobserver variability of 11 μm , the semi-automated method's variability was at the level of image resolution (3 μm). Although it appears that the system may require some further refining in image acquisition protocol or image analysis, the results of this initial study are promising and may one day give the practitioner more objective data to help guide IIH management.

Evan Price

THE IMPLICATIONS OF HORNER SYNDROME IN CERVICAL ARTERY DISSECTION

Lyrer PA, Brandt T, Metso TM, Metso AJ, Kloss M, Debette S, Leys D, Caso V, Pezzini A, Bonati LH, Thijs V, Bersano A, Touzé E, Gensicke H, Martin JJ, Lichy C, Tatlisumak T, Engelter ST, Grond-Ginsbach C; Cervical Artery Dissection and Ischemic Stroke Patients (CADISP) Study Group. Clinical import of Horner syndrome in internal carotid and vertebral artery dissection. *Neurology* 2014;82:1653–1659.

In this observational study of the patients in the CADISP (Cervical Artery Dissection and Ischemic Stroke Patients) database, the authors sought to identify the association of Horner syndrome (HS) with various characteristics of internal carotid artery dissection (ICAD) or vertebral artery dissection (VAD). A total of 765 patients were included in the study, and patients were enrolled both prospectively and retrospectively. HS was noted in 38.5% of ICAD patients and 13.4% of VAD patients. HS+ ICAD patients less frequently presented with stroke/Transient ischemic attack (TIA) or had bilateral ICA or ICAD occlusion than HS– patients. They also had fewer severe strokes, were less likely to have a new stroke within the 3-month follow-up period, and were more likely to have a favourable functional 3-month outcome based on modified Rankin scale score. On the other hand, HS+ VAD patients more often had occlusive arterial dissection, though no other parameters were statistically different as compared with HS– VAD patients. It is reasoned that this difference in outcomes between ICAD and VAD patients with HS is indicative of different pathophysiological mechanisms in the two conditions. Whereas HS in ICAD is thought to result from pressure on the pericarotid sympathetic plexus, HS in VAD is often a sign of brainstem stroke. The authors note that no patients in their study group died and that severely debilitated patients were unable to give informed consent, which may tend to make their findings less applicable to more severely

affected cervical artery dissection patients. Nevertheless, this large study is helpful in elucidating the prognostic implications of HS in ICAD and VAD patients.

Evan Price

JOB SECURITY FOR NEURO-OPHTHALMOLOGISTS?

Martínez-Lapiscina EH, Fraga-Pumar E, Pastor X, Gómez M, Conesa A, Lozano-Rubí R, Sánchez-Dalmau B, Alonso A, Villoslada P. Is the incidence of optic neuritis rising? Evidence from an epidemiological study in Barcelona (Spain), 2008–2012. *J Neurol* 2014;261:759–767.

The incidence of optic neuritis (ON) has previously been estimated to be 0.56–5.1/100,000 person-years. The authors of this study examined records from a 300,000-person cohort in Spain between 2008 and 2012. They found 82 cases of optic neuritis. Of these, 48 were in the setting of multiple sclerosis (MS), 26 idiopathic, 2 neuromyelitis optica (NMO), and 2 chronic relapsing ischemic optic neuropathy (CRION). Vasculitis, sarcoid, human immunodeficiency virus (HIV), and other systemic inflammatory diseases were also represented. Female/male ratio was 2.4, with an average age of 36. The mean annual incidence rate in this study was 5.36/100,000 person-years. One reason that prior studies' incidence rates may have been lower than the present report is the exclusion of NMO or other systemic disease in previous epidemiological studies of optic neuritis. The presence of new and powerful treatments of MS compared with what was available at the time of these earlier studies may also have produced an increased awareness of ON and stronger reason to diagnose and report. Finally, electronic health records probably allow better tracking of true incidences of disease, a tool which wasn't previously available. One unanswered question that remains to be answered is whether or not the actual incidence of optic neuritis is increasing, and if so, why? Studies like this are important because regional epidemiology helps bring representation to certain groups in regards to disease description. They may also help to provide clues as to aetiology of disease, if epidemiological differences are uncovered.

John H. Pula

DIFFERENTIATING MILD PAPHILLOEDEMA FROM BURIED DISC DRUSEN WITH SD-OCT

Kulkarni KM, Pasol J, Rosa PR, Lam BL. Differentiating mild papilledema and buried optic nerve head drusen using spectral domain optical coherence tomography. *Ophthalmology* 2014;121:959–963.

The authors studied 16 eyes of 9 patients with ultrasound-proven buried optic nerve head drusen (ONHD) using spectral domain optical coherence tomography (SD-OCT). Twelve eyes of 6 patients had less than or equal to Frisén grade 2 papilloedema from idiopathic intracranial hypertension. Two normal fellow eyes of patients with buried ONHD were included. Five clinicians trained and masked to the underlying diagnosis independently reviewed the SD-OCT images and found that SD-OCT is not clinically reliable in differentiating buried ONHD and mild papilloedema.

Michael Vaphiades

TEMPORAL ARTERITIS

Lee AW, Chen C, Cugati S. Temporal arteritis. *Neurol Clin Pract* 2014;4:106–112.

The authors review temporal arteritis and discuss clinical presentation, investigations, temporal artery biopsy, histopathology, laboratory parameters, noninvasive vascular imaging, therapy, and length of steroid therapy and adjuvant steroid sparing agents. They stress that high-dose steroids should be begun the moment the diagnosis is considered and only withdrawn once it has been excluded and a gradual tapering of the steroid dose should occur

over at least 1 year. Ophthalmologists in clinical practice will potentially diagnose and treat this disorder, and thus should read this important review.

Michael Vaphiades

INCREASED OPTIC NERVE RADIOSENSITIVITY FOLLOWING OPTIC NEURITIS

Sanda N, Heran F, Daly-Schveitzer N, Sahel JA, Safran AB. Increased optic nerve radiosensitivity following optic neuritis. *Neurology* 2014;82:1474–1475.

The authors report a patient with radiation-induced optic neuropathy following radiation therapy delivered at a dose widely considered as innocuous, occurring on a background of previous isolated optic neuritis. Increased brain sensitivity to radiation has been observed in patients with multiple sclerosis, but this case is unique in that it solely involves the optic nerves.

Michael Vaphiades

MY EYES ARE TURNED OUTSIDE

Al-Zubidi N, Chevez-Barrios P, Foroozan R, Bhatti MT. “My eyes are turned outside”. *Surv Ophthalmol* 2014;59:354–360.

The authors report a 78-year-old white woman with worsening headache and horizontal and vertical diplopia of 2 months’ duration. Giant cell arteritis (GCA) was suspected and corticosteroids prescribed. An initial temporal artery biopsy (TAB) was negative for GCA, yet a second TAB was positive. In time there was complete resolution of the ophthalmoplegia. The case highlights that diplopia may be a manifestation of GCA and if the first TAB is negative, and the clinical suspicion is high, obtain a second biopsy.

Michael Vaphiades

INFILTRATION OF THE OPTIC CHIASM, NERVE, AND DISC BY GLIOMATOSIS CEREBRI

Traynis I, Singer S, Winterkorn J, Rosenblum M, Dinkin M. Infiltration of the optic chiasm, nerve, and disc by gliomatosis cerebri. *J Neuroophthalmol* 2014;34:44–46.

The authors report an 18-year-old man with gliomatosis cerebri originating in the right frontal lobe, which progressed despite treatment to involve the optic chiasm and right optic nerve including the optic disc. Tumour invasion of the optic nerve head is extremely rare in this entity.

Michael Vaphiades

RACE/ETHNICITY DOES ASSOCIATE WITH VISUAL OUTCOME

Moss HE, Gao W, Balcer LJ, Joslin CE. Association of race/ethnicity with visual outcomes following acute optic neuritis: an analysis of the optic neuritis treatment trial. *JAMA Ophthalmol* 2014;132:421–427.

Although prior data from the Optic Neuritis Treatment Trial (ONTT) indicated that there was no association between race/ethnicity and visual outcomes, a retrospective study of optic neuritis in Atlanta demonstrated worse visual outcomes in patients of black race/ethnicity compared with individuals of white race/ethnicity. To circumvent the limitation of cross-sectional analysis in the ONTT, the authors conducted a secondary analysis of

the ONTT data set through application of longitudinal data analysis techniques. They found race/ethnicity was significantly associated with outcomes in contrast sensitivity and visual acuity during a 15-year period following acute optic neuritis, with black race/ethnicity being associated with worse scores for both.

An-Guor Wang

SLO MAY HELP MONITOR TREATMENT EFFICACY IN PSEUDOTUMOR CEREBRI

Moss HE, Treadwell G, Wanek J, Deleon S, Shahidi M. Retinal vessel diameter assessment in papilledema by semi-automated analysis of SLO images: feasibility and reliability. *Invest Ophthalmol Vis Sci* 2014;55:2049–2054.

Currently, we use lumbar puncture to assess the intracranial pressure (ICP) in patients with idiopathic intracranial hypertension (IIH). Identifying a reliable non-invasive marker reflecting the ICP status is critical for future clinical management of IIH. The authors report the feasibility and reliability of a semi-automated scanning laser ophthalmoscopy (SLO) image analysis method for retinal vessel diameter measurements in subjects with papilloedema before and after treatment. They found retinal vein diameter in papilloedema subjects was larger than in pseudopapilloedema and normal subjects. In addition, papilloedema subjects had a decrease in retinal vein diameter following treatment for and resolution of papilloedema. The clinical utility of retinal vein diameter measurements as a marker for diagnosis and treatment of papilloedema is promising.

An-Guor Wang

IN VIVO MANIPULATION OF RETINAL ATP

Gkotsi D, Begum R, Salt T, Lascaratos G, Hogg C, Chau KY, Schapira AH, Jeffery G. Recharging mitochondrial batteries in old eyes. Near infra-red increases ATP. *Exp Eye Res* 2014;122C:50–53.

Aging changes in tissues with high metabolic demand is accompanied by mitochondrial DNA mutations, decreased adenosine triphosphate (ATP) production, and increasing reactive oxygen species output. Near infra-red (670nm) exposure, presumably absorbed by cytochrome-c oxidase (COX), may improve mitochondrial membrane potentials and ameliorate retinal inflammation in aged eyes. The authors use mouse model to demonstrate that exposure of near infra-red may increase ATP in old mice. In the retina, this was associated with increased COX and reduced acrolein expression. This is the first report of ATP manipulation in vivo and may provide a new non-invasive method to rejuvenate mitochondrial function.

An-Guor Wang

TELEMEDICINE: NEURO-OPHTHALMOLOGY FOR THE FUTURE?

Maamari RN, Keenan JD, Fletcher DA, Margolis TP. A mobile phone-based retinal camera for portable wide field imaging. *Br J Ophthalmol* 2014;98:438–441.

The authors describe a potentially useful tool for telemedicine—a mobile retinal camera that allows images to be captured onto smartphones, and directly transmitted via wireless telecommunication system for remote evaluation. Mobile retinal cameras are available commercially—though barriers to its use for most commercially available cameras include cost and the need for a separate computer to view and transmit images. The authors created a portable device from off-the-shelf components, coupled to an unmodified, commercially available mobile phone (the one named after a fruit). The authors show high-quality images, and make a valuable contribution to this developing field; there are also a few other researchers working on using a smartphone to obtain fundus images—perhaps this accessible technology may be of interest to neuro-ophthalmologists “on-the-move”, or as telemedicine for the future?

Sui Wong