Published in final edited form as:

Glob Heart. 2016 September ; 11(3): 365–367. doi:10.1016/j.ghheart.2016.09.002.

Charting the Future for Ethnicity and Health Research: Clinical and Population Science Insights from the Multi-Ethnic Study of Atherosclerosis (MESA)

George A. Mensah, MD, FACC^{1,2}

¹ Center for Translation Research and Implementation Science; National Heart, Lung, and Blood Institute; National Institutes of Health, Bethesda, Maryland, USA

² Division of Cardiovascular Sciences; National Heart, Lung, and Blood Institute; National Institutes of Health, Bethesda, Maryland, USA

Abstract

The Multi-Ethnic Study of Atherosclerosis (MESA) has been highly successful in investigating the prevalence, characteristics, and progression of subclinical cardiovascular disease (CVD) in a multi-ethnic American cohort of adult men and women free of cardiovascular disease at baseline. MESA has also championed the use of novel biomarkers and emerging imaging techniques for the assessment of subclinical CVD and has created an extensive set of data that continues to fuel dozens of ongoing analyses. Insights from MESA include the first demonstration of ethnic differences in coronary artery calcification and its association with sub-clinical disease progression and incident CVD. Other findings include ethnic differences in the prevalence of pharmacological, behavioral, and lifestyle interventions for the primary prevention of CVD. MESA has also shown the association between residential neighborhood characteristics and behavioral and biomedical risk factors for CVD. This vast amount of data documenting ethnic differences in progression of subclinical CVD, diabetes, kidney disease, and pulmonary disease contrasts sharply with the relative scarcity of specific that can pave the way for the elimination of racial and ethnic disparities. Intervention research however goes beyond the original objectives of MESA and other observational studies. The time has now come to build on the legacy of MESA by supporting rigorous intervention research that informs clinical and public health strategies as well as policy

---

Address for Correspondence Center for Translation Research and Implementation Science (CTRIS), National Heart, Lung, and Blood Institute; National Institutes of Health, One Rockledge Center, 6705 Rockledge Drive, Suite 6070, Bethesda, MD 20892, Telephone: 301-496-1051; Fax: 301-402-1051; George.Mensah@nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest Disclosure: None

Disclosure:

Dr. Mensah is a staff member of the National Heart, Lung and Blood Institute (the source of funding for the MESA study) and acting director of the Division of Cardiovascular Sciences where the MESA study is overseen.

Disclaimer:

The views expressed in this article are those of the author and do not necessarily represent the views of the National Heart, Lung, and Blood Institute; National Institutes of Health; or the United States Department of Health and Human Services.

and environmental changes for eliminating racial and ethnic disparities in CVD and other chronic diseases and advancing the health of multi-ethnic communities.

Introduction

By most standards of assessment, the Multi-Ethnic Study of Atherosclerosis (MESA) has been highly productive and continues to successfully address its core research mission. Initiated in July 2000 to investigate the epidemiological characteristics and progression of subclinical cardiovascular disease (CVD) in a population-based sample of 6500 men and women aged 45-84 years,¹ MESA successfully recruited 6814 participants with a rich multi-ethnic representation of whom 38 percent are white, 28 percent African-American, 22 percent Hispanic, and 12 percent Asian, predominantly of Chinese descent.² Through the creative use of ancillary studies and the collaborative efforts of hundreds of investigators and many active scientific working groups, MESA has also successfully built an extensive set of cohort data that is the basis of dozens of ongoing analytic studies.³ These data include novel markers of CVD risk (including coronary artery calcification (CAC) using computed tomography;⁴⁻⁷ ventricular structure and function using magnetic resonance imaging;^{5, 8-10} and carotid intima-media thickness (CIMT) using ultrasonography.^{4, 5, 11-14} Additionally, MESA has amassed a wealth of phenotypic data as well as an extensive database of geographic, meteorological, and other physical and social characteristics of residential neighborhood environments of its participants.¹⁵⁻¹⁷

A Rich Source of Insights and Clinical Evidence in Ethnicity and Health

As a testament to the scientific productivity of MESA investigators and collaborators, nearly 1100 papers have been published or were in press as of mid-August 2016.³ This extensive source of clinical and public health insights includes the first demonstration of ethnic differences in CAC;¹⁸ that CAC score is a stronger predictor of incident coronary heart disease (CHD) that shows no major differences in its predictive ability among racial and ethnic groups;¹⁹ and that CAC provided superior discrimination and risk reclassification in comparison with any other risk marker in individuals at intermediate risk for incident CHD.²⁰ MESA also demonstrated that among persons with diabetes or metabolic syndrome, absence of CAC conferred a very low risk for incident CHD whereas in the presence of CAC scores greater than 400, incident CHD risk increased more than 10-fold.²¹ Importantly, MESA also showed significant ethnic differences in the prevalence of pharmacological, behavioral, and lifestyle interventions for the primary prevention of CHD. Other evidence coming out of MESA relate to ethnic differences in dietary intake, physical activity, smoking, blood pressure, body mass index, visceral fat, and their associations with incident CHD and diabetes.²²⁻²⁷ More recently, MESA is beginning to shed light on whether the significant Genome-Wide Association Study findings in predominantly European-descent populations are also present and associated with markers of subclinical atherosclerosis and disease progression in the multi-ethnic cohort of MESA.²⁸⁻³⁰

MESA has also provided important insights from social and environmental determinants of subclinical CVD and disease progression. For example, we now have a better understanding

of the association between ambient air pollution and increased levels of atherosclerosis (as measured by CIMT) and especially how that association differs by ethnicity in the MESA communities.¹⁵ In addition, MESA has provided a deeper understanding of the drivers of risk exposure that may inform policy and environmental interventions for CVD prevention. In a study of 5921 White, Black, Hispanic, and Chinese adults across 6 US cities between 2000 and 2002, Jones et al.¹⁶ demonstrated that air pollution exposures were significantly higher in majority Hispanic neighborhoods compared to majority White neighborhoods and that neighborhood-level segregation was importantly associated with markers of ambient air pollution.¹⁶ More recently, a 12-year follow-up of 5950 adult participants of MESA demonstrated that social and physical neighborhood characteristics such as the density of recreational facilities, density of healthy food stores, and measures of the availability of healthy foods, walking environment, and social environment were associated with changes in CAC.³¹ The authors concluded that a “greater access to neighborhood healthy food resources may slow the development of coronary atherosclerosis in middle-aged and older adults.”³¹

The Need for Types 2 and 3 Evidence in Ethnicity and Health

As an observational cohort study, MESA has produced a wealth of evidence that has significantly improved our understanding of the important associations between traditional and novel markers of risk and the development and progression of sub-clinical atherosclerosis and eventual development of overt clinical CVD and diabetes, especially in the non-white ethnic populations in the study. Rychetnik et al.³² refer to this category of evidence as Type 1 evidence or findings from “research that describes risk-disease relations, and identifies the magnitude, severity, and preventability of public health problems.” Thus, Type 1 evidence in ethnicity and health identifies important associations that suggests that “*something should be done*”.³² In contrast, Type 2 evidence identifies “the relative effectiveness of specific interventions aimed at addressing a problem” or “*what* should be done”; while Type 3 evidence informs us on *how* it should be done.³²

Relative to the wealth of Type 1 evidence available, there is a real scarcity of Types 2 and 3 evidence in ethnicity and health. For example, Type 1 evidence from MESA and other observational cohort studies on the documented burden of subclinical and overt clinical CVD in several racial and ethnic populations is quite compelling. Although limited Type 2 evidence exists to inform us of which interventions are effective in controlling traditional risk factors such as hypertension and LDL-cholesterol and preventing progression of subclinical atherosclerosis,^{33, 34} very few of these interventions actually demonstrate efficacy or effectiveness in eliminating racial and ethnic disparities in CVD. For example, Mueller et al.³⁵ identified 39 interventions and several policy initiatives at the state and national levels that addressed reducing racial and ethnic disparities in blood pressure control. The predominant racial/ethnic group in most interventions was African-American. Out of the completed interventions, 27 demonstrated some improvement in blood pressure control or related process measures and 7 showed no improvement.³⁵ Of the 6 completed studies that specifically examined disparities, 3 reduced, 2 increased, and 1 had no effect on disparities.³⁵

Conclusions

As demonstrated in the articles in this special issue of *Global Heart*, the MESA investigators have made very important progress. As a result, we know more about the development, characteristics, and progression of subclinical atherosclerosis and incident CVD, especially in racial and ethnic minority populations in the United States. However, as Goethe eloquently stated, “*Knowing is not enough; we must apply.*” The time has now come for us to apply the insights we have learned from MESA and design the appropriately rigorous and sustainable interventions that pave the way for how we eliminate racial and ethnic disparities in cardiovascular health. This challenge should not fall on the shoulders of MESA investigators and their collaborators because intervention research goes beyond the original objectives of observational cohort studies. However, observational studies will continue to play an important role using evolving state-of-the-art methodologies to make new discoveries, but all of public health should be asking when do we take action to intervene to reduce preventable death and disability and eliminate racial and ethnic disparities. Answers to these questions will be crucial in advancing ethnicity and health research, especially now as we enter the implementation phase of the NHLBI Strategic Vision for the next decade.³⁶

Reference

- (1). Bild DE, Bluemke DA, Burke GL, et al. Multi-Ethnic Study of Atherosclerosis: objectives and design. *Am J Epidemiol*. Nov 1; 2002 156(9):871–81. [PubMed: 12397006]
- (2). Multi-Ethnic Study of Atherosclerosis Coordinating Center. About MESA. MESA. 2016. Available at: URL: <https://www.mesa-nhlbi.org/aboutMESA.aspx>
- (3). Multi-Ethnic Study of Atherosclerosis Coordinating Center. MESA Website. MESA. 2016. Available at: URL: <https://www.mesa-nhlbi.org/>
- (4). Gepner AD, Young R, Delaney JA, et al. Comparison of coronary artery calcium presence, carotid plaque presence, and carotid intima-media thickness for cardiovascular disease prediction in the Multi-Ethnic Study of Atherosclerosis. *Circ Cardiovasc Imaging*. Jan.2015 8(1):e002262. [PubMed: 25596139]
- (5). Sharma RK, Donekal S, Rosen BD, et al. Association of subclinical atherosclerosis using carotid intima-media thickness, carotid plaque, and coronary calcium score with left ventricular dyssynchrony: the multi-ethnic Study of Atherosclerosis. *Atherosclerosis*. Apr; 2015 239(2):412–8. [PubMed: 25682041]
- (6). Ahmed HM, Blaha MJ, Nasir K, et al. Low-risk lifestyle, coronary calcium, cardiovascular events, and mortality: results from MESA. *Am J Epidemiol*. Jul 1; 2013 178(1):12–21. [PubMed: 23733562]
- (7). Budoff MJ, Young R, Lopez VA, et al. Progression of coronary calcium and incident coronary heart disease events: MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol*. Mar 26; 2013 61(12):1231–9. [PubMed: 23500326]
- (8). Edvardsen T, Rosen BD, Pan L, et al. Regional diastolic dysfunction in individuals with left ventricular hypertrophy measured by tagged magnetic resonance imaging--the Multi-Ethnic Study of Atherosclerosis (MESA). *Am Heart J*. Jan; 2006 151(1):109–14. [PubMed: 16368301]
- (9). Liu CY, Liu YC, Wu C, et al. Evaluation of age-related interstitial myocardial fibrosis with cardiac magnetic resonance contrast-enhanced T1 mapping: MESA (Multi-Ethnic Study of Atherosclerosis). *J Am Coll Cardiol*. Oct 1; 2013 62(14):1280–7. [PubMed: 23871886]
- (10). Newton N, Opdahl A, Choi EY, et al. Left ventricular global function index by magnetic resonance imaging--a novel marker for assessment of cardiac performance for the prediction of cardiovascular events: the multi-ethnic study of atherosclerosis. *Hypertension*. Apr; 2013 61(4):770–8. [PubMed: 23424238]

- (11). Chen LY, Leening MJ, Norby FL, et al. Carotid Intima-Media Thickness and Arterial Stiffness and the Risk of Atrial Fibrillation: The Atherosclerosis Risk in Communities (ARIC) Study, Multi-Ethnic Study of Atherosclerosis (MESA), and the Rotterdam Study. *J Am Heart Assoc.* May.2016 5(5):e002907. [PubMed: 27207996]
- (12). Tattersall MC, Gassett A, Korcez CE, et al. Predictors of carotid thickness and plaque progression during a decade: the Multi-Ethnic Study of Atherosclerosis. *Stroke.* Nov; 2014 45(11):3257–62. [PubMed: 25213342]
- (13). Polak JF, Szklo M, Kronmal RA, et al. The value of carotid artery plaque and intima-media thickness for incident cardiovascular disease: the multi-ethnic study of atherosclerosis. *J Am Heart Assoc.* Apr 8.2013 2(2):e000087. [PubMed: 23568342]
- (14). Polak JF, Wong Q, Johnson WC, et al. Associations of cardiovascular risk factors, carotid intima-media thickness and left ventricular mass with inter-adventitial diameters of the common carotid artery: the Multi-Ethnic Study of Atherosclerosis (MESA). *Atherosclerosis.* Oct; 2011 218(2): 344–9. [PubMed: 21726862]
- (15). Jones MR, Diez-Roux AV, O'Neill MS, et al. Ambient air pollution and racial/ethnic differences in carotid intima-media thickness in the Multi-Ethnic Study of Atherosclerosis (MESA). *J Epidemiol Community Health.* Dec; 2015 69(12):1191–8. [PubMed: 26142402]
- (16). Jones MR, Diez-Roux AV, Hajat A, et al. Race/ethnicity, residential segregation, and exposure to ambient air pollution: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Public Health.* Nov; 2014 104(11):2130–7. [PubMed: 25211756]
- (17). Pang Y, Jones MR, Tellez-Plaza M, et al. Association of Geography and Ambient Air Pollution with Urine Metal Concentrations in Six US Cities: The Multi-Ethnic Study of Atherosclerosis. *Int J Environ Res Public Health.* Mar 15.2016 13(3):E324. [PubMed: 26999173]
- (18). Bild DE, Detrano R, Peterson D, et al. Ethnic differences in coronary calcification: the Multi-Ethnic Study of Atherosclerosis (MESA). *Circulation.* Mar 15; 2005 111(10):1313–20. [PubMed: 15769774]
- (19). Detrano R, Guerci AD, Carr JJ, et al. Coronary calcium as a predictor of coronary events in four racial or ethnic groups. *N Engl J Med.* Mar 27; 2008 358(13):1336–45. [PubMed: 18367736]
- (20). Yeboah J, McClelland RL, Polonsky TS, et al. Comparison of novel risk markers for improvement in cardiovascular risk assessment in intermediate-risk individuals. *JAMA.* Aug 22; 2012 308(8):788–95. [PubMed: 22910756]
- (21). Malik S, Budoff MJ, Katz R, et al. Impact of subclinical atherosclerosis on cardiovascular disease events in individuals with metabolic syndrome and diabetes: the multi-ethnic study of atherosclerosis. *Diabetes Care.* Oct; 2011 34(10):2285–90. [PubMed: 21844289]
- (22). Sanchez DR, Diez Roux AV, Michos ED, et al. Comparison of the racial/ethnic prevalence of regular aspirin use for the primary prevention of coronary heart disease from the multi-ethnic study of atherosclerosis. *Am J Cardiol.* Jan; 2011 107(1):41–6. [PubMed: 21146684]
- (23). Joseph JJ, Echouffo-Tcheugui JB, Golden SH, et al. Physical activity, sedentary behaviors and the incidence of type 2 diabetes mellitus: the Multi-Ethnic Study of Atherosclerosis (MESA). *BMJ Open Diabetes Res Care.* Jun 23.2016 4(1):e000185.
- (24). Joseph JJ, Echouffo-Tcheugui JB, Carnethon MR, et al. The association of ideal cardiovascular health with incident type 2 diabetes mellitus: the Multi-Ethnic Study of Atherosclerosis. *Diabetologia.* Sep; 2016 59(9):1893–903. [PubMed: 27272340]
- (25). Shah AD, Kandula NR, Lin F, et al. Less favorable body composition and adipokines in South Asians compared with other US ethnic groups: results from the MASALA and MESA studies. *Int J Obes (Lond).* Apr; 2016 40(4):639–45. [PubMed: 26499444]
- (26). McClelland RL, Jorgensen NW, Post WS, Szklo M, Kronmal RA. Methods for estimation of disparities in medication use in an observational cohort study: results from the Multi-Ethnic Study of Atherosclerosis. *Pharmacoepidemiol Drug Saf.* May; 2013 22(5):533–41. [PubMed: 23382107]
- (27). Oza-Frank R, Chan C, Liu K, Burke G, Kanaya AM. Incidence of type 2 diabetes by place of birth in the Multi-Ethnic Study of Atherosclerosis (MESA). *J Immigr Minor Health.* Oct; 2013 15(5):918–24. [PubMed: 22833256]

- (28). Pankow JS, Decker PA, Berardi C, et al. Circulating cellular adhesion molecules and risk of diabetes: the Multi-Ethnic Study of Atherosclerosis (MESA). *Diabet Med.* Jul; 2016 33(7):985–91. [PubMed: 26937608]
- (29). Vargas JD, Manichaikul A, Wang XQ, et al. Common genetic variants and subclinical atherosclerosis: The Multi-Ethnic Study of Atherosclerosis (MESA). *Atherosclerosis.* Feb.2016 245:230–6. doi: 10.1016/j.atherosclerosis.2015.11.034. Epub@2015 Dec 8.:230-6. [PubMed: 26789557]
- (30). Bielinski SJ, Berardi C, Decker PA, et al. P-selectin and subclinical and clinical atherosclerosis: the Multi-Ethnic Study of Atherosclerosis (MESA). *Atherosclerosis.* May; 2015 240(1):3–9. [PubMed: 25744700]
- (31). Wing JJ, August E, Adar SD, et al. Change in Neighborhood Characteristics and Change in Coronary Artery Calcium: A Longitudinal Investigation in the MESA (Multi-Ethnic Study of Atherosclerosis) Cohort. *Circulation.* Aug 16; 2016 134(7):504–13. [PubMed: 27528645]
- (32). Rychetnik L, Hawe P, Waters E, Barratt A, Frommer M. A glossary for evidence based public health. *J Epidemiol Community Health.* Jul; 2004 58(7):538–45. [PubMed: 15194712]
- (33). Howard BV, Roman MJ, Devereux RB, et al. Effect of lower targets for blood pressure and LDL cholesterol on atherosclerosis in diabetes: the SANDS randomized trial. *JAMA.* Apr 9; 2008 299(14):1678–89. [PubMed: 18398080]
- (34). Howard WJ, Russell M, Fleg JL, et al. Prevention of atherosclerosis with LDL-C lowering - lipoprotein changes and interactions: The SANDS Study. *J Clin Lipidol.* Oct 1; 2009 3(5):322–31. [PubMed: 20161568]
- (35). Mueller M, Purnell TS, Mensah GA, Cooper LA. Reducing racial and ethnic disparities in hypertension prevention and control: what will it take to translate research into practice and policy? *Am J Hypertens.* Jun; 2015 28(6):699–716. [PubMed: 25498998]
- (36). National Heart, Lung, and Blood Institute. Charting the Future Together: The NHLBI Strategic Vision. NHLBI; Bethesda, MD: 2016.