Received:
20 April 2016

Revised:
27 June 2016

Accepted:
30 June 2016

<http://dx.doi.org/10.1259/bjr.20160353>

Cite this article as:

Bray TJP, Amies T, Vendhan K, Humphries P, Sen D, Ioannou Y, et al. Discordant inflammatory changes in the apophyseal and sacroiliac joints: serial observations in enthesitis-related arthritis. *Br J Radiol* 2016; **89**: 20160353.

FULL PAPER

Discordant inflammatory changes in the apophyseal and sacroiliac joints: serial observations in enthesitis-related arthritis

¹TIMOTHY J P BRAY, MB BChir, MA, ¹THOMAS AMIES, MB BS, BSc, ¹KANIMOZHI VENDHAN, MB BS, MD, ¹PAUL HUMPHRIES, MB BS, BSc, ²DEBAJIT SEN, MB BS, ²YIANNIS IOANNOU, MB BS, PhD and ¹MARGARET A HALL-CRAGGS, MB BS, MD

¹Centre for Medical Imaging, University College London, London, UK

²Arthritis UK Centre for Adolescent Rheumatology, University College London, London, UK

Address correspondence to: Dr Timothy J P Bray

E-mail: t.bray@ucl.ac.uk

Objective: To determine the extent to which inflammation of the sacroiliac joints (SIJs) and apophyseal joints (AJs) changes concordantly after treatment in enthesitis-related arthritis (ERA).

Methods: A retrospective study was performed with institutional review board approval. 31 young patients with ERA who had been scanned between March 2009 and November 2014 were included. All patients had post-contrast imaging of the SIJs and lumbar spine and short tau inversion-recovery (STIR) images of the SIJs. The severity of sacroiliitis was scored using a modification of an established technique, and inflammation of the AJs was evaluated using a recently described grading system. The changes in SIJ and AJ scores after treatment were classified as either concordant or discordant, and the proportion of scan pairs in these

groups was recorded. In addition, the correlation between change in SIJ STIR score ($\Delta n\text{fla}$) and change in AJ score (ΔAJ) was assessed using Spearman's correlation coefficient.

Results: Of a total of 43 scan pairs, the changes in inflammation were concordant in 16 scan pairs and discordant in 27 scan pairs. There was no significant correlation between $\Delta n\text{fla}$ and ΔAJ ($R = 0.14$, $p = 0.37$).

Conclusion: Inflammatory changes in the SIJs and AJs are often discordant. This may be a reason why patients experience ongoing back pain despite apparent improvement in one or the other site.

Advances in knowledge: Inflammation may behave differently at different anatomical sites. The SIJs and AJs should both be imaged in patients with ERA with back pain.

INTRODUCTION

Enthesitis-related arthritis (ERA) is a juvenile-onset spondyloarthritis and a subtype of juvenile idiopathic arthritis (JIA), as defined by the International League of Associations for Rheumatology classification criteria for childhood arthritis.¹ The most commonly affected joints at diagnosis are the sacroiliac joints (SIJs), knees, ankles and hips.² Axial inflammation is a hallmark of ERA and up to 35–48% of children with ERA have clinical or radiographic evidence of sacroiliitis.^{3–6} A subset of children with sacroiliitis will progress to spondylitis as adults, which is characterized by back pain, stiffness and eventual fusion of the vertebra.³ Early treatment in spondyloarthritis may have a disease-modifying effect and consequently improve outcomes;⁷ but, if treatment is inadequate, then outcomes are poor in terms of physical health, pain and physical activity.⁴

MRI is commonly used in ERA and has traditionally focused on the early detection of inflammation at the SIJs.^{8,9}

However, a recent cross-sectional study described coexisting inflammation of the lumbar apophyseal joints (AJs) and interspinous ligaments on MRI in a significant proportion of patients.¹⁰ The extent to which AJ and SIJ inflammation are responsible for lower back pain in patients with ERA is unknown. Although it has previously been argued that spinal involvement can be “inferred” from the presence of sacroiliitis (and that routine imaging of the lumbar spine is therefore unnecessary), this suggestion remains unproven.¹¹ Furthermore, it is unknown whether AJ inflammation causes AJ fusion in patients with ERA, although studies in adult spondyloarthritis suggest close relationships between inflammation, excessive bone formation and subsequent fusion.^{12–14} Fusion is known to contribute to impairment of spinal mobility and increased fracture risk.^{15,16}

To our knowledge, there are no published data on the relationship between inflammation of the SIJs and

inflammation of the AJs in patients with ERA. In particular, it is unclear whether SIJ and AJ inflammation improve or deteriorate simultaneously or whether inflammation at these two sites behaves differently. This issue is clinically important because it influences whether imaging of the SIJs alone is sufficient or whether both the SIJs and lumbar spine should be imaged in patients with ERA with lower back pain.

This study aimed to determine the extent to which inflammation at the SIJs and AJs changes concordantly over serial MRI scans. We hypothesized that improvements in inflammation in the SIJs would generally be associated with improvements in the AJs, since both joints might be subject to similar fluctuations in immune/inflammatory activity over time (including when patients are treated).

METHODS AND MATERIALS

We performed a retrospective review of patients suspected of having ERA, who were attending a specialist outpatient adolescent rheumatology clinic at our institution. The study was performed with institutional review board approval. Informed consent was obtained for review of all clinical investigations.

Subjects

A local database was used to identify young people (aged 11–23 years) who had MRI scans between March 2009 and November 2014. Patients were recruited for the study if they met the International League of Associations for Rheumatology criteria for ERA¹ and had complete MRI of the lumbar spine and SIJs, with post-contrast imaging at both sites (as specified below).

MRI technique

MRI of the lumbar spine and SIJs was performed using a 1.5-T system (Avanto; Siemens, Germany). Sequence information is provided in Table 1 [it should be noted that the short tau inversion-recovery (STIR) images were acquired using turbo spin-echo sequences, which are referred to as turbo inversion recovery magnitude in the sequence information]. The contrast agent used was gadoterate meglumine, which was administered as an i.v. bolus at a dose of 0.2 mmol kg⁻¹ using an infusion pump. Following the contrast bolus, the SIJ images were acquired first (axial then coronal), followed by the lumbar spine images. This regime has been designed such that the majority of images are acquired in the early equilibrium phase.

Table 1. Scan parameters

Sequence	Plane	Parameters
Sacroiliac joints		
T_1 weighted TSE	Coronal	TR/TE 610/11 ms, slices 15, slice thickness 4 mm, FOV 200 mm, matrix size 256 × 256, ETL 7, series time 3 min
T_1 weighted TSE	Axial	TR/TE 475/11 ms, slices 15, slice thickness 5 mm, FOV 200 mm, matrix size 256 × 256, ETL 7, series time 5 min
TIRM	Axial	TR/TE 6070/83 ms, inversion time 150 ms, flip angle 150°, slices 15, slice thickness 5 mm, FOV 200 mm, matrix size 256 × 256, ETL 13, series time 5 min
TIRM	Coronal	TR/TE 4340/83 ms, inversion time 150 ms, flip angle 150°, slices 15, slice thickness 4 mm, FOV 200 mm, matrix size 256 × 256, ETL 13, series time 8 min
Post-contrast T_1 weighted TSE with fat saturation	Axial	TR/TE 619/11 ms, slices 15, slice thickness 5 mm, FOV 200 mm, matrix size 256 × 256, ETL 7, series time 5 min
Post-contrast T_1 weighted TSE with fat saturation	Coronal	T_1 TSE fat sat coronal—TR/TE 795/11 ms, slices 15, slice thickness 4 mm, FOV 200 mm, matrix size 256 × 256, ETL 7, series time 3 min
Lumbar spine		
T_1 weighted TSE	Sagittal	TR/TE 400/10 ms, slice thickness 4 mm, slices 15, FOV 300 mm, matrix size 512 × 512, ETL 7, series time 4 min
TIRM	Sagittal	TR/TE 4000/74 ms, inversion time 150 ms, slice thickness 4 mm, slices 15, flip angle 150°, FOV 300 mm, matrix size 256 × 256, ETL 13, series time 4 min
Post-contrast T_1 weighted TSE with fat saturation	Sagittal	TR/TE 510/10 ms, slice thickness 4 mm, slices 15, matrix size 512 × 512, ETL 7, series time 4 min

ETL, echo train length; FOV, field of view; TE, echo time; TIRM, turbo inversion recovery magnitude; TR, repetition time; TSE, turbo spin echo. Series times represent the total interval between each series being acquired inclusive of imaging reconstruction time and time for contrast administration.

Image analysis

All images were reviewed by two consultant radiologists with more than 20 years and 8 years' of experience, respectively, in musculoskeletal MRI and expertise in adolescent spinal imaging. Anonymized images were presented at random to each reader, who was blinded to clinical data. Each observer graded the SIJs and AJs as described below. A reference sheet with examples of the various grades of inflammation was available to each reader to ensure consistency. The reference sheets were produced using a previously described grading system.^{10,17} One of the two observers was involved in conceiving the use of the reference sheets, but was not involved in the choice of specific images or preparation of the sheets themselves. The majority of patients had an initial scan compared with a single follow-up scan, but in patients with more than one follow-up scan, second scans were compared with a third scan and third scans were compared with a fourth scan. Comparison was always with respect to the most recent previous scan. The scoring data were recorded on a score sheet and the two reader scores were averaged.

Apophyseal joint grading

The AJs were graded using the sagittal STIR, T_1 weighted and post-contrast T_1 weighted images, according to the adapted grading method described by Vendhan et al.¹⁰ The grades of severity for joint inflammation are: Grade 0 = normal, Grade 1 = high signal confined to the joint capsule, Grade 2 = intra-articular and periarticular signal abnormality and Grade 3 = features of Grade 2 plus bone marrow oedema of articular process. To account for differences in the "burden" of inflammation associated with each grade of inflammation, we assigned the following weighted scoring system: Grade 0 = Score 0, Grade 1 = Score 0.5, Grade 2 = Score 1 and Grade 3 = Score 2. Levels were scored from T11/12 to L5/S1 (seven vertebral levels). Right and left joints were scored separately and the whole lumbar spine could therefore score a maximum of 28 (*i.e.* 7 vertebral levels; scored 0.5, 1 or 2; left and right). Each scorer recorded additional observations such as erosions or pars defects separately.

Sacroiliac joint grading

The SIJs were scored for inflammation according to the method described by the Spondyloarthritis Research Consortium of Canada¹⁷—an established, validated scoring method for grading sacroiliitis in ankylosing spondyloarthritis in adults. This scoring method takes six coronal slices through the joint from anterior to posterior. For every slice, each joint is divided into quadrants and each quadrant is given a point if high signal is present on the STIR images. An extra point is given if any quadrant on the slice demonstrates high signal extending >1 cm from the articular surface or when the signal is particularly intense. Each slice can therefore score a maximum of 12 points and the whole joint can score 72 points.

Statistical analysis

Assessment of concordance

Changes in score were classified as concordant if the SIJs and AJs both improved or both worsened and as discordant if either the SIJ and AJ score changed in opposite directions or one remained stable while the other changed. We recorded the number of patients in whom inflammatory changes were concordant/discordant and for

each group, we determined the median change in inflammatory scores for subjects who improved and for subjects who worsened.

In order to assess for any dependence of concordance on the interval between scans, the interval between scans for each scan pair was compared between concordant and discordant groups using a Mann–Whitney–Wilcoxon test. Spearman's correlation was used to assess the correlation between change in SIJ score (Δ SIJ) and change in AJ score (Δ AJ) between scans. If patients underwent more than two scans, the change in scores was assessed for each consecutive scan pair (*e.g.* first to second scan, second to third scan etc.) for both the SIJs and AJs.

Repeatability

Interobserver variation in SIJ and AJ scoring was assessed using Bland–Altman plots (95% limits of agreement) and intraclass correlation coefficient (absolute agreement between measurements).

RESULTS

Subjects

There were a total of 31 patients with ERA who had at least 2 MR scans between March 2009 and November 2014, with 74 scans available for analysis. The subject mean age at the time of the first scan was 16.1 years (range 11.5–23 years). This included 24 males with a mean age of 16.7 years (range 12.7–23 years) and 7 females with a mean age of 14.2 years (range 11.5–17.3 years). All patients had at least two scans; eight patients had three scans and two patients had four scans. This allowed for direct comparison between 43 pairs of scans. The mean interval between the first and the last scan was 1.7 years (range 0.3–5.7 years). The mean interval between successive follow-up scans was 1.3 years (range 0.3–3 years).

The number of patients receiving treatment at each scan is provided in Table 2. Between the initial scans and the first follow-up scan, the medication regime was reduced in 1 patient; 14 patients remained on the same therapy and 15 patients required an escalation in therapy. In those patients who had a third scan, five patients remained on the same therapy and four patients required an escalation; in the one patient who had a fourth scan, their medication did not change between the third and fourth scans.

Sacroiliac joint and apophyseal joint scores

On the initial scan, 28 (90%) patients had sacroiliitis and 23 (74%) patients had AJ inflammation. In 21 (68%) patients, there was coexisting sacroiliitis and AJ inflammation. 2 (6%) patients had AJ inflammation on the initial scan but no sacroiliitis. The most severe sacroiliitis scored 64 points (out of a maximum of 72 points) and the most severe AJ inflammation in the spine scored 24 points (out of a maximum of 28 points). 26 (84%) patients had SIJ erosions on their initial scan.

SIJ inflammation improved in 23 scan pairs (21 patients), with a median score reduction of 10 (range 1–33). SIJ inflammation progressed in 17 scan pairs (15 patients), with a median score increase of 5 (range 1–37). In three scan pairs, there was no change in SIJ inflammation.

Table 2. Medication information

Medication	Scan 1	Scan 2	Scan 3	Scan 4
DMARD	12	20	4	2
TNFi	4	9	6	2
NSAID	9	2	0	0
Oral steroid	1	0	0	0
Patients on medication	26	26	8	2
No medication	5	5	0	0

DMARD, disease-modifying antirheumatic drug; NSAID, non-steroidal anti-inflammatory drug; TNFi, tumour necrosis factor inhibitors. The number of patients receiving DMARDs, TNFi, NSAIDs and oral steroids are recorded for each successive scan.

AJ inflammation improved in 17 scan pairs (16 patients), with a median score reduction of 3 (range 1–6). AJ inflammation progressed in 14 scan pairs (14 patients), with a median score increase of 3 (range 1–8). In 12 scan pairs (12 patients), there was no change in AJ inflammation.

Assessment of concordance

Inflammatory changes were concordant (*i.e.* the inflammation either got worse, remained stable or improved at both sites) in 16/43 (37%) scan pairs. An example of a subject's scans showing concordant improvement is shown in Figure 1. Of the concordant scan pairs, inflammation became more severe at both the SIJs and the AJs in six pairs (median increase in inflammation score: SIJ +8, AJ +2), remained stable at both joints in two pairs and improved at both sites in eight scan pairs (median decrease in inflammation score: SIJ –16, AJ –3).

Inflammatory changes were discordant (*i.e.* changes in inflammation at either the SIJs or the AJs did not correspond with the same change at the other joint) in 27/43 (63%) scan pairs. An example of a subject's scans showing discordance is shown in Figure 2. Of the discordant scan pairs, inflammation worsened at the SIJs and improved/remained stable at the AJs in 11 pairs (median increase in inflammation score: SIJ +5, AJ –1). Inflammation improved at the SIJs and worsened/remained stable at the AJs in 15 pairs (median increase in inflammation score: SIJ –5, AJ +2). In one scan pair, inflammation remained stable at the SIJs but increased at the AJs (AJ +1).

The mean interval between scans for each scan pair was 1.23 years for concordant scan pairs and 1.20 years for discordant scan pairs; there was no significant difference between these groups ($p = 0.80$).

Figure 1. Concordant improvement in both sacroiliac joints (SIJs) and apophyseal joints (AJs) (change in SIJ score –23, change in AJ score –6). The AJ images are T_1 weighted post-gadolinium and the SIJs are shown on short tau inversion-recovery images. Arrows indicate areas of bone marrow oedema (inflammation).

Figure 2. Discordance: in this case, the apophyseal joint (AJ) inflammation has worsened on the follow-up scan (change in AJ score +9), while the sacroiliac joint (SIJ) inflammation has improved (change in SIJ score –33). The AJ images are T_1 weighted post-gadolinium and the SIJs are shown on short tau inversion-recovery images. Arrows indicate areas of bone marrow oedema (inflammation).

Details of inflammatory changes in discordant cases are given in Tables 3 and 4. Concordant and discordant cases are also highlighted in Figure 3, which shows a scatter plot demonstrating the relationship between Δ SIJ and Δ AJ. There was no significant correlation between Δ SIJ and Δ AJ ($R = 0.14$, $p = 0.37$).

Repeatability

For the SIJ scores, the Bland–Altman 95% limits of agreement for the two observers were ± 13 across a range of values from 0 to 66. The intraclass correlation coefficient was 0.90.

For the AJ scores, the Bland–Altman 95% limits of agreement for the two observers were ± 8.5 across a range of values from 0 to 25. The intraclass correlation coefficient was 0.57.

DISCUSSION

Inflammation of the lumbar AJs is a recently described phenomenon which has been proposed as a potential cause for back pain in patients with ERA.¹⁰ However, to our knowledge, there is no existing literature describing the relationship between AJ and SIJ inflammation over successive scans. The results of this study indicate a surprising degree of discordance between AJ and SIJ

Table 3. Discordant cases: sacroiliac joint (SIJ) inflammatory scores improving vs apophyseal joint (AJ) score worsening (scans are first scan vs second scan unless otherwise stated)

Patient number	SIJ mean scores	Δ SIJ	AP mean scores	Δ AJ
6	14 to 12	–2	6 to 13	+7
7	41 to 27	–14	0 to 3	+3
13	4 to 2	–2	1 to 4	+2
15	52 to 25	–27	0 to 1	+1
18	50 to 17	–33	16 to 24	+8
25	27 to 25	–2	0 to 4	+4
29	5 to 0	–5	1 to 4	+3

Δ AJ, change in AJ score; Δ SIJ, change in SIJ score.

Table 4. Discordant cases: sacroiliac joint (SIJ) inflammatory score worsening vs apophyseal joint (AJ) score improving (scans are first scan vs second scan unless otherwise stated)

Patient number	SIJ mean scores	Δ SIJ	AJ mean scores	Δ AJ
3	0 to 1	+1	8 to 4	-4
4	27 to 64	+37	2 to 1	-1
5	25 to 41	+16	1 to 0	-1
6 ^a	12 to 27	+13	13 to 10	-3
6 ^b	27 to 32	+5	10 to 6	-4
14	27 to 31	+4	4 to 3	-1
26 ^b	10 to 11	+1	3 to 2	-1
29 ^a	0 to 1	+1	4 to 3	-1
31 ^a	4 to 14	+10	12 to 11	-1

Δ AJ, change in AJ score; Δ SIJ, change in SIJ score.

^aSecond vs third scan.

^bThird vs fourth scan.

inflammation. This suggests that imaging of the SIJs alone may be insufficient for diagnosis in patients with ERA with ongoing back pain, since an absence of SIJ inflammation does not necessarily imply an absence of AJ inflammation. Our data also indicate that AJ inflammation can sometimes precede sacroiliitis, which is consistent with previous studies.¹⁸

Importantly, several authors have recently questioned the need for gadolinium-enhanced scans when imaging young people with lower back pain, arguing that imaging the SIJs alone is

Figure 3. Scatter plot showing change in inflammation in the sacroiliac joints (SIJs) and apophyseal joints (AJs). Each point represents the change between one pair of scans. There was no significant correlation between SIJ change and AJ change ($R = 0.14$, $p = 0.37$).

sufficient.^{19–21} This study reinforces previous suggestions that imaging of the lumbar spine (in addition to the SIJs) is necessary for the complete assessment of young people with lower back pain.

There is an ongoing debate about the pathological mechanisms underlying inflammatory enthesitis, but it is possible that the discordance between AJ and SIJ scores may arise because the pathological processes underlying inflammation at these sites are different.¹² Interestingly, a recent inception cohort study has suggested that enthesitis often persists after treatment in ERA, even in patients who have no active joints (as determined by clinical assessment).²²

Previous studies¹⁰ have suggested that contrast-enhanced images are more sensitive than STIR images for the detection of synovitis. For this reason, we based our scoring on the T_1 sagittal contrast-enhanced images. We employed a weighted scoring system of 0.5, 1 and 2 for the three grades of inflammation, thereby giving greater importance to the presence of bone marrow oedema, which is known to be strongly associated with symptoms.²³ Grade 1 inflammation is, by definition, confined to the joint capsule and therefore arguably reflects only a small inflammatory burden. However, there is clearly scope for further research to determine the clinical significance of different grades of AJ inflammation.

In general, the medication requirements of the patients in our cohort increased over the duration of our study. This is consistent with existing data describing the natural history of ERA and disease progression.²⁴ Compared with other subgroups of JIA, ERA is associated with worse function, quality of life and pain²⁵ as well as a smaller likelihood of attaining inactive disease 1 year after treatment initiation.²⁶

We found that erosions of the SIJs were common in our patients and were present in 81% (25/31) of cases on the initial scan. Other authors studying ankylosing spondylitis suggest that

erosions should be given more attention in quantitative MRI assessment scores of SIJs.²⁷ They conclude that erosions, not bone marrow oedema or contrast enhancement, are the most disease-specific measurable imaging finding in SIJs.

There are some limitations in our study. Although the mean scan interval was 12 months, there was considerable variation. This is an inherent problem when studying an adolescent population. Routinely, these patients are scanned at 6–12-month intervals as part of their normal follow-up. However, appointments were invariably missed, cancelled or rescheduled to allow for events such as school examinations. Specialist nurses maintained contact with patients between outpatient appointments and if symptoms progressed, patients were scanned earlier than the standard interval time. The patient numbers in this study are modest, although ERA is an uncommon subtype affecting 2–10% of all children with JIA (which has an overall prevalence in the UK of 1–2 per 1000 children^{28,29}). Therefore, this study does represent a sizable population and is comparable with other studies. It would also be of interest to correlate measures of inflammatory back pain and function to MRI changes within the SIJ and AJ to ascertain which of these sites is most associated to pain and functional compromise.

A further limitation of this work is that there is scope for improvement in terms of interobserver variation, particularly when scoring AJ inflammation. Descriptions of AJ inflammation have emerged relatively recently and more work is required to increase the repeatability of this score. It is possible that scoring only “definite” inflamed joints may improve this.

In this study, we have shown that in patients with ERA, sacroiliitis and AJ inflammation can change independently and consequently, back pain could be due to the persistence of one site of inflammation, even though the other is responding to therapy. Full assessment of back pain in ERA should include imaging of both the SIJs and the spine with post-contrast imaging of the AJs.

FUNDING

This work was undertaken at the University College London Hospitals (UCLH)/University College London, which receives funding from the Department of Health's the National Institute for Health Research (NIHR) Biomedical Research Centre (BRC) funding scheme. MH-C, TJPB and YI are supported by the NIHR UCLH BRC and YI is also supported by Arthritis Research UK Grant 20164.

REFERENCES

- Petty RE, Southwood TR, Manners P, Baum J, Glass DN, Goldenberg J, et al. International League of Associations for Rheumatology classification of juvenile idiopathic arthritis: second revision, Edmonton, 2001. *J Rheumatol* 2004; **31**: 390–2.
- Weiss PF, Klink AJ, Behrens EM, Sherry DD, Finkel TH, Feudtner C, et al. Enthesitis in an inception cohort of enthesitis-related arthritis. *Arthritis Care Res (Hoboken)* 2011; **63**: 1307–12. doi: <http://dx.doi.org/10.1002/acr.20508>
- Weiss PF. Diagnosis and treatment of enthesitis-related arthritis. *Adolesc Health Med Ther* 2012; **2012**: 67–74. doi: <http://dx.doi.org/10.2147/AHMT.S25872>
- Flatø B, Hoffmann-Vold AM, Reiff A, Førre Ø, Lien G, Vinje O. Long-term outcome and prognostic factors in enthesitis-related arthritis: a case-control study. *Arthritis Rheum* 2006; **54**: 3573–82.
- Weiss P, Klink AH, Behrens EM. Prevalence of enthesitis in pediatric patients with enthesitis-related arthritis. *Arthritis Rheum* 2010; **62**: S98–9.
- Rachlis AC, Babyn PS, Lobo-Mueller E, Benseler SM, Stimec J, Anderson M, et al. Whole body magnetic resonance imaging in juvenile spondyloarthritis: will it provide vital information compared to clinical exam alone?. *Arthritis Rheum* 2011; (63 Suppl. 1): 749.
- Maksymowych WP, Dougados M, van der Heijde D, Sieper J, Braun J, Citera G, et al. Clinical and MRI responses to etanercept in early non-radiographic axial spondyloarthritis: 48-week results from the EMBARK study. *Ann Rheum Dis* 2016; **75**: 1328–35. doi: <http://dx.doi.org/10.1136/annrheumdis-2015-207596>
- Rudwaleit M, Jurik AG, Hermann KG, Landewé R, van der Heijde D, Baraliakos X, et al. Defining active sacroiliitis on magnetic resonance imaging (MRI) for classification of axial spondyloarthritis: a consensual approach by the ASAS/OMERACT MRI group. *Ann Rheum Dis* 2009; **68**: 1520–7. doi: <http://dx.doi.org/10.1136/ard.2009.110767>
- Fisher C, Ioannou Y, Hall-Craggs MA, Sen D. Enthesitis related arthritis: a new era of understanding. *Ann Paediatr Rheum* 2012; **1**: 8–16.
- Vendhan K, Sen D, Fisher C, Ioannou Y, Hall-Craggs MA. Inflammatory changes of the lumbar spine in children and adolescents with enthesitis-related arthritis: magnetic resonance imaging findings. *Arthritis Care Res (Hoboken)* 2014; **66**: 40–6. doi: <http://dx.doi.org/10.1002/acr.22201>
- Bray TP, Vendhan K, Fisher C, Sen D, Ioannou Y, Hall-Craggs MA. Low back pain in adolescents with inflammatory arthritis can be due to lumbar spine apophyseal joint inflammation, and this requires contrast enhancement for adequate assessment: comment on the article by Weiss et al. *Arthritis Rheumatol* 2015; **68**: 263. doi: <http://dx.doi.org/10.1002/art.39452>
- Miceli-Richard C. Enthesitis: the clue to the pathogenesis of spondyloarthritis? *Joint Bone Spine* 2015; **82**: 402–5. doi: <http://dx.doi.org/10.1016/j.jbspin.2015.02.018>
- Ball J. Enthesopathy of rheumatoid and ankylosing spondylitis. *Ann Rheum Dis* 1971; **30**: 213–23. doi: <http://dx.doi.org/10.1136/ard.30.3.213>
- Lories RJ, Luyten FP, de Vlam K. Progress in spondylarthritis. Mechanisms of new bone formation in spondyloarthritis. *Arthritis Res Ther* 2009; **11**: 221. doi: <http://dx.doi.org/10.1186/ar2642>
- Machado P, Landewé R, Braun J, Hermann KG, Baker D, van der Heijde D. Both structural damage and inflammation of the spine contribute to impairment of spinal mobility in patients with ankylosing spondylitis. *Ann Rheum Dis* 2010; **69**: 1465–70. doi: <http://dx.doi.org/10.1136/ard.2009.124206>
- Vosse D, Landewé R, van der Heijde D, van der Linden S, van Staa TP, Geusens P. Ankylosing spondylitis and the risk of fracture: results from a large primary care-based nested case-control study. *Ann Rheum Dis* 2009; **68**: 1839–42. doi: <http://dx.doi.org/10.1136/ard.2008.100503>

17. Maksymowych WP, Inman RD, Salonen D, Dhillon SS, Williams M, Stone M, et al. Spondyloarthritis research Consortium of Canada magnetic resonance imaging index for assessment of sacroiliac joint inflammation in ankylosing spondylitis. *Arthritis Care Res* 2005; **53**: 703–9. doi: <http://dx.doi.org/10.1002/art.21445>
18. Stoll ML, Bhole R, Dempsey-Robertson M, Punaro M. Spondyloarthritis in a pediatric population: risk factors for sacroiliitis. *J Rheumatol* 2010; **37**: 2402–8. doi: <http://dx.doi.org/10.3899/jrheum.100014>
19. Weiss PF, Xiao R, Biko DM, Johnson AM, Chauvin NA. Detection of inflammatory sacroiliitis in children with magnetic resonance imaging: is gadolinium contrast enhancement necessary? *Arthritis Rheumatol* 2015; **67**: 2250–6. doi: <http://dx.doi.org/10.1002/art.39159>
20. Bray TP, Vendhan K, Fisher C, Sen D, Ioannou Y, Hall-Craggs MA. Low back pain in adolescents with inflammatory arthritis can be due to lumbar spine apophyseal joint inflammation, and this requires contrast enhancement for adequate assessment: comment on the article by Weiss et al. *Arthritis Rheumatol* 2016; **68**: 263. doi: <http://dx.doi.org/10.1002/art.39452>
21. Weiss PF, Vossough A, Chauvin NA. Reply. *Arthritis Rheumatol* 2016; **68**: 263–4. doi: <http://dx.doi.org/10.1002/art.39456>
22. Oen K, Duffy CM, Tse SM, Ramsey S, Ellsworth J, Chedeville G, et al. Early outcomes and improvement of patients with juvenile idiopathic arthritis enrolled in a Canadian multicenter inception cohort. *Arthritis Care Res (Hoboken)* 2010; **62**: 527–36. doi: <http://dx.doi.org/10.1002/acr.20044>
23. Flatø B, Aasland A, Vinje O, Førre O. Outcome and predictive factors in juvenile rheumatoid arthritis and juvenile spondyloarthritis. *J Rheumatol* 1998; **25**: 366–75.
24. Fisher C, Ioannou Y, Hall-Craggs MA, Sen D. Enthesitis related arthritis: a new era of understanding. *Ann Paediatr Rheum* 2012; **1**: 8–16. doi: <http://dx.doi.org/10.5455/apr.011520120733>
25. Weiss PF, Schanberg L, Kimura Y, Colbert RA; CARRAnet investigators BT. *Enthesitis is a significant predictor of decreased quality of life, function, and arthritis-specific pain across juvenile idiopathic arthritis (JIA) categories: preliminary analyses from the CARRAnet registry.* ACR/ARHP Annual Scientific Meeting; 2011. p. 280.
26. Donnithorne KJ, Cron RQ, Beukelman T. Attainment of inactive disease status following initiation of TNF- α inhibitor therapy for juvenile idiopathic arthritis: enthesitis-related arthritis predicts persistent active disease. *J Rheumatol* 2011; **38**: 2675–81. doi: <http://dx.doi.org/10.3899/jrheum.110427>
27. Wick MC, Weiss RJ, Jaschke W, Klauser AS. Erosions are the most relevant magnetic resonance imaging features in quantification of sacroiliac joints in ankylosing spondylitis. *J Rheumatol* 2010; **37**: 622–7. doi: <http://dx.doi.org/10.3899/jrheum.090602>
28. Adib N, Silman A, Thomson W. Outcome following onset of juvenile idiopathic inflammatory arthritis: I. frequency of different outcomes. *Rheumatology (Oxford)* 2005; **44**: 995–1001. doi: <http://dx.doi.org/10.1093/rheumatology/keh620>
29. Oen K, Malleson PN, Cabral DA, Rosenberg AM, Petty RE, Cheang M. Disease course and outcome of juvenile rheumatoid arthritis in a multicenter cohort. *J Rheumatol* 2002; **29**: 1989–99.