What Does the Cochrane Collaboration Say about Exercise after Breast Cancer?

Ezzo J, Manheimer E, McNeely ML, et al. Manual lymphatic drainage for lymphedema following breast cancer treatment. *Cochrane Database Syst Rev.* 2015;(5):CD003475. <http://dx.doi.org/10.1002/14651858.CD003475.pub2>. Medline:25994425

Khan F, Amatya B, Ng L, et al. Multidisciplinary rehabilitation for follow-up of women treated for breast cancer. *Cochrane Database Syst Rev.* 2012;(12):CD009553. <http://dx.doi.org/10.1002/14651858.CD009553.pub2>. Medline:23235677

Markes M, Brockow T, Resch KL. Exercise for women receiving adjuvant therapy for breast cancer. *Cochrane Database Syst Rev.* 2006;(4):CD005001. <http://dx.doi.org/10.1002/14651858.CD005001.pub2>. Medline:17054230

McNeely ML, Campbell K, Ospina M, et al. Exercise interventions for upper-limb dysfunction due to breast cancer treatment. *Cochrane Database Syst Rev.* 2010;(6):CD005211. <http://dx.doi.org/10.1002/14651858.CD005211.pub2>. Medline:20556760

Stuiver MM, ten Tusscher MR, Agasi-Idenburg CS, et al. Conservative interventions for preventing clinically detectable upper-limb lymphoedema in patients who are at risk of developing lymphoedema after breast cancer therapy. *Cochrane Database Syst Rev.* 2015;(2):CD009765. <http://dx.doi.org/10.1002/14651858.CD009765.pub2>. Medline:25677413

The Cochrane Collaboration is an international not-for-profit and independent organization dedicated to making up-to-date, accurate information about the effects of health care readily available worldwide. It produces and disseminates systematic reviews of health care interventions and promotes the search for evidence in the form of clinical trials and other studies of interventions. For more information, visit <http://www.cochrane.org>.

DOI:10.3138/ptc.68.4.cochrane