Published in final edited form as:

Epidemiology. 2017 January ; 28(1): e1. doi:10.1097/EDE.0000000000000560.

Re. Modeling the Effects of E-Cigarettes on Smoking Behavior: Implications for Future Adult Smoking Prevalence

Samir Soneji,

Dartmouth College

Brian Primack,

University of Pittsburgh

John Pierce,

University of California, San Diego

Hai-Yen Sung, and

University of California, San Francisco

James Sargent

Dartmouth College

To the Editor

Cherng et al. (2016) recently published a simulation-based study that estimated the effect of electronic cigarette (e-cigarette) use on the future prevalence of combustible, or traditional, cigarette smoking.¹ We appreciate that the authors addressed this topic of growing global public health importance. However, we disagree with the authors' choice of a key parameter in their primary analysis: the probability of cigarette smoking initiation among youth e-cigarette users. Their choice of this parameter is inconsistent with a growing body of scientific evidence and limits the usefulness of their overall analysis.

Cherng et al. assumed e-cigarette use among youth does not affect the probability of cigarette smoking initiation. Yet, three recently published longitudinal studies involving a total of 5,562 adolescents and young adults all concluded e-cigarette use among never cigarette smokers significantly increased the probability of subsequent cigarette smoking initiation.²⁻⁴ This estimated increase for each of the three studies was 4.7%, 14.1%, and 27.9%, respectively. Using random-effects meta-analysis, we calculated that the pooled odds ratio of cigarette smoking initiation equaled 2.5 (95% confidence interval: 1.5 to 4.0) for e-cigarette users, compared to non-e-cigarette users, adjusting for multiple known demographic, psycho-social, and behavioral risk factors.

Based on the 2014 intercensal population estimate and nationally representative data from the National Youth Tobacco Survey and National Health Interview Survey, 3.1 million

Address correspondence to: Samir Soneji, The Dartmouth Institute for Health Policy, Dartmouth College, One Medical Center Drive, Lebanon NH, 03756. Telephone: 603-650-0398. samir.soneji@dartmouth.edu.

Conflict of Interest Disclosures: None reported.

adolescents and young adults ages 12–24 had never smoked cigarettes and had ever used e-cigarettes in 2014. Applying the increased probability of cigarette smoking initiation—estimated from the three longitudinal studies—to this population, between 148,000 and 876,000 *more* adolescents and young adults would initiate cigarette smoking. Although the sensitivity analysis performed by Cherng et al. varied the increased probability of cigarette smoking initiation among e-cigarette users, the prevalence of cigarette smoking remained virtually unchanged because the denominator of prevalence is the size of the entire US population of 18–75 year olds (225 million people in 2014). Yet, it is crucial not to ignore these several hundred thousand more young cigarette smoking initiators each year. Each young cigarette smoking initiator will lose a dozen or more years of life, on average, if he or she becomes a lifelong nicotine-addicted cigarette smoker.⁵

In conclusion, the assumptions of Cherng et al. around cigarette smoking initiation among youth e-cigarette users substantially undermines the usefulness of their simulation model and its results. When we measure the harms of e-cigarette use based on the most up-to-date scientific evidence, it will be possible to directly measure the balance of benefits and harms of e-cigarette use.

Acknowledgments

We thank the following individuals for their review of and feedback on the manuscript: Valerie Lewis, PhD.

Financial Support: This study was supported by grant R21-CA197912 from the National Cancer Institute (Dr. Soneji), R01-CA077026 from the National Cancer Institute (Dr. Sargent), grants R01-CA140150 and R21-CA185767 from the National Cancer Institute (Dr. Primack), and grant R01-CA190347 from the National Cancer Institute (Dr. Pierce).

REFERENCES

1. Cherng ST, Tam J, Christine PJ, Meza R. Modeling the Effects of E-Cigarettes on Smoking Behavior: Implications for Future Adult Smoking Prevalence. *Epidemiol.* 2016; 28:xxx.
2. Leventhal AM, Strong DR, Kirkpatrick MG, et al. Association of electronic cigarette use with initiation of combustible tobacco product smoking in early adolescence. *JAMA.* 2015; 314(7):700–707. [PubMed: 26284721]
3. Primack BA, Soneji S, Stoolmiller M, Fine MJ, Sargent JD. Progression to traditional cigarette smoking after electronic cigarette use among US adolescents and young adults. *JAMA Pediatr.* 2015 Sep.:1–7.
4. Wills TA, Knight R, Sargent JD, Gibbons FX, Pagano I, Williams RJ. Longitudinal study of e-cigarette use and onset of cigarette smoking among high school students in Hawaii. *Tob Control.* 2016 Jan. tobaccocontrol-2015-052705.
5. Jha P, Ramasundarahettige C, Landsman V, et al. 21st-Century Hazards of Smoking and Benefits of Cessation in the United States. *N Engl J Med.* 2013; 368(4):341–350. [PubMed: 23343063]