Critical role of ATP-induced ATP release for Ca^{2+} signaling in nonsensory cell networks of the developing cochlea

Federico Ceriani^{a,b,c,1}, Tullio Pozzan^{d,e,2}, and Fabio Mammano^{a,b,c,f,2}

^aDepartment of Biomedical Sciences, Institute of Cell Biology and Neurobiology, Italian National Research Council, 00015 Monterotondo (RM), Italy; ^bDepartment of Physics and Astronomy, University of Padua, 35131 Padua, Italy; ^cConnexin Structure and Function Unit, Venetian Institute of Molecular Medicine, Foundation for Advanced Biomedical Research, 35129 Padua, Italy; ^dDepartment of Biomedical Sciences, Institute of Neuroscience (Padua Section), Italian National Research Council, 35121 Padua, Italy; ^eDepartment of Biomedical Sciences, University of Padua, 35121 Padua, Italy; and ^fLaboratory of Phenotypic Screening, Shanghai Institute for Advanced Immunochemical Studies, ShanghaiTech University, Shanghai 201210, China

Contributed by Tullio Pozzan, October 1, 2016 (sent for review July 5, 2016; reviewed by Gary D. Housley and Luc Leybaert)

Spatially and temporally coordinated variations of the cytosolic free calcium concentration ($[\text{Ca}^{2+}]_c$) play a crucial role in a variety of tissues. In the developing sensory epithelium of the mammalian cochlea, elevation of extracellular adenosine triphosphate concentration ($[\text{ATP}]_e$) triggers $[\text{Ca}^{2+}]_c$ oscillations and propagation of intercellular inositol 1,4,5-trisphosphate (IP_3)-dependent Ca^{2+} waves. What remains uncertain is the relative contribution of gap junction channels and connexin hemichannels to these fundamental mechanisms, defects in which impair hearing acquisition. Another related open question is whether $[\text{Ca}^{2+}]_c$ oscillations require oscillations of the cytosolic IP_3 concentration ($[\text{IP}_3]_c$) in this system. To address these issues, we performed Ca^{2+} imaging experiments in the lesser epithelial ridge of the mouse cochlea around postnatal day 5 and constructed a computational model in quantitative adherence to experimental data. Our results indicate that $[\text{Ca}^{2+}]_c$ oscillations are governed by Hopf-type bifurcations within the experimental range of $[\text{ATP}]_e$ and do not require $[\text{IP}_3]_c$ oscillations. The model replicates accurately the spatial extent and propagation speed of intercellular Ca^{2+} waves and predicts that ATP-induced ATP release is the primary mechanism underlying intercellular propagation of Ca^{2+} signals. The model also uncovers a discontinuous transition from propagating regimes (intercellular Ca^{2+} wave speed $> 11 \mu\text{m}\cdot\text{s}^{-1}$) to propagation failure (speed = 0), which occurs upon lowering the maximal ATP release rate below a minimal threshold value. The approach presented here overcomes major limitations due to lack of specific connexin channel inhibitors and can be extended to other coupled cellular systems.

inositol trisphosphate | calcium waves | calcium oscillations | cochlear nonsensory cells | connexins

In the mammalian cochlea, sensory hair cells perform mechanotransduction, i.e., the conversion of sound-evoked mechanical stimuli into electrical signals that are conveyed to the central nervous system along the fibers of the auditory nerve (1). Unlike normal human newborns that hear at birth, most rodents start hearing at postnatal days 10–14 (P10–P14, where P0 is day of birth) and achieve adult-level auditory thresholds by the third postnatal week (2). Preceding the acquisition of hearing, the highly specialized and polarized sensory epithelium of the murine cochlea comprises the greater epithelial ridge, from which inner hair cells and medial nonsensory cells originate, and the adjacent lesser epithelial ridge, which is presumed to generate outer hair cells and lateral nonsensory cells (3, 4). Several of these cells express a glial marker (5).

Purinergic signaling, which is key to cochlear physiology and pathology (6–8), promotes cytosolic free calcium concentration ($[\text{Ca}^{2+}]_c$) oscillations and intercellular Ca^{2+} waves in the matrix of gap-junction-coupled nonsensory cells (9) that embed the sensory inner and outer hair cells of the organ of Corti (2). In this cellular system, a host of experimental data (10, 11) indicate that extracellular ATP binding to G-protein-coupled P2Y receptors activates a canonical transduction

cascade whereby phospholipase C (PLC)-dependent hydrolysis of phosphatidylinositol 4,5-bisphosphate (PIP_2) generates the second-messenger inositol 1,4,5-trisphosphate (IP_3) and diacylglycerol (12). When IP_3 binds to and opens intracellular receptors (IP_3R), it triggers Ca^{2+} release from the endoplasmic reticulum (ER) and raises the $[\text{Ca}^{2+}]_c$ (12). Phosphatidylinositol phosphate kinase type 1 γ ($\text{PIP}1\gamma$) is the enzyme that is primarily responsible for the synthesis of the IP_3 precursor PIP_2 in the cochlea (13).

Connexins participate in this sequence of events as follows: (i) as gap junction channels, composed of connexins 26 and 30 (14–17), which enable IP_3 movement between coupled cells of the organ of Corti (18); (ii) as connexin hemichannels in the cell plasma membrane (19, 20), which mediate paracrine signaling by opening in response to a raised level of the $[\text{Ca}^{2+}]_c$ (21–23); and (iii) these hemichannels release intracellular ATP to the extracellular milieu, whereas ATP degradation by ectonucleotidases terminates this purinergic signaling (19, 20). Altogether, these concerted molecular mechanisms promote encoding of signals by $[\text{Ca}^{2+}]_c$ oscillations (24, 25) and convey crucial biochemical information to every district of the cochlear sensory epithelium via intercellular Ca^{2+} waves (2, 26).

Significance

This study dissects the mechanisms underlying the occurrence of ATP- and inositol 1,4,5-trisphosphate (IP_3)-dependent intracellular cytosolic free calcium concentration $[\text{Ca}^{2+}]_c$ oscillations and intercellular Ca^{2+} waves in the syncytium formed by nonsensory cells in the postnatal mouse cochlea. The findings are significant with regard to development of the cochlear sensory epithelium, injury signaling in the cochlea, and pathophysiology around connexinopathies that dominate prelingual deafness. On a broader frame, this work provides an accurate, quantitative description of the mode of propagation of extracellular ATP-mediated paracrine signaling in epithelial cells. Critically, the modeling brings together a synthesis of quantitative data on the key elements concerning the signaling molecules (ATP and IP_3) and propagation mechanisms from a broad range of prior work.

Author contributions: F.C., T.P., and F.M. designed research; F.C. performed research; F.C. analyzed data; and T.P. and F.M. wrote the paper.

Reviewers: G.D.H., University of New South Wales; and L.L., Ghent University.

The authors declare no conflict of interest.

Freely available online through the PNAS open access option.

¹Present address: Department of Biomedical Science, University of Sheffield, Sheffield S10 2TN, United Kingdom.

²To whom correspondence may be addressed. Email: tullio.pozzan@unipd.it or fabio.mammano@cnr.it.

This article contains supporting information online at www.pnas.org/lookup/suppl/doi:10.1073/pnas.1616061113/-DCSupplemental.

Prior work with mouse cochlear organotypic cultures indicates that $[Ca^{2+}]_c$ oscillations and Ca^{2+} waves in cochlear nonsensory cells of both the lesser (10, 11) and the greater epithelial ridge (27, 28) (Fig. S1) share the same PLC- and IP_3R -dependent signal transduction cascade activated by ATP (13). This signaling cascade has been implicated in the sensing of noise-induced hearing loss (10), as well as in developmental defects that impair hearing acquisition due to genetic mutations in key components of the cascade (13, 18, 28). However, the relative contribution of connexin hemichannels and intercellular gap junction channels has not been determined, due to lack of specific inhibitors acting selectively on one of the two types of channels made by connexins (29). Another related open question that is difficult to address experimentally in this native tissue is whether oscillations in IP_3 concentration ($[IP_3]_c$) are an obligatory component of $[Ca^{2+}]_c$ oscillations (30–33).

In this study, we present a data-driven computational model designed to address these unresolved issues. We show that the occurrence of $[Ca^{2+}]_c$ oscillations within a well-defined range of extracellular adenosine triphosphate concentration ($[ATP]_e$) (10, 11) can be described in terms of Hopf-type bifurcations (34). In addition, we quantify precisely the contribution of IP_3 diffusion through gap junction channels and paracrine signaling mediated by ATP release through connexin hemichannels to the propagation of intercellular Ca^{2+} signals. Our experimental results are consistent with a model whereby Ca^{2+} oscillations do not require IP_3 oscillations and can develop in the presence of an elevated and stable background of IP_3 .

Results

$[Ca^{2+}]_c$ Oscillations Do Not Require $[IP_3]_c$ Oscillations and Are Governed by Hopf-Type Bifurcations. It is well known that $[Ca^{2+}]_c$ can be modulated by a variety of mechanisms, including Ca^{2+} release from various intracellular stores and influx from the extracellular space (25). However, in the present as well as the majority of our previously published work, we focused on purinergic signaling at the endolymphatic surface of the sensory epithelium around P5. Endolymph, which fills the cochlear duct (scala media) and bathes the apical surface of both sensory and nonsensory cells, is an unusual extracellular fluid characterized by a positive potential, known as the endocochlear potential (35). Moreover, endolymph contains high levels of K^+ and Cl^- , low levels of Na^+ (36), and an extremely low extracellular free Ca^{2+} concentration ($[Ca^{2+}]_o$) (37). In mice at P5, the endocochlear potential is still negligible, whereas the endolymphatic ion concentrations reach almost adult levels (38). Our experiments with cochlear organotypic cultures obtained from P5 mice were conducted in low, endolymph-like $[Ca^{2+}]_o$ (20 μM), a condition in which Ca^{2+} influx is reduced to a minimum. In fact, $[Ca^{2+}]_c$ oscillations evoked by extracellular ATP in cochlear nonsensory cells of the lesser epithelial ridge (10, 11, 13, 19, 20), as well as spontaneous Ca^{2+} transients in the greater epithelial ridge (13, 28), persisted for tens of minutes in 20 μM $[Ca^{2+}]_o$ or even in a divalent-free extracellular medium. Therefore, in constructing the model represented in Fig. 1 and described in *Supporting Information*, we neglected Ca^{2+} exchange through the plasma membrane [including the basolateral plasma membrane that in vivo is exposed to ordinary $[Ca^{2+}]_o$ levels and contains elements of the Ca^{2+} entry pathway, such as TRPC channels (39)], effectively reducing our stereotyped cell to a closed compartment in which the total intracellular $[Ca^{2+}]_o$ is constant. In addition, based on the insensitivity of cochlear nonsensory cells to caffeine (11) and akin to similar models of Ca^{2+} dynamics in nonexcitable cells (32, 33), we considered IP_3 -dependent Ca^{2+} release from the ER as the only mechanism responsible for the observed $[Ca^{2+}]_c$ increase. Finally, to fine-tune model parameters (Table S1) we performed a variety of Ca^{2+} imaging experiments in apical cochlear cultures from P5 mice and iteratively compared model predictions to experimental results, as detailed hereafter.

Fig. 1. Components of the computational model of Ca^{2+} dynamics in cochlear nonsensory cells. (A) Schematic representation of the model. Letters J and K indicate fluxes through channels/pumps and rate of production/degradation, respectively (all in $\mu M s^{-1}$). (B) Four-state model of a connexin hemichannel. (C) Hemichannel open probability as a function of $[Ca^{2+}]_c$. Black squares, data from ref. 22; blue solid line, fit generated by the model in B.

In a first set of experiments (Fig. 2), we stimulated cochlear nonsensory cells of the lesser epithelial ridge by the steady pressure application of ATP from a glass micropipette. $[Ca^{2+}]_c$ oscillations were detected only within a limited range of ATP concentrations, consistent with previous reports (10, 11). Specifically, $[Ca^{2+}]_c$ responses failed to exhibit an oscillatory character at $[ATP]_e < 10$ nM (two cultures for each concentration). Small-amplitude, damped $[Ca^{2+}]_c$ oscillations were detected between ~ 20 nM and 50 nM $[ATP]_e$ (three cultures for each concentration). Sizeable, self-sustained $[Ca^{2+}]_c$ oscillations could be reliably elicited in nine of nine organotypic cultures for $[ATP]_e$ in the concentration range from 50 nM to 1 μM . Above this $[ATP]_e$ value, oscillations displayed again a damped (or overdamped) character and $[Ca^{2+}]_c$ returned to baseline in 40–110 s, probably due to a combination of P2Y receptor desensitization (40) and limiting factors affecting G proteins or PLC function.

$[Ca^{2+}]_c$ responses in the computational model replicated experimental data (Fig. 2A) and fitted the ATP dose-response curve for the peak ATP-evoked $[Ca^{2+}]_c$ over the whole range of tested $[ATP]_e$ (Fig. 2B). Analysis of the model results suggests that (i) for $[ATP]_e$ between approximately 50 nM and 1 μM , self-sustained $[Ca^{2+}]_c$ oscillations in cochlear nonsensory cells can develop in the presence of an elevated and stable background of $[IP_3]_c$; (ii) for $[ATP]_e$ in excess of ~ 1 μM , after an initial $[IP_3]_c$ surge, P2Y receptor desensitization combines with IP_3 degradation, causing an exponential fall of $[IP_3]_c$ over time. We performed a computational analysis based on bifurcation theory (41) and determined that the appearance and disappearance of $[Ca^{2+}]_c$ oscillations are governed by supercritical Hopf bifurcations (34) occurring at ~ 0.055 μM and ~ 0.745 μM $[ATP]_e$ (Fig. 3).

In our computational model, the occurrence of $[Ca^{2+}]_c$ oscillations (blue traces in Fig. 2A) is due to the interplay of (i) IP_3 -dependent Ca^{2+} release from the ER; (ii) $[Ca^{2+}]_c$ regulation of IP_3 receptors

To corroborate this conclusion we used a Ca^{2+} imaging assay based on eliciting $[\text{Ca}^{2+}]_c$ oscillations by ATP application followed by photostimulation with caged IP_3 , to increase the $[\text{IP}_3]_c$ at a specific time point (32). An increase in the oscillation frequency is expected if $[\text{Ca}^{2+}]_c$ oscillations do not depend on $[\text{IP}_3]_c$ oscillations. Conversely, a delay in the occurrence of the first $[\text{Ca}^{2+}]_c$ peak after IP_3 uncaging is expected if $[\text{Ca}^{2+}]_c$ oscillations require (i.e., are driven by) $[\text{IP}_3]_c$ oscillations through feedback regulation of IP_3 production or degradation (32). To perform this assay, we coloaded cochlear organotypic cultures with Fluo-4 and caged IP_3 and pressure-applied ATP (200 nM) from a glass micropipette positioned above the sensory epithelium while imaging Fluo-4 fluorescence. Forty seconds after starting ATP application, we activated a 365-nm light-emitting diode for 400 ms to stimulate all cells in the field of view by the uncaging of IP_3 .

These results confirm that $[\text{Ca}^{2+}]_c$ oscillations in this system do not require $[\text{IP}_3]_c$ oscillations (32), in agreement with our model predictions (Fig. 2A). However, it might be of some interest to predict what would happen in the case that $[\text{Ca}^{2+}]_c$ oscillations had to rely on $[\text{IP}_3]_c$ oscillations (Figs. S2–S4). Changing three parameters (amplitude, frequency, and tonic component of $[\text{IP}_3]_c$ oscillations) in a nonsystematic way showed that, at low frequency (0.1 Hz), $[\text{Ca}^{2+}]_c$ oscillations would follow faithfully $[\text{IP}_3]_c$ oscillations, albeit with a systematic delay (Fig. S2). Upon raising the frequency (0.5 Hz), the $[\text{Ca}^{2+}]_c$ signal would track the $[\text{IP}_3]_c$ phasic component only provided $[\text{IP}_3]_c$ oscillations started from $[\text{IP}_3]_c = 0$. In the unlikely event that $[\text{IP}_3]_c$ oscillated at a frequency of 2 Hz, the $[\text{Ca}^{2+}]_c$ signal would be unable to keep the pace. Indeed, $[\text{IP}_3]_c$ oscillations at both 0.5 Hz (Fig. S3) and 2 Hz (Fig. S4) would drive $[\text{Ca}^{2+}]_c$ oscillations with variable degrees of damping and a frequency close to 0.1 Hz, which is the natural oscillation frequency of $[\text{Ca}^{2+}]_c$ oscillations in the model.

Waves initiated by IP₃ uncaging. Having characterized and modeled Ca²⁺ dynamics at the single-cell level, we tackled the problem of intercellular signaling (26). Also in this case we used cochlear cultures coloaded with the AM ester forms of caged IP₃ and of Fluo-4. However, for these experiments Ca²⁺ responses in the lesser epithelial ridge were evoked by a brief (170 ms) and focalized pulse of UV light. The irradiated area comprised a central cell and its six nearest neighbors from which radial Ca²⁺ waves propagated to 18 ± 1 cells of the culture ($n = 4$ experiments in three cultures) (Fig. 5). Similar waves were also evoked by photostimulation with caged IP₃ in the greater epithelial ridge (13). To compare model predictions to experimental results, we reconstructed cell network topology by laser scanning confocal imaging (Fig. S1). All cells in the model were attributed a volume $V = 3,900 \mu\text{m}^3$ (average estimate from our confocal images) whereas the unitary permeability to IP₃ was set at $p_u = 72 \times 10^{-3} \mu\text{m}^3 \cdot \text{s}^{-1}$ from ref. 44. Then, based on realistic cell network topology, we imposed an *in silico* sudden increase of [IP₃]_i in a group of neighboring cells (5 μM in one cell and 0.7 μM in its nearest neighbors, reflecting the different intensities of laser

Fig. 4. Ca^{2+} imaging assay to confirm that Ca^{2+} oscillations do not require IP₃ oscillations. (A) Representative fluorescence trace from a cochlear nonsensory cell during the application of the protocol detailed in ref. 32. After a baseline recording of 20 s, ATP (200 nM) was applied for the duration of the experiment. The solid arrow indicates IP₃ photoactivation by a pulse of UV light. (B) Box plot summarizing the results obtained from $n = 29$ cells using the protocol in A. An 88% increase in the mean oscillation frequency was detected after photoactivation of caged IP₃.

irradiation in the target cell and those adjacent to it). Finally, we evaluated the number of cells (N_{cells}) in which $[\text{Ca}^{2+}]_c$ reached an arbitrary threshold value (10% of the peak in the flashed cell). Model simulations replicated experimental responses (Fig. 5) with a number of gap junction channels $N_{\text{ch}} = 935$ coupling each pair of neighboring cells (Fig. S5).

Waves initiated by extracellular ATP puffs. In cochlear organotypic cultures, Ca^{2+} waves can be triggered also by focal application of brief ATP puffs (4 μM , 50 ms) from a glass micropipette placed in close proximity to a nonsensory cell (10, 11, 13, 19). We simulated these conditions by setting the model $[\text{ATP}]_e$ initial value to 4 μM in a focal spot of 2 μm diameter for 50 ms and compared model responses to experimental results (Movie S1). Wave speed depends on both the maximal ATP release (v_{HC}) and the ATP degradation rate ($r_{\text{deg}}^{\text{ATP}}$). For $300 \mu\text{M}\cdot\text{s}^{-1} \leq v_{\text{HC}} \leq 2,000 \mu\text{M}\cdot\text{s}^{-1}$ and low values of $r_{\text{deg}}^{\text{ATP}}$ ($< 0.1 \text{ s}^{-1}$), the model generated wave speeds in the range from $13 \mu\text{m}\cdot\text{s}^{-1}$ to $17 \mu\text{m}\cdot\text{s}^{-1}$ determined experimentally

(10, 11, 13, 19) (Fig. 6A). For $r_{\text{deg}}^{\text{ATP}} > 0.1 \text{ s}^{-1}$, Ca^{2+} waves failed to propagate irrespective of the value attributed to v_{HC} in the above range because rapid degradation of the ATP released by any given cell hampered wave progression. Lowering v_{HC} (at constant $r_{\text{deg}}^{\text{ATP}} < 0.1 \text{ s}^{-1}$) produced model responses, with progressively lower speed, which eventually lost their self-regenerative character (epitomized by a constant speed of propagation, Fig. 6B). At $v_{\text{HC}} = 0$, Ca^{2+} signals invaded only first-order cells (Fig. 6B), consistent with experiments in which the extracellular purinergic pathway was abrogated pharmacologically by exposure to 200 μM suramin (figure 4 of ref. 18). Note that, in agreement with experimental results, all traces in Fig. 6B superimpose for distances $< 20 \mu\text{m}$ from the ATP source, as in this distance range cellular responses depend on direct activation by ATP diffusing out of the puff pipette. For some combinations of v_{HC} and $r_{\text{deg}}^{\text{ATP}}$, the model generated oscillatory Ca^{2+} waves (Fig. S6), which were occasionally observed in cochlear nonsensory cells (e.g., figure 9 of ref. 20).

Altogether, these data-driven simulations indicate that a self-regenerative ATP-induced ATP-release mechanism sustains wave propagation at constant speed over a vast range of distances (in excess of 200 μm from the source). The transition from propagating to nonpropagating regimes is abrupt (Fig. 6A and C), indicating that (for each $r_{\text{deg}}^{\text{ATP}}$ value) the system becomes self-regenerative only above a minimal threshold value of the v_{HC} parameter. At the opposite extreme, increasing v_{HC} results in propagation speeds that saturate at a value that depends on the ATP diffusion constant D in the extracellular medium (Fig. 6D), indicating that this parameter sets the ultimate limit to the cell-to-cell signal propagation (diffusion-limited rate).

Besides paracrine signaling mediated by extracellular ATP, our computational model also included IP₃ diffusion between adjoining cells through gap junction channels (Fig. 1 and Supporting Information). However, the relative contribution to the propagation mechanism of gap junction channels (transferring IP₃) and connexin hemichannels (releasing ATP) is unknown. Pharmacological isolation of the two components is difficult to achieve due to lack of connexin hemichannels blockers that do not also affect gap junction channels (29). Currently, hemichannel block without gap junction block is possible for connexin 43 (45) but not for connexin 26 and connexin 30. Computationally, gap junction

Fig. 5. Focal photoactivation of caged IP₃: comparison between experiments and simulations. (A) Cochlear cultures were loaded with the photoactivatable precursor of IP₃. $[\text{Ca}^{2+}]_c$ oscillations and the propagation of Ca^{2+} signals between adjacent cells were elicited by illuminating a single cell, indicated by the white arrow, with a brief pulse of UV light (365 nm). Photolytic release of IP₃ was reproduced in silico by a step increase of intracellular IP₃ concentration. (Scale bar: 20 μm .) (B) Time course of $[\text{Ca}^{2+}]_c$ signals in experiments (Top) and simulations (Middle) from the regions of interest shown in A. The bottom set of traces is the corresponding model $[\text{IP}_3]_c$ signals.

Fig. 6. Parametric analysis of radial Ca^{2+} wave propagation triggered by focal application of ATP. Experimental conditions were simulated by setting the model $[\text{ATP}]_e$ initial value to $4 \mu\text{M}$ in a focal spot of $2 \mu\text{m}$ diameter for 50 ms . (A) Heat map showing the dependence of Ca^{2+} wave speed on the maximal ATP release rate, v_{HC} , and the ATP degradation constant, $r_{\text{deg}}^{\text{ATP}}$. Values in the blue region of the graph prevented ATP-induced Ca^{2+} wave propagation. (B) The maximal radius of the wave is plotted against time for different values of v_{HC} ($0.0 \mu\text{M}\cdot\text{s}^{-1}$, $140 \mu\text{M}\cdot\text{s}^{-1}$, $200 \mu\text{M}\cdot\text{s}^{-1}$, and $1,000.0 \mu\text{M}\cdot\text{s}^{-1}$) at constant $r_{\text{deg}}^{\text{ATP}} = 0.001 \text{ s}^{-1}$. (C) Wave speed as a function of the ATP release rate v_{HC} , at constant $r_{\text{deg}}^{\text{ATP}} = 0.001 \text{ s}^{-1}$; note discontinuity at $v_{\text{HC}} = 150 \mu\text{M}\cdot\text{s}^{-1}$, a value below which waves failed to propagate and above which waves propagate with speed $> 11 \mu\text{m}\cdot\text{s}^{-1}$. (D) The same as C, but with values plotted on a larger, linear scale; for large values of v_{HC} , wave speed saturates to an asymptotic value that depends on the ATP diffusion coefficient D : black circles, $D = 363 \mu\text{m}^2\cdot\text{s}^{-1}$ (correct value); red squares, $D = 726 \mu\text{m}^2\cdot\text{s}^{-1}$ (hypothetical value, to make the point). Note that also the value of v_{HC} at which the discontinuity arises depends on the diffusion coefficient D , as higher values cause faster diffusion of ATP away from the endolymphatic surface of the cells, and thus more ATP release is required to generate Ca^{2+} waves. (E and F) Ca^{2+} wave radius vs. time for $v_{\text{HC}} = 1,000.0 \mu\text{M}\cdot\text{s}^{-1}$ and $r_{\text{deg}}^{\text{ATP}} = 0.01 \text{ s}^{-1}$ in the presence (triangles) or absence (squares) of gap-junction-mediated IP_3 diffusion. The fraction of ATP-releasing supporting cells is 100% in E and 50% in F. Data points for $N_{\text{ch}} = 935$ are the same in E and F.

channels blockade can be simulated by setting $N_{\text{ch}} = 0$ while leaving all other parameters unchanged (and thus also hemichannel function). When all model cells were allowed to release ATP, abrogation of gap junction coupling resulted in negligible changes to the Ca^{2+} wave speed (Fig. 6E). The contribution of IP_3 diffusion through gap junction channels became appreciable only upon reducing the number of model cells that were allowed to release ATP (Fig. 6F). Therefore, our computational results single out ATP-induced ATP release through connexin hemichannels as the primary mechanism responsible for long-range propagation of Ca^{2+} signals in the developing mouse cochlea.

Discussion

The results of the assay we performed based on ref. 32 confirm that $[\text{IP}_3]_e$ oscillations are not an obligatory component of $[\text{Ca}^{2+}]_e$ oscillations (30–32) in cochlear nonsensory cells. Accordingly, we constructed a computational model for the occurrence of $[\text{Ca}^{2+}]_e$ oscillations based on sequential positive and negative feedback of Ca^{2+} on the IP_3 receptor. As has been previously described in models of this type (32), Ca^{2+} oscillations also may develop at constant $[\text{IP}_3]_e$. Of note, our model (i) captures the range of values of $[\text{ATP}]_e$ that evoke $[\text{Ca}^{2+}]_e$ oscillations in nonsensory cells, (ii) accounts for the dose–response relationship between $[\text{Ca}^{2+}]_e$ and $[\text{ATP}]_e$, and (iii) links the occurrence of $[\text{Ca}^{2+}]_e$ oscillations in a well-defined range of $[\text{ATP}]_e$ to supercritical Hopf bifurcations (34).

The model highlights also another critical behavior, namely the abrupt (discontinuous) transition from propagating regimes (intercellular Ca^{2+} wave speed $> 11 \mu\text{m}\cdot\text{s}^{-1}$) to propagation failure (speed = 0) that occurs upon lowering the maximal ATP release rate (v_{HC} parameter, Table S1) below a minimal threshold value, which depends on ATP degradation rate ($r_{\text{deg}}^{\text{ATP}}$ parameter, Table S1).

Previous experiments with ATP biosensor cells demonstrated that photostimulation with caged IP_3 releases ATP to the extracellular medium at the endolymphatic surface of the lesser epithelial ridge and triggers intercellular Ca^{2+} wave propagation (19). Similar waves were also evoked by photostimulation with caged IP_3 in the greater epithelial ridge (13). Leybaert and coworkers showed that (i) an IP_3 -mediated increase in the $[\text{Ca}^{2+}]_e$ is sufficient to trigger hemichannel opening and (ii) hemichannel open probability has a bell-shaped relationship with $[\text{Ca}^{2+}]_e$ (21–23). The hemichannel scheme in our model, although oversimplified, captures these critically important experimental results. However, a bell-shaped dependence on $[\text{Ca}^{2+}]_e$ has been demonstrated for connexins 32 and 43 (21–23). Whether this applies also to connexins 26 and 30, although very likely, awaits direct experimental confirmation. Provided sufficient ATP is released through connexin hemichannels, its binding to P_2Y receptors (10, 11) activates PLC-dependent IP_3 production in a population of nearby cells (13). This in turn promotes Ca^{2+} release from intracellular stores, raising $[\text{Ca}^{2+}]_e$, which, up to $\sim 500 \text{ nM}$, increases the hemichannel open probability (21–23), fostering further ATP-dependent ATP release in a self-regenerative cascade of biochemical reactions that sustains Ca^{2+} wave propagation at constant speed across the cell network.

It is important to note that the four parameters of our simplified hemichannel model would still fit the experimental data if they were all scaled by the same value, thus making the opening kinetics faster or slower. With the particular set of parameters we used, the hemichannel open probability peaks in $\sim 200 \text{ ms}$ following a step-like increase of the $[\text{Ca}^{2+}]_e$, which seems reasonable given that the effect of intracellular Ca^{2+} on hemichannel gating is likely an indirect one (23). A slower response seems unlikely, and faster kinetics would not affect much the propagation speed of Ca^{2+} waves, because IP_3 production and Ca^{2+} release are the rate-limiting factors in this process.

We also ran some simulations using a two-state hemichannel model (open and closed states) in which hemichannel inactivation at high Ca^{2+} is not present (Fig. S7). The results we obtained for the ATP-evoked propagation of Ca^{2+} waves resemble those generated by the four-state model. Thus, inactivation by high Ca^{2+} is not required for the propagation of Ca^{2+} waves and for the associated ATP-induced ATP release. We expect that similar results can be obtained with other types of hemichannel models (e.g., a three-state model, with an open, a closed, and an inactive state) as long as the open probability is different from 0 at physiological Ca^{2+} levels. For all these reasons, we believe that our conclusions are robust and independent of the particular hemichannel models or kinetics.

We have previously implicated IP_3 synthesis deficits (13) as well as gap junction channel IP_3 permeability defects (18) in the

pathogenesis of hereditary hearing disorders. Furthermore, using a knock-in mouse model, we have linked defects in the spontaneous release of ATP from connexin hemichannels in the developing cochlea to disease phenotype in adulthood (28). The unitary permeability of gap junction channels to IP_3 (p_u , Table S1) is an important parameter in our numerical simulations of IP_3 -evoked intercellular Ca^{2+} waves. It has been proposed that junctions in the murine cochlea are composed of connexin 26/30 heteromeric connexons (14, 15), although this may not be the case for the human cochlea (46). As the p_u of heterotypic/heteromeric channels is not known, we used the value we previously determined for homomeric/homotypic connexin 26 channels in transfected HeLa cells (44). Another critical model parameter that influences cell coupling is the average number of channels (N_{ch}) shared by each cell pair in the syncytium. Using voltage imaging, we previously estimated that nonsensory cells in the apical coil of the cochlea are coupled by 910–1,260 open channels per cell pair (47). This range of values is compatible with the estimate of $N_{\text{ch}} = 935$ obtained here, by a completely different approach, to account for the propagation range of Ca^{2+} signals evoked by photolytic release of IP_3 .

Estimating the number of open hemichannels that release ATP is more difficult than estimating the number of gap junction channels between supporting cells, mainly because the unitary permeability of hemichannels to ATP is not known; we provide an educated guess in *Supporting Information*. In the light of the present analysis, ATP release through connexin hemichannels of the cochlear sensory epithelium, and subsequent diffusion through endolymph (19), appears to influence intercellular Ca^{2+} signaling more drastically than the cell-to-cell diffusion of IP_3 through gap junction channels (18). On the other hand, permeation of these two signaling molecules, which are both highly negatively charged and have similar size (Movie S2), is most likely correlated (48). Therefore, a point mutation that affects IP_3 passage through the channel pore may affect also the passage of ATP and vice versa.

The propagation of ATP-dependent intercellular Ca^{2+} waves has been described in glial cells (which express connexins 40, 43, 45, and 46) (49) as well as in a variety of other connexin-expressing epithelial tissues, including (but not limited to) airway epithelia (connexin 32) (50), vascular endothelium (connexins 37, 40, 43, and 45) (51), keratinocytes (connexins 26, 30, 30.3, 31.1, 37, and 43) (52), renal glomerular endothelial cells (connexin 40) (53), and corneal endothelial cells (connexin 43) (54). We believe that the combination of experimental measurements and computational modeling presented here is of general interest as it (i) overcomes major limitations due to lack of specific connexin channel inhibitors and (ii) can be readily extended (by suitable changes to the set of parameters in Table S1) to other cellular systems, including (but not limited to) those mentioned above.

Finally, the results of this modeling offer critical insight into the unique operating conditions of connexin hemichannels at the endolymphatic surface of the cochlear sensory epithelium. All hemichannels are activated upon depolarization and are effectively closed by hyperpolarization; however, voltage dependence, kinetic properties, and sensitivity to $[\text{Ca}^{2+}]_o$, which acts as an open pore blocker (55), vary significantly between hemichannels composed of different connexin isoforms (56). At transmembrane potentials (ΔV_m , inside minus outside) ≤ -30 mV, connexin 30 hemichannels have zero open probability also in zero $[\text{Ca}^{2+}]_o$ (57). Under the same conditions, connexin 26 hemichannels transit frequently between open and closed states (58, 59). This peculiar behavior has been attributed to the so-called “loop gate,” which remains constitutively active at all voltages even in zero $[\text{Ca}^{2+}]_o$ (58). The same behavior is not observed in hemichannels formed by other connexins (e.g., connexins 46 and 50), which remain stably open in low $[\text{Ca}^{2+}]_o$ and close only if subjected to robust hyperpolarization (60, 61).

Depolarization promotes hemichannel opening, and therefore the cell resting potential (and input resistance of the plasma membrane) can be completely run down in the presence of strong hemichannel activity (56). However, the resting potential of nonsensory cells in cochlear organotypic cultures exposed to endolymph-like $[\text{Ca}^{2+}]_o$ (20 μM) is ≤ -60 mV (47). We conclude that, at resting $[\text{Ca}^{2+}]_o$ and in 20 μM $[\text{Ca}^{2+}]_o$, hemichannels of nonsensory cells of the lesser epithelial ridge are only modestly active and transient hemichannel opening promotes ATP release that is constantly neutralized by the action of ectonucleotidases.

Consistent with this conclusion, nonsensory cells of the lesser epithelial ridge remain quiescent if not perturbed by stimuli (e.g., extracellular ATP or intracellular IP_3) that promote hemichannel opening by elevating the $[\text{Ca}^{2+}]_o$. However, they develop spontaneous Ca^{2+} oscillations when exposed to an ectonucleotidase inhibitor, such as ARL67156 (19). As already mentioned, the same PLC- and IP_3R -dependent signal transduction cascade activated by extracellular ATP governs $[\text{Ca}^{2+}]_o$ oscillations and Ca^{2+} waves in cochlear nonsensory cells of the greater epithelial ridge (13). Therefore, the occurrence of ATP-dependent spontaneous Ca^{2+} signaling activity in the greater epithelial ridge (13, 27, 28) is likely due to a constitutively lower ectonucleotidase activity at the endolymphatic surface of the tall columnar cells that populate that portion of the developing sensory epithelium.

In vivo, the ΔV_m across the apical plasma membrane of all cells exposed to endolymph depends on the series combination of the cell resting potential and the endocochlear potential (1). In mice, the endocochlear potential remains below 15 mV between P1 and P5 and increases rapidly after P7, reaching levels of 80–100 mV around P16 (38). In adult mice, we measured endocochlear potentials as large as 117 mV (62). In adult nonsensory cells of the rodent organ of Corti, the cell resting potential can be as negative as -90 mV (63); therefore the ΔV_m across their apical plasma membrane can exceed -200 mV (inside minus outside). Extrapolating available data, which cover a range from $+60$ mV to -100 mV (57, 59), suggests that a ΔV_m around -200 mV should be more than enough to keep connexin 26 (and certainly connexin 30) hemichannels closed in endolymphatic $[\text{Ca}^{2+}]_o$. Hemichannel opening may still occur under conditions that promote a reduction of the ΔV_m (64). Acting synergistically with ATP-gated ion channels (7, 8), the release of ATP into endolymph through connexin hemichannels of the adult cochlea may then foster adaptation to elevated sound levels.

Materials and Methods

Mathematical Methods and Model Parameters. A schematic representation of the model is shown in Fig. 1. The model comprises three main subsystems (Ca^{2+} , IP_3 , and ATP) as described in detail in *Supporting Information*. Model parameters are summarized in Table S1.

Preparation of Cochlear Organotypic Cultures. Animal work was approved by the Ethics committee of the University of Padua (Comitato Etico di Ateneo per la Sperimentazione Animale, protocol no. 104230, October 12, 2013). Cochleae were dissected from P5 mouse pups in ice-cold Hepes-buffered (10 mM, pH 7.2) HBSS, placed onto glass coverslips coated with 185 $\mu\text{g}/\text{mL}$ of Cell Tak, and incubated overnight at 37 °C in DMEM/F12 supplemented with 5% (vol/vol) FBS.

Ratiometric Ca^{2+} Imaging. In this study, we used membrane-permeable AM ester derivatives of Fura-2 (Thermo Fisher Scientific, catalog no. F1221), Fluo-4 (Thermo Fisher Scientific, catalog no. F14217), and caged inositol (3,4,5) trisphosphate [iso-Ins(1,4,5)P3/PM caged, Enzo Life Sciences, catalog no. ALX307071]. Pluronic F-127 (catalog no. P2443) and sulphinpyrazone (catalog no. S9509) were purchased from Sigma-Aldrich. All experiments were performed at room temperature (22–25 °C). Cochlear cultures were transferred to the stage of an upright wide-field fluorescence microscope (BX51; Olympus) and continually superfused with EXM, an extracellular medium containing 138 mM NaCl, 5 mM KCl, 2 mM CaCl_2 , 0.3 mM NaH_2PO_4 , 0.4 mM KH_2PO_4 , 10 mM Hepes-NaOH, and 6 mM d-glucose (pH 7.2, 300 mOsm). Cochlear cultures were incubated for 40 min at 37 °C in DMEM supplemented

with Fura-2 AM (16 μM). The incubation medium contained also pluronic F-127 (0.1%, wt/vol) and sulfinpyrazone (250 μM) to prevent dye sequestration and secretion. Cultures were then transferred onto the stage of an upright microscope (BX51; Olympus) and perfused in EXM for 20 min at 2 mL/min to allow for deesterification. For recording, EXM was substituted by ECM, a medium obtained by replacing 2 mM Ca^{2+} in EXM with an endolymph-like concentration (20 μM). Fura-2 fluorescence was excited using alternatively two light-emitting diodes (LEDs) (center wavelengths 365 nm and 385 nm; M365L and M385L, Thorlabs) passing through an FF01-360/12-2 and an FF01-387-11 filter, respectively (Semrock), and directed onto the sample through a dichromatic mirror (T400LP; Chroma). Fluorescence emission was selected by a BA495-540HQ filter (Olympus) to form fluorescence images on a scientific-grade CCD camera (SensiCam, PCO AG), using a 60 $\times$ water immersion objective (N.A. 1.0, Fluor; Nikon). Image sequences were acquired using software developed in the laboratory, stored on a disk, and processed offline using the Matlab software package (The MathWorks). Signals were measured as dye emission ratio changes, $\Delta R = R(t) - R(0)$, where t is time and $R(t)$ is emission intensity excited at 365 nm divided by the intensity excited at 385 nm, and $R(0)$ indicates prestimulus ratio. To directly compare fluorescence measurements to Ca^{2+} concentrations computed with computer simulations, we converted fluorescence intensity ratios R to $[\text{Ca}^{2+}]_c$ values by the Grynkiewicz equation (65)

$$[\text{Ca}^{2+}]_c = K_D \frac{R - R_{\min}}{R_{\max} - R} \frac{F_{\max}}{F_{\min}},$$

where $K_D = 287.3$ nM is the Fura-2 dissociation constant at 25 $^{\circ}\text{C}$ (66); $R_{\min}$ and $R_{\max}$ are ratio values in Ca^{2+} -free conditions and in saturating Ca^{2+} conditions, respectively; and $F_{\min}/F_{\max}$ is the ratio of the fluorescence intensity after excitation at 385 nm in free and saturating Ca^{2+} conditions, respectively.

$R_{\max}$ and $F_{\max}$ were measured by perfusing cochlear cultures with an extracellular solution containing 10 μM ionomycin and 5 mM Ca^{2+} for 2 min. $R_{\min}$ and $F_{\min}$ were measured after perfusion with a solution containing 100 μM EGTA and 10 μM ionomycin for 30 min. The overall $[\text{Ca}^{2+}]_c$ of the cell was computed by averaging R values in a region of interest centered on it.

For ATP stimulation experiments, ATP dissolved in ECM was applied by pressure, using glass microcapillaries (puff pipettes) that were pulled to a tip of 2–10 μm on a vertical puller (PP80; Narishige) and placed near the target cell. Pressure was applied at the back of the pipette by delivering a transistor–transistor logic (TTL) pulse of carefully controlled duration to a Pneumatic PicoPump (PV800; World Precision Instruments) under software control. All cells tested responded to ATP, whereas no response was detected when ATP (or other P2YR agonists) was omitted from the solution used to fill the puff pipette, indicating that accidental mechanical activation of the cells was negligible.

Photostimulation with Caged IP_3 . Cochlear cultures were incubated for 30 min at 37 $^{\circ}\text{C}$ in DMEM supplemented with the Ca^{2+} dye Fluo-4 AM (16 μM), caged IP_3 AM (5 μM), pluronic F-127 (0.1%, wt/vol), and sulfinpyrazone (250 μM) and thereafter perfused in EXM for 10 min at 2 mL/min to allow for deesterification. Fluorescence emission was selected by a ET535/30M filter (Chroma), centered around a 535-nm wavelength, to form fluorescence images on a scientific-grade CCD camera (SensiCam; PCO AG) using a 20 $\times$ water immersion objective (N.A. 0.95, LumPlanFI; Olympus) connected to a microscope (BX51; Olympus) and illuminated by a collimated 470-nm light-emitting diode (M470L2; Thorlabs) directed onto the sample through a dichromatic mirror (505 dcxr; Chroma). For focal photostimulation with caged IP_3 , the output of a TTL-controlled semiconductor laser module (20 mW, 379 nm, part no. FBB-375-020-FS-FS-1-1; RGBLase LLC) was injected into a UV-permissive fiber-optic cable (multimode step index 0.22 N.A., 105- μm core, part no. AFS105/125YCUSTOM; Thorlabs GmbH). Fiber output was projected onto the specimen plane by an aspheric condenser lens (20-mm effective focal length, part no. ACL2520; Thorlabs) and the recollimated beam was directed onto a dichromatic mirror (400 dclp; Chroma) placed at 45 $^{\circ}$ just above the objective lens of the microscope. By carefully adjusting the position of the fiber in front of the aspheric lens we projected a sharp image of the illuminated fiber core (spot) onto the specimen focal plane selected by the (infinity-corrected) objective lens. Under these conditions, the fiber-optic diameter determined accurately the laser-irradiated area. The size of the irradiated area was estimated by measuring the dimensions of the spot carved by the focused laser into a thin film of black ink deposited on a microscope coverslip located at the front focal plane of the objective. On average, this area comprised a central cell and its six nearest neighbors (Ca^{2+} signal generators). Baseline (prestimulus) fluorescence emission (F_0) was recorded for 2 s, and thereafter a UV laser pulse of 170 ms was applied to release IP_3 and fluorescence emission was monitored for up to 60 s. Signals were measured as relative changes of fluorescence emission intensity ($\Delta F/F_0$), where F is fluorescence at poststimulus time t and $\Delta F = F - F_0$.

Statistical Analysis. Means are quoted $\pm$ SEM and P values are indicated by the letter P . Statistical comparisons were made using the Mann–Whitney u test and $P < 0.05$ was selected as the criterion for statistical significance.

ACKNOWLEDGMENTS. We thank Francesco Zonta for providing [Movie S2](#) and Giulia Crispino for immunofluorescence staining and laser scanning confocal imaging of fixed cochlear organotypic cultures ([Fig. S1](#)). This work was supported by Fondazione Telethon (Grant GGP13114 to F.M.), the Italian National Research Council (Project DSB.AD009.001.004/INVECCHIAMENTO IBCN to F.M.) and the Italian Ministry of University and Research (Grant FIRB-RBAP11X42L to T.P.). F.C. was supported by a junior postdoctoral fellowship from the University of Padua (Grant CPDR132235 to F.M.).

- Fettiplace R, Kim KX (2014) The physiology of mechanoelectrical transduction channels in hearing. *Physiol Rev* 94(3):951–986.
- Mammano F (2013) ATP-dependent intercellular Ca^{2+} signaling in the developing cochlea: Facts, fantasies and perspectives. *Semin Cell Dev Biol* 24(1):31–39.
- Lim D, Rueda J (1992) Structural development of the cochlea. *Development of Auditory and Vestibular Systems - 2*, ed Romand R (Elsevier, New York), 1st Ed, pp 33–58.
- Eggston AA, Wolff D (1947) Embryology of the ear. *Histopathology of the Ear, Nose, and Throat* (Williams and Wilkins, Baltimore), pp 37–64.
- Rio C, Dikkes P, Liberman MC, Corfas G (2002) Glial fibrillary acidic protein expression and promoter activity in the inner ear of developing and adult mice. *J Comp Neurol* 442(2):156–162.
- Housley GD, Bringmann A, Reichenbach A (2009) Purinergic signaling in special senses. *Trends Neurosci* 32(3):128–141.
- Housley GD, et al. (2013) ATP-gated ion channels mediate adaptation to elevated sound levels. *Proc Natl Acad Sci USA* 110(18):7494–7499.
- Yan D, et al. (2013) Mutation of the ATP-gated P2X(2) receptor leads to progressive hearing loss and increased susceptibility to noise. *Proc Natl Acad Sci USA* 110(6):2228–2233.
- Kikuchi T, Kimura RS, Paul DL, Takasaka T, Adams JC (2000) Gap junction systems in the mammalian cochlea. *Brain Res Brain Res Rev* 32(1):163–166.
- Gale JE, Piazza V, Ciubotaru CD, Mammano F (2004) A mechanism for sensing noise damage in the inner ear. *Curr Biol* 14(6):526–529.
- Piazza V, Ciubotaru CD, Gale JE, Mammano F (2007) Purinergic signalling and intercellular Ca^{2+} wave propagation in the organ of Corti. *Cell Calcium* 41(1):77–86.
- Berridge MJ (2009) Inositol trisphosphate and calcium signalling mechanisms. *Biochim Biophys Acta* 1793(6):933–940.
- Rodriguez L, et al. (2012) Reduced phosphatidylinositol 4,5-bisphosphate synthesis impairs inner ear Ca^{2+} signaling and high-frequency hearing acquisition. *Proc Natl Acad Sci USA* 109(35):14013–14018.
- Forge A, et al. (2003) Gap junctions in the inner ear: Comparison of distribution patterns in different vertebrates and assessment of connexin composition in mammals. *J Comp Neurol* 467(2):207–231.
- Ahmad S, Chen S, Sun J, Lin X (2003) Connexins 26 and 30 are co-assembled to form gap junctions in the cochlea of mice. *Biochem Biophys Res Commun* 307(2):362–368.
- Ortolano S, et al. (2008) Coordinated control of connexin 26 and connexin 30 at the regulatory and functional level in the inner ear. *Proc Natl Acad Sci USA* 105(48):18776–18781.
- Zonta F, Polles G, Zanotti G, Mammano F (2012) Permeation pathway of homomeric connexin 26 and connexin 30 channels investigated by molecular dynamics. *J Biomol Struct Dyn* 29(5):985–998.
- Beltramello M, Piazza V, Bukauskas FF, Pozzan T, Mammano F (2005) Impaired permeability to $\text{Ins}(1,4,5)\text{P}_3$ in a mutant connexin underlies recessive hereditary deafness. *Nat Cell Biol* 7(1):63–69.
- Anselmi F, et al. (2008) ATP release through connexin hemichannels and gap junction transfer of second messengers propagate Ca^{2+} signals across the inner ear. *Proc Natl Acad Sci USA* 105(48):18770–18775.
- Majumder P, et al. (2010) ATP-mediated cell-cell signaling in the organ of Corti: The role of connexin channels. *Purinergic Signal* 6(2):167–187.
- Leybaert L, et al. (2003) Connexin channels, connexin mimetic peptides and ATP release. *Cell Commun Adhes* 10(4–6):251–257.
- De Vuyst E, et al. (2006) Intracellular calcium changes trigger connexin 32 hemichannel opening. *EMBO J* 25(1):34–44.
- De Vuyst E, et al. (2009) Ca^{2+} regulation of connexin 43 hemichannels in C6 glioma and glial cells. *Cell Calcium* 46(3):176–187.
- Uhlén P, Fritz N (2010) Biochemistry of calcium oscillations. *Biochem Biophys Res Commun* 396(1):28–32.
- Parekh AB (2011) Decoding cytosolic Ca^{2+} oscillations. *Trends Biochem Sci* 36(2):78–87.
- Leybaert L, Sanderson MJ (2012) Intercellular Ca^{2+} waves: Mechanisms and function. *Physiol Rev* 92(3):1359–1392.
- Tritsch NX, Yi E, Gale JE, Glowatzki E, Bergles DE (2007) The origin of spontaneous activity in the developing auditory system. *Nature* 450(7166):50–55.
- Schütz M, et al. (2010) The human deafness-associated connexin 30 T5M mutation causes mild hearing loss and reduces biochemical coupling among cochlear non-sensory cells in knock-in mice. *Hum Mol Genet* 19(24):4759–4773.

29. Sáez JC, Leybaert L (2014) Hunting for connexin hemichannels. *FEBS Lett* 588(8):1205–1211.
30. Li W, Llopis J, Whitney M, Zlokarnik G, Tsien RY (1998) Cell-permeant caged InsP₃ ester shows that Ca²⁺ spike frequency can optimize gene expression. *Nature* 392(6679):936–941.
31. Matsuura T, et al. (2006) Cytosolic inositol 1,4,5-trisphosphate dynamics during intracellular calcium oscillations in living cells. *J Cell Biol* 173(5):755–765.
32. Sneyd J, et al. (2006) A method for determining the dependence of calcium oscillations on inositol trisphosphate oscillations. *Proc Natl Acad Sci USA* 103(6):1675–1680.
33. Gaspers LD, et al. (2014) Hormone-induced calcium oscillations depend on cross-coupling with inositol 1,4,5-trisphosphate oscillations. *Cell Reports* 9(4):1209–1218.
34. Strogatz SH (2001) *Nonlinear Dynamics and Chaos* (Westview, Boulder, CO).
35. Nin F, et al. (2008) The endocochlear potential depends on two K⁺ diffusion potentials and an electrical barrier in the stria vascularis of the inner ear. *Proc Natl Acad Sci USA* 105(5):1751–1756.
36. Anniko M, Wróblewski R (1986) Ionic environment of cochlear hair cells. *Hear Res* 22:279–293.
37. Bosher SK, Warren RL (1978) Very low calcium content of cochlear endolymph, an extracellular fluid. *Nature* 273(5661):377–378.
38. Yamasaki M, Komune S, Shimozono M, Matsuda K, Haruta A (2000) Development of monovalent ions in the endolymph in mouse cochlea. *ORL J Otorhinolaryngol Relat Spec* 62(5):241–246.
39. Wong AC, Birnbaumer L, Housley GD (2013) Canonical transient receptor potential channel subtype 3-mediated hair cell Ca²⁺ entry regulates sound transduction and auditory neurotransmission. *Eur J Neurosci* 37(9):1478–1486.
40. Erb L, Weisman GA (2012) Coupling of P2Y receptors to G proteins and other signaling pathways. *Wiley Interdiscip Rev Membr Transp Signal* 1(6):789–803.
41. Arnold VI (1994) *Bifurcation Theory and Catastrophe Theory* (Springer, Berlin).
42. Bezprozvanny I, Watras J, Ehrlich BE (1991) Bell-shaped calcium-response curves of Ins(1,4,5)P₃- and calcium-gated channels from endoplasmic reticulum of cerebellum. *Nature* 351(6329):751–754.
43. Adkins CE, Taylor CW (1999) Lateral inhibition of inositol 1,4,5-trisphosphate receptors by cytosolic Ca²⁺. *Curr Biol* 9(19):1115–1118.
44. Hernandez VH, et al. (2007) Unitary permeability of gap junction channels to second messengers measured by FRET microscopy. *Nat Methods* 4(4):353–358.
45. Wang N, et al. (2013) Selective inhibition of Cx43 hemichannels by Gap19 and its impact on myocardial ischemia/reperfusion injury. *Basic Res Cardiol* 108(1):309.
46. Liu W, et al. (2016) Super-resolution structured illumination fluorescence microscopy of the lateral wall of the cochlea: The Connexin26/30 proteins are separately expressed in man. *Cell Tissue Res* 365(1):13–27.
47. Ceriani F, Mammano F (2013) A rapid and sensitive assay of intercellular coupling by voltage imaging of gap junction networks. *Cell Commun Signal* 11:78.
48. Harris AL (2007) Connexin channel permeability to cytoplasmic molecules. *Prog Biophys Mol Biol* 94(1–2):120–143.
49. Guthrie PB, et al. (1999) ATP released from astrocytes mediates glial calcium waves. *J Neurosci* 19(2):520–528.
50. Isakson BE, Evans WH, Boitano S (2001) Intercellular Ca²⁺ signaling in alveolar epithelial cells through gap junctions and by extracellular ATP. *Am J Physiol Lung Cell Mol Physiol* 280(2):L221–L228.
51. Lohman AW, Billaud M, Isakson BE (2012) Mechanisms of ATP release and signalling in the blood vessel wall. *Cardiovasc Res* 95(3):269–280.
52. Tsutsumi M, Denda S, Inoue K, Ikeyama K, Denda M (2009) Calcium ion gradients and dynamics in cultured skin slices of rat hindpaw in response to stimulation with ATP. *J Invest Dermatol* 129(3):584–589.
53. Toma I, et al. (2008) Connexin 40 and ATP-dependent intercellular calcium wave in renal glomerular endothelial cells. *Am J Physiol Regul Integr Comp Physiol* 294(6):R1769–R1776.
54. Gomes P, Srinivas SP, Vereecke J, Himpens B (2005) ATP-dependent paracrine intercellular communication in cultured bovine corneal endothelial cells. *Invest Ophthalmol Vis Sci* 46(1):104–113.
55. Bennett BC, et al. (2016) An electrostatic mechanism for Ca²⁺-mediated regulation of gap junction channels. *Nat Commun* 7:8770.
56. Fasciani I, et al. (2013) Regulation of connexin hemichannel activity by membrane potential and the extracellular calcium in health and disease. *Neuropharmacology* 75:479–490.
57. Valiunas V, Weingart R (2000) Electrical properties of gap junction hemichannels identified in transfected HeLa cells. *Pflügers Arch* 440(3):366–379.
58. Sanchez HA, Slavi N, Srinivas M, Verselis VK (2016) Syndromic deafness mutations at Asn 14 differentially alter the open stability of Cx26 hemichannels. *J Gen Physiol* 148(1):25–42.
59. Sanchez HA, Villone K, Srinivas M, Verselis VK (2013) The D50N mutation and syndromic deafness: Altered Cx26 hemichannel properties caused by effects on the pore and intersubunit interactions. *J Gen Physiol* 142(1):3–22.
60. Trexler EB, Bennett MV, Bargiello TA, Verselis VK (1996) Voltage gating and permeation in a gap junction hemichannel. *Proc Natl Acad Sci USA* 93(12):5836–5841.
61. Srinivas M, Kronengold J, Bukauskas FF, Bargiello TA, Verselis VK (2005) Correlative studies of gating in Cx46 and Cx50 hemichannels and gap junction channels. *Biophys J* 88(3):1725–1739.
62. Crispino G, et al. (2011) BAAV mediated GJB2 gene transfer restores gap junction coupling in cochlear organotypic cultures from deaf Cx26^{lox10} mice. *PLoS One* 6(8):e23279.
63. Oesterle EC, Dallos P (1990) Intracellular recordings from supporting cells in the guinea pig cochlea: DC potentials. *J Neurophysiol* 64(2):617–636.
64. Thorne PR, Muñoz DJ, Housley GD (2004) Purinergic modulation of cochlear partition resistance and its effect on the endocochlear potential in the Guinea pig. *J Assoc Res Otolaryngol* 5(1):58–65.
65. Grynkiewicz G, Poenie M, Tsien RY (1985) A new generation of Ca²⁺ indicators with greatly improved fluorescence properties. *J Biol Chem* 260(6):3440–3450.
66. Larsson D, Larsson B, Lundgren T, Sundell K (1999) The effect of pH and temperature on the dissociation constant for fura-2 and their effects on [Ca²⁺]_i in enterocytes from a poikilothermic animal, Atlantic cod (*Gadus morhua*). *Anal Biochem* 273(1):60–65.
67. De Young GW, Keizer J (1992) A single-pool inositol 1,4,5-trisphosphate-receptor-based model for agonist-stimulated oscillations in Ca²⁺ concentration. *Proc Natl Acad Sci USA* 89(20):9895–9899.
68. De Pittà M, Goldberg M, Volman V, Berry H, Ben-Jacob E (2009) Glutamate regulation of calcium and IP₃ oscillating and pulsating dynamics in astrocytes. *J Biol Phys* 35(4):383–411.
69. Agulhon C, et al. (2008) What is the role of astrocyte calcium in neurophysiology? *Neuron* 59(6):932–946.
70. Lytton J, Westlin M, Burk SE, Shull GE, MacLennan DH (1992) Functional comparisons between isoforms of the sarcoplasmic or endoplasmic reticulum family of calcium pumps. *J Biol Chem* 267(20):14483–14489.
71. Li YX, Rinzel J (1994) Equations for InsP₃ receptor-mediated [Ca²⁺]_i oscillations derived from a detailed kinetic model: A Hodgkin-Huxley like formalism. *J Theor Biol* 166(4):461–473.
72. Lemon G, Gibson WG, Bennett MR (2003) Metabotropic receptor activation, desensitization and sequestration-I: Modelling calcium and inositol 1,4,5-trisphosphate dynamics following receptor activation. *J Theor Biol* 223(1):93–111.
73. Sneyd J, Wetton BT, Charles AC, Sanderson MJ (1995) Intercellular calcium waves mediated by diffusion of inositol trisphosphate: A two-dimensional model. *Am J Physiol* 268(6 Pt 1):C1537–C1545.
74. Hubley MJ, Locke BR, Moerland TS (1996) The effects of temperature, pH, and magnesium on the diffusion coefficient of ATP in solutions of physiological ionic strength. *Biochim Biophys Acta* 1291(2):115–121.