Published in final edited form as:

Environ Res. 2017 January ; 152: 221–225. doi:10.1016/j.envres.2016.09.026.

E-CIGARETTES AS A SOURCE OF TOXIC AND POTENTIALLY CARCINOGENIC METALS

Catherine Ann Hess^{a,b}, Pablo Olmedo^b, Ana Navas-Acien^{b,c}, Walter Goessler^d, Joanna E. Cohen^c, and Ana Maria Rule^b

^aUniversity of California, Berkeley, School of Public Health Prevention Research Center, 180 Grand Ave., Ste. 1200, Oakland, CA 94612.

^bDepartment of Environmental Health Sciences, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe St. Baltimore, MD 21205.

^cInstitute for Global Tobacco Control, Johns Hopkins Bloomberg School of Public Health, 615 N. Wolfe St. Baltimore, MD 21205.

^dKarl-Franzens-Universität Graz, Graz, Institute of Chemistry, Universitätsplatz 1, 8010 Graz, Austria

Abstract

Background and Aims—The popularity of electronic cigarette devices is growing worldwide. The health impact of e-cigarette use, however, remains unclear. E-cigarettes are marketed as a safer alternative to cigarettes. The aim of this research was the characterization and quantification of toxic metal concentrations in five, nationally popular brands of cig-a-like e-cigarettes.

Methods—We analyzed the cartomizer liquid in 10 cartomizer refills for each of five brands by Inductively Coupled Plasma Mass Spectrometry (ICP-MS).

Results—All of the tested metals (cadmium, chromium, lead, manganese and nickel) were found in the e-liquids analyzed. Across all analyzed brands, mean (SD) concentrations ranged from 3.20 (0.454) to 1960 (1540) µg/L for lead, 53.9 (6.95) to 2110 (5220) µg/L for chromium and 115 (49.4) to 22600 (24400) µg/L for nickel. Manganese concentrations ranged from 28.7 (9.79) to 6919 (12200) µg/L. We found marked variability in nickel and chromium concentration within and between brands, which may come from nichrome heating elements.

Conclusion—Additional research is needed to evaluate whether e-cigarettes represent a relevant exposure pathway for toxic metals in users.

Corresponding Author: Catherine Ann Hess, PhD, 180 Grand Avenue, Suite 1200, Oakland, CA 94612, chess@prev.org, 510-883-5731.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Contributions: CAH and PO developed and implemented experimental methods and analysis; AR and ANA contributed to experimental design; CAH conducted data analysis and wrote the manuscript; PO, ANA, AR and JEC contributed to the preparation of the manuscript.

Competing Interests: None

Keywords

Electronic nicotine delivery devices; carcinogens; non-cigarette tobacco products

1. Introduction

E-cigarettes are increasing in popularity in the United States with sales in 2015 exceeding \$3.5 billion.¹ There is great controversy surrounding e-cigarettes and some evidence showing that e-cigarettes are not harmless, although less so than cigarettes and may have long-term health implications for the user.^{2,3} Many of the active smokers switch to e-cigarettes, and never smokers that start using them, do so in the belief that these devices are completely safer than combustible tobacco.^{4,5}

Cig-a-likes, the rechargeable or fully disposable devices commonly sold at convenience and liquor stores, are sometimes referred to as “first-generation” devices, implying that these e-cigarettes are waning in popularity.⁶ We chose to analyze cig-a-likes because as of 2015, cig-a-likes still maintained a strong market share, despite falling in popularity compared to “second-generation” devices.⁷ Surveys of e-cigarette users report that 99 % of adult users are former or current smokers.^{4,8} Over 80 % of e-cigarette users are former tobacco smokers (defined as no longer smoking any tobacco cigarettes).^{4,9} In the US, e-cigarette use is increasing among teenagers who have never used tobacco.¹⁰⁻¹³

Regulation of e-cigarettes varies across countries although at the time this research was conducted, cig-a-likes were unregulated in the US. Recently however, the US Food and Drug Administration (FDA) has announced new deeming regulations that bring e-cigarettes under the same regulations as tobacco.¹⁴ Scheduled to come into effect as of August 2016, the rules require FDA approval for all e-cigarette products which entered the market after 2007. This move may have a substantial impact on the e-cigarette market and could potentially increase the market share of cig-a-like devices in the US, as many of these devices are produced by established tobacco companies who may be better positioned to afford the high cost of FDA product approval than smaller, independent device and e-liquid producers.¹⁵ The European Union (EU) has also recently implemented regulations on e-cigarettes.¹⁶ These regulations include new labeling requirements and advertising restrictions.

Cig-a-like devices work by heating a liquid mixture of propylene glycol, glycerin, nicotine and flavorings. When heated with a metal coil, the mixture is aerosolized into a “vapor”, which is inhaled by the user. The commonly held belief among consumers of e-cigarettes is that they are a safer alternative to cigarettes.^{5,17,18} However, based on investigations including our own, there is strong evidence to suggest that these devices may be a source of toxic chemical exposure for users, particularly substances with known carcinogenic properties.¹⁹⁻²⁴

Very little research has evaluated the potential of e-cigarettes to be a source of toxic metal exposure, including metals with known carcinogenic properties. To date, few published studies have investigated metal concentrations in US e-cigarette brands.^{25,26} Goniewicz et al. investigated 12 Polish and British cig-a-like e-cigarettes and identified only nickel, cadmium

and lead in cig-a-like aerosol, and in concentrations similar to that of a commercially available nicotine inhaler.²⁵ Concentrations ranged from 0.11 to 0.29 $\mu\text{g}/\text{e-cigarette}$ (150 puffs) for nickel and 0.03 μg to 0.57 $\mu\text{g}/\text{e-cigarette}$ for lead. That study did not report chromium or manganese in any brand. Williams et al. analyzed metal concentration in both liquid and aerosol and report the presence of nickel, chromium and lead, but not cadmium.²⁶ Reported concentrations were 0.005 $\mu\text{g}/10$ puffs for nickel, 0.007 $\mu\text{g}/10$ puffs for chromium and 0.017 $\mu\text{g}/10$ puffs for lead.²⁶ The aim of this study was to analyze metal concentrations in the liquid of popular brands of e-cigarettes.

2. Materials and Methods

We selected five popular brands of rechargeable “cig-a-like” devices available in the United States. The retail environment and sales of cig-a-likes are difficult to determine. Brands increase and decrease in popularity rapidly as cig-a-like manufacturers bring new products to market.²⁷ We chose five brands based on national market share. Three of the brands we tested comprised 71 % of the market share of cig-a-likes in 2015.²⁸ Three of the brands are manufactured by tobacco companies and two are not, but all brands are available nationally in the US at big-box retail outlets, convenience stores, and online. All brands contained nicotine in concentrations of approximately 1.6 to 1.8 mg/mL, as stated by the manufacturer on the cartridge packaging.

Cartridges from each brand were purchased at retail outlets or online. All cartridges from a given brand were bought together and, where batch numbers were given, cartridges from the same batch were used. The liquid from 10 cartridges from each brand were analyzed. For each cartridge, we aimed to obtain enough liquid sample (approximately 400 μL) for two replicates. In the end we had a total of 48 liquid samples instead of 50 because two samples from Brand C did not yield enough liquid for analysis and those two samples were excluded. We only selected one flavor for each brand and flavor choice was determined by retail availability at the time of purchase. We found that total volume of liquid per cartridge varied significantly by brand and ranged from 300 to 600 μL . For this reason we chose not to measure per-cartridge metal content but instead report metal concentrations in $\mu\text{g}/\text{L}$, which allows for consistency in reporting across brands.

The end caps of each cartomizer were removed with standard pliers and the pad, free of the heating coil, was removed from the cartridge using polypropylene forceps. Pads were centrifuged for 10 minutes at 1,540 RCF. Two aliquots of 250 μL were collected from each sample for Brand A, Brand B, Brand D and Brand E, and 150 μL for Brand C and diluted to 5 mL final volume with 1 % HNO_3 and 0.5% HCl (Fisher Optima Trace Element Grade) in ultra-pure MilliQ water and vortexed prior to analysis. Cd, Cr, Pb, Mn, and Ni were analyzed using inductively coupled plasma mass spectrometry (ICPMS, Agilent 7500ce Octopole ICP-MS, Agilent Technologies, Santa Clara, USA). Method limits of detection (MLD) were calculated using procedural blanks and are reported in Table 1. Accuracy was successfully tested using NIST traceable Certified Reference Material TMDW-B (High Purity Standards, Charleston, SC). We estimated the intra-class correlation coefficient (ICC) for the two aliquots from the same sample (intra-laboratory ICC) and given the high reliability (Table 1), we calculated and used in the analysis the mean metal concentration of

the two replicates for each e-cigarette liquid sample. We also conducted a duplicate analysis in a random subset of four e-cigarette liquid samples at the Trace Element Laboratory of the Institute of Chemistry Analytical Chemistry, Graz University (Graz, Austria), showing high comparability between laboratories (inter-laboratory ICC, Table 1).

3. Results

We found high levels of metals in the liquids of some brands. Cd, Cr, Pb, Mn and Ni were detected in all liquids analyzed. Metal concentrations per brand are given in Table 1 and Figure 1 (cadmium was not included in the figure as the concentrations were markedly lower in most brands compared to the other metals). Brand A had the highest mean concentrations of all metals investigated. Brand B had the second highest mean concentrations of Cr, Mn and Ni. Mean (*SD*) Ni concentration in Brand A was 22,600 (24,400) µg/L and was nearly 400 times that of the lowest Ni concentration of 58.0 (22.4) µg/L measured in liquid from Brand D. Mean Cr concentration in Brand A was 2,110 (5,220) µg/L, 39 times that of the lowest Cr concentration of 53.9 (6.95). Mean (*SD*) Mn concentration in Brand A was 1,960 (1,450) µg/L, 240 times that of the lowest Mn concentration, measured in Brand E. Cd levels were fairly low, except in Brand A. Pb concentrations were fairly low in Brand C and Brand D and highly variable in other brands.

Intra class correlation coefficients were calculated for sample repeats for inter- and intra-laboratory results. ICCs for all elements are > 0.96, indicating high reliability of analytical results. Variation in and distribution of metal concentrations within some brands was high, particularly in Brand A for all metals and Brand C and Brand E brands for Pb (Figure 1).

4. Discussion

This analysis of cig-a-like e-cigarette liquid found marked variability in nickel and chromium, manganese and lead concentrations within and between brands. For cadmium, the concentrations were comparatively low, except for Brand A. To date, few studies have investigated metal concentrations in e-cigarettes liquid. Comparisons with previous studies are difficult because of differences in the type of sample analyzed (e-cigarette liquid vs. aerosol), sampling protocol and reporting methods across studies. We have reported metal concentrations in µg/L, compared to a per-cartridge concentration, in part, because we found variation in total cartridge liquid volume both within and between brands.

The concentrations of nickel, chromium and manganese in some brands warrant further detailed investigation into metal concentrations in e-cigarette liquid and in aerosol. Nickel is a Group 1 carcinogen and has been associated with chronic bronchitis and lung cancer in occupationally exposed populations.^{29,30} In animal models, inhaled nickel can enter the lymphatic system inducing lymph node damage and reducing acquired immunity.²⁹ Inhaled chromium has been associated with emphysema and chronic lung infection and reduced lung function in humans.³¹ More generally, nickel is a known respiratory and skin irritant.³² It is estimated that the prevalence of nickel contact allergy is approximately 12 percent in the North American population, with recent evidence that the prevalence is increasing.³³ Nickel allergy may be higher among younger individuals and women. Nickel allergy is also

associated with cigarette smoking, as tobacco is a significant source of nickel.³² Effects of inhaled nickel can include, rhinitis, chronic sinusitis and bronchitis and allergic asthma.²⁹ Chronic dermal exposure from vaping can occur around the peri-oral area could potentially result in contact dermatitis from e-cigarettes containing nickel.

Recent research has highlighted the potential harmful effects of even small concentrations of chromium (III), indicating the potential for the oxidization of chromium (III) into carcinogenic chromium (VI) at the cellular level.³⁴ There is growing evidence that chromium (III) is genotoxic³⁵, highlighting the importance of measuring total chromium. In our study, we could not measure the valence state of chromium. It is possible that the nickel and chromium concentrations stem from the use of nickel and chromium (nichrome) in the heating elements of most devices.³⁶ The origin of lead and manganese is unclear, but it could be present due to contamination during the production of the heating coil. Concerns over the health risks of metals in cig-a-likes have been debated, however the high toxicity of these metals justifies the further study of their concentrations in e-cigarette devices, and are high enough to cause concern for user health.³⁷ Lead and manganese, though measured at lower concentrations in our study, are both highly toxic when inhaled. Lead is of particular concern as it affects multiple organs and systems, even at low exposure levels, and inhaled lead is more readily absorbed into the blood stream compared to other routes.³⁸ Manganese is a potent neurotoxicant, and exposure to inhaled manganese is associated with neurological symptoms which resemble Parkinson's Disease, tremor, and muscle spasms as well as inflammation of the lungs.³⁹⁻⁴¹

Direct translation of these results into a quantified level of exposure for users is complicated and beyond the scope of this paper. Electronic cigarettes do not produce side-stream aerosol in the same way as a tobacco cigarette produces side-stream smoke. Because the aerosol is only generated when the user activates the battery through inhalation, a significant portion of the aerosol generated is inhaled into the lungs. The data presented do show the potential for high concentrations of metals in the aerosols produced across the life of one cartridge. While it is unknown how much of the metal in the liquid is aerosolized, even if only a fraction of these metals were aerosolized and transferred into the lung, the concentrations and the variability presented in this paper warrant caution and additional research. More research is needed to evaluate metal exposure in the generated aerosol, including the relatively high concentrations of toxic metals in some brands of e-cigarettes but not others, and the variability within brands. Research is also needed measuring metal concentrations in biospecimens of e-cigarette users. Limits for inhaled metals are generally set for occupational exposure and measured in mg/m³ over a set period of time. A user exposed to the total metal concentrations present in these liquids could exceed NIOSH recommended exposure limits as well as the more conservative ATSDR (Agency For Toxic Substances And Disease Registry) Maximum Recommended Limit (MRL) in one cartridge, particularly for nickel, chromium and lead.^{42,43}

From a consumer standpoint, the variability in metal concentrations makes it difficult to determine which brands or devices may be less harmful than others with regards to toxic metal exposure. More critically, from a quality control perspective, high variability within brands and batches makes safety testing of these devices more difficult for both

manufacturers and regulatory agencies. We did not analyze the metal heating coil, however previous studies in both the US and Japan have reported nichrome heating coils in cig-a-likes.^{26,44} Additionally, nichrome, along with kanthal, an iron/chromium/aluminum alloy, is among the most commonly used alloys for resistance heating components. When in use, the heating coil comes in direct contact with e-liquid, and at higher temperatures could result in some leaching of the coil metals into the liquid. Given the likelihood that the source of some of these metals are the device components themselves, it appears that the existing screening of the liquid for metals prior to assembly of the device is insufficient. While the concentrations of metals in e-cigarette liquid are higher than would be expected in aerosol, and may be lower than in tobacco, the metals and concentrations reported here indicate that these devices are a source of toxic metal exposure. This exposure may be of particular concern in the case of non-smokers who use e-cigarettes, a demographic which is predominantly adolescents.

This study does have limitations. Firstly, it is difficult to translate these findings into delivered dose estimates. This is primarily due to uncertainty in vaping topography and subsequently, in estimating metals exposure from “typical” vaping behavior. Secondly, we did not quantify nickel or chromium species in cig-a-likes, however this is an important subsequent step in determining more precise health risks associated with the element concentrations reported here.

The implications of these findings are particularly relevant in light of increased regulation of e-cigarette manufacturing. New FDA deeming rules may bring about change and may result in more stringent quality control regarding product constituents as well as greater transparency for consumers. The regulations require that manufacturers of electronic cigarettes and e-liquids are required to submit both ingredient lists as well as information on harmful or potentially harmful constituents (HPHC), which includes nickel, lead and chromium and cadmium.¹⁴ A more thorough investigation of the mechanical components of e-cigarettes is needed, as is greater chemical monitoring of e-cigarette liquids after prolonged contact with the device itself as well as monitoring of the final aerosol. For cig-a-likes, hazard reduction may take the form of a shift away from nichrome heating components and greater scrutiny of the materials used in device components.

Acknowledgements

The authors thank Walter Goessler at the Institute for Chemistry at Karl-Franzens-Universität Graz, Austria for conducting the inter-laboratory quality control evaluation in a subset of the samples and Maria Grau Pérez for her assistance in preparing the figure for the manuscript.

Funding: This study was funded by the Institute for Global Tobacco Control, Johns Hopkins School of Public Health (Grant # 118402); NIEHS Training Grant T32ES007141-31A1; NIAAA Training Grant T32-AA014125 and the Alfonso Martín Escudero Foundation.

References

1. Herzog, B. National Association of Tobacco Outlets E-cig and Vape Panel: Key trends facing the industry. Wells Fargo Securities; Apr 22. 2015 2015
2. Rom O, Pecorelli A, Valacchi G, Reznick AZ. Are E-cigarettes a safe and good alternative to cigarette smoking? *Ann N Y Acad Sci.* 2015; 1340:65–74. [PubMed: 25557889]

3. Grana R, Benowitz N, Glantz SA. E-Cigarettes a scientific review. *Circulation*. 2014; 129(19):1972–1986. [PubMed: 24821826]
4. Etter J-F, Bullen C. Electronic cigarette: users profile, utilization, satisfaction and perceived efficacy. *Addiction*. 2011; 106(11):2017–2028. [PubMed: 21592253]
5. Goniewicz ML, Lingas EO, Hajek P. Patterns of electronic cigarette use and user beliefs about their safety and benefits: An Internet survey. *Drug and Alcohol Review*. 2013; 32(2):133–140. [PubMed: 22994631]
6. Lechner WV, Meier E, Wiener JL, et al. The comparative efficacy of first- versus second-generation electronic cigarettes in reducing symptoms of nicotine withdrawal. *Addiction*. 2015; 110(5):862–867. [PubMed: 25639148]
7. Herzog, B.; Gerber, J. E-cigs revolutionizing the tobacco industry. Wells Fargo Securities LLC, Equity Research Department; 2013.
8. Etter J. Electronic cigarettes: a survey of users. *BMC Public Health*. 2010; 10:231. [PubMed: 20441579]
9. Piñeiro B, Correa JB, Simmons VN, et al. Gender differences in use and expectancies of e-cigarettes: Online survey results. *Addictive Behaviors*. 2016; 52:91–97. [PubMed: 26406973]
10. McCarthy M. Cigarette, alcohol, and illicit drug use down among US teens but e-cigarette use common, survey finds. *BMJ*. 2014; 349:7735.
11. McCarthy M. “Alarming” rise in popularity of e-cigarettes is seen among US teenagers as use triples in a year. *BMJ*. 2015; 350:2083.
12. Wills TA, Knight R, Williams RJ, Pagano I, Sargent JD. Risk factors for exclusive e-cigarette use and dual e-cigarette use and tobacco use in adolescents. *Pediatrics*. 2015; 135(1):e43–e51. [PubMed: 25511118]
13. Gilreath TD, Leventhal A, Barrington-Trimis JL, et al. Patterns of alternative tobacco product use: Emergence of hookah and e-cigarettes as preferred products amongst youth. *Journal of Adolescent Health*. 2016; 58(2):181–185. [PubMed: 26598059]
14. US Food and Drug Administration. 81 FR 28973:28973–29106.
15. Yandle B, Meiners RE, Adler JH, Morris AP. Bootleggers, Baptists, and E-Cigarettes. 2015
16. [September 27, 2016] Directive 2014/40/EU of the European Parliament and of the Council of 3 April 2014 on the approximation of the laws, regulations and administrative provisions of the Member States concerning the manufacture, presentation and sale of tobacco and related products and repealing Directive 2001/37/EC.. *Official Journal of the European Union*. Available at: http://ec.europa.eu/health/tobacco/docs/dir_201440_en.pdf.
17. Dockrell M, Morrison R, Bauld L, McNeill A. E-cigarettes: prevalence and attitudes in Great Britain. *Nicotine Tob Res*. 2013; 15(10):1737–1744. [PubMed: 23703732]
18. Farsalinos KE, Romagna G, Tsiapras D, Kyrzopoulos S, Voudris V. Characteristics, perceived side effects and benefits of electronic cigarette use: A worldwide survey of more than 19,000 consumers. *Int J Environ Res Public Health*. 2014; 11(4):4356–4373. [PubMed: 24758891]
19. Chervona Y, Arita A, Costa M. Carcinogenic metals and the epigenome: understanding the effect of nickel, arsenic, and chromium. *Metallomics*. 2012; 4(7)
20. Cheng T. Chemical evaluation of electronic cigarettes. *Tobacco Control*. 2014; 23:11–17.
21. Lerner CA, Sundar IK, Watson RM, et al. Environmental health hazards of e-cigarettes and their components: Oxidants and copper in e-cigarette aerosols. *Environmental pollution (Barking, Essex : 1987)*. 2015; 198:100–107.
22. Tokar EJ, Benbrahim-Tallaa L, Waalkes MP. Metal ions in human cancer development. *Met Ions Life Sci*. 2011; 8:375–401. [PubMed: 21473387]
23. Varlet V, Farsalinos K, Augsburger M, Thomas A, Etter J-F. Toxicity Assessment of Refill Liquids for Electronic Cigarettes. *Int J Environ Res Public Health*. 2015; 12(5):4796. [PubMed: 25941845]
24. Barrington-Trimis JL, Samet JM, McConnell R. Flavorings in electronic cigarettes an unrecognized respiratory health hazard? *JAMA*. 2014; 312(23):2493–2494. [PubMed: 25383564]
25. Goniewicz ML, Knysak J, Gawron M, et al. Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. *Tobacco Control*. 2014; 23(2):133–139. [PubMed: 23467656]

26. Williams M, Villarreal A, Bozhilov K, Lin S, Talbot P. Metal and silicate particles including nanoparticles are present in electronic cigarette cartomizer fluid and aerosol. *PLoS One*. 2013; 8(3)
27. Zhu SH, Sun JY, Bonnevie E, et al. Four hundred and sixty brands of e-cigarettes and counting: implications for product regulation. *Tobacco Control*. 2014; 23:3–9.
28. Craver R. Vuse overwhelming e-cig competition. *Richmond Times-Dispatch*. Jun 1.2015 2015.
29. ATSDR. Toxicological profile for Nickel. U.S. Department of Health and Human Services PHS; Atlanta, GA: 2005.
30. Arsenic, metals, fibres and dusts. International Agency for Research on Cancer; Lyon (FR): 2012. IARC Working Group on the Evaluation of Carcinogenic Risk to Humans..
31. ATSDR. Toxicological Profile for Chromium. U.S. Department of Health and Human Services; Atlanta, GA: 2012.
32. Thyssen JP, Linneberg A, Menné T, Johansen JD. The epidemiology of contact allergy in the general population – prevalence and main findings. *Contact Dermatitis*. 2007; 57(5):287–299. [PubMed: 17937743]
33. Admani S, Jacob SE. Allergic contact dermatitis in children: Review of the past decade. *Current Allergy and Asthma Reports*. 2014; 14(4):1–11.
34. Wu LE, Levina A, Harris HH, et al. Carcinogenic chromium(VI) compounds formed by intracellular oxidation of chromium(III) dietary supplements by adipocytes. *Angew Chemie International Edition*. 2016; 55(5):1742–1745.
35. Fang Z, Zhao M, Zhen H, Chen L, Shi P, Huang Z. Genotoxicity of tri- and hexavalent chromium compounds and their modes of action on DNA damage. *PLoS One*. 2014; 9(8)
36. Brown CJ, Cheng JM. Electronic cigarettes: product characterisation and design considerations. *Tobacco Control*. 2014; 23(suppl 2):ii4–ii10. [PubMed: 24732162]
37. Farsalinos KE, Voudris V, Poulas K. Are metals emitted from electronic cigarettes a reason for health concern? A risk-assessment analysis of currently available literature. *Int J Environ Res Public Health*. 2015; 12(5):5215–5232. [PubMed: 25988311]
38. ATSDR. Toxicological Profile for Lead. Department of Health and Human Services; Atlanta, Georgia: 2007. Atlanta, GA
39. Mergler D, Baldwin M, Belanger S, et al. Manganese neurotoxicity, a continuum of dysfunction: Results from a community based study. *NEUROTOXICOLOGY*. 1999; 20(2-3):327–342. [PubMed: 10385894]
40. Han J, Lee J-S, Choi D, et al. Manganese (II) induces chemical hypoxia by inhibiting HIF-prolyl hydroxylase: Implication in manganese-induced pulmonary inflammation. *Toxicol Appl Pharmacol*. 2009; 235(3):261–267. [PubMed: 19263519]
41. O'Neal SL, Zheng W. Manganese toxicity upon overexposure: a decade in review. *Current environmental health reports*. 2015; 2(3):315–328. [PubMed: 26231508]
42. ATSDR. Minimal Risk Levels for Hazardous Substances. 2015
43. NIOSH. Pocket Guide to Chemical Hazards. US Department of Health and Human Services, Public Health Service, Centers for Disease Prevention, National Institute for Occupational Safety and Health; Cincinnati, OH: 1997.
44. Bekki K, Uchiyama S, Ohta K, Inaba Y, Nakagome H, Kunugita N. Carbonyl Compounds generated from electronic cigarettes. *Int J Environ Res Public Health*. 2014; 11(11):11192. [PubMed: 25353061]

Highlights

- Certain brands of cig-a-like e-cigarettes contain high levels of nickel and chromium
- Cig-a-likes contain low levels of cadmium, compared to tobacco cigarettes
- Nickel and chromium in the e-liquid of cig-a-likes may come from nichrome heating coils

Figure 1.

Distribution of metal concentration within brands. Horizontal lines within boxes indicate medians; boxes, interquartile ranges; error bars, values within 1.5 times the interquartile range; solid circles, outlying data points.

Table 1

Metal concentrations in five commercial brands of cig-a-like e-cigarettes (µg/L).

Brand	N	Cadmium			Chromium			Lead			Manganese			Nickel		
		Mean (SD)	Median	Range	Mean (SD)	Median	Range	Mean (SD)	Median	Range	Mean (SD)	Median	Range	Mean (SD)	Median	Range
Brand A (µg/L)	10	205 (318)	12.40	322-735	21180 (5220)	213	98.6-16900	1970 (1450)	1630	500-4870	6910 (12200)	918	541-31500	22600 (24400)	15400	2040-72700
Brand B (µg/L)	10	1.17 (1.09)	0.796	0.469-1.63	788 (284)	726	550-1130	58.1 (79.4)	18.5	3.53-218	670 (283)	627	476-1200	13400 (4540)	13100	4560-20500
Brand C (µg/L)	8	1.57 (1.30)	1.17	0.157-4.18	231 (71.6)	205	162-381	5.83 (1.80)	5.15	2.70 -3.96	200 (33.9)	187	154-258	463 (132)	491	316-639
Brand D (µg/L)	10	0.982 (0.802)	0.502	0.249-2.23	76.1 (11.0)	75.6	60.2 -92.7	4.89 (0.893)	4.98	3.17-5.89	41.50 (13.9)	44.4	11.8-65.5	58.7 (22.4)	58.1	13.7-85.4
Brand E (µg/L)	10	0.415 (0.38)	0.204	0.137-1.23	53.9 (6.95)	56.7	41.5-60.79	93.4 (80.5)	69.3	7.94-233	28.7 (9.79)	26.1	15.5-48.23	114 (49.3)	134	39.3-175
LOD (µg/L) *		0.04		0.1	0.02		0.08	0.1								
Intra-laboratory ICC	48×2	0.965		0.999	0.997		1.000	1.000								
Inter-laboratory ICC	4×2			0.997			0.993			0.997			0.988			0.988

ICC: intraclass correlation coefficient. The intra-laboratory ICC was calculated from duplicate aliquots from the same e-cigarette liquid sample. Mean concentration was calculated by taking the mean of 2 duplicate samples from the same e-cigarette. The inter-laboratory ICC was calculated from duplicate analyses conducted in a subset of 4 e-cigarette liquid samples conducted at Graz University (Graz, Austria).

* LOD are calculated to a 1:20 dilution factor.