Published in final edited form as:

Addict Behav. 2017 January ; 64: 70–77. doi:10.1016/j.addbeh.2016.08.015.

Descriptive and Injunctive Network Norms Associated with Non Medical Use of Prescription Drugs among Homeless Youth

Anamika Barman-Adhikari, PhD¹, Alia Al Tayyib, PhD^{2,3}, Stephanie Begun, MSW¹, Elizabeth Bowen, PhD⁴, and Eric Rice, PhD⁵

¹Graduate School of Social Work, University of Denver, Denver, CO, USA

²Denver Public Health, Denver Health and Hospital Authority, Denver, CO, USA

³Department of Epidemiology, Colorado School of Public Health, Aurora, CO, USA

⁴School of Social Work, State University of New York, Buffalo, NY, USA

⁵School of Social Work, University of Southern California, Los Angeles, CA, USA

Abstract

Background—Nonmedical use of prescription drugs (NMUPD) among youth and young adults is being increasingly recognized as a significant public health problem. Homeless youth in particular are more likely to engage in NMUPD compared to housed youth. Studies suggest that network norms are strongly associated with a range of substance use behaviors. However, evidence regarding the association between network norms and NMUPD is scarce. We sought to understand whether social network norms of NMUPD are associated with engagement in NMUPD among homeless youth.

Methods—1,046 homeless youth were recruited from three drop-in centers in Los Angeles, CA and were interviewed regarding their individual and social network characteristics. Multivariate logistic regression was employed to evaluate the significance of associations between social norms (descriptive and injunctive) and self-reported NMUPD.

Results—Approximately 25% of youth reported past 30-day NMUPD. However, more youth (32.28%) of youth believed that their network members engage in NMUPD, perhaps suggesting some pluralistic ignorance bias. Both descriptive and injunctive norms were associated with self-reported NMUPD among homeless youth. However, these varied by network type, with presence of NMUPD engaged street-based and home-based peers (descriptive norm) increasing the likelihood of NMUPD, while objections from family-members (injunctive norm) decreasing that likelihood.

Conclusions—Our findings suggest that, like other substance use behaviors, NMUPD is also influenced by youths' perceptions of the behaviors of their social network members. Therefore,

Corresponding author: Anamika Barman-Adhikari (anamika.barmanadhikari@du.edu).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors of this manuscript report no financial interests or potential conflicts of interest.

prevention and interventions programs designed to influence NMUPD might benefit from taking a social network norms approach.

Keywords

social network norms; nonmedical use of prescription drugs; homeless youth; substance use

1. Introduction

1.1. NMUPD among youth and young adults

The nonmedical use of prescription drugs (NMUPD) is a serious public health problem in the United States (White House Office of National Drug Control Policy, 2011). NMUPD is defined as using medications that were not prescribed or were taken only for the experience or feeling that they caused, with the most commonly misused drugs being opioid pain relievers, stimulants, tranquilizers, and sedatives. The most recent data from the National Survey on Drug Use and Health indicates that an estimated 655,000, or 2.6% of adolescents' aged 12 to 17 reported NMUPD in 2014. For those aged 18 to 25, an estimated 1.6 million, or 4.4% of young adults, reported NMUPD in the same year (Center for Behavioral Health Statistics and Quality, 2015). These numbers are concerning, particularly given that NMUPD is associated with a variety of negative health correlates for young adults, such as sexual risk behaviors, mental health concerns, and drug overdose (Drazdowski, 2016; Zullig & Divin, 2012).

1.2. Social norms of NMUPD

Perception of social norms is an important aspect of drug use. Norms are defined as perceived rules or properties of a group that typify specific beliefs around what behaviors are considered acceptable or common within that group (Kincaid, 2004). Perceived social norms form a crucial component of many common theories of health behaviors (Bandura, 1977; Fishbein, Middlestadt, & Hitchcock, 1994; Fisher & Fisher, 1992), and have been linked to both behavior and behavioral intentions (Barrington et al., 2009). Perceived norms have been generally characterized as descriptive or injunctive norms. Descriptive norms indicate the perceived prevalence of a behavior in a group, whereas injunctive norms refer to perceived approval or disapproval of a behavior (Davey-Rothwell & Latkin, 2007).

Social network analysis provides a fruitful means through which one can assess the prevalence of both descriptive and injunctive norms in naturally occurring social groups (Latkin et al., 2009). There is an important distinction between "social norms" and "network norms". For example, most studies assessing social norms have used survey data, in which participant respond to a "question regarding the level of drug use their peers engage in" and those responses are used to assess perceived social norms. Researchers have found this to be a little problematic because of the ambiguity regarding who these peers are (i.e., the specificity of the reference group. In a recent meta-analysis, Borsari and Carey (2003) reported that the degree to which youth were able to correctly estimate their peers' drug use depended on the degree of familiarity and proximity of the relationship. Youth generally tend to know better what their close friends are doing compared to a hypothetical or typical peer group (Pape 2012). The use of network methods partially mitigates this issue because

respondents are asked about a very specific network (i.e. people they regularly communicate or interact with) instead of a hypothetical group of people.

Extant research shows that network norms are strongly associated with substance use behavior. However, most of the research describing the association between network norms and drug use behaviors has tended to focus either on college students, especially in relationship to alcohol and marijuana use (Barrett, Darredeau, Bordy, & Pihl, 2005; McCabe, Knight, Teter, & Wechsler, 2006; Rabiner et al., 2009) or regarding adult persons who inject drugs (Knowlton, Hua & Latkin, 2005; Kottiri, Friedman, Neaigus, Curtis, & Des Jarlais, 2002; Latkin, Kuramoto, Davey-Rothwell, & Tobin, 2010).

Additionally, studies have also found that norms are not one-dimensional by nature. More specifically, different kinds of referent groups contribute different kinds of norms surrounding both risk and protective behaviors. For example, while one's family members might promote norms around abstinence, peer groups might promote norms promoting substance use behaviors (Latkin et al., 2009). Conventional wisdom dictated that homeless youth have strained family relationships and their social networks are only comprised of other similarly situated street peers (Whitbeck, 2009). However, evidence from various studies suggests that homeless youth's relationships are not confined to street associations alone (Rice, Milburn & Rotheram-Borus, 2007; Wenzel et al., 2012; Whitbeck & Hoyt, 1999). Homeless youth relationships involve parents, other relatives, home-based peers (peers to whom homeless youths were connected before they became homeless), and social service program staff members (de la Haye et al., 2012; Ennett, Bailey & Federman, 1999; Rice, Milburn, & Monro, 2011). To the best of our knowledge, no study has yet network norms regarding NMUPD vary based on different referent groups in the homeless youth population.

Though less commonly researched, some studies have explored social norms regarding NMUPD. A recent systematic review examined risk and protective factors associated with NMUPD among youth aged 14 to 24 from a social ecological perspective and included 7 longitudinal studies, 36 cross-sectional studies, and 7 reviews (Nargiso et al., 2015). In the interpersonal domain related to peers, the review found that peer approval of substance use (an injunctive norm) and NMUPD were strongly associated, and that a close friend's use of substances (a descriptive norm) was one of the strongest and most consistent risk factors for NMUPD (Nargiso et al., 2015). Several of the studies included in the review analyzed data from the National Survey on Drug Use and Health and the majority of the others used convenience samples from schools and colleges.

1.3. NMUPD among homeless youth: individual correlates and the role of social norms

Much less is known about the prevalence of NMUPD among homeless youth and perceived social norms related to nonmedical use of prescription drugs in this population. To the best of our knowledge, this is the first study to investigate associations between social network norms and NMUPD among homeless youth. Extant data from smaller studies of homeless or street-involved youth have reported that 9% to 22% of homeless youth engage in NMUPD (Al-Tayyib, Rice, Rhoades, & Riggs, 2014; Hadland et al. 2014; Rhoades, Winetrobe, & Rice, 2014), which is several times the rate reported in the general population of youth and

young adults. One study of 16 to 24 year olds, in which 60% of participants were currently homeless, found that 69% of participants reported lifetime nonmedical use of prescription opioids, tranquilizers, and stimulants (Lankenau et al. 2012).

A vast majority of the literature on NMUPD has been focused on school or college going housed youth (Berenson & Rahman, 2011, 2011; Zullig & Divin, 2012). Fewer studies have tried to assess NMUPD specifically among homeless youth (Al-Tayyib, Rice, Rhoades, & Riggs, 2014; Rhoades, Winetrobe, & Rice, 2014). Among youth in the general population, NMUPD has been associated with depression (Zullig & Divin, 2012), post-traumatic stress disorder (PTSD) (Berenson & Rahman, 2011) and use of other substances such as heroin, marijuana and methamphetamine (Lankenau et al., 2012). In samples of homeless youth, NMUPD has been associated with hard drug use, unprotected sex, suicidal ideation and history of foster care (Al-Tayyib, Rice, Rhoades, & Riggs, 2014; Rhoades, Winetrobe, & Rice, 2014). However, beyond these two studies, not much is known about NMUPD among homeless youth.

Notably, interventions attempting to reduce drug use have successfully utilized social networks to disseminate and reinforce behavioral norms that are supportive of protective behaviors related to drug use, especially among hard-to-reach populations (Barrington et al., 2009). Unfortunately, little is known about social norms of NMUPD among homeless youth, leaving researchers and service providers with a dearth of evidence for how to design effective network interventions. Given the paucity of information related to social norms surrounding NMUPD among homeless youth, we sought to understand how social norms regarding NMUPD is associated with self-reported NMUPD in a particularly vulnerable population.

2. Methods

2.1. Procedures

As part of a social network panel study, homeless youth ages 13–24 in Los Angeles ($N=1,046$) were surveyed between October 2011 and June 2013. Participants were recruited from three homeless youth drop-in centers in Hollywood and Santa Monica across 4 different waves. Each wave of data collection was approximately 6 months apart. Youth were invited to participate at multiple waves, but only their baseline survey data were used in the current study. Any client receiving services at each respective agency was eligible to participate; this included youth living outdoors or in places not meant for human habitation, youth in shelters or transitional housing programs, and youth living with family, friends, or relatives but spending most of their time on the street. Recruitment was conducted for 19 days at each agency during each wave; during each recruitment period, researchers were present at the agency to approach youth for the duration of service provision hours. Signed voluntary informed consent was obtained from each youth, with caveats that child abuse and suicidal and homicidal intentions would be reported to appropriate authorities. Informed consent was obtained from youth 18 years of age and older and informed assent was obtained from youth 13 to 17 years old. The Institutional Review Board waived parental consent, as homeless youth under 18 years are unaccompanied minors who may not have a parent or adult guardian from whom to obtain consent.

The study consisted of two parts: a computerized self-administered survey and a social network interview. The social network interview was conducted using a free-recall name generator (Rice, Barman-Adhikari, Milburn, & Monro, 2012) to collect names of participants' network members, defined as someone with whom the youth had face-to-face, phone, or Internet contact in the last month. The social network-mapping interview conducted face-to-face by a trained research staff member. The research team used an iPad application to create the network map.). Respondents are then asked to list as many people as they can. Additionally, 17 follow-up questions were asked about specific types of interactions with each network member, and their perceptions of these youths' behaviors. All participants received \$20 in cash or gift cards as compensation for their time. The Institutional Review Board at the [redacted for review] approved all survey items and procedures. More detailed information about the study's design and data collection procedures may be found elsewhere (blinded for review).

2.2. Measures

2.2.1. NMUPD and Substance Use Behaviors—Recent NMUPD was assessed by asking how many times a participant took a prescription drug without a doctor's prescription, used more of the drug than what was prescribed, or took the drug more often than prescribed within the last 30 days (Eaton et al., 2012; Youth Risk Behavior Survey [YRBS] [CDC, 2012]). Prescription drugs included of sedatives, stimulants and opioids, however youth were not asked to separately specify which type of prescription drug that they had use nonmedically. Similarly, using items adapted from the YRBS (Eaton et al., 2012), we assessed for past 30-day use of binge drinking and marijuana, heroin, and injection drug use. Binge drinking was defined as having 5 or more drinks of alcohol in a row, that is, within a couple of hours.

2.2.2 Sociodemographic Characteristics—Demographics included age, time spent homeless (in years), gender (male vs. female), ethnicity (White, Black, Latino/a, and other and mixed race), and sexual orientation (heterosexual vs. lesbian, gay, bisexual, or queer [LGBQ]). Youth experiencing literal homelessness were defined as those who indicated that they were currently staying in a shelter (emergency or temporary), a stranger's home, hotel, motel, street, beach, tent or campsite, abandoned building, car, or bus (vs. those living with their biological family, foster family, relative, friend, group home, sober living facility, transitional living program, or own apartment but still spent considerable time on the streets).

2.2.3 Health Service Use—The frequency of past month health service utilization was assessed with the question; "I have gone to a place(s) for medical or health care services." Responses ranged from "every day or almost every day" to "not at all this month," and then were dichotomized to indicate health service use versus non-use.

2.2.4 Depression—Depressive symptoms were assessed by the 10-item Center for Epidemiological Studies Depression Scale (CES-D; Kohout, Berkman, Evans, & Cornoni-Huntley, 1993).

2.2.5 Social Norms of NMUPD—We assessed both descriptive as well as injunctive norms. In order to assess descriptive and injunctive norms regarding NMUPD, after youths finished nominating their network members, they were asked: “Out of the people you nominated, how many of them engage in NMUPD, would encourage and object you engaging in NMUPD?” This was calculated as the number of people specified by relationship type (i.e., street peers, home-based peers, family members (which included both biological and foster family, extended relatives and siblings, and finally caseworkers/staff members) whom respondents thought engaged in or perceived would object to or encourage them to engage in NMUPD. These network members are all mutually exclusive. However, every youth could have more than one network member type in their total network (i.e. youth who have a family member engaging in NMUPD could also have a street-peer engaging in NMUPD).

To account for varying sized networks, we created network member proportion variables who either engaged in NMUPD, objected to or encourage NMUPD using the total number of network members nominated in the last 90 days as the denominator. However, these proportion variables were skewed as well. Typically, in the social network literature, the way skewness has been handled is to recode proportion into categorical variables (Barrington et al., 2009; Davey-Rothwell & Latkin, 2007; Latkin et al., 2010; Tucker et al., 2011; Tyler, 2013; Valente & Auerswald, 2013). The median is used to create a threshold for measures that are not uniformly distributed (Wang, Fan & Wilson, 1996). Based on the median, the descriptive norm and the injunctive norms (encourage and object NMUPD) were consequently dichotomized as either no network member (coded as 0) or at least one or more network members (coded as 1) who participants believed engaged in NMUPD or would encourage or object to them engaging in NMUPD.

2.3. Analytic approach

Data analyses were conducted using SAS Version 9.2 (SAS Institute Inc., 2008). Analyses examined bivariate associations between socio-demographic variables and perceived NMUPD social norms and self-reported NMUPD, using chi-square statistics for categorical variables and t-tests for continuous variables. Two-sided significance from Pearson’s chi square tests was reported for all categorical variables and t-tests were used for continuous level variables. The degree of freedom for all binary variables was 1. Bivariate tests were not conducted for any of the service-related network variables because of the sparse nature of the cell sizes. Additionally, because these service-related networks could not be independently included in bivariate and multivariate logistic regression models, they were dropped from the subsequent bivariate and multivariate logistic analyses.

To evaluate the significance of the independent associations between perceived NMUPD norms and self-reported NMUPD, logistic regression was conducted. In order to preserve statistical power and degrees of freedom (because of the large number of variables being examined), bivariate analyses were first conducted to examine unadjusted associations between study variables and outcome measures. Any variable that was significantly associated with the outcome at $p < .10$ was retained in the multivariate model. Multivariate models were then constructed based on these analyses (Hosmer & Lemeshow, 2004).

Sociodemographic characteristics (i.e. age, gender, sexual orientation, race/ethnicity) were controlled for regardless of their significance in bivariate analysis. Significance levels were adjusted within each hypothesis using the step-down Bonferroni (Holm) procedure (SAS, 9.2.). The benefit of using step-down methods is that the tests are made more powerful (smaller adjusted p -values) while, in most cases, maintaining strong control of the familywise error rate (Holm, 1979).

Since we were interested in whether or not youth had initiated use and not on dependence or abuse, a dichotomous variable was created to indicate 0 = “no NMUPD use” versus 1 = “recent NMUPD use.” It is important to note that dichotomization does not significantly reduce statistical power and yields meaningful effect sizes (i.e. odds ratios) (Farrington & Loeber, 2000). Individual ORs were interpreted in accordance to Chen, Cohen, and Chen’s (2010) guidelines for small (OR = 1.46), medium (OR = 2.49), and large (OR = 4.14) effect sizes when predicted outcomes are present in at least or more than 10% of the general population. Tjur R-square (Tjur, 2009) was also used to assess overall strength of association. Tjur R² was .28, which represents a moderate-to-strong effect size (Ferguson, 2009). Furthermore, multivariate analyses were restricted to participants without missing data for the variables included in the models. Therefore, the sample size for the multivariate model is smaller than the study’s total sample size ($n=969$).

3. Results

3.1. Descriptive Statistics

Socio demographic characteristics are presented in Table 1. The sample was predominantly male (70.27%) and heterosexual (75.10%). Average age of participants was 21.34 years. In terms of race/ethnicity, Whites represented the largest group (38.24%). A total of 247 youth (23.61%) reported engaging in NMUPD in the past 30 days. Participants differed significantly in their nonmedical use of prescription drugs by variables including race/ethnicity; health care access in the past 30 days; use of heroin, marijuana, or injection drugs in the past 30 days; and depressive symptoms. Participants who reported recent NMUPD compared to those who did not were more likely to be White (44.53% vs. 36.30%, $p < 0.05$) and less likely to be African-American (14.75% vs. 26.03%, $p < .0001$). Youth who accessed health care in the past 30 days were also more likely to report current NMUPD (51.74% vs. 39.41%, $p < .01$). Additionally, use of other substances was significantly associated with self-reported NMUPD, with increased rates of NMUPD for participants who reported using heroin (45.75% vs. 5.51%, $p < .0001$), marijuana (64.78% vs. 54.19%, $p < .01$), or injection drug use (43.72% vs. 4.63%, $p < .0001$). On average, participants who reported recent NMUPD had higher depression scores on the CES-D than those who did not report NMUPD ($M = 18.79$ vs. 17.02 , $p < .01$).

3.2. Network Norm Characteristics

Social network norm characteristics are also presented in Table 1. A total of 360 participants (32.28%) reported having any network members (aggregated across all network types) that they perceived would engage in NMUPD. There were variations when this was stratified by network type. More youth thought that their street-based peers were more likely to engage in

NMUPD (18.99%) compared to home-based peers (9.02%), family-based network members (3.89%) and service-related network members (0.38%) engaging in NMUPD. In regards to injunctive norms, more youth believed that their network members would object to them engaging in NMUPD relative to perceptions of receiving encouragement from their network members.

Bivariate statistics indicated that participants differed significantly in their self-reported NMUPD based on these social norms especially by network type. Youth were more likely to engage in NMUPD if they had at least one street-based peer (32.68% vs. 17.27%, $p < .0001$) or home-based peer (16.93 vs. 6.51%, $p < .0001$) who engage in NMUPD. Similarly, youth were more likely to endorse recent NMUPD if they had at least one street-based peer (26.88% vs. 10.64%, $p < .0001$) or home-based peer (9.84 vs. 2.88 %, $p < .0001$) who encouraged them to engage in NMUPD. On the contrary, youth who had at least one family member who objected to NMUPD, youth were less likely to engage in NMUPD (45.25% vs. 58.69%, $p < .0001$).

3.3. Bivariate and Multivariate Logistic Regression Analyses

Bivariate and multivariate statistics are presented in Table 2. As noted above, only variables significant in the bivariate model at the $p < .10$ level were included in the bivariate model. In the multivariate model, having accessed health care in the last 30 days, recent heroin, marijuana, injection drug use, and NMUPD. Youth who accessed health care in the past 30 days were 1.91 times more likely to engage in NMUPD ($p < .001$). Youth who reported engaging in recent heroin, marijuana, and injection drug use were 4.50 times ($p < .0001$), 2.64 times ($p < .0001$) and 6.85 times ($p < .0001$) more likely to engage in NMUPD, respectively. Youth who perceived that at least one street-based network member engaged in NMUPD were 1.49 times more likely to report engaging in NMUPD themselves ($p < .001$). Alternatively, youth who perceived that at all family-based people in their network would object to them engaging in NMUPD were 31% less likely to engage in NMUPD ($p < .001$). When the final model was rerun using the more conservative Bonferroni step-down corrected alpha-levels, all variables remained significantly associated with recent NMUPD.

4. Discussion

The current study advances our understanding of NMUPD among homeless youth, a high-risk and especially hard-to-reach population, in many ways. First, this study, to the best of our knowledge, is the first to demonstrate an association between perceived norms of NMUPD and self-reported NMUPD among homeless youth. Both descriptive and injunctive norms were strongly and significantly associated with NMUPD, a finding consistent with previous research on social norms of NMUPD, although previous studies focused on housed and college attending populations (Meisel & Goodie, 2015).

While less than one-quarter of youth indicated that they had engaged in NMUPD, a greater percentage of youth (32.28%) believed that their network members (aggregating across all network relationships) engaged in NMUPD. This finding indicates that there might be an “overestimation” of the prevalence of NMUPD among homeless youth, perhaps suggesting a “pluralistic ignorance” bias, in which individuals tend to misperceive their peers’ drug use,

regardless of their own status as users or nonusers (Henry, Kobus, & Schoeny, 2011). Several studies have found evidence consistent with this phenomenon (Bourgeois & Bowen 2001; Henry et al., 2011). The social norms intervention approach, however, is designed to correct the demonstrated inconsistency between actual and perceived behaviors (Ott & Doyle, 2005), and could provide an appropriate approach to address NMUPD with homeless youth. A few interventions have in fact been successful in demonstrating reduced misperceptions of norms of NMUPD (Schinke et al., 2009). The current study was not specifically designed to understand the discrepancy between actual and perceived NMUPD, but given these preliminary findings, it would be prudent to explore this issue further.

Second, our study's focus on injunctive norms furthers our understanding of perceived approval and disapproval as they relate to NMUPD. Injunctive norms continued to be significantly associated with NMUPD even after controlling for (i.e. holding constant) descriptive norms in the multivariate model. Extant research has found that descriptive and injunctive norms have an independent association with behavior across a number of behaviors, including drug use (McMillan & Conner, 2003), engagement in sexual risk (Peterson, Rothenberg, Kraft, Beeker, & Trotter, 2009), physical activity (Rhodes & Courneya, 2003), and violence (Norman, Clark, & Walker, 2005).

Injunctive norms, as noted before, refer to both perceived approval and disapproval of a behavior. What is encouraging about our findings is that while there is significant perceived disapproval of NMUPD use among homeless youths' network members, there is relatively little perceived approval of it. Specifically, in this study, we found that only 7.50% of youth believed that their street-based peers and 4.56% believed that their home-based peers would encourage them to engage in NMUPD. However, a number of preventive interventions have based on the theory that there is active pressure on youth (especially in the form of peer encouragement) to use substances; therefore, interventions have generally sought to provide the young person with social and interpersonal skills that would enable them to resist such proactive pressure (Coggans & McKellar, 1994; Denscombe, 2001). More studies are needed to understand whether the absence of such peer pressure is a matter of perception, or if peer pressure works in other, subtle ways not captured by survey data (Denscombe, 2001). It is possible that observational studies or qualitative studies could better capture the nuances of how such influence is exerted.

Perhaps the most compelling aspects of these data is the finding that the role of descriptive and injunctive norms on NMUPD varies depending on the nature of the referent (Borsari & Carey, 2003; Cho, 2006). Our findings indicate that while the presence of NMUPD engaged street-based and home-based peers increased the likelihood of youth reporting recent NMUPD, objections from family-members decreasing that likelihood. These findings support the "Social Identity Theory" (Turner & Tajfel, 1986), which proposes that norms are linked to specific social groups, and identification with these groups determines what will be regarded as normative. It can be assumed that homeless youth who are connected to family members have more role models to rely on in understanding what can be considered normative behavior. In addition, it is also conceivable that the conversations that youth are having with their family members might act as sources of informational support, which is known, to influence the expression of norms and consequently engagement in behavior.

However, it is important to mention that this study did not collect any data on the nature of these interactions, more specifically, what was discussed. A future study that collects detailed network level information on ties and the content of interactions across those ties would do much to increase our understanding of these processes, which could be critical in informing clinical interventions with youth and families.

In addition to network norm characteristics, our study found associations between other sociodemographic and behavioral characteristics and self-reported NMUPD. What is perhaps most compelling is the association between recent health care use and engagement in NMUPD. Nearly half (42%) of participants reported accessing health care in the past 30 days. Emerging research has indicated that one of the principal factors connected to increased NMUPD among youth and young adults is increased access to these drugs (McCabe, Cranford, Boyd & Teter, 2007). This has been in part attributed to less than conservative prescribing practices of doctors of opioid painkillers and changing medical guidelines (Twombly & Holtz, 2008). However, this also underscores the critical role that physicians can play in screening for and communicating about the negative effects of nonmedical use of prescription drugs with their patients. This may be particularly important for homeless youth, as building trusting relationships with health care providers is one of the key avenues through which substance abusing homeless youth are typically able to access treatment and resources (Hudson, Nyamathi, & Sweat, 2008).

Similar to other studies of NMUPD, we found a significant association between NMUPD and heroin use (Compton, Jones, & Baldwin, 2016). However, much of the existing literature describing the relationship between NMUPD and heroin use is in the adult population. Our findings add to the much smaller body of evidence suggesting that the association between NMUPD and heroin is present at a younger age. Though we did not capture information related to the specific route of administration for heroin, we also found a strong association between NMUPD and injection drug use.

In addition, youth who reported engaging in marijuana use were more likely to engage in NMUPD. This study reinforces previous evidence (Arria et al., 2008; Catalano et al., 2011; Haddox et al., 2014; McCabe et al., 2005), which has found that marijuana and NMUPD are linked especially among college students. This study provides some preliminary evidence that marijuana could prove to be a gateway to NMUPD even among homeless youth. Given that these substance use behaviors are clustered, community-based interventions that address substance misuse more universally could provide a viable means of addressing NMUPD among youth populations.

4.1. Limitations and conclusions

Certain study limitations must be noted. First, the study includes cross-sectional data, thereby reducing the ability to draw causal conclusions. Thus, it is plausible that substance-using youth are more likely to perceive that their peers also use substances, and/or would approve of their substance use. Future research would benefit from longitudinal investigations of how network dynamics function over time to better discern the causal pathways through which network attributes facilitate the formation of substance use norms among homeless youth. Additionally, all behavioral study variables are self-reported, which

could introduce bias. It is perhaps inevitable that some behaviors are under- or over-reported due to social desirability, despite reminders given to participants that data were confidential and that they could skip questions while completing the questionnaire. Further, the study's use of a computer-assisted self-interview modality, which has been shown to diminish threats of social desirability and impression management (Schroder, Carey, & Venable, 2003), is likely to generate less biased responses. Moreover, norms variables were based on youths' perceptions and, as aforementioned, may not be accurately reflective of reality. Additional research regarding social norms would benefit from further incorporating independent confirmations of norms. Also, we did not assess for perceived availability of prescription drugs in our study. It is possible that perceived norms may have been confounded with availability/access of prescription drugs via network members. However, the social network interview was separate from the individual survey and was interviewer administered. We believe that knowing that these are separate questions assessing their perceptions of their network members' and having an interviewer clarify it for them might have reduced the confusion noted above. Last, but not the least, respondents were not asked to identify the type of prescription drug (i.e., stimulants, sedatives, and/or opioids) that they had use nonmedically. Previous studies (Kelly et al., 2014) have found that young adults who engage in NMUPD have different patterns and clusters of use and therefore are important to assess.

4.2. Conclusion

Our findings help further our understanding of the relationship between perceived norms of NMUPD and self-reported NMUPD among homeless youth. Given that NMUPD may be associated with youths' perceived prevalence of their peers' use (even if an overestimation of that use), and also by their perceived peers' approval or disapproval of NMUPD, such an understanding can help inform development of network interventions that specifically target prevention of NMUPD among homeless youth.

Acknowledgments

Funded by NIMH R01 MH093336

References

- Al-Tayyib AA, Rice E, Rhoades H, Riggs P. Association between prescription drug misuse and injection among runaway and homeless youth. *Drug and Alcohol Dependence*. 2014; 134:406–409. [PubMed: 24300900]
- Arria AM, O'Grady KE, Caldeira KM, Vincent KB, Wish ED. Nonmedical use of prescription stimulants and analgesics: Associations with social and academic behaviors among college students. *Pharmacotherapy*. 2008; 38:1045–1060.
- Bandura A. Self-efficacy: toward a unifying theory of behavioral change. *Psychological Review*. 1977; 84(2):191–215. [PubMed: 847061]
- Barrett SP, Darredeau C, Bordy LE, Pihl RO. Characteristics of methylphenidate misuse in a university student sample. *Canadian Journal of Psychiatry*. 2005; 50(8):457–461. [PubMed: 16127963]
- Barrington C, Latkin C, Sweat MD, Moreno L, Ellen J, Kerrigan D. Talking the talk, walking the walk: Social network norms, communication patterns, and condom use among the male partners of female sex workers in la Romani, Dominican Republic. *Social Science & Medicine*. 2009; 68(11):2037–2044. [PubMed: 19356834]

- Berenson AB, Rahman M. Prevalence and correlates of prescription drug misuse among young, low-income women receiving public healthcare. *Journal of addictive diseases*. 2011; 30(3):203–215. [PubMed: 21745043]
- Borsari B, Carey KB. Descriptive and injunctive norms in college drinking: A meta-analytic integration. *Journal of Studies on Alcohol*. 2003; 64:331–341. [PubMed: 12817821]
- Bourgeois MJ, Bowen A. Self-organization of alcohol-related attitudes and beliefs in a campus housing complex: An initial investigation. *Health Psychology*. 2001; 20(6):434–437. [PubMed: 11714185]
- Catalano RF, White HR, Fleming CB, Haggerty KP. Is nonmedical prescription opiate use a unique form of illicit drug use? *Addictive Behaviors*. 2011; 36:79–86. [PubMed: 20864261]
- Center for Behavioral Health Statistics and Quality. Behavioral health trends in the United States: Results from the 2014 National Survey on Drug Use and Health. 2015. HHS Publication No. SMA 15-4927, NSDUH Series H-50 Retrieved from: <http://www.samhsa.gov/data/>
- Chen H, Cohen P, Chen S. How big is a big odds ratio? Interpreting the magnitudes of odds ratios in epidemiological studies. *Communications in Statistics—Simulation and Computation*. 2010; 39(4):860–864.
- Cho H. Influences of norm proximity and norm types on binge and non-binge drinkers: Examining the under-examined aspects of social norms interventions on college campuses. *Journal of Substance Use*. 2006; 11:417–429.
- Coggans N, McKellar S. Drug use amongst peers: Peer pressure or peer preference? *Drugs: Education, Prevention and Policy*. 1994; 1(1):15–26.
- Compton WM, Jones CM, Baldwin GT. Relationship between Nonmedical Prescription-Opioid Use and Heroin Use. *New England Journal of Medicine*. 2016; 374(2):154–163. [PubMed: 26760086]
- Davey-Rothwell MA, Latkin CA. Gender differences in social network influence among injection drug users: Perceived norms and needle sharing. *Journal of Urban Health*. 2007; 84(5):691–703. [PubMed: 17682947]
- de la Haye K, Green HD Jr, Kennedy DP, Zhou A, Golinelli D, Wenzel SL, Tucker JS. Who is supporting homeless youth? Predictors of support in personal networks. *Journal of Research on Adolescence*. 2012; 22:604–616. [PubMed: 23204810]
- Denscombe M. Peer group pressure, young people and smoking: New developments and policy implications. *Drugs: Education, Prevention and Policy*. 2001; 8(1):7–32.
- Drazdowski TK. A systematic review of the motivations for the nonmedical use of prescription drugs in young adults. *Drug and Alcohol Dependence*. 2016; Advance online publication. doi: 10.1016/j.drugalcdep.2016.01.011
- Eaton DK, Kann L, Kinchen S, Shanklin S, Flint KH, Hawkins J, Centers for Disease Control and Prevention (CDC). Youth risk behavior surveillance – United States, 2011. *Morbidity and Mortality Weekly Report*. 2012; 61(4):1–162.
- Ennett ST, Bailey SL, Federman EB. Social network characteristics associated with risky behaviors among runaway and homeless youth. *Journal of Health and Social Behavior*. 1999; 40:63–78. [PubMed: 10331322]
- Farrington DP, Loeber R. Some benefits of dichotomization in psychiatric and criminological research. *Criminal Behaviour and Mental Health*. 2000; 10(2):100–122.
- Ferguson CJ. An effect size primer: A guide for clinicians and researchers. *Professional Psychology: Research and Practice*. 2009; 40(5):532–538.
- Fiellin LE, Tetrault JM, Becker WC, Fiellin DA, Hoff RA. Previous use of alcohol, cigarettes, and marijuana and subsequent abuse of prescription opioids in young adults. *Journal of Adolescent Health*. 2013; 52(2):158–163. [PubMed: 23332479]
- Fishbein, M.; Middlestadt, SE.; Hitchcock, PJ. Preventing AIDS. Springer; 1994. Using information to change sexually transmitted disease-related behaviors; p. 61-78.
- Fisher JD, Fisher WA. Changing AIDS-risk behavior. *Psychological Bulletin*. 1992; 111(3):455–474. [PubMed: 1594721]
- Haddox JD, Weiler RM, Pealer LN, Barnett TE. Early initiation of alcohol or marijuana use and nonmedical use of prescription drugs. *Drug & Alcohol Dependence*. 2014; 140:e77.

- Hadland SE, DeBeck K, Kerr T, Feng C, Montaner JS, Wood E. Prescription opioid injection and risk of hepatitis C in relation to traditional drugs of misuse in a prospective cohort of street youth. *BMJ Open*. 2014; 4(7) e005419-2014-005419. doi: 10.1136/bmjopen-2014-005419
- Henry DB, Kobus K, Schoeny ME. Accuracy and bias in adolescents' perceptions of friends' substance use. *Psychology of Addictive Behaviors*. 2011; 25(1):80–89. [PubMed: 21244119]
- Holm S. A simple sequentially rejective multiple test procedure. *Scandinavian journal of statistics*. 1979:65–70.
- Hosmer, DW., Jr; Lemeshow, S. *Applied logistic regression*. John Wiley & Sons; 2004.
- Hudson AL, Nyamathi A, Sweat J. Homeless youths' interpersonal perspectives of health care providers. *Issues in Mental Health Nursing*. 2008; 29(12):1277–1289. [PubMed: 19052945]
- Kelly BC, Wells BE, Pawson M, LeClair A, Parsons JT. Combinations of prescription drug misuse and illicit drugs among young adults. *Addictive Behaviors*. 2014; 39(5):941–944. [PubMed: 24462348]
- Kincaid DL. From innovation to social norm: Bounded normative influence. *Journal of Health Communication*. 2004; 9(S1):37–57. [PubMed: 14960403]
- Knowlton A, Hua W, Latkin C. Social support networks and medical service use among HIV-positive injection drug users: Implications to intervention. *AIDS Care*. 2005; 17(4):479–492. [PubMed: 16036234]
- Kohout FJ, Berkman LF, Evans DA, Cornoni-Huntley J. Two shorter forms of the CES-D (Center for Epidemiological Studies Depression) depression symptoms index. *Journal of Aging and Health*. 1993; 5(2):179–193. [PubMed: 10125443]
- Kottiri BJ, Friedman SR, Neaigus A, Curtis R, Des Jarlais DC. Risk networks and racial/ethnic differences in the prevalence of HIV infection among injection drug users. *Journal of Acquired Immune Deficiency Syndromes* (1999). 2002; 30(1):95–104. [PubMed: 12048369]
- Lankenau SE, Schrager SM, Silva K, Kecojevic A, Bloom JJ, Wong C, Iverson E. Misuse of prescription and illicit drugs among high-risk young adults in Los Angeles and New York. *Journal of Public Health Research*. 2012; 1(1):22–30. [PubMed: 22798990]
- Latkin CA, Donnell D, Metzger D, Sherman S, Aramrattana A, Davis-Vogel A, Celentano DD. The efficacy of a network intervention to reduce HIV risk behaviors among drug users and risk partners in Chiang Mai, Thailand and Philadelphia, USA. *Social Science & Medicine*. 2009; 68:740–748. [PubMed: 19070413]
- Latkin C, Kuramoto S, Davey-Rothwell M, Tobin K. Social norms, social networks, and HIV risk behavior among injection drug users. *AIDS and Behavior*. 2010; 14(5):1159–1168. [PubMed: 19466537]
- McCabe SE, Knight JR, Teter CJ, Wechsler H. Non-medical use of prescription stimulants among US college students: Prevalence and correlates from a national survey. *Addiction*. 2005; 100(1):96–106. [PubMed: 15598197]
- McCabe SE, Teter CJ, Boyd CJ, Knight JR, Wechsler H. Nonmedical use of prescription opioids among US college students: Prevalence and correlates from a national survey. *Addictive behaviors*. 2005; 30(4):789–805. [PubMed: 15833582]
- McCabe SE, Cranford JA, Boyd CJ, Teter CJ. Motives, diversion and routes of administration associated with nonmedical use of prescription opioids. *Addictive behaviors*. 2007; 32(3):562–575. [PubMed: 16843611]
- McMillan B, Conner M. Applying an extended version of the theory of planned behavior to illicit drug use among Students1. *Journal of Applied Social Psychology*. 2003; 33(8):1662–1683.
- Meisel MK, Goodie AS. Predicting prescription drug misuse in college students' social networks. *Addictive Behaviors*. 2015; 45:110–112. [PubMed: 25661988]
- Nargiso JE, Ballard EL, Skeer MR. A systematic review of risk and protective factors associated with nonmedical use of prescription drugs among youth in the United States: A social ecological perspective. *Journal of Studies on Alcohol and Drugs*. 2015; 76(1):5–20. [PubMed: 25486389]
- Norman P, Clark T, Walker G. The theory of planned behavior, descriptive norms, and the moderating role of group identification. *Journal of Applied Social Psychology*. 2005; 35(5):1008–1029.
- Ott CH, Doyle LH. An evaluation of the small group norms challenging model: Changing substance use misperceptions in five urban high schools. *The High School Journal*. 2005; 88(3):45–55.

- Pape H. Young people's overestimation of peer substance use: an exaggerated phenomenon? *Addiction*. 2012; 107(5):878–884. [PubMed: 22324656]
- Peterson JL, Rothenberg R, Kraft JM, Beeker C, Trotter R. Perceived condom norms and HIV risks among social and sexual networks of young African American men who have sex with men. *Health Education Research*. 2009; 24(1):119–127. DOI: 10.1093/her/cyn003 [PubMed: 18281710]
- Rabiner DL, Anastopoulos AD, Costello EJ, Hoyle RH, McCabe SE, Swartzwelder HS. The misuse and diversion of prescribed ADHD medications by college students. *Journal of Attention Disorders*. 2009; 13(2):144–153. DOI: 10.1177/1087054708320414 [PubMed: 19448150]
- Rhoades H, Winetrobe H, Rice E. Prescription drug misuse among homeless youth. *Drug and Alcohol Dependence*. 2014; 138:229–233. [PubMed: 24613220]
- Rhodes RE, Courneya KS. Investigating multiple components of attitude, subjective norm, and perceived control: An examination of the theory of planned behaviour in the exercise domain. *British Journal of Social Psychology*. 2003; 42(1):129–146. [PubMed: 12713760]
- Rice E, Milburn NG, Rotheram-Borus MJ. Pro-social and problematic social network influences on HIV/AIDS risk behaviours among newly homeless youth in Los Angeles. *AIDS Care*. 2007; 19:697–704. [PubMed: 17505933]
- Rice E, Milburn NG, Monro W. Social networking technology, social network composition, and reductions in substance use among homeless adolescents. *Prevention Science*. 2011; 12:80–88. [PubMed: 21194011]
- Rice E, Barman-Adhikari A, Milburn NG, Monro W. Position-specific HIV risk in a large network of homeless youths. *American Journal of Public Health*. 2012; 102(1):141–147. [PubMed: 22095350]
- Schinke SP, Fang L, Cole KC. Substance use among early adolescent girls: Risk and protective factors. *Journal of Adolescent Health*. 2008; 43(2):191–194. [PubMed: 18639794]
- Schroder KE, Carey MP, Venable PA. Methodological challenges in research on sexual risk behavior: I. Item content, scaling, and data analytical options. *Annals of Behavioral Medicine*. 2003; 26(2):76–103. [PubMed: 14534027]
- Tjur T. Coefficients of determination in logistic regression models—A new proposal: The coefficient of discrimination. *The American Statistician*. 2009; 63(4):366–372.
- Tucker JS, Hu J, Golinelli D, Kennedy DP, Green HD, Wenzel SL. Social network and individual correlates of sexual risk behavior among homeless young men who have sex with men. *Journal of Adolescent Health*. 2012; 51(4):386–392. [PubMed: 22999840]
- Turner JC, Tajfel H. The social identity theory of intergroup behavior. *Psychology of intergroup relations*. 1986:7–24.
- Tyler KA. Homeless youths' HIV risk behaviors with strangers: Investigating the importance of social networks. *Archives of sexual behavior*. 2013; 42(8):1583–1591. [PubMed: 23613136]
- Twombly EC, Holtz KD. Teens and the misuse of prescription drugs: Evidence-based recommendations to curb a growing societal problem. *The journal of primary prevention*. 2008; 29(6):503–516. [PubMed: 19015990]
- Valente AM, Auerswald CL. Gender differences in sexual risk and sexually transmitted infections correlate with gender differences in social networks among San Francisco homeless youth. *Journal of Adolescent Health*. 2013; 53(4):486–491. [PubMed: 23871131]
- Wang L, Fan X, Willson VL. Effects of nonnormal data on parameter estimates and fit indices for a model with latent and manifest variables: An empirical study. *Structural Equation Modeling: A Multidisciplinary Journal*. 1996; 3(3):228–247.
- Wenzel S, Holloway I, Golinelli D, Ewing B, Bowman R, Tucker J. Social networks of homeless youth in emerging adulthood. *Journal of Youth and Adolescence*. 2012; 41(5):561–571. [PubMed: 21863378]
- White House Office of National Drug Control Policy. Epidemic: Responding to America's prescription drug abuse crisis. 2011. Retrieved from: https://www.whitehouse.gov/sites/default/files/ondcp/policy-and-research/rx_abuse_plan.pdf
- Whitbeck, LB.; Hoyt, DR. Nowhere to grow: Homeless and runaway adolescents and their families. Transaction Publishers; 1999.

Zullig KJ, Divin AL. The association between nonmedical prescription drug use, depressive symptoms, and suicidality among college students. *Addictive Behaviors*. 2012; 37(8):890–899. [PubMed: 22541802]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Highlights

- 1.** 23.61% of homeless youth (n=1046) reported non-medical use of prescription drugs (NMUPD).
- 2.** Significant associations were found between perceived network norms (descriptive and injunctive) and NMUPD among this at-risk population.
- 3.** The associations between network norms and NMUPD varied by network type, with presence of NMUPD engaged street-based and home-based peers increasing the likelihood of youth reporting recent NMUPD, while objections from family-members decreasing that likelihood.
- 4.** Findings also indicate associations between other drug use (marijuana, heroin and injection drug use) and NMUPD.
- 5.** Interventions designed to target social network norms of NMUPD might be a viable strategy to reduce NMUPD among homeless youth.

Table 1Sample Characteristics of Homeless Youth in Los Angeles ($n=1,046$)

	Full Sample		Recent NMUPD		No NMUPD	
	<i>n</i>	%	<i>n</i> = 247	% = 23.61	<i>n</i> = 799	
Gender						
Male	735	70.27	165	66.80	570	
Female	311	29.73	82	33.20	229	
Sexual Orientation ($n = 1,010$)						
Heterosexual	759	75.10	159	72.27	600	
LGBQ	251	24.90	61	27.73	190	
Race/Ethnicity						
African-American	244	23.33	36	14.57	208	
Latino	139	13.29	31	12.55	108	
Other Race	212	20.26	42	4.02	170	
White	400	38.24	110	44.53	290	
Traveler						
Yes	377	36.00	99	34.79	278	
Accessed Health Care in Past 30 Days ($n=980$)						
Yes	411	41.90	104	51.74	307	
Alcohol Use (Binge Drinking) in Past 30 Days						
Yes	936	89.50	225	91.09	711	
Heroin Use in Past 30 Days						
Yes	157	15.00	113	45.75	134	
Marijuana Use in Past 30 Days						
Yes	593	56.70	160	64.78	433	
Injection Drug Use in Past 30 Days						
Yes	145	13.90	108	43.72	37	
Descriptive Norms						
All Alters Using Prescription Drugs						
Any	360	32.28	131	48.99	229	
Street-Based Peer Networks engage in NMUPD						

	Full Sample		Recent NMUPD		No NMUPD	
	n	%	n =247	%=23.61	n =799	
Any	221	20.99	83	32.68	138	
Home-Based Peer Networks engage in NMUPD						
Any	95	9.02	43	16.93	52	
Family-Based Networks engage in NMUPD						
Any	41	3.89	16	4.00	25	
Service-Related Networks engage in NMUPD						
Any	4	0.38	—	—	—	
Injunctive Norms (Encouragement)						
All Alters Encouraging Prescription Drug Use						
Any	156	14.53	70	26.88	86	
Street-Based Peer Networks encourage NMUPD						
Any	79	7.50	38	14.96	41	
Home-Based Peer Networks encourage NMUPD						
Any	48	4.56	25	9.84	23	
Family-Based Networks encourage NMUPD						
Any	28	2.66	8	3.00	20	
Proportion of Service Related Networks encourage NMUPD						
Any	2	0.19	—	—	—	
Injunctive Norms (Object)						
All Alters Objecting to Prescription Drug Use						
Any	776	73.69	158	70.16	618	
Street-Based Peer Networks Object NMUPD						
Any	509	48.34	111	43.70	398	
Home-Based Peer Networks Object NMUPD						
Any	502	47.67	108	42.52	394	
Family-Based Networks Object NMUPD						
Any	577	54.80	138	45.25	439	
Service Related Networks Object NMUPD						
Any	4	0.38	—	—	—	

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

	Full Sample		Recent NMUPD		No NMUPD	
	<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>	<i>Mean</i>	<i>SD</i>
Age	21.34	2.16	21.46	2.12	21.31	
Time Homeless (0–7)	3.63	1.35	3.96	1.60	3.52	
CESD Score (0–30)	18.37	9.74	18.79	9.00	17.02	

* p<0.05.
** p<0.01,
*** p<0.001,
**** p<0.0001

Note: Bivariate tests not conducted for service related networks because of the sparse nature of cell sizes

Table 2
Multivariate Logistic Regression of Social Network Norms and Prescription Drug Misuse (PDM)

	Prescription Drug Misuse			
	Unadj OR	95% CI	p	Adj OR
Demographics				
Age	1.03	0.96	1.11	1.04
Male	1.05	0.76	1.46	1.05
LGBQ (Ref= Heterosexual)	1.21	0.87	1.70	1.23
Race (Ref=White)				
African-American	0.47	0.32	0.69	*** 0.72
Latino	0.89	0.58	1.36	0.62
Other Race	0.73	0.51	1.06	0.68
Behavioral/Situational Variables				
Traveler	0.61	0.44	0.84	** —
Accessed Health Care in last 30 days	2.34	1.75	3.12	*** 1.91
CESD Score (Depression)	1.02	1.01	1.04	* 1.01
Length of Time Homeless				—
Substance Use (Past 30 Days)				
Alcohol Use (Binge drinking)	1.31	0.80	2.13	—
Heroin Use	15.37	10.39	22.72	*** 4.50
Marijuana Use	1.44	1.08	1.92	** 2.64
Injection Drug Use	17.04	11.28	25.73	*** 6.85
Perceived Social Norms in Network				
<i>Descriptive Norms</i>				
Street-Based Peer Networks engage in NMUPD	1.62	1.35	1.93	*** 1.49
Home-Based Peer Networks engage in NMUPD	1.45	1.15	1.82	*** 1.09
Family-Based Networks engage in NMUPD	1.28	0.80	2.03	—
<i>Injunctive Norms (Encouragement)</i>				
Street-Based Peer Networks encourage NMUPD	1.42	1.21	1.68	*** 1.12
Home-Based Peer Networks encourage NMUPD	1.62	1.22	2.16	*** 1.04
Family-Based Networks encourage NMUPD	1.50	0.88	2.55	—

Prescription Drug Misuse				
	<i>Unadj OR</i>	<i>95% CI</i>	<i>p</i>	<i>Adj OR</i>
<i>Injunctive Norms (Object)</i>				
Street-Based Peer Networks Object NMUPD	0.96	0.91 1.00	—	—
Home-Based Peer Networks Object NMUPD	0.95	0.89 1.01	—	—
Family-Based Networks Object NMUPD	0.27	0.09 0.80	****	0.69
<i>n</i>				969
AIC				736.50
SC				813.37
−2 log likelihood				704.50
Wald chi-square				147.48****
Tjur R2				0.28

Note. LGBQ = lesbian, gay, bisexual, or questioning; OR = odds ratio; CI = confidence interval; AIC = Akaike information criterion; SC = Schwarz criterion; df = degrees of freedom

* p<0.05.

** p<0.01,

*** p<0.001,

**** p<0.0001

Note: Only variables significant in the bivariate model at p<.10 included in the multivariate model