Published in final edited form as:

Behav Modif. 2016 September ; 40(5): 663–677. doi:10.1177/0145445515610315.

Distress Tolerance Among Students Referred for Treatment Following Violation of Campus Cannabis Use Policy: Relations to Use, Problems, and Motivation

Julia D. Buckner¹, Emily R. Jeffries¹, Meredith A. Terlecki², and Anthony H. Ecker¹

¹Louisiana State University, Baton Rouge, USA

²University of East London, UK

Abstract

Students referred to treatment after violating campus drug policies represent a high-risk group. Identification of factors related to these students' cannabis use could inform prevention and treatment efforts. Distress tolerance (DT) is negatively related to substance-related behaviors and may be related to high-risk cannabis use vulnerability factors that can impact treatment outcome. Thus, the current study tested whether DT was related to cannabis use frequency, cannabis-related problems, and motivation to change cannabis use among 88 students referred for treatment after violating campus cannabis policies. DT was robustly, negatively related to cannabis use and related problems. DT was also significantly, negatively correlated with coping, conformity, and expansion motives. DT was directly and indirectly related to cannabis problems via coping (not conformity or expansion) motives. Motives did not mediate the relation of DT to cannabis use frequency. DT may be an important target in treatment with students who violate campus cannabis policies.

Keywords

mandated students; cannabis; marijuana; distress tolerance; motivation

Over one third of college students endorse current cannabis use, a rate comparable with non-college same-age peers (Johnston, O'Malley, Bachman, Schulenberg, & Miech, 2014). Furthermore, nearly one fourth of past-year cannabis using first-year students meet *Diagnostic and Statistical Manual of Mental Disorders* (4th ed.; *DSM-IV*; American Psychiatric Association, 1994) criteria for a cannabis use disorder (Caldeira, Arria, O'Grady, Vincent, & Wish, 2008), and frequent cannabis use among college students is associated

Reprints and permissions: sagepub.com/journalsPermissions.nav

Corresponding Author: Julia D. Buckner, Department of Psychology, Louisiana State University, 236 Audubon Hall, Baton Rouge, LA 70803, USA. jbuckner@lsu.edu.

Authors' Note

The National Institute on Drug Abuse had no further role in study design; in the collection, analysis, and interpretation of data; in the writing of the manuscript; or in the decision to submit the manuscript for publication.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

with greater problems academically, as well as poorer physical health and psychosocial functioning (Buckner, Ecker, & Cohen, 2010; Caldeira et al., 2008). Despite the high rates of cannabis use and use-related problems among college students, the vast majority of students with cannabis-related impairment are not interested in treatment to help them manage their cannabis use (Buckner et al., 2010; Caldeira et al., 2008). Thus, the campus judicial process has become a point of intervention for many students who misuse alcohol and/or drugs.

Yet, little is known about the students that are referred for treatment after violating campus cannabis use policies. This is unfortunate given that students who violate campus drug and alcohol policies tend to engage in more substance use and experience more substance-related problems (Clements, 1999; Merrill, Carey, Lust, Kalichman, & Carey, 2014; O'Hare, 1997) relative to other students. Identification of cognitive vulnerability factors could inform campus-wide and personalized prevention and treatment efforts. One such vulnerability factor that may be especially relevant is distress tolerance (DT). DT is a capacity to withstand negative emotional states. Lower DT is associated with more cannabis-related problems (Buckner, Keough, & Schmidt, 2007; Bujarski, Norberg, & Copeland, 2012). Consistent with the notion that cannabis users with low DT may use cannabis to decrease distress, cannabis users with lower DT report more coping motives for cannabis use (Bujarski et al., 2012; Simons & Gaher, 2005; Zvolensky et al., 2009). Yet, such use is problematic, as coping motives are robustly related to cannabis-related impairment among college students (Buckner, 2013). In fact, in a community sample of non-treatment-seeking cannabis users, coping (but not conformity) motives mediated the association between DT and cannabis-related problems (Bujarski et al., 2012).

The current study tested whether DT would be related to cannabis use frequency, use-related problems, and motivation to change cannabis use behaviors among students referred for treatment following a violation of campus cannabis use policies. Identification of whether DT is related to baseline predictors of treatment outcome among students referred for treatment following violation of campus drug and alcohol policies is important given that low DT is related to substance use disorder treatment attrition (Daughters, Lejuez, Bornoalova, et al., 2005; Daughters, Lejuez, Kahler, Strong, & Brown, 2005) including pre-treatment attrition (MacPherson, Stipelman, Duplinsky, Brown, & Lejuez, 2008), shorter abstinence duration (Daughters, Lejuez, Kahler, et al., 2005), and greater lapse following smoking cessation (Brown, Lejuez, Kahler, & Strong, 2002; Brown et al., 2009). To test whether DT was robustly related to these cannabis-related outcomes, we also examined whether these relations remained after controlling for other substance use (smoking and alcohol use) and gender, given that these variables are related to DT and cannabis (e.g., Buckner, Keough, & Schmidt, 2007; Leyro, Bernstein, Vujanovic, McLeish, & Zvolensky, 2011; Stinson, Ruan, Pickering, & Grant, 2006). To further understand DT's relations with these cannabis vulnerability factors, we also tested whether relevant cannabis use motives (e.g., coping motives) mediated the relations of DT to these cannabis-related outcomes. As per prior work with non-treatment seeking community adults (Bujarski et al., 2012), it was hypothesized that coping motives would mediate the relationship between DT and cannabis-related problems.

Method

Participants and Procedures

Undergraduates were invited to participate in an ongoing study of the utility of brief motivational interventions for students referred by the university for violation of campus policies regarding cannabis use. Students who were observed violating the university's policies regarding cannabis use were informed of the study by staff in the university's Office of Student Advocacy and Accountability or the Office of Residential Life. Students who engaged in "predatory dealing" (i.e., selling cannabis to students other than one's friends) were not referred to the intervention study, as these students were usually expelled from the university. Inclusion criteria were (a) having received a campus disciplinary referral following a recent cannabis policy violation, (b) being at least 18 years of age, (c) being a current student at the university, and (d) endorsing lifetime cannabis use. Students who refused to participate were re-referred to their referring office to arrange an alternate treatment program. Participants were charged US\$60 total for the baseline intake and treatment appointments. They were not compensated for study participation. The study was approved by the university's institutional review board. The confidentiality of research data was assured with a Certificate of Confidentiality from the U.S. Department of Health and Human Services.

Of the 118 students who were referred for treatment, 18 refused to schedule an intake appointment, 7 dropped out after scheduling an intake appointment, 4 did not respond to attempts to contact, and 1 was ineligible due to not being a current student at the university. Thus, 88 completed the baseline appointment and were included in the current study. The racial/ethnic composition of the sample was: 80.7% Caucasian, 10.2% African American, 4.5% Asian, and 2.3% Native American, and 2.3% Hispanic/Latino/a. Participants were ethnically representative of the university during recruitment which included 76% Caucasian students. Compared with the larger university sample, participants were more likely to be male (88% vs. 49%). The majority (65.9%) lived in their own residence, with 17.0% in dorms, 11.4% with parents, and 5.7% in fraternity housing. The mean age was 19.5 ($SD = 3.1$), and 40.9% were employed.

The majority (81.8%) endorsed past-month drinking, with 44.3% endorsing at least weekly drinking and 2.3% endorsing daily drinking. Although most (65.9%) endorsed lifetime tobacco use, 25.0% endorsed current smoking. The majority (69.3%) endorsed at least monthly cannabis use, 46.6% endorsed at least weekly use, and 9.1% endorsed daily use. Offenses leading the treatment referral were as follows: cannabis possession (77.3%), cannabis paraphernalia possession (40.9%), and "other" (13.6%). Examples of other charges include being in the presence of cannabis, resisting arrest, trespassing, synthetic cannabis possession, and obstruction of justice. The majority (68.2%) were charged with one offense, 28.4% with two offenses, and 2.3% with three or more offenses.

Measures

Distress intolerance was assessed with the 15-item Distress Tolerance Scale (DTS; Simons & Gaher, 2005). Participants rated items concerning participants' perceived ability to

withstand negative psychological states from 1 (*strongly agree*) to 5 (*strongly disagree*). Thus, lower scores indicated greater intolerance of distress. The DTS has shown good psychometric properties in prior work (Simons & Gaher, 2005; Zvolensky et al., 2009) and in the present sample ($\alpha = .95$)

Typical frequency of cannabis and alcohol use was assessed per the Core Institute's Campus Assessment of Alcohol and Other Drug Norms assessed. Participants were asked to rate how often they typically use cannabis and alcohol. Response options ranged from 0 (*never*) to 4 (*2–3 times per month*) to 8 (*daily*). Participants were also asked to report whether they ever and currently smoke cigarettes.

Cannabis problems were assessed via the Marijuana Problems Scale (MPS; Stephens, Roffman, & Curtin, 2000), a 19-item list of negative consequences related to cannabis use (e.g., memory loss, financial difficulties, and legal and medical problems) in the past 90 days. Participants rated cannabis use problems on a 0 (*no problem*) to 2 (*serious problem*) scale. Endorsed items (i.e., scores of 1 or 2 on each item) were summed to create an index of total number of cannabis-related problems. The MPS has demonstrated adequate internal consistency in prior work (Buckner et al., 2010; Lozano, Stephens, & Roffman, 2006; Stephens et al., 2000; Stephens et al., 2004) and in the present sample ($\alpha = .84$).

Cannabis use motives were assessed with the *Marijuana Motives Measure* (MMM; Simons, Correia, Carey, & Borsari, 1998), a 25-item measure assessing the following cannabis use motives: enhancement (e.g., to get high), coping (e.g., to forget my worries), social (e.g., to be more sociable), conformity (e.g., to fit in with a group I like), and expansion (e.g., to expand my awareness). Participants indicated from 1 (*almost never/never*) to 5 (*almost always/always*) the degree to which they have smoked cannabis for particular reasons. MMM subscales have demonstrated good internal consistency in prior work (Chabrol, Ducongé, Casas, Roura, & Carey, 2005) and in the present sample (conformity, $\alpha = .72$; enhancement, $\alpha = .91$; social, $\alpha = .84$; coping, $\alpha = .81$; and expansion, $\alpha = .94$).

Motivation was assessed using the *Importance/Confidence Form* (ICF) adapted from Miller and Rollnick's (2002) importance/confidence rulers. The first item asked, "On a scale of 0–10, rate how important it is for you to change your marijuana use" in which 0 = *not at all important* and 10 = *most important*. The second item asked, "On a scale of 0–10, rate how confident you are that you can change your marijuana use" in which 0 = *not at all confident* and 10 = *most confident*. Similar scales correspond with changes in cannabis use (Gates, Norberg, Copeland, & Digiusto, 2012) and to increase as a result of a motivation enhancement intervention (Buckner & Schmidt, 2009).

Data Analyses

First, bivariate correlations were conducted to examine relations between study variables and to determine whether DT was related to cannabis factors (frequency of use, problems, motivation to change, use motives). Second, to test the robustness of the observed relations between DT and cannabis criterion variables, a series of hierarchical linear regression models were conducted. Separate regressions were conducted for each relevant cannabis criterion variable. Predictor variables were as follows—Step 1: gender, past-month drinking

frequency, tobacco smoking status, cannabis use frequency (for the cannabis problems model); and Step 2: DT. This strategy ensured that effect at Step 2 cannot be attributed to variance shared with variables in Step 1 (Cohen & Cohen, 1983). Third, mediational analyses were conducted using PROCESS, a macro used with SPSS 22.0 that utilizes an ordinary least squares regression-based path analytical framework to test for both direct and indirect effects (Hayes, 2013) using bootstrap analyses with 10,000 resamples from which bias-corrected 95% confidence intervals (CIs) were estimated (Hayes, 2009; Preacher & Hayes, 2004, 2008).

Results

Table 1 presents means, standard deviations, and correlations among study variables. DT was significantly correlated with frequency of typical cannabis use, number of cannabis-related problems, and coping, conformity, and expansion motives. The correlation between DT and confidence to change cannabis use was small to medium ($p = .058$). DT was unrelated to importance to change cannabis use behaviors or to social or enhancement motives.

Next, the robustness of the relations between DT and cannabis variables was tested. As evidenced in Table 2, covariates accounted for 4.6% of the variance in cannabis use frequency. After accounting for this variance, DT accounted for an additional 4.8% of the variance. Covariates accounted for 11.8% of the variance in cannabis-related problems. After accounting for this variance, DT accounted for an additional 10.9% of the variance.

Cannabis-Related Problems

Coping, Conformity, and Expansion motives were the only motives correlated with cannabis problems and DT and were thus evaluated as mediators. Smoking and drinking variables, cannabis frequency, and gender were included as covariates.¹ For the relation between DT and cannabis-related problems, the total effects model accounted for significant variance ($R^2 = .230$, $df = 5, 82$, $F = 4.90$, $p < .001$) and the full model with coping motives accounted for significant variance ($R^2 = .539$, $df = 6, 81$, $F = 5.52$, $p = .0001$). The direct effect of DT and cannabis-related problems remained significant after controlling for coping motives ($B = -0.09$, $SE = 0.03$, $p = .001$). The indirect effect was estimated and revealed that DT was predictive of more cannabis problems indirectly through greater coping motivated use ($B = -0.02$, $SE = 0.012$, 95% CI = $[-.052, -.004]$).

For the analyses concerning conformity motives, the total effects model accounted for significant variance ($R^2 = .480$, $df = 5, 82$, $F = 4.90$, $p < .001$) and the full model with conformity motives accounted for significant variance ($R^2 = .244$, $df = 6, 81$, $F = 4.36$, $p < .001$). The direct effect of DT and cannabis-related problems remained significant after controlling for conformity motives ($B = -0.08$, $SE = 0.03$, $p = .009$). The indirect effect was estimated and revealed that DT was not indirectly related to cannabis problems through greater conformity motivated use ($B = -0.01$, $SE = 0.01$, 95% CI = $[-.041, .007]$).

¹A similar pattern was obtained when analyses were conducted without these covariates.

For the analyses concerning expansion motives, the total effects model accounted for significant variance ($R^2 = .230$, $df = 5$, 82 , $F = 4.90$, $p < .001$) and the full model with expansion motives accounted for significant variance ($R^2 = .233$, $df = 6$, 81 , $F = 4.11$, $p = .001$). The direct effect of DT and cannabis-related problems remained significant after controlling for enhancement motives ($B = -0.0$, $SE = 0.03$, $p = .002$). The indirect effect was estimated and revealed that DT was not indirectly related to cannabis problems through greater enhancement motivated use ($B = -0.002$, $SE = 0.006$, $95\% \text{ CI} = [-.023, .004]$).

Cannabis Use Frequency

Coping motives were the only motives significantly correlated with cannabis use frequency and were thus tested as a putative mediator of the relations between DT and cannabis use frequency. Smoking and drinking variables and gender were included as covariates see Note 1. For the relation between DT and cannabis use frequency, the total effects model accounted for significant variance ($R^2 = .112$, $df = 4$, 83 , $F = 2.62$, $p = .040$) and the full model with coping motives was marginally significant ($R^2 = .117$, $df = 5$, 82 , $F = 2.18$, $p = .065$). The direct effect of DT and cannabis frequency was no longer significant after controlling for coping motives ($B = -0.04$, $SE = 0.02$, $p = .115$). The indirect effect was estimated and revealed that DT was not predictive of more frequent cannabis use indirectly through greater coping motivated use ($B = -0.005$, $SE = 0.008$, $95\% \text{ CI} = [-.024, .008]$).

Discussion

The current study is the first known study to identify personality factors related to cannabis use and use-related problems among students referred by the university for treatment after violating campus cannabis use policies. Consistent with prior work with volunteer undergraduate students who endorsed lifetime cannabis use (Simons & Gaher, 2005) and young adult community current cannabis users (Zvolensky et al., 2009), DT among these students was negatively related to coping motives and unrelated to enhancement motives. Also consistent with young adult cannabis users, DT was negatively correlated with conformity motives (Zvolensky et al., 2009). Consistent with prior work with volunteer undergraduates, DT was negatively related to cannabis-related problems (Buckner, Keough, & Schmidt, 2007). Contrary to studies with young adult cannabis users and volunteer undergraduates (Buckner, Keough, & Schmidt, 2007; Zvolensky et al., 2009), low DT among students referred for treatment for violating campus cannabis use policies was related to more frequent cannabis use.

The current findings further extend the extant literature in several key ways. First, the current study extends prior work (Simons & Gaher, 2005; Zvolensky et al., 2009) by determining that DT is significantly related to cannabis problems indirectly through coping motives (but not conformity or expansion motives). Thus, students referred for treatment after violating campus cannabis policies may benefit from cognitive-behavioral skills (see Steinberg et al., 2002) to help them manage negative affectivity using more adaptive emotion regulation skills. Notably, DT was no longer related to cannabis use frequency after controlling for coping motives, although DT was not indirectly related to frequency via coping motives. These data suggest that the relation of DT to cannabis use frequency is not robust, which is

consistent with prior work (Buckner, Keough, & Schmidt, 2007; Potter, Vujanovic, Marshall-Berenz, Bernstein, & Bonn-Miller, 2011; Zvolensky et al., 2009). Taken together, this pattern of findings is consistent with prior work suggesting that individuals with higher levels of emotional reactivity (e.g., those with elevated social anxiety; Buckner, Bonn-Miller, Zvolensky, & Schmidt, 2007; Buckner et al., 2010; Buckner & Schmidt, 2008; Ecker, Richter, & Buckner, 2014) may not be using cannabis more frequently than other individuals, but something about the way in which they are using is placing them at risk for experiencing more problems related to their use.

Unexpectedly, DT was unrelated to importance to change cannabis use and the relation of DT to confidence to change cannabis was of a small-to-medium effect, suggesting that DT's relations to these two components of motivation were not strong. This is somewhat inconsistent with data from other substance use treatment samples in which lower DT was associated with proxy measures of motivation such as greater perceived barriers to smoking cessation (Kraemer, McLeish, Jeffries, Avallone, & Luberto, 2013) and substance use disorder treatment attrition (Daughters, Lejuez, Kahler, et al., 2005; MacPherson et al., 2008). However, a unique feature of our sample is that the patients were referred for treatment by the university following a violation of campus cannabis use policies. Thus, our sample may have been more extrinsically motivated to change their cannabis use than prior samples which may have impacted DT's relation to motivation. This line of research could benefit from testing whether motivation to change cannabis use among these students is related to more objective measures of DT (i.e., paced auditory serial addition task; Lejuez, Kahler, & Brown, 2003), which have been associated with proxy measures of motivation in prior work (i.e., treatment attrition; Daughters, Lejuez, Bornovalova, et al., 2005) and which have not been found to correlate with self-report measures of DT (McHugh et al., 2011).

The present study should be considered in light of limitations that can inform future work in this area. First, the sample was comprised solely of students referred for treatment following a violation of campus cannabis use policies and future work comparing these students with cannabis using students who have not been caught violating campus policies or other cannabis using populations will be an important next step. Second, a large proportion of students referred for treatment following a violation of campus policies did not present for intake. Given data finding low DT to be related to greater pre-treatment attrition (MacPherson et al., 2008), future work could benefit from testing whether pre-treatment drop-outs differ on DT from students who do comply with treatment recommendations. Third, data were self-report and future work could benefit from multi-method (e.g., biological verification of cannabis use, behavioral measures of DT, prospective designs such as ecological momentary assessment) and multi-informant (e.g., collateral reports of cannabis use and problems) approaches. Fourth, the sample was predominantly male. Although this probably reflects that cannabis use is greater among college men than women (Johnston et al., 2014), future work could benefit from inclusion of more women to examine whether results generalize to women or whether the relations of DT to cannabis vary as a function of gender.

In sum, the current study identified a cognitive vulnerability factor related to more frequent cannabis use and cannabis-related problems among students referred by the university for

treatment following violations of campus cannabis use policies. DT was robustly associated with more frequent cannabis use and cannabis-related problems. Thus, future work testing whether targeting DT directly during treatment improves outcomes for these high-risk students will be an important next step.

Acknowledgments

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: Funding for this study was provided in part by Grant 1R34DA031937-01A1 from the National Institute on Drug Abuse.

References

- American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 4. Washington, DC: Author; 1994.
- Brown RA, Lejuez CW, Kahler CW, Strong DR. Distress tolerance and duration of past smoking cessation attempts. *Journal of Abnormal Psychology*. 2002; 111:180–185. DOI: 10.1037/0021-843X.111.1.180 [PubMed: 11866171]
- Brown RA, Lejuez CW, Strong DR, Kahler CW, Zvolensky MJ, Carpenter LL, ... Price LH. A prospective examination of distress tolerance and early smoking lapse in adult self-quitters. *Nicotine & Tobacco Research: Official Journal of the Society for Research on Nicotine and Tobacco*. 2009; 11:493–502. DOI: 10.1093/ntr/ntp041 [PubMed: 19372572]
- Buckner JD. College cannabis use: The unique roles of social norms, motives, and expectancies. *Journal of Studies on Alcohol and Drugs*. 2013; 74:720–726. [PubMed: 23948531]
- Buckner JD, Bonn-Miller MO, Zvolensky MJ, Schmidt NB. Marijuana use motives and social anxiety among marijuana-using young adults. *Addictive Behaviors*. 2007; 32:2238–2252. DOI: 10.1016/j.addbeh.2007.04.004 [PubMed: 17478056]
- Buckner JD, Ecker AH, Cohen AS. Mental health problems and interest in marijuana treatment among marijuana-using college students. *Addictive Behaviors*. 2010; 35:826–833. DOI: 10.1016/j.addbeh.2010.04.001 [PubMed: 20483200]
- Buckner JD, Keough ME, Schmidt NB. Problematic alcohol and cannabis use among young adults: The roles of depression and discomfort and distress tolerance. *Addictive Behaviors*. 2007; 32:1957–1963. DOI: 10.1016/j.add-beh.2006.12.019 [PubMed: 17258398]
- Buckner JD, Schmidt NB. Marijuana effect expectancies: Relations to social anxiety and marijuana use problems. *Addictive Behaviors*. 2008; 33:1477–1483. DOI: 10.1016/j.addbeh.2008.06.017 [PubMed: 18694625]
- Buckner JD, Schmidt NB. A randomized pilot study of motivation enhancement therapy to increase utilization of cognitive-behavioral therapy for social anxiety. *Behaviour Research and Therapy*. 2009; 47:710–715. DOI: 10.1016/j.brat.2009.04.009 [PubMed: 19467647]
- Bujarski SJ, Norberg MM, Copeland J. The association between distress tolerance and cannabis use-related problems: The mediating and moderating roles of coping motives and gender. *Addictive Behaviors*. 2012; 37:1181–1184. DOI: 10.1016/j.addbeh.2012.05.014 [PubMed: 22698896]
- Caldeira KM, Arria AM, O'Grady KE, Vincent KB, Wish ED. The occurrence of cannabis use disorders and other cannabis-related problems among first-year college students. *Addictive Behaviors*. 2008; 33:397–411. DOI: 10.1016/j.addbeh.2007.10.001 [PubMed: 18031940]
- Chabrol H, Ducongé E, Casas C, Roura C, Carey KB. Relations between cannabis use and dependence, motives for cannabis use and anxious, depressive and borderline symptomatology. *Addictive Behaviors*. 2005; 30:829–840. DOI: 10.1016/j.addbeh.2004.08.027 [PubMed: 15833585]
- Clements R. Prevalence of alcohol-use disorders and alcohol-related problems in a college student sample. *Journal of American College Health*. 1999; 48:111–118. DOI: 10.1080/07448489909595682 [PubMed: 10584445]

- Cohen, J.; Cohen, P. Applied multiple regression/correlation analysis for the behavioral sciences. Hillsdale, NJ: Lawrence Erlbaum; 1983.
- Daughters SB, Lejuez CW, Bornovalova MA, Kahler CW, Strong DR, Brown RA. Distress tolerance as a predictor of early treatment dropout in a residential substance abuse treatment facility. *Journal of Abnormal Psychology*. 2005; 114:729–734. DOI: 10.1037/0021-843x.114.4.729 [PubMed: 16351393]
- Daughters SB, Lejuez CW, Kahler CW, Strong DR, Brown RA. Psychological distress tolerance and duration of most recent abstinence attempt among residential treatment-seeking substance abusers. *Psychology of Addictive Behaviors*. 2005; 19:208–211. [PubMed: 16011392]
- Ecker AH, Richter AA, Buckner JD. Short communication: Cannabis-related impairment: The impacts of social anxiety and misconceptions of friends' cannabis-related problems. *Addictive Behaviors*. 2014; 39:1746–1749. DOI: 10.1016/j.addbeh.2014.07.004 [PubMed: 25119421]
- Gates PJ, Norberg MM, Copeland J, Digiusto E. Randomized controlled trial of a novel cannabis use intervention delivered by telephone. *Addiction*. 2012; 107:2149–2158. DOI: 10.1111/j.1360-0443.2012.03953.x [PubMed: 22632139]
- Hayes AF. Beyond Baron and Kenny: Statistical mediation analysis in the new millennium. *Communication Monographs*. 2009; 76:408–420. DOI: 10.1080/03637750903310360
- Hayes, AF. Introduction to mediation, moderation, and conditional process analysis: A regression-based approach. New York, NY: Guilford Press; 2013.
- Johnston, LD.; O'Malley, PM.; Bachman, JG.; Schulenberg, JE.; Miech, RA. Monitoring the future: National survey results on drug use 1975–2013: Volume 2, college students & adults ages 19–55. Ann Arbor: Institute for Social Research, The University of Michigan; 2014.
- Kraemer KM, McLeish AC, Jeffries ER, Avallone KM, Luberto CM. Distress tolerance and perceived barriers to smoking cessation. *Substance Abuse*. 2013; 34:277–282. DOI: 10.1080/08897077.2013.771597 [PubMed: 23844959]
- Lejuez CW, Kahler CW, Brown RA. A modified computer version of the Paced Auditory Serial Addition Task (PASAT) as a laboratory-based stressor. *The Behavior Therapist*. 2003; 26:290–293.
- Leyro TM, Bernstein A, Vujanovic AA, McLeish AC, Zvolensky MJ. Distress Tolerance Scale: A confirmatory factor analysis among daily cigarette smokers. *Journal of Psychopathology and Behavioral Assessment*. 2011; 33:47–57. DOI: 10.1007/s10862-010-9197-2 [PubMed: 23935238]
- Lozano BE, Stephens RS, Roffman RA. Abstinence and moderate use goals in the treatment of marijuana dependence. *Addiction*. 2006; 101:1589–1597. DOI: 10.1111/j.1360-0443.2006.01609.x [PubMed: 17034438]
- MacPherson L, Stipelman BA, Duplinsky M, Brown RA, Lejuez CW. Distress tolerance and pre-smoking treatment attrition: Examination of moderating relationships. *Addictive Behaviors*. 2008; 33:1385–1393. DOI: 10.1016/j.addbeh.2008.07.001 [PubMed: 18706768]
- McHugh R, Daughters S, Lejuez CW, Murray H, Hearon B, Gorka S, Otto M. Shared variance among self-report and behavioral measures of distress intolerance. *Cognitive Therapy and Research*. 2011; 35:266–275. DOI: 10.1007/s10608-010-9295-1 [PubMed: 23894216]
- Miller, WR.; Rollnick, S. Motivational interviewing: Preparing people for change. 2. New York, NY: Guilford Press; 2002.
- Merrill JE, Carey KB, Lust SA, Kalichman SC, Carey MP. Do students mandated to intervention for campus alcohol-related violations drink more than nonmandated students? *Psychology of Addictive Behaviors*. 2014; 28:1265–1270. [PubMed: 25180559]
- O'Hare TM. Measuring excessive alcohol use in college drinking contexts: The drinking context scale. *Addictive Behaviors*. 1997; 22:469–477. DOI: 10.1016/s0306-4603(96)00050-0 [PubMed: 9290857]
- Potter CM, Vujanovic AA, Marshall-Berenz EC, Bernstein A, Bonn-Miller MO. Posttraumatic stress and marijuana use coping motives: The mediating role of distress tolerance. *Journal of Anxiety Disorders*. 2011; 25:437–443. DOI: 10.1016/j.janxdis.2010.11.007 [PubMed: 21146357]
- Preacher KJ, Hayes AF. SPSS and SAS procedures for estimating indirect effects in simple mediation models. *Behavior Research Methods, Instruments, & Computers*. 2004; 36:717–731. DOI: 10.3758/BF03206553

- Preacher KJ, Hayes AF. Asymptotic and resampling strategies for assessing and comparing indirect effects in multiple mediator models. *Behavior Research Methods*. 2008; 40:879–891. DOI: 10.3758/BRM.40.3.879 [PubMed: 18697684]
- Simons JS, Correia CJ, Carey KB, Borsari BE. Validating a five-factor marijuana motives measure: Relations with use, problems, and alcohol motives. *Journal of Counseling Psychology*. 1998; 45:265–273. DOI: 10.1037/0022-0167.45.3.265
- Simons JS, Gaher RM. The Distress Tolerance Scale: Development and validation of a self-report measure. *Motivation and Emotion*. 2005; 29:83–102. DOI: 10.1007/s11031-005-7955-3
- Steinberg KL, Roffman RA, Carroll KM, Kabela E, Kadden R, Miller M, Duresky D. Tailoring cannabis dependence treatment for a diverse population. *Addiction*. 2002; 97:135–142. DOI: 10.1046/j.1360-0443.97.s01.5.x [PubMed: 12460135]
- Stephens RS, Roffman RA, Curtin L. Comparison of extended versus brief treatments for marijuana use. *Journal of Consulting and Clinical Psychology*. 2000; 68:898–908. DOI: 10.1037/0022-006X.68.5.898 [PubMed: 11068976]
- Stephens RS, Roffman RA, Fearer SA, Williams C, Picciano JF, Burke RS. The Marijuana Check-up: Reaching users who are ambivalent about change. *Addiction*. 2004; 99:1323–1332. DOI: 10.1111/j.1360-0443.2004.00832.x [PubMed: 15369571]
- Stinson FS, Ruan WJ, Pickering R, Grant BF. Cannabis use disorders in the USA: Prevalence, correlates and co-morbidity. *Psychological Medicine*. 2006; 36:1447–1460. DOI: 10.1017/S0033291706008361 [PubMed: 16854249]
- Zvolensky MJ, Marshall EC, Johnson K, Hogan J, Bernstein A, Bonn-Miller MO. Relations between anxiety sensitivity, distress tolerance, and fear reactivity to bodily sensations to coping and conformity marijuana use motives among young adult marijuana users. *Experimental and Clinical Psychopharmacology*. 2009; 17:31–42. DOI: 10.1037/a0014961 [PubMed: 19186932]

Biographies

Julia D. Buckner, PhD, is an Associate Professor in the Department of Psychology at Louisiana State University and director of the Anxiety and Addictive Behaviors Laboratory and Clinic. Her research focuses on factors related to the etiology, maintenance, and treatment of anxiety and substance use disorders.

Emily R. Jeffries, BA, is a graduate student in the Clinical Psychology Training Program at Louisiana State University. Her research focuses on transdiagnostic cognitive vulnerability factors related to anxiety and substance use disorders.

Meredith A. Terlecki, PhD, is a Senior Lecturer in the School of Psychology at the University of East London, United Kingdom. Her research is focused broadly on the identification and prevention of alcohol and substance use disorders among high-risk young adults.

Anthony H. Ecker, MA, is a graduate student in the Clinical Psychology Training Program at Louisiana State University. He is currently on predoctoral internship at the West Haven VA Healthcare System. His research interests include anxiety disorders and the comorbidity of anxiety disorders and substance use, and substance-related problems.

Table 1

Means, Standard Deviations, and Correlations Among Study Variables.

	1	2	3	4	5	6	7	8	9	10	11	12
Distress tolerance												
Cannabis frequency	-.24**											
Cannabis problems	-.42***	.26**										
Importance	.01	-.06	.30***									
Confidence	.20*	.05	-.20*	.25**								
Social motives	-.02	.30***	.01	-.13	.10							
Coping motives	-.36***	.22**	.42***	-.05	-.10	.39***						
Enhancement motives	-.14	.36***	.21*	-.07	.11	.57***	.41***					
Conformity motives	-.31***	-.07	.22**	.09	-.20*	.17	.26**	.02				
Expansion motives	-.24**	.34***	.22**	-.18*	-.14	.40***	.53***	.43***	.28***			
Alcohol frequency	-.18	.19*	.24**	.20*	.21*	.19*	.28***	.40***	.08	.24**		
Smoking status	.15	.08	-.10	-.08	.12	.12	.05	-.15	.27**	.20	.02	
<i>M (SD)</i>	55.0 (13.5)	4.0 (2.7)	3.7 (3.5)	5.2 (3.6)	8.7 (2.7)	10.5 (4.4)	8.7 (3.7)	15.8 (6.3)	6.1 (2.0)	9.8 (5.4)	2.0 (1.3)	25.0%

Note. Importance refers to importance to change cannabis use. Confidence refers to confidence to change cannabis use.

* $p < .10$.

** $p < .05$.

*** $p < .01$.

Table 2

Hierarchical Linear Models Testing the Robustness of Distress Tolerance in the Prediction of Cannabis-Related Variables.

	<i>B</i>	<i>SE</i>	β	<i>t</i>	<i>p</i>	<i>R</i> ²
DV: Cannabis use frequency.						
Step 1						
Gender	−0.083	0.851	−.010	−0.097	.923	.046
Smoker status	0.557	0.647	.093	0.861	.392	
Drinking frequency	0.357	0.213	.181	1.677	.097	
Step 2						
Distress tolerance	−0.045	0.021	−.227	−2.10	.038	.048
DV: Cannabis-related problems						
Step 1						
Gender	1.224	1.097	.115	1.115	.268	.118
Smoker status	−0.306	0.838	−.038	−0.365	.716	
Drinking frequency	0.520	0.279	.197	1.860	.066	
Cannabis use frequency	0.303	0.141	.228	2.155	.034	
Step 2						
Distress tolerance	−0.092	0.027	−.350	−3.405	.001	.109

Note. Separate models were analyzed for each variable in Step 2. DTS = Distress Tolerance Scale (Simons & Gaher, 2005), DV = dependent variable.