Published in final edited form as:

Neuropharmacology. 2017 February ; 113(Pt A): 502–510. doi:10.1016/j.neuropharm.2016.11.001.

Dysregulated corticostriatal activity in open-field behavior and the head-twitch response induced by the hallucinogen 2,5-dimethoxy-4-iodoamphetamine

Claudia Rangel-Barajas^{a,1}, Ana María Estrada-Sánchez^{a,b,1}, Scott J. Barton^a, Robert R. Luedtke^c, and George V. Rebec^{a,*}

^aIndiana University Bloomington, Department of Psychological and Brain Sciences, Program in Neuroscience, 1101 E. 10th St. Bloomington, IN 47405

^bIntellectual and Developmental Disabilities Research Center, Semel Institute, Department of Psychiatry & Biobehavioral Sciences, David Geffen School of Medicine, University of California Los Angeles, USA

^cUniversity of North Texas Health Science Center, the Department of Pharmacology and Neuroscience, 3500 Camp Bowie Boulevard, Fort Worth, Texas 76107

Abstract

The substituted amphetamine, 2,5-dimethoxy-4-iodoamphetamine (DOI), is a hallucinogen that has been used to model a variety of psychiatric conditions. Here, we studied the effect of DOI on neural activity recorded simultaneously in the primary motor cortex (M1) and dorsal striatum of freely behaving FvB/N mice. DOI significantly decreased the firing rate of individually isolated neurons in M1 and dorsal striatum relative to pre-drug baseline. It also induced a bursting pattern of activity by increasing both the number of spikes within a burst and burst duration. In addition, DOI increased coincident firing between simultaneously recorded neuron pairs within the striatum and between M1 and dorsal striatum. Local field potential (LFP) activity also increased in coherence between M1 and dorsal striatum after DOI in the low frequency gamma band (30–50 Hz), while corticostriatal coherence in delta, theta, alpha, and beta activity decreased. We also assessed corticostriatal LFP activity in relation to the DOI-induced head-twitch response (HTR), a readily identifiable behavior used to assess potential treatments for the conditions it models. The HTR was associated with increased delta and decreased theta power in both M1 and dorsal striatum. Together, our results suggest that DOI dysregulates corticostriatal communication and that the HTR is associated with this dysregulation.

*Corresponding Author: Department of Psychological and Brain Sciences, Program in Neuroscience, Indiana University, 1101 E. 10th St. Bloomington, IN 47405-7007. (812) 855-4832, rebec@indiana.edu (G.V. Rebec).

¹Contributed equally to this work.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

twitch response; DOI; 5-HT_{2A} serotonin receptor; *in vivo* electrophysiology

1. INTRODUCTION

Cortical pyramidal neurons (CPNs) are a major source of glutamate-mediated excitatory drive on medium-sized spiny neurons (MSNs), which account for >90% of the neuronal population in dorsal striatum (Gerfen and Bolam, 2010; Deng et al, 2015). Corticostriatal connectivity plays a key role in cognition, learning, behavioral control and movement (Pennartz et al, 2009; Santos et al., 2014; Shepherd, 2013). This connectivity is monosynaptic such that glutamatergic CPNs send projections to striatum, but striatum communicates only indirectly with cortex through subcortical networks (Harris and Shepherd, 2015). Disruptions in this connectivity have been linked to multiple neurological and neuropsychiatric conditions including obsessive-compulsive disorder, Tourette's syndrome and schizophrenia (Estrada-Sánchez et al, 2015; Shepherd *et al*, 2013). Interestingly, the substituted amphetamine, 2,5-dimethoxy-4-iodoamphetamine (DOI), a hallucinogen that binds primarily to serotonin 5HT_{2A} receptors, has been used as a tool for studying these conditions. Although DOI does not mimic the entire symptom profile in each case, the drug induces a rapid, lateral movement of the head, known as the head-twitch response (HTR), which has been shown to have predictive validity for assessing potential pharmacological treatments (Canal and Morgan, 2012). For example, we reported that highly selective D₃ dopamine receptor ligands as well as antipsychotic drugs, which are known to block D₂-like dopamine receptors, attenuate the DOI-induced HTR (Rangel-Barajas et al, 2014; 2015). However, despite ample information about DOI-induced changes in consciousness and sensory perception leading to hallucinations (Nichols, 2004), behavior-related changes in corticostriatal activity produced by DOI are largely unknown.

We assessed three major aspects of DOI-induced behavioral modulation of corticostriatal processing. First, we analyzed the effects of DOI in freely behaving animals on the overall firing rate and pattern of individually isolated spike trains, which allowed us to assess the extent of correlated activity between simultaneously recorded neuron pairs, a process known to be altered by neurodegenerative disease and drugs of abuse (Ball et al, 2010; Miller et al., 2008; Walker et al, 2008). We also assessed changes in burst firing as a measure of information transmission and neuroplasticity (Izhikevich et al, 2003). Second, we simultaneously recorded local field potentials (LFPs), which, by capturing the activity of thousands of neurons in the vicinity of the recording electrode, integrate synaptic processes that cannot be measured by observing spike activity alone (Einevoll et al., 2013). Thus, LFPs provide information about the aggregated activity of relatively localized neuronal populations, including synaptic potentials and other low frequency events (Buzsaki et al, 2012). We assessed the overall LFP phase relationship between M1 and striatum to determine the effects of DOI administration on corticostriatal connectivity. Third, to examine behavior-related modulation of corticostriatal connectivity, we also assessed corticostriatal LFP oscillations in relation to the DOI-induced HTR itself. Collectively, our

results indicate that DOI modulates multiple aspects of corticostriatal processing to alter behavioral output.

2. MATERIALS AND METHODS

2.1 Animals

Data were obtained from male FvB/N mice supplied by Jackson Laboratories (Bar Harbor, Maine). All mice ranged in age from 20–39 weeks; they were kept under controlled conditions of humidity and temperature, with water and food available *ad libitum* and with a 12 hr day/night cycle. All experiments were conducted during the light phase of the diurnal cycle. All animal procedures were carried out in accord with the conditions set forth by the Institutional Animal Care and Use Committee of Indiana University based on guidelines established by the National Institutes of Health.

2.2 Behavioral Analysis

We assessed quiet rest (QR) periods before and after DOI treatment. QR was defined as the lack of overt motor activity. The HTR analysis, performed as previously described (Rangel-Barajas et al, 2014; 2015), was defined as a rapid left to right (or right to left) movement of the head, without involvement of the front paws. Any HTR that occurred immediately before or after grooming behavior was not considered in the analysis. Two observers counted the number of head-twitches by visual examination. The number of head-twitches was recorded for 5-min intervals; data are presented as the mean values obtained by two observers.

2.3 Surgical procedure

Some animals were anesthetized with a mixture of chloral hydrate (170 mg/kg) and sodium pentobarbital (40 mg/kg) in an intraperitoneal (i.p.) injection volume of 0.4 ml/100 mg. Mice were placed in a stereotaxic frame, two micro-electrode bundles, each consisting of four insulated stainless steel wires (25 µm diameter) and one uninsulated stainless steel ground wire (50 µm diameter), were unilaterally placed. One bundle was positioned in M1 (coordinates: +0.5 mm anterior, ±1.4 lateral from bregma and –0.5mm from dura) and one in dorsal striatum (+0.8mm anterior, ±2.2 lateral from bregma and –3.2mm from dura) according to Paxinos and Franklin (2001). The recording apparatus was secured with dental acrylic and two small stainless steel screws. All animals were allowed a week for recovery before the recording sessions began.

2.4 Electrophysiological recording

All electrophysiological recordings and data collection were performed as described previously (Estrada-Sánchez et al, 2015). Briefly, a lightweight wire harness equipped with field-effect transistors for unity gain was attached to the skull-mounted electrode assembly. The harness was attached to a swiveling commutator that permitted freedom of movement. We simultaneously recorded LFPs and extracellular spikes in M1 and dorsal striatum in mice behaving spontaneously in an open-field arena (26 × 18 cm) located in a soundproof and electrically shielded chamber. LFP activity and spike trains were analyzed in conjunction with the Multichannel Acquisition Processor system (Plexon, Dallas, TX, USA).

Electrophysiological signals were recorded for analysis for 30 min under basal conditions (untreated) followed by 30 min after DOI (5 mg/kg, ip).

Spikes were first identified by oscilloscope at a voltage threshold 2.5 times background noise. Approximately 800 waveform samples were sorted and collected to define templates based on principal component analysis. Spike-2 software (Cambridge Electronic Design Cambridge, England) was used to confirm the accuracy of signal isolation. Classification of presumed CPNs and MSNs followed previously established criteria (McCormick et al., 1985; Wilson et al, 1993). Presumed interneurons, identified as fast spiking units (10 Hz) with short-duration (<1 ms) waveforms, were not considered for analysis.

Videotapes of behavior were analyzed both before (basal) and after DOI treatment. To assess behavior-related changes in electrophysiological activity, we focused on LFP activity during periods of QR and DOI-induced HTR. The time at which both behaviors occurred were identified from the videotape and analyzed in conjunction with LFP activity.

2.5 Drug

2,5-dimethoxy-4-iodoamphetaminehydrochloride (DOI) was purchased from Sigma Aldrich Research Biochemicals Inc. A dose of 5 mg/kg was selected based on the intensity and frequency of DOI-induced HTR as described in our previous work (Rangel-Barajas et al, 2014; 2015). Control animals received vehicle injections (ip).

2.6 Histology

Upon completion of recording, mice were deeply anesthetized and a current (30 μ A) was applied through each micro-wire to mark the recording location. Animals were transcardially perfused and brains were removed and preserved as previously reported (Miller et al, 2008). Coronal sections (50 μ m) were analyzed for electrode placement confirmation.

2.7 Data Analysis

Single-unit spike and LFP activity were analyzed with NeuroExplorer 4.13 (Nex Technologies Littleton, MA, USA) in conjunction with written MATLAB scripts (Matworks, Natick, MA, USA). Firing rate was calculated as spikes/s. We also measured the mean inter-spike interval (ISI) and the coefficient of variation (CV) of the ISIs (CV ISIs) as previously reported (Miller et al., 2008, 2011). Burst firing consists of a sudden onset of a cluster of action potentials occurring at a fast rate (Homayoun et al., 2005). We analyzed bursting activity using the Poisson surprise method (Legendy and Salcman, 1985). This method defines a burst as a group of spikes in which several successive ISIs do not follow a Poisson distribution (Boraud et al., 2002; Miller et al., 2008). The surprise value is defined as the negative natural logarithm of the probability that a series of spikes in a given time interval is significantly different from that expected from a Poisson process with the same mean firing rate, providing an estimate of the statistical significance of each burst in the spike train. In accord with previous studies, we used a surprise value of five, which estimates that a burst occurs approximately 150 times more often ($p < 0.007$) than would be expected with the same mean firing rate (Homayoun et al., 2005; Miller et al., 2008). Therefore, the surprise value is

an index of how intense or “surprising” the ISI of a particular burst is in relation to other ISIs in the same spike train. This method can detect not only burst rate but several parameters of bursting including: the percentage of spikes in a burst, the mean number of spikes within a burst, mean burst duration, mean burst surprise, and mean inter-burst interval. The surprise method is well established for detecting bursts in presumed striatal MSNs (Stanford et al., 2007; Miller et al., 2008).

To assess correlated activity between two spike-trains, cross-correlation histograms were constructed for each pairwise comparison (Miller et al, 2008). Synchronous firing between two spike-trains was identified by 99% confidence intervals. Statistically significant peaks indicate correlated activity. Cross-correlation histograms were performed within and between cortical and striatal spike-trains.

LFP power spectral densities were analyzed in relation to instances of specific behavioral events (QR and HTR). LFPs were assessed by frequency decomposition using Fourier transforms as previously described (Hong et al, 2012). Start and stop times for each behavioral episode were marked and corresponding LFPs were analyzed. We constructed mean power spectral density (PSD) histograms and peri-event spectrograms for QR and HTR averaged across repeated instances of the same behavior and the results were imported to MATLAB for further analysis. The area under the curve (AUC) from the normalized PSD was calculated for individual frequency bands. We focused on delta (1–4 Hz), theta (4–7 Hz), alpha (7–13 Hz), beta (13–30 Hz), and low gamma (30–50 Hz) oscillations.

Further analysis was performed on the cross-power spectra from M1 and dorsal striatum to obtain coherence and relative phase values across frequency bands. Coherence values ranged from 0 to 1; a value of 1 represents perfect synchrony between M1 and dorsal striatum. Relative phase values ranged from -180° to 180° , where 0° relative phase indicates an in-phase corticostriatal relationship and $\pm 180^\circ$ indicates an anti-phase pattern. Since M1 was used as a reference signal, a positive phase value indicates that M1 leads, while negative value indicates that striatum leads.

2.8 Statistical Analysis

Statistical analysis was performed with Prism 6.0a GraphPad Software (San Diego, CA, USA). Significance for all statistical comparisons was set at $p = 0.05$. The nonparametric Friedman test followed by Dunn’s test was used to analyze the HTR over time. Firing rate and burst analysis (burst rate, mean spikes in burst, mean inter-burst interval, percentage of spikes in a burst, mean burst duration and mean burst surprise) and coherent LFP frequency oscillations were analyzed with the matched-paired Wilcoxon test. Cross-correlation histograms were analyzed by Chi-square in conjunction with Fisher’s test to calculate the exact p value. The AUC of the power spectral density histograms was calculated using the trapezoidal method and frequency oscillations were analyzed by the Kruskal-Wallis test followed by Dunn’s multiple comparisons test.

3. RESULTS

We first confirmed that 5 mg/kg DOI could elicit a robust dataset of HTRs in FvB/N mice because the frequency of the DOI-induced HTR varies with mouse strain (Canal et al., 2010; Canal and Morgan, 2012; Rangel-Barajas et al., 2014). We then recorded electrophysiological signals in M1 and dorsal striatum in additional animals to determine: (1) how this dose alters the basal firing patterns of presumed CPN and MSN neurons as mice behave spontaneously; and (2) how M1 and striatal processing, as measured by LFP activity, changes during spontaneous behavior and during the HTR.

3.1 The DOI-induced HTR in FvB/N mice

Administration of DOI (5 mg/kg ip) in FvB/N mice (n=8) induced 75.8 ± 1.30 HTRs over a 30-min recording period (Figure 1A). In line with our data from other mouse strains (Rangel-Barajas et al, 2014; 2015), the DOI-induced HTR in FvB/N mice peaked between 5–10 min after injection and decreased over time (Figure 1B). Furthermore, the number of head-twitches was equivalent to the number reported previously for Swiss albino mice (Darmani et al, 1990). Thus, our data establish that 5 mg/kg DOI is sufficient to elicit a robust HTR dataset in the FvB/N strain

3.2 Firing patterns of presumed CPN and MSN units in spontaneously behaving mice before and after DOI

We recorded 63 individually isolated units in M1 and 50 in dorsal striatum in a separate group of 16 FvB/N mice (4 with vehicle and 12 with DOI) that behaved spontaneously in an open field. Both M1 and striatal neurons showed spike waveforms (Figure 2A, H) and pre-drug firing rates of 5.3 ± 0.7 and 6.4 ± 1.3 spikes/s consistent with presumed CPN and MSN activity, respectively (Miller et al., 2008; Walker et al, 2008).

For the 30-min post-injection period, DOI significantly decreased overall firing in both M1 ($p<0.001$) and striatum ($p<0.0001$) relative to pre-drug activity (Figure 2B, I). Although no changes were found in the CV ISI in either location (Figure 2C, J), the mean ISI was significantly increased in M1 ($p<0.05$) and striatum ($p<0.01$) (Figure 2D, K). The lack of change in the CV ISI suggests that DOI induced an overall decrease in activity rather than a change in spike-train variability.

Further analysis showed that DOI also disrupted the pattern of spike activity. There was a significant increase in the percentage of spikes within a burst in both M1 ($p<0.05$) and striatum ($p<0.05$) relative to the pre-drug baseline (Figure 2E, L). The mean burst duration was also increased after DOI administration in both M1 ($p<0.0001$) and striatum ($p<0.001$) (Figure 2F, M), while the mean burst surprise significantly increased in M1 ($p<0.01$) but not in striatum (Figure 2G, N). Other burst parameters (burst rate, mean number of spikes within a burst, and mean inter-burst interval) remained unchanged by DOI (data not shown).

To assess the extent of correlated activity between simultaneously recorded neuron pairs, we constructed cross-correlation histograms for all recorded units. Representative histograms are shown in Figure 3 (**right panel**); peaks at or above the 99% confidence interval (broken lines) indicate synchronous firing. After DOI treatment the number of correlated neurons

increased (Figure 3 **left panel**) in striatum (from 10 to 27 neurons; $\chi^2=13$ df=1, $p<0.001$) and between M1 and the striatum (from 6 to 19 neurons; $\chi^2=8.2$ df=1, $p<0.01$). A trend in the same direction was found in M1 (from 9 to 18 neurons; $\chi^2=4$ df=1, $p=0.07$).

Vehicle had no effect on firing rate or any of the bursting patterns examined (data not shown).

3.3 Effect of DOI on LFP activity

We analyzed LFP activity to assess behaviorally relevant synaptic processes that cannot be determined by spike data alone. We first examined DOI-induced changes in LFP coherence between M1 and striatum throughout the period of spontaneous behavior (Figure 4). DOI significantly decreased the coherence value of delta, theta, alpha, and beta oscillations ($p<0.001$), while low gamma oscillations were significantly increased ($p<0.001$) (Figure 4A **and inset**). In addition, coherence phase was significantly decreased by DOI for delta ($p<0.05$), alpha, and beta ($p<0.001$), but increased ($p<0.001$) for low gamma oscillations (Figure 4B, C). The negative phase value with M1 as the reference signal indicates that striatum leads M1 in the DOI-induced disruption of LFP activity.

When we focused specifically on HTRs, we found a marked increase in low frequency power in dorsal striatum and a similar though less dramatic trend in M1. This change was most evident during the first 10 min after DOI injection when the number of DOI-induced HTRs is at its peak. Representative LFP spectrograms from M1 and striatum for the 30-min periods before (Figure 5A, C) and after DOI treatment (Figure 5B, D) show that major changes in power occur in striatum (Figure 5D) in parallel with the HTR time course.

To assess this relationship in more detail, we evaluated LFP activity during 10 randomly selected QR and HTR episodes from each animal before and after DOI. Sample LFP traces from one recording are shown for episodes of QR before and after DOI and for DOI-induced HTR (Figure 5E); boxes indicate the 2 s window around the time when the behavioral event occurred. Mean PSD histograms were constructed and the AUC was calculated for delta, theta, alpha, and beta oscillations in each brain region. There was a significant increase in delta activity during QR episodes that occurred after DOI administration in both M1 and dorsal striatum ($p<0.05$ and $p<0.001$, respectively). This effect was significantly enhanced during the HTR ($p<0.05$ and $p<0.01$ respectively) (Figure 5J). In contrast, QR-related theta activity was significantly decreased by DOI treatment in M1 and dorsal striatum ($p<0.01$) without further changes during the HTR (Figure 5K). No changes were observed in alpha or beta oscillations (Figure 5L, M).

Because dorsal striatal neurons exhibit entrainment to high-voltage spindle oscillations (Berke et al., 2004), which mainly occur during periods of immobility or activation of vibrissae and disappear immediately after initiation of body movements (Shaw and Liao, 2005), we searched for signs of this paroxysmal activity (mu rhythm between 5–12 Hz), but found none. In fact, our analysis of the mu rhythm did not show significant differences during QR either in the basal condition or after DOI treatment (data not shown). These oscillations, however, may be more likely to occur when there is no motivation to act or move, such as when the environment has already been explored (Berke et al., 2004). Other

factors, such as age and strain of animal, also may affect the occurrence of high-voltage spindles (Buzsaki et al, 1988).

Peri-event spectrograms constructed from LFPs revealed no change in M1 power during QR (Figure 5F) or DOI-induced HTR (Figure 5G). In dorsal striatum, however, an increase in power occurred during QR (Figure 5H), and dysregulated delta oscillations were found during the DOI-induced HTR (Figure 5I). Thus, the DOI-induced disruption of LFP activity is more prominent in dorsal striatum than M1 during both QR and the HTR.

4. DISCUSSION

Three interesting observations emerge from our results: first, DOI dysregulates spike activity in both M1 and striatum by increasing burst firing and the correlated firing between simultaneously recorded neuron pairs; second, DOI disrupts LFP activity between M1 and striatum; and third, delta oscillations increase during the DOI-induced HTR in both areas but consistently high delta power is correlated with the HTR in striatum.

4.1 Dysregulated spike activity

The effects of DOI on neuronal activity in forebrain areas related to behavioral control and known to be involved in psychiatric conditions have been assessed *in vitro* and in anesthetized preparations. *In vitro* studies indicate that DOI increases excitatory postsynaptic currents in layer V CPNs of rat prefrontal cortex (Aghajanian and Marek, 1999). In contrast, studies of anesthetized rats found that DOI inhibited neuronal firing in orbital cortex and striatum (Rueter et al, 2000), whereas in prefrontal cortex both excitatory and inhibitory effects were recorded (Celada et al, 2008). Our studies extend this research to behaving animals by assessing the effects of DOI in the corticostriatal motor circuit.

Interestingly, DOI decreased the overall firing rate in both M1 and striatum by increasing the mean ISI without changing variability in the pattern of spikes. Thus, the decrease in firing appeared to be relatively uniform, uncomplicated by increased variability in the timing of individual spikes. Changes in the CV ISI usually reflect a complex spike-train pattern that often indicates increased bursting (Miller et al., 2011). DOI, however, did not change the CV ISI but increased the mean ISI, indicating that the decrease in firing rate was likely due to a consistently longer interval between spikes. A lack of effect of DOI on CV ISI is consistent with a corresponding lack of change in the rate of bursting. Importantly, however, other parameters of burst activity (e.g., the percentage of spikes in a burst and the mean burst duration) were increased by DOI in both M1 and the striatum, while the mean burst surprise was altered only in M1. Because the Poisson surprise algorithm allows for differentiation between irregular firing, single spikes, and bursting patterns within the same spike-train (Jackson et al, 2004), the mean burst surprise effect in M1 might reflect changes in bursting not captured in burst rate. For instance, even though DOI decreased spike rate, individual spikes occurred more frequently in a burst, which was denoted by increases in the number of spikes within a burst and the mean burst duration in both M1 (Figure 2E, F) and striatum (Figure 2L, M). In contrast, only M1 showed a change in mean burst surprise, suggesting that DOI rearranged the burst pattern to include unexpected or “surprising” burst clusters. Changes in cortical firing and bursting activity may alter downstream connections.

Accordingly, when we assessed the activity of simultaneously recorded pairs of neurons, we found that DOI significantly increased correlated firing in striatum and produced a similar trend in M1. Correlated firing between M1 and striatum was also increased, indicating that DOI alters neuronal communication between these two brain regions. Interestingly, we previously found increased correlated neuronal activity in response to another hallucinogen, the ($\pm$)3,4-methylenedioxymethamphetamine (MDMA) (Ball et al, 2010). Corticostriatal connectivity plays an important role in the initiation of conscious movements through the basal ganglia motor loop, which effectively propagates information downstream (Kimpo et al, 2003). Thus, the increase in correlated activity may represent an altered form of information transmission induced by hallucinogenic psychoactive drugs.

4.2 Disrupted communication between M1 and striatum

Synchronized corticostriatal LFP activity has been reported in relation to cognitive tasks (DeCoteau et al, 2007) and motor processing (von Nicolai et al, 2014). DOI, in contrast, has the opposite effect as indicated by disrupted corticostriatal LFP coherence. The decrease in synchrony and phase coherence of delta, theta, alpha and beta oscillations but increase in the low gamma range (Figure 4) is consistent with a report of DOI-induced decreased EEG oscillations in frontal cortex and striatum of rats with the exception of the alpha band (Dimpfel et al, 1989). Delta oscillations also have been reported to decrease in response to DOI in frontal cortex (Celada et al, 2008). In addition, DOI has been reported to increase high gamma oscillations (130–180 Hz) to a greater extent than low gamma oscillations (40–60 Hz) in the nucleus accumbens of freely moving rats (Goda et al, 2013). Interestingly, an increase in low gamma power in striatum also has been found in response to movement initiation (Masimore et al, 2005), while high gamma oscillations predominated during running (von Nicolai et al, 2014). The dynamic interplay of low and high gamma oscillations may play a role in motor states. Accordingly, it has been suggested that gamma oscillations help to organize ensembles of synchronously firing neurons (Laurent, 2002). Thus, the increased coherence of low gamma oscillations induced by DOI could be linked to the increased correlated spike activity we found in M1 and striatum.

4.3 The HTR and low frequency LFP oscillations

Rhythmic oscillations in LFP activity play critical roles in various aspects of brain processing, including sensory binding, cognition, as well as movement preparation and initiation (Buzsaki et al., 2012). In the corticostriatal system, beta, theta, and gamma are dominant in sensorimotor processing, but their exact function is difficult to define and likely depends on environmental context and behavioral state (Fries, 2015). Our analysis of the time course of LFP activity indicated a peak increase in power during the first 5 min after DOI administration, the same time as the peak number of head twitches, suggesting that the HTR might be related to changes in LFP power. The LFP changes, moreover, predominate at delta, alpha, and beta frequencies, consistent with evidence of low frequency oscillations during movement (Masimore et al, 2005). In fact, DOI not only increased delta oscillations in both M1 and dorsal striatum during QR, but further increased delta activity during the HTR. In addition, our LFP coherence analysis suggests that striatum leads M1 (Figure 4A). Effective unidirectional connectivity from cortex to striatum has been reported for sleep states and anesthesia (David *et al*, 2008), while effective connectivity has been shown to

become bidirectional in animals engaged in voluntary movement (Nakhnikian et al, 2014). Interestingly, in the delta band M1 led striatum during basal conditions, whereas after DOI administration striatum led M1 (Figure 4C), suggesting that DOI affects the direction of corticostriatal communication.

Delta oscillations have been linked to the activation of T-type calcium channels (Ernst et al, 2009), which are involved in mediating calcium influx into cells after an action potential (Perez-Reyes et al, 2003). Delta activity can be eliminated by genetic deletion of the most conserved pore-subunit (CaV3.1) of this channel (Kim et al, 2001). Overexpression of this subunit in mice results in absence epilepsy, characterized by intermittent delta activity (Ernst et al, 2009; Gelisse et al., 2011). In a further link to the HTR-related increase in delta activity, T-type calcium channels have been shown to participate in motor function (Yang et al, 2014) and blocking these channels improves locomotor symptoms in a Parkinson's disease model (Tai et al, 2011).

Apart from a motor component, the HTR might underlie perceptual alterations, delusions, or hallucinations (Nichols, 2004). Interestingly, delta wave disruptions also have been associated with altered mental states in schizophrenia (Venables et al, 2009), and the HTR is blocked by antipsychotics (Rangel-Barajas et al, 2014), including those that act as T-type calcium channel antagonists (Enyeart et al., 1992; Santi et al, 2002; Choi and Rhim, 2010).

Collectively, these data suggest that elevated delta activity in corticostriatal circuitry is an important component of the DOI-induced HTR. DOI could modulate cellular calcium equilibrium mediated by receptor-linked intracellular signaling pathways or through modulation of T-type calcium channels, increasing delta oscillations observed during DOI treatment and the HTR. Further studies are needed to identify the cellular mechanism by which DOI changes LFP activity. Our behavioral assessments, moreover, did not include potential cognitive and emotional changes induced by DOI, which could also impact the contribution of specific LFP rhythms.

4.4 DOI Pharmacology and Receptors

DOI is a partial agonist of the 5-HT₂ receptor family, which includes 5-HT_{2A}, 5-HT_{2B} and 5-HT_{2C} receptors. This family is positively coupled to phospholipase C (PLC), which increases accumulation of inositol phosphates and therefore increases intracellular calcium (Barnes and Sharp, 1999). Most atypical antipsychotics, hallucinogens, and some antidepressants, bind to these receptors. Although DOI exhibits the highest affinity for 5-HT_{2A} and 5-HT_{2C} receptors (Rangel-Barajas et al, 2014), ample evidence suggests that 5-HT_{2A} receptors are required for the DOI-induced HTR. For example, M100907, a 5-HT_{2A} receptor antagonist, blocked the DOI-induced HTR in mice (Kehne et al, 1996), and mice lacking 5-HT_{2A} receptors do not exhibit the HTR, while selective expression of 5-HT_{2A} receptors in cortical neurons restored this behavior (Gonzalez-Maeso et al, 2007). Electrophysiological studies have reported that the changes in high frequency gamma oscillations were fully reversed by a 5-HT_{2A} receptor antagonist and were not affected by a 5-HT_{2C} receptor antagonist, suggesting that 5-HT_{2A} receptors are the main site of action of DOI (Goda et al, 2013). In addition, the DOI-induced inhibition of neuronal firing in orbito-

frontal cortex and striatum is not impaired in 5-HT_{2C} receptor knockout mice (Rueter et al, 2000), further implicating 5-HT_{2A} receptors in expression of the HTR.

The effect of DOI in corticostriatal connectivity can be explained by the localization of 5-HT_{2A} receptors. It has been shown that 5-HT_{2A} receptors are expressed in M1 CPNs (Weber and Andrade, 2010), and these receptors can modulate the activity of the corticostriatal intratelencephalic tract (Shepherd, 2013), which preferentially targets MSNs. This connection is part of the direct pathway of the basal ganglia (Reiner *et al*, 2010), which is involved in the initiation of movement.

5-HT_{2A} receptors show enriched expression in dorsal striatum and play an important role in sensorimotor-related connections (Ward and Dorsa, 1996; Lopez-Gimenez et al., 2002; Crittenden and Graybiel, 2011). In fact, alterations in 5-HT_{2A} receptors have been linked to motor deficits. In Parkinsonian patients, for example, the 5-HT_{2A} antagonist, ritanserin, improved akinesia and gait (Henderson et al, 1992), while quetiapine, another 5-HT_{2A} antagonist, inhibited levodopa-induced motor complications in monkeys (Oh et al, 2002) and improved extrapyramidal movement disorders in schizophrenic patients (Vesely et al, 2000). Interestingly, although DOI binds to 5-HT_{2A} but not dopamine D2-like receptors, D2 selective antagonists significantly decreased the DOI-induced HTR (Rangel-Barajas et al, 2014), suggesting a functional interaction between D2 and 5-HT_{2A} receptors. It is likely, therefore, that there is an interaction between dopamine and serotonin receptors, probably by intracellular signaling mechanisms or by receptor heterodimer formation.

5. CONCLUSIONS

DOI disrupted the pattern of corticostriatal neuronal firing by decreasing firing rate, altering bursting activity, and increasing corticostriatal correlated firing. Corticostriatal LFP activity was also disrupted by DOI, while the HTR was accompanied by an increase in delta oscillations. Dysregulated corticostriatal communication, perhaps involving dopamine-serotonin receptor interactions, may disrupt downstream information flow that could contribute to the DOI-induced HTR.

Acknowledgments

This work was supported by National Institute of Health (P50 DA05312 and R01 DA 012964). The authors would like to thank Dr. Maninder Malik for her valuable assistance in the collection of behavioral data.

ABBREVIATIONS

CPN	Cortical Pyramidal Neuron
CV	Coefficient of Variation
DOI	2,5-dimethoxy-4-iodoamphetamine
HTR	Head Twitch Response
ISI	Inter-spike Interval

LFP	Local Field Potential
MSN	Medium-sized Spiny Neuron
QR	Quiet Rest

REFERENCES

- Aghajanian GK, Marek GJ. Serotonin, via 5-HT_{2A} receptors, increases EPSCs in layer V pyramidal cells of prefrontal cortex by an asynchronous mode of glutamate release. *Brain Res.* 1999; 825:161–171. [PubMed: 10216183]
- Ball KT, Wellman CL, Miller BR, Rebec GV. Electrophysiological and structural alterations in striatum associated with behavioral sensitization to (+/-)-3,4-methylenedioxymethamphetamine (ecstasy) in rats: Role of drug context. *Neuroscience.* 2010; 171:794–811. [PubMed: 20875842]
- Barnes NM, Sharp T. A review of central 5-HT receptors and their function. *Neuropharmacology.* 1999; 38:1083–1152. [PubMed: 10462127]
- Berke JD, Okatan M, Skurski J, Eichenbaum HB. Oscillatory entrainment of striatal neurons in freely moving rats. *Neuron.* 2004; 43:883–896. [PubMed: 15363398]
- Boraud T, Bezard E, Bioulac B, Gross CE. From single extracellular unit recording in experimental and human Parkinsonism to the development of a functional concept of the role played by the basal ganglia in motor control. *Progress in Neurobiology.* 2002; 66:265–283. [PubMed: 11960681]
- Buzsaki G, Anastassiou CA, Koch C. The origin of extracellular fields and currents--EEG, ECoG, LFP and spikes. *Nat Rev Neurosci.* 2012; 13:407–420. [PubMed: 22595786]
- Buzsaki G, Bickford RG, Armstrong DM, Ponomareff G, Chen KS, Ruiz R, Thal LJ, Gage FH. Electric activity in the neocortex of freely moving young and aged rats. *Neuroscience.* 1988; 26:735–744. [PubMed: 3200427]
- Canal CE, Morgan D. Head-twitch response in rodents induced by the hallucinogen 2,5-dimethoxy-4-iodoamphetamine: A comprehensive history, a re-evaluation of mechanisms, and its utility as a model. *Drug Test Anal.* 2012; 4:556–576. [PubMed: 22517680]
- Canal CE, Olaghere da Silva UB, Gresch PJ, Watt EE, Sanders-Bush E, Airey DC. The serotonin 2C receptor potentially modulates the head-twitch response in mice induced by a phenethylamine hallucinogen. *Psychopharmacology (Berl).* 2010; 209:163–174. [PubMed: 20165943]
- Celada P, Puig MV, Diaz-Mataix L, Artigas F. The hallucinogen DOI reduces low-frequency oscillations in rat prefrontal cortex: Reversal by antipsychotic drugs. *Biol Psychiatry.* 2008; 64:392–400. [PubMed: 18436196]
- Choi KH, Rhim H. Inhibition of recombinant $\text{Ca}_v3.1$ ($\alpha 1G$) T-type calcium channels by the antipsychotic drug clozapine. *Eur J Pharmacol.* 2010; 626:123–130. [PubMed: 19782679]
- Crittenden JR, Graybiel AM. Basal ganglia disorders associated with imbalances in the striatal striosome and matrix compartments. *Front Neuroanat.* 2011; 5:59. [PubMed: 21941467]
- Darmani NA, Martin BR, Glennon RA. Withdrawal from chronic treatment with (+/-)-DOI causes super-sensitivity to 5-HT₂ receptor-induced head-twitch behaviour in mice. *Eur J Pharmacol.* 1990; 186:115–118. [PubMed: 2282932]
- David O, Guillemain I, SAILLET S, Rey S, Deransart C, Segebarth C, Depaulis A. Identifying neural drivers with functional MRI: An electrophysiological validation. *PLoS Biol.* 2008; 6:2683–2697. [PubMed: 19108604]
- DeCoteau WE, Thorn C, Gibson DJ, Courtemanche R, Mitra P, Kubota Y, Graybiel AM. Oscillations of local field potentials in the rat dorsal striatum during spontaneous and instructed behaviors. *J Neurophysiol.* 2007; 97:3800–3805. [PubMed: 17329629]
- Deng Y, Lanciego J, Goff LK, Coulon P, Salin P, Kachidian P, Lei W, Del Mar N, Reiner A. Differential organization of cortical inputs to striatal projection neurons of the matrix compartment in rats. *Front Syst Neurosci.* 2015; 9:51. [PubMed: 25926776]
- Dimpfel W, Spuler M, Nichols DE. Hallucinogenic and stimulatory amphetamine derivatives: Fingerprinting DOM, DOI, DOB, MDMA, and MBDB by spectral analysis of brain field

- potentials in the freely moving rat (tele-stereo-EEG). *Psychopharmacology (Berl)*. 1989; 98:297–303. [PubMed: 2568652]
- Einevoll GT, Kayser C, Logothetis NK, Panzeri S. Modelling and analysis of local field potentials for studying the function of cortical circuits. *Nat Rev Neurosci*. 2013; 14:770–785. [PubMed: 24135696]
- Enyeart JJ, Biagi BA, Mlinar B. Preferential block of T-type calcium channels by neuroleptics in neural crest-derived rat and human C cell lines. *Mol Pharmacol*. 1992; 42:364–372. [PubMed: 1355264]
- Ernst WL, Zhang Y, Yoo JW, Ernst SJ, Noebels JL. Genetic enhancement of thalamocortical network activity by elevating alpha 1g-mediated low-voltage-activated calcium current induces pure absence epilepsy. *J Neurosci*. 2009; 29:1615–1625. [PubMed: 19211869]
- Estrada-Sanchez AM, Burroughs CL, Cavaliere S, Barton SJ, Chen S, Yang XW, Rebec GV. Cortical efferents lacking mutant huntingtin improve striatal neuronal activity and behavior in a conditional mouse model of huntington's disease. *J Neurosci*. 2015; 35:4440–4451. [PubMed: 25762686]
- Fries P. Rhythms for cognition: communication through coherence. *Neuron*. 2015; 88:220–235. [PubMed: 26447583]
- Gelisse P, Serafini A, Velizarova R, Genton P, Crespel A. Temporal intermittent delta activity: A marker of juvenile absence epilepsy? *Seizure*. 2011; 20:38–41. [PubMed: 21044851]
- Gerfen, CR.; Bolam, JP. The Neuroanatomical organization of the Basal Ganglia. In: Steiner, H.; Tseng, KY., editors. *Handbook of Basal Ganglia Structure and Function*. San Diego; 2010. p. 3-28.
- Goda SA, Piasecka J, Olszewski M, Kasicki S, Hunt MJ. Serotonergic hallucinogens differentially modify gamma and high frequency oscillations in the rat nucleus accumbens. *Psychopharmacology (Berl)*. 2013; 228:271–282. [PubMed: 23525524]
- Gonzalez-Maeso J, Weisstaub NV, Zhou M, et al. Hallucinogens recruit specific cortical 5-HT(2A) receptor-mediated signaling pathways to affect behavior. *Neuron*. 2007; 53:439–452. [PubMed: 17270739]
- Harris KD, Shepherd GM. The neocortical circuit: Themes and variations. *Nat Neurosci*. 2015; 18:170–181. [PubMed: 25622573]
- Henderson J, Yiannikas C, Graham JS. Effect of ritanserin, a highly selective 5-HT2 receptor antagonist, on parkinson's disease. *Clin Exp Neurol*. 1992; 29:277–282. [PubMed: 1343870]
- Homayoun H, Jackson ME, Moghaddam B. Activation of metabotropic glutamate 2/3 receptors reverses the effects of NMDA receptor hypofunction on prefrontal cortex unit activity in awake rats. *J Neurophysiol*. 2005; 93:1989–2001. [PubMed: 15590730]
- Hong SL, Cossyleon D, Hussain WA, Walker LJ, Barton SJ, Rebec GV. Dysfunctional behavioral modulation of corticostriatal communication in the R6/2 mouse model of huntington's disease. *PLoS One*. 2012; 7:e47026. [PubMed: 23056565]
- Izhikevich EM, Desai NS, Walcott EC, Hoppensteadt FC. Bursts as a unit of neural information: Selective communication via resonance. *Trends Neurosci*. 2003; 26:161–167. [PubMed: 12591219]
- Jackson ME, Homayoun H, Moghaddam B. NMDA receptor hypofunction produces concomitant firing rate potentiation and burst activity reduction in the prefrontal cortex. *Proc Natl Acad Sci U S A*. 2004; 101:8467–8472. [PubMed: 15159546]
- Kehne JH, Baron BM, Carr AA, et al. Preclinical characterization of the potential of the putative atypical antipsychotic MDL 100,907 as a potent 5-HT2A antagonist with a favorable CNS safety profile. *J Pharmacol Exp Ther*. 1996; 277:968–981. [PubMed: 8627580]
- Kim D, Song I, Keum S, Lee T, Jeong MJ, Kim SS, McEnery MW, Shin HS. Lack of the burst firing of thalamocortical relay neurons and resistance to absence seizures in mice lacking alpha(1G) T-type Ca^{2+} channels. *Neuron*. 2001; 31:35–45. [PubMed: 11498049]
- Kimpo RR, Theunissen FE, Doupe AJ. Propagation of correlated activity through multiple stages of a neural circuit. *J Neurosci*. 2003; 23:5750–5761. [PubMed: 12843279]
- Laurent G. Olfactory network dynamics and the coding of multidimensional signals. *Nat Rev Neurosci*. 2002; 3:884–895. [PubMed: 12415296]
- Legendary CR, Salzman M. Bursts and recurrences of bursts in the spike trains of spontaneously active striate cortex neurons. *J Neurophysiol*. 1985; 53:926–939. [PubMed: 3998798]

- Lopez-Gimenez JF, Tecott LH, Palacios JM, Mengod G, Vilaro MT. Serotonin 5-HT_{2C} receptor knockout mice: Autoradiographic analysis of multiple serotonin receptors. *J Neurosci Res.* 2002; 67:69–85. [PubMed: 11754082]
- McCormick DA, Connors BW, Lighthall JW, Prince DA. Comparative electrophysiology of pyramidal and sparsely spiny stellate neurons of the neocortex. *J Neurophysiol.* 1985; 54:782–806. [PubMed: 2999347]
- Masimore B, Schmitzer-Torbert NC, Kakalios J, Redish AD. Transient striatal gamma local field potentials signal movement initiation in rats. *Neuroreport.* 2005; 16:2021–2024. [PubMed: 16317346]
- Miller BR, Walker AG, Barton SJ, Rebec GV. Dysregulated neuronal activity patterns implicate corticostriatal circuit dysfunction in multiple rodent models of huntington's disease. *Front Syst Neurosci.* 2011; 5:26. [PubMed: 21629717]
- Miller BR, Walker AG, Shah AS, Barton SJ, Rebec GV. Dysregulated information processing by medium spiny neurons in striatum of freely behaving mouse models of huntington's disease. *J Neurophysiol.* 2008; 100:2205–2216. [PubMed: 18667541]
- Nakhnikian A, Rebec GV, Grasse LM, Dwiel LL, Shimono M, Beggs JM. Behavior modulates effective connectivity between cortex and striatum. *PLoS One.* 2014; 9:e89443. [PubMed: 24618981]
- Nichols DE. Hallucinogens. *Pharmacol Ther.* 2004; 101:131–181. [PubMed: 14761703]
- Pennartz CM, Berke JD, Graybiel AM, Ito R, Lansink CS, van der Meer M, Redish AD, Smith KS, Voorn P. Corticostriatal interactions during learning, memory processing, and decision making. *J Neurosci.* 2009; 29:12831–12838. [PubMed: 19828796]
- Perez-Reyes E, Cribbs LL, Daud A, Lacerda AE, Barclay J, Williamson MP, Fox M, Rees M, Lee JH. Molecular characterization of a neuronal low-voltage-activated T-type calcium channel. *Nature.* 1998; 391:896–900. [PubMed: 9495342]
- Rangel-Barajas C, Malik M, Mach RH, Luedtke RR. Pharmacological modulation of abnormal involuntary DOI-induced head twitch response movements in male DBA/2J mice: II. effects of D3 dopamine receptor selective compounds. *Neuropharmacology.* 2015; 93:179–190. [PubMed: 25698528]
- Rangel-Barajas C, Malik M, Vangveravong S, Mach RH, Luedtke RR. Pharmacological modulation of abnormal involuntary DOI-induced head twitch response in male DBA/2J mice: I. effects of D2/D3 and D2 dopamine receptor selective compounds. *Neuropharmacology.* 2014; 83C:18–27.
- Reiner A, Hart NM, Lei W, Deng Y. Corticostriatal projection neurons - dichotomous types and dichotomous functions. *Front Neuroanat.* 2010; 4:142. [PubMed: 21088706]
- Rueter LE, Tecott LH, Blier P. In vivo electrophysiological examination of 5-HT_{2C} responses in 5-HT_{2C} receptor mutant mice. *Naunyn Schmiedeberg's Arch Pharmacol.* 2000; 361:484–491. [PubMed: 10832601]
- Santi CM, Cayabyab FS, Sutton KG, McRory JE, Mezeyova J, Hamming KS, Parker D, Stea A, Snutch TP. Differential inhibition of T-type calcium channels by neuroleptics. *J Neurosci.* 2002; 22:396–403. [PubMed: 11784784]
- Santos L, Opris I, Hampson R, Godwin DW, Gerhardt G, Deadwyler S. Functional dynamics of primate cortico-striatal networks during volitional movements. *Front Syst Neurosci.* 2014; 8:27. [PubMed: 24653682]
- Shaw FZ, Liao YF. Relation between activities of the cortex and vibrissae muscles during high-voltage rhythmic spike discharges in rats. *J Neurophysiol.* 2005; 93:2435–2448. [PubMed: 15625092]
- Shepherd GM. Corticostriatal connectivity and its role in disease. *Nat Rev Neurosci.* 2013; 14:278–291. [PubMed: 23511908]
- Stanford JA, Gerhardt GA. Age-related changes in striatal function of freely moving F344 rats. *Neurobiol Aging.* 2001; 22:659–669. [PubMed: 11445266]
- Tai CH, Yang YC, Pan MK, Huang CS, Kuo CC. Modulation of subthalamic T-type Ca²⁺ channels remedies locomotor deficits in a rat model of parkinson disease. *J Clin Invest.* 2011; 121:3289–3305. [PubMed: 21737877]
- Venables NC, Bernat EM, Sponheim SR. Genetic and disorder-specific aspects of resting state EEG abnormalities in schizophrenia. *Schizophr Bull.* 2009; 35:826–839. [PubMed: 18381357]

- Vesely C, Kufferle B, Brucke T, Kasper S. Remission of severe tardive dyskinesia in a schizophrenic patient treated with the atypical antipsychotic substance quetiapine. *Int Clin Psychopharmacol*. 2000; 15:57–60. [PubMed: 10836289]
- von Nicolai C, Engler G, Sharott A, Engel AK, Moll CK, Siegel M. Corticostriatal coordination through coherent phase-amplitude coupling. *J Neurosci*. 2014; 34:5938–5948. [PubMed: 24760853]
- Walker AG, Miller BR, Fritsch JN, Barton SJ, Rebec GV. Altered information processing in the prefrontal cortex of huntington's disease mouse models. *J Neurosci*. 2008; 28:8973–8982. [PubMed: 18768691]
- Ward RP, Dorsa DM. Colocalization of serotonin receptor subtypes 5-HT_{2A}, 5-HT_{2C}, and 5-HT₆ with neuropeptides in rat striatum. *J Comp Neurol*. 1996; 370:405–414. [PubMed: 8799865]
- Weber ET, Andrade R. Htr2a gene and 5-HT(2A) receptor expression in the cerebral cortex studied using genetically modified mice. *Front Neurosci*. 2010; 4 10.3389/fnins.2010.00036. eCollection 2010.
- Wilson CJ. The generation of natural firing patterns in neostriatal neurons. *Prog Brain Res*. 1993; 99:277–297. [PubMed: 8108553]
- Yang YC, Tai CH, Pan MK, Kuo CC. The T-type calcium channel as a new therapeutic target for Parkinson's disease. *Pflugers Arch*. 2014; 466:747–755. [PubMed: 24531801]

HIGHLIGHTS

- DOI dysregulates corticostriatal firing and bursting activity.
- DOI increases corticostriatal correlated firing.
- DOI decreases theta oscillations during quiet rest and head-twitch response.
- Delta oscillations increase during the DOI-induced head-twitch response.

Figure 1. DOI-induced HTR in FvB/N mice

A: Mean ($\pm$ SEM) number of head twitches in FvB/N mice ($n = 8$) recorded for 30 min after DOI administration. **B:** Time course of the DOI-induced HTR. The number of head twitches decreased progressively from the peak recorded at 5 min post injection. The nonparametric Friedman test followed by Dunn's multiple comparisons test indicated significant differences from the 5-min peak (* $p < 0.05$; ** $p < 0.01$).

Figure 2. Multiple measures of neuronal activity before and after DOI administration in M1 and dorsal striatum

A and H Putative cortical pyramidal and medium-sized spiny neuron waveforms recorded from M1 and dorsal striatum. These waveforms are representative of all M1 ($n = 51$) and striatal ($n = 42$) neurons used for data analysis. Scale indicates $25 \mu\text{V}$ (vertical) and $300 \mu\text{s}$ (horizontal). **B and I:** Firing rate. DOI significantly decreases firing rate in both M1 and dorsal striatum (*** $p < 0.001$ and **** $p < 0.0001$, respectively). **C and J:** CV ISI. No significant effect induced by DOI on CV ISI was noted in either M1 or striatum ($p = 0.7$ and $p = 0.9$, respectively). **D and K:** Mean ISI. DOI significantly increased the mean ISI in both M1 and dorsal striatum (* $p < 0.05$ and ** $p < 0.01$ respectively). **E and L:** Percentage of spikes that occur in bursts. There was a significant effect of DOI in both M1 and dorsal striatum (* $p < 0.05$). **F and M:** Mean burst duration. DOI significantly increased the mean burst duration in both M1 and dorsal striatum (**** $p < 0.0001$ and ** $p < 0.01$ respectively). **G and N:** Mean burst surprise. DOI significantly increased mean burst surprise in M1 (** $p < 0.01$) but not dorsal striatum ($p = 0.1$). For these and all subsequent Tukey box-and-whisker plots, the line within the box is the median score, the edges of the box represent the 25th and 75th quartile, and the whiskers indicate standard deviation above and below the mean. Statistical data are based on the matched-pair Wilcoxon test.

M1 **Basal** **DOI** **Striatum** **M1-Striatum**

Figure 3. Coincident firing in M1 and dorsal striatum before and after DOI administration

Right: Representative cross-correlation histograms derived from pairs of simultaneously recorded neurons. Broken lines indicate 99% confidence intervals in M1 (top), dorsal striatum (middle), and between M1 and dorsal striatum (bottom) used to identify a significant peak. **Left:** Distribution of correlated pairs assessed by Chi-square (χ^2); Fisher's test was used to calculate an exact p value. Significant differences before and after DOI were found in dorsal striatum ($\chi^2=13$, df=1, ***p<0.001; middle) and between M1 and dorsal striatum ($\chi^2=8.2$ df=1, **p<0.01; bottom), but only a trend in M1 ($\chi^2=4$ df=1 p=0.07; top).

Figure 4. Coherence of LFP activity between M1 and striatum before and after DOI administration

A: Mean coherence value of LFP activity before (black) and after (gray) DOI treatment.

Inset: Tukey box-and-whisker plots of the changes induced by DOI at each frequency band analyzed. **B:** Mean relative phase. **C:** Coherence phase for each frequency band. Statistical differences were analyzed by the matched-pair Wilcoxon test (* $p < 0.05$ and *** $p < 0.001$).

was further enhanced during the HTR (* $p < 0.05$ and ** $p < 0.01$), while theta power was decreased after DOI treatment without a further change during the HTR in either M1 or striatum (* $p < 0.05$). No significant differences were found in the alpha and beta bands.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript