another coexistent disease, she needed urgent relief of the obstruction. Chronic progressive liver disease and even metastatic cancer are not emergencies; obstruction of the common bile duct is.

The general practitioner faced two communication challenges. Firstly, what to tell the patient? Respondents agreed they would tell her she has a liver problem that may be serious and she needs more tests. The second challenge was for the general practitioner to be the patient's advocate and negotiate prompt assessment in secondary care.

Mrs Prior's general practitioner worked the patient into his schedule, identified her problems, ordered key diagnostic tests, followed her course, recognised the

urgent need for intervention, and helped the patient and her family understand the process. What more can you hope for? Only that the consultant system meets this patient's needs.

Competing interests: None declared.

- 1 Heathcote J. Abnormal liver function found after an unplanned consultation: case outcome. *BMJ* 2004;329:500.
- 2 Orzano AJ, Gregory PM, Nutting PA, Werner JJ, Flocke SA, Stange KC. Care of the secondary patient in family practice. A report from the Ambulatory Sentinel Practice Network. *J Fam Pract* 2001;50:113-6.
- 3 Heathcote J. Abnormal liver function found after an unplanned consultation: case progression. *BMJ* 2004;329:342.
- 4 Beckingham IJ, Ryder SD. ABC of diseases of liver, pancreas, and biliary system: investigation of liver and biliary disease. *BMJ* 2001;322:33-6.
- 5 Beckingham IJ. ABC of diseases of liver, pancreas, and biliary system: gallstone disease. *BMJ* 2001;322:91-4.

Commentary: The gap between doctors' and patients' perceptions

Kieran Walsh

BMJ Learning,
BMA House,
London WC1H 9JR
Kieran Walsh
editorial registrar
kmwalsh@
bmjgroup.com

During her husband's consultation Mrs Prior mentioned three symptoms of her own: itching, cystitis, and a rash.¹ She mentioned them as a "by the way," and her general practitioner saw her and did some blood tests. Would you be so patient? Or would you ask her to come back? There is no right answer to this question but, reading between the lines, perhaps the primary purpose of the consultation was to discuss her problems. Respondents on *bmj.com* thought that this was common and that the reasons for it included not wanting to have to make a separate appointment or to wait and, in some countries, not wanting to have to pay. Maybe she thought her problems were trivial or maybe she was afraid that they were not and wanted to downplay them to herself and her husband and her doctor.

Whatever her reasons, the literature suggests that doctors could be better at finding out why patients come to see them. One study showed that in routine consultations 45% of patient's concerns are not elicited.² In half of visits the doctor and the patient do not agree on the nature of the presenting problem.² And doctors often assume that patients' first complaint is their chief one, but in fact the order in which patients present their problems is often unrelated to their clinical importance.³ These are all strong arguments for longer appointment times in primary care.

Readers were also asked what they would tell the patient about the results of her tests. The responses varied: most people said they would tell her that she had abnormal liver function and that she needed further investigations. Some said that they would tell her that she had septicaemia or a bug or something nasty. It is striking that nobody discussed how to tell her this news, how much information to give her, or how much she would be able to take in at this stressful time. Respondents were expansive about the differential diagnosis but clammed up when talking to the patient. Maybe it's not surprising then that, while most patients say they want more information than they receive, doctors vastly overestimate the time they spend on explanation and planning for the future.^{4 5}

Competing interests: None declared.

- 1 Heathcote J. Abnormal liver function found after an unplanned consultation case outcome. *BMJ* 2004;329:500.
- 2 Kurtz S, Silverman J, Draper J. *Teaching and learning communication skills in medicine*. Abingdon: Radcliffe Medical Press, 1998.
- 3 Beckman HB, Frankel RM. The effect of physician behavior on the collection of data. *Ann Intern Med* 1984;101:692-6.
- 4 Waitzkin H. Doctor-patient communication. Clinical implications of social scientific research. *JAMA* 1984;252:2441-6.
- 5 Makoul G, Arntson P, Schofield T. Health promotion in primary care: physician-patient communication and decision making about prescription medications. *Soc Sci Med* 1995;41:1241-54.

Commentary: "Double loop" learning

Ed Peile

Division of Medical
Education, Warwick
Medical School,
University of
Warwick, Coventry
CV4 7AL
Ed Peile
professor of medical
education
ed.peile@
warwick.ac.uk

As well as sympathising with the patient, many respondents to this case commentary found themselves engaging in the difficulties faced by the general practitioner.¹ Donald Schon, in *Educating the Reflective Practitioner*, described a "swampy lowland" wherein lie the problems of greatest human concern and where "messy, confusing problems defy technical solution."² Schon contrasts beautifully the technical and rational

approaches of the theoretical high ground with the reflective approaches we often need in professional practice.

In interactive case reports, readers are asked how they would react to a given situation. The usual response for professionals is to delineate what Schon and his co-author, American psychologist Chris Argyris, call our "theory of action."³ They describe this