Published in final edited form as:

Transpl Int. 2016 December ; 29(12): 1286–1295. doi:10.1111/tri.12855.

Liver transplant recipients with portal vein thrombosis receiving an organ from a high-risk donor are at increased risk for graft loss due to hepatic artery thrombosis

Jonathan G. Stine, MD¹, Curtis K. Argo, MD¹, Shawn J. Pelletier, MD², Daniel G. Maluf, MD², and Patrick G. Northup, MD¹

¹Center for the Study of Coagulation Disorders in Liver Disease, Division of Gastroenterology & Hepatology, Department of Medicine, University of Virginia

²Division of Transplant, Department of Surgery, University of Virginia

Abstract

Problem—We hypothesize that recipients with pre-transplant portal vein thrombosis (PVT) receiving organs from high-risk donors (HRD) are at increased risk of HAT.

Methods—Data on all liver transplants in the United States from February 2002-March 2015 were analyzed. Recipients were sorted into two groups: those with PVT and those without. HRDs were defined by Donor Risk Index (DRI) >1.7. Multivariable logistic regression models were constructed to assess independent risk factors for HAT with resultant graft loss 90 days from transplantation.

Results—60,404 candidates underwent liver transplantation; of those recipients, 623 (1.0%) had HAT, of which 66.0% (n=411) received organs from HRDs compared to 49.3% (n=29,473) in recipients without HAT (p<0.001). 2,250 (3.7%) recipients had pre-transplantation PVT and received organs from HRDs. On adjusted multivariable analysis, PVT with a HRD organ was the most significant independent risk factor (OR 3.56, 95% CI 2.52–5.02, p<0.001) for the development of HAT.

Conclusions—Candidates with pre-transplant PVT who receive an organ from a HRD are at the highest risk for post-operative HAT independent of other measurable factors. Recipients with pre-

Corresponding Author: Patrick G. Northup, MD MHS, Division of Gastroenterology and Hepatology, JPA and Lee Street, MSB 2145, PO Box 800708, University of Virginia, Charlottesville, VA 22908-0708, Office: 434.243.2718, Fax: 434.244.9454, northup@virginia.edu.

Meeting: This research was presented as a podium presentation at AASLD section of Digestive Diseases Week 2016, San Diego, CA. This research was also presented as a podium presentation at the Young Investigators Forum during the Controversies in Transplantation Conference, 2016, Breckenridge, CO.

Conflicts of interest/disclosures: The authors of this manuscript have no conflicts of interest to disclose.

Authorship statement: All authors participated in the listed roles below.

JS- planning/conducting study, collecting and/or interpreting data, drafting manuscript, final approval

SP- drafting manuscript, final approval

CA- drafting manuscript, final approval

DM- drafting manuscript, final approval

PN- planning/conducting study, collecting and/or interpreting data, drafting manuscript, final approval

Guarantor of the article: Patrick G. Northup MD MHS

transplant PVT would benefit from careful donor selection and possibly anticoagulation perioperatively.

Keywords

Hepatology; Coagulopathy; Cirrhosis; Portal hypertension; Outcomes

Introduction

Hepatic artery thrombosis (HAT) is an uncommon complication after liver transplantation with serious clinical implications including graft loss and increased recipient mortality.[1–3] Surgical risk factors including organ cold ischemia time (CIT), surgical technique, delay in reperfusion and anatomic abnormalities all likely play a role in outcomes.[4, 5] Less is understood about recipient-specific risk factors; however, prior liver transplantation, inherited thrombophilia, primary sclerosing cholangitis, acute intermittent porphyria and onset of diabetes post-transplantation may be associated with HAT.[6–12] We have previously shown that liver transplant recipients with pre-transplantation portal vein thrombosis (PVT) are at increased odds of HAT and that donor risk index (DRI) is predictive of HAT.[3] PVT is commonly associated with increased hepatic decompensation and mortality complication in patients with cirrhosis.[13, 14] [15] Both PVT and increased DRI in liver transplantation recipients are associated with a higher rate of graft loss due to primary non-function, in many cases leading to re-transplantation.[16] While careful donor selection may be important in order to prevent post-operative HAT in recipients with pre-transplantation PVT, a widely accepted cutoff has not yet been established. Originally proposed by Feng et al in 2006,[17] DRI is the most widely used system to evaluate donor risk and assist the transplant team in decision-making regarding graft utilization.[18] A DRI cutoff of greater than 1.7 has been proposed to define “high-risk” and validated by multiple studies showing this value is predictive of poorer post-transplantation outcomes including recipient survival.[19, 20] In this retrospective cross-sectional study of liver transplant recipients in the United States, we aimed to examine HAT risk factors in liver transplant recipients. Owing to their presumed hypercoagulable nature, we hypothesize that recipients with pre-transplant PVT who receive organs from high-risk donors (HRD) are at increased risk of HAT.

METHODS

Study design and recipient characteristics

Data on all liver transplantations between February 1, 2002 and March 31, 2015 in the United States were analyzed from the Organ and Transplantation Network (OPTN) with permission from the United Network for Organ Sharing (UNOS). This cross-sectional nationwide database has been previously validated to analyze HAT and PVT in recipients undergoing liver transplantation.[3, 13, 21–23] Only recipients who were transplanted at or above 18 years of age were included in our analysis. All living donor transplants, re-transplants, status one candidates, multi-visceral transplants, acute liver failure transplants and recipients with transjugular intrahepatic portosystemic shunts (TIPS) were excluded. Due to the higher rate of non-thrombotic complications, the analysis was performed both

with and without donation after cardiac death (DCD) recipients and the fundamental conclusions of the statistical analysis were not changed, therefore DCD organ recipients were also excluded. In the OPTN/UNOS dataset, there are ten values for the cause of graft loss variable: “recurrent disease, infection, chronic rejection, acute rejection, de novo hepatitis, recurrent hepatitis, primary nonfunction, biliary, vascular thrombosis and other (write-in field).” Based on previously validated methodology,[3, 24, 25] utilizing the “vascular thrombosis” and “other” category with the search terms “hepatic artery thrombosis,” HAT was defined by these two values provided there was resultant graft loss within 90 days of liver transplantation. Recipients who experienced HAT >90 days post transplantation were included in the non HAT group (n= 2). Missing data were handled initially by comparing the patients with known HAT status to those with unknown status. This comparison yielded one major difference; an increased percentage of recipients with underlying chronic hepatitis C (HCV) which we have previously shown to not be associated with an increased or decreased risk of HAT[3] and thus not a clinically relevant difference. Given the similarities in the two cohorts, the decision was made to include all recipients with unknown HAT status and to consider them as having “no HAT” in order to prevent inducing allocation bias.

Baseline recipient, operative and donor characteristics were analyzed. Recipient characteristics included the etiology of liver disease [hepatitis B, HCV, NASH/cryptogenic, autoimmune, cholestatic, hepatocellular carcinoma (HCC), alcoholic, and other liver diseases, which included recipients receiving an organ for any other reason than those above], severity of liver disease based on laboratory Model for End-Stage Liver Disease (MELD) score at transplantation and portal hypertensive complications including encephalopathy, ascites and PVT. Operative factors reviewed included locoregional or national organ sharing, CIT and parenteral heparin use at the time of cross clamping. Donor variables included organ steatosis content, age, ethnicity, cause of death, desmopressin (DDAVP) use for bleeding complications, cytomegalovirus status (IgG) and DRI which was dichotomized into high-risk and normal-risk (NRD) based on a previously validated cutoff of >1.7. [19, 20] The OPTN/UNOS dataset does not contain sufficient information regarding anticoagulant use for pre-existing venous thromboembolic disease nor does it contain variables specifying the presence of an inherited thrombophilic state.

Statistics

Using univariate logistic regression modeling, recipients with pre-transplantation PVT were compared statistically to those without PVT in an effort to compare recipient, operative and donor characteristics. Similarly, candidates receiving an organ from a HRD were compared to recipients transplanted with an organ from a standard risk donor (SRD). The primary outcome was graft loss secondary to HAT within the first 90 days of transplantation. Multivariable logistic regression models were then constructed to assess statistical associations and risk factors for the development of HAT utilizing maximum likelihood estimates. Variables were included in the final model only if they have previously been shown to be clinically relevant to the development of post-transplantation HAT or were statistically significant by univariate analysis ($p < 0.20$). [26, 27] Final variables included in the logistic regression model included recipient age at transplantation, gender, BMI, African

American race, diabetes, HCC, HCV, cholestatic liver disease, autoimmune hepatitis, encephalopathy (which was divided into those with encephalopathy score >2), ascites (similarly divided by score > 2), laboratories at transplantation (INR, bilirubin, creatinine, albumin, sodium), PVT], operative (heparin use at cross clamp, and donor factors (age, gender, HRD, DDAVP). Interaction terms for PVT and HRD, HCV and donor age, NASH and BMI and NASH and diabetes were included in the final model. Possible covariates for the PVT-HRD interaction variable included PVT with HRD, no PVT with HRD, as well as PVT with SRD. These were compared to a reference of no PVT with SRD. A p value of less than or equal to 0.05 was considered statistically significant, and all tests were two sided. Data imputation was not performed. SAS (version 9.4, Cary, NC) was utilized for all analysis and dataset manipulation. The OPTN/UNOS dataset is de-identified, thus, institutional review board approval was not required. No transplantations involving prisoners were included in our analysis.

RESULTS

60,404 recipients underwent liver transplantation from February 27, 2002, through March 31, 2015 and met our inclusion criteria; of these, 623 (1.0%) had HAT leading to early graft loss within 90 days of LT, which is similar to the accepted incidence of post-transplantation HAT. [3] Overall PVT prevalence was 7.5%, similar to previously published rates using the OPTN/UNOS dataset.[3] Of the recipients with post-transplantation HAT, 66.0% (n=411) received organs from HRDs compared to 49.3% (n=29,473) in recipients without HAT (p<0.001). Pre-transplant PVT was found in 13.5% (n=84) of recipients with post-transplantation HAT versus 7.5% (n=4,471) in those without HAT (p<0.001). 2,250 (3.7%) recipients had pre-transplantation PVT and received organs from HRDs.

On univariate analysis, recipient characteristics including demographics, etiology of liver disease, and severity of liver disease (both portal hypertension and laboratory values), were statistically similar or within marginal clinically important differences for patients with and without pre-transplantation PVT (Table 1) and when comparing HRD to NRD (Table 2), with several exceptions. Recipients with PVT were more likely to have pre-transplantation diabetes mellitus (30.2% versus 22.8%, p<0.001), NASH (17.4% versus 10.9%, p<0.001), HCC (26.4% versus 21.7%, p<0.001), and grade 3–4 ascites (32.2% versus 27.8%, p<0.001). Recipients with PVT were less likely to have underlying chronic HCV (25.3% versus 30.2%, p<0.001). In terms of surgical factors, heparin use at aortic cross-clamp was more common in recipients with pre-transplantation PVT (89.8% versus 84.8%, p<0.001) and CIT was slightly longer in the PVT group (6.95 +/- 3.34 hours versus 6.81 +/- 3.43 hours, p=0.015).

In comparing recipients receiving HRD versus SRD organs, several notable differences were observed in baseline characteristics. Candidates who received an organ from a HRD were less likely to be male (64.6% versus 69.6%, p<0.001), be on dialysis at the time of liver transplantation (8.3% versus 12.6%, p<0.001), have underlying chronic HCV (27.7% versus 31.8%, p<0.001), have lower mean BMI (27.0 +/- 5.66 kg/m² versus 28.4 +/- 10.2 kg/m², p<0.001), and have lower mean MELD scores (20.3 +/- 10.0 versus 22.1 +/- 10.6, p<0.001) at transplantation with corresponding differences in the individual MELD covariates (serum

bilirubin 7.33 $\pm$ 9.97 mg/dL versus 8.57 $\pm$ 11.2 mg/dL, $p < 0.001$; INR 1.83 $\pm$ 1.23 versus 1.92 $\pm$ 1.28, $p < 0.001$; creatinine 1.45 $\pm$ 1.24 g/dL versus 1.64 $\pm$ 1.48 g/dL, $p < 0.001$). Interestingly, the HRD group had a lower rate of anoxic donor death (11.5% versus 25.6%, $p < 0.001$). Heparin use at aortic cross-clamp was less likely as well (81.8% versus 88.5%, $p < 0.001$) as was DDAVP use (18.5% versus 24.0%, $p < 0.001$). HRD organ recipients were more likely to receive organs from older donors (mean age 52.5 $\pm$ 14.5 years versus 30.9 $\pm$ 11.1 years, $p < 0.001$), female donors (53.0% versus 29.4%, $p < 0.001$), donors with a cerebrovascular attack as the cause of death (57.2% versus 23.0%, $p < 0.001$), organ sharing both regionally (25.9% versus 16.0%, $p < 0.001$) and nationally (9.2% versus 0.5%, $p < 0.001$) and longer CIT (7.18 $\pm$ 4.08 hours versus 6.48 $\pm$ 2.58 hours, $p < 0.001$). CMV donor positivity was also more common in the HRD organ recipients (71.0% versus 60.6%, $p < 0.001$).

Multivariable regression analysis (Table 3) of risk factors for HAT with resultant graft loss within 90 days of liver transplantation demonstrated that in the presence of pre-transplantation PVT, using an organ from a HRD was statistically significantly associated with increased risk of post-transplantation HAT with OR 3.56, 95% CI 2.52–5.012 $p < 0.001$. Recipients with pre-transplantation PVT who received an organ from a SRD still had an increased odds of post-transplantation HAT (OR 2.25, 95% CI 1.48–3.42, $p < 0.001$) as did recipients without pre-transplantation PVT who received a HRD organ (OR 1.71, 95% CI 1.36–2.14, $p < 0.001$). Other factors protective against HAT included creatinine values at transplantation (OR 0.91, 95% CI 0.83–0.99, $p = 0.035$), heparin use at cross-clamp (OR 0.61, 95% CI 0.47–0.79, $p < 0.001$), INR values at transplantation (OR 0.87, 95% CI 0.76–0.98, $p = 0.027$) and receiving an organ from a male donor (OR 0.53, 95% CI 0.44–0.64, $p < 0.001$). These recipient, surgical and donor factors were all independently associated with a lower risk of HAT and early graft loss. While significant on univariate analysis, diabetes was not found to be independently predictive with multivariable regression modeling (OR 0.88, 95% CI 0.68–1.14, $p = 0.318$). Recipient age at transplantation was statistically significant on both univariate and multivariable analysis and for each year of age a recipient's risk of HAT decreased by 2% (OR 0.98, 95% CI 0.97–0.99, $p < 0.001$).

DISCUSSION

Based on a large U.S. based national liver transplantation database and after adjusting for known donor, recipient and surgical risk factors, we have found that liver transplant recipients with pre-transplant PVT who receive an organ from a HRD (DRI > 1.7) are at increased odds of post-transplantation HAT. These findings lend credence to the consideration of more than just surgical technique and CIT in determining a recipient's risk for HAT, especially in the setting of pre-transplantation coagulation abnormalities, including PVT, which by itself is associated with higher rates of primary graft nonfunction and re-transplantation.[16] With the increasing utilization of HRDs to meet the organ supply and demand issue,[28] it appears we can expect more issues with early graft loss from post-transplantation thrombosis and we would suggest this patient population as a potential target recipient for more aggressive post-operative care and prevention, certainly in the presence of pre-transplantation venothromboembolic disease.

Our findings also raise the question of what is the most appropriate pre-, peri- and postoperative management strategy for anticoagulation? Efficacy data on routine administration of antiplatelet agents (namely aspirin) for the prevention of post-operative HAT is mixed and complicated by retrospective study limitations including varying time definitions of early and late HAT; however, aspirin does appear safe as bleeding events are generally infrequent.[29–31] We are unaware of any studies looking at dual antiplatelet therapy as prophylaxis. Our Coagulation in Liver Disease Study Group recently published a series of 39 patients with cirrhosis comparing the safety of utilizing direct oral anticoagulants (DOAC) (factor Xa inhibitors apixiban and rivaroxaban) to traditional anticoagulants warfarin and low-molecular weight heparin (LMWH) and found similar safety profiles in terms of bleeding and no episodes of drug-induced liver injury.[32] The bleeding rate of 5% with DOAC and the absence of fatal bleeding was similar to the pooled incidence from a recent meta-analysis of 16 studies by Qi et al. who found a pooled rate of 3.3% for LMWH or warfarin use, 95% CI 1.1–6.7%.[33] The widespread use of DOACs may be limited by the lack of currently available reversal agents, although a recent study randomized placebo-controlled study of 101 healthy older adults, none of whom had liver disease, by Siegal et al[34] utilizing intravenous andexanet alpha holds promise. While the exact role of DOACs in the management of peri-operative coagulopathy unique to liver transplantation recipients has yet to be firmly established, these data are nonetheless promising and provide a starting point for future prospective study. Given that there are significant hemostatic abnormalities in the peri- and post-operative periods that may interact to create a hypercoagulable milieu[35] it seems that DOAC use in recipients at high-risk for post-transplantation thrombotic complications could be considered at least until the coagulability equilibrium shifts away from a pro-thrombotic state, the timing of which could be aided by the routine use of post-transplantation thromboelastography (TEG), a method for determining the real-time viscous and elastic properties of blood and blood clot.

While PVT is a common complication in patients with cirrhosis[13, 14] and is associated with both increased hepatic decompensation and mortality, [15] prospective studies with a direct comparator group examining both pharmacologic prevention and treatment of PVT is lacking and consensus guidelines on pre-operative management of PVT have yet to be disseminated. Nonetheless, several unblinded single center studies have found both regression of liver disease with mitigation of portal hypertensive complications as well as a long-term survival benefit with either daily prophylactic (the equivalent of 40mg/day) [36] or therapeutic dosing (1 mg/kg every twelve hours or 1.5mg/kg daily) [37] of low molecular weight heparin. Following these studies, multi-center investigation of the safety and efficacy of prevention of PVT with pharmacologic anticoagulation is currently underway with anticipated results within the next five years.

A recent multi-center Canadian experience with 118 liver transplant recipients published by Seal et al[38] found that intraoperative use of tissue plasminogen activator (tPA) directly into the donor hepatic artery either before hepatic artery anastomosis or portal vein anastomosis (5–10 minutes before portal reperfusion) reduced complications from ischemic biliary strictures and led to superior one- and three-year overall and graft survival rates in recipients receiving DCD organs and significantly lower re-transplantation rates in the absence of increased bleeding. While the authors did not look specifically at HAT and others have

argued that the inherent profound activation of fibrinolysis with withdrawal of life support is enough to prevent significant downstream ischemic thrombotic biliary complications,[39] these findings are nonetheless thought provoking given that no deaths or graft loss was attributable to HAT in the tPA group. These findings combined with our findings that intraoperative administration of heparin at the time of aortic cross-clamping is associated with a lower risk of HAT, lend importance to further consideration of perioperative anticoagulation when using organs from HRDs.

Our study has several weaknesses worth noting. Despite aggressive verification by auditors and data technicians, large datasets are dependent on diagnostic coding accuracy to preclude induction of information bias and also suffer from bias due to missing data, and the OPTN/UNOS database is no exception to this rule.[40] Our method of handling the missing data provided similar incidence rates of HAT when compared to those published by other study groups who utilized the UNOS database to investigate HAT. The OPTN/UNOS database also does not contain information on pre-operative anticoagulant use nor does it provide information on inherited thrombophilia testing. However, previous study has found the incidence of inherited thrombophilia to be similar when comparing recipients who experience post transplantation HAT to those who do not.[1] The database also does not contain a description of the extent of pre-transplantation PVT, including which specific vessels are involved and whether or not the clot is partial or complete nor does it contain information regarding the method of surgical reconstruction including arterial reconstruction which is arguably the most technically challenging aspect of liver transplant surgery and has been implicated in post-transplantation complications including HAT.[5, 6, 41–44] Information on the extent of hepatic hilar manipulation during portal vein reconstruction is also lacking. This may predispose to hepatic artery injury or rethrombosis of the portomesenteric veins.[45] Interestingly, a recent single center experience of ten patients with grade four PVT suggests that intraoperative renoportal bypass combined with direct measurements of portal vein and hepatic artery flow may be used successfully to mitigate post-transplantation complications including primary graft non-function and mortality associated with pre-transplantation PVT.[46] Creating renoportal or cavoportal anastomoses in transplant recipients with pre-transplantation PVT may also decrease the risk of post-transplantation vascular complications.[47]

Conclusions

Liver transplant candidates with pre-transplant PVT who receive an organ from a HRD are at the highest risk for post-operative HAT independent of other measurable recipient, surgical and donor factors. Utilizing a DRI cutoff of 1.7 in order to prevent post-operative vascular thrombotic complications may be useful. Recipients with pre-transplant PVT would benefit from careful donor selection and consideration of aggressive perioperative anticoagulation to improve patient centered outcomes.

Acknowledgments

Financial support: Research reported in this publication was supported by the National Institute of Diabetes and Digestive and Kidney Diseases of the National Institutes of Health under award number T32DK007769.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

This work was supported in part by Health Resources and Services Administration contract 234-2005-370011C. The content is the responsibility of the authors alone and does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

Abbreviations

BMI	body mass index
CIT	cold ischemia time
CMV	cytomegalovirus
DCD	donation after cardiac death
DDAVP	desmopressin
DOAC	direct oral anticoagulant
DRI	donor risk index
HAT	hepatic artery thrombosis
HRD	high-risk donor
HCC	hepatocellular carcinoma
HCV	hepatitis C virus
INR	international normalized ratio
LT	liver transplant
NASH	non-alcoholic steatohepatitis
OPTN	organ procurement and transplantation network
PVT	portal vein thrombosis
TEG	thromboelastography
TIPS	transjugular intrahepatic portosystemic shunt
TPA	tissue plasminogen activator
UNOS	united network for organ sharing

References

1. Ayala R, Martinez-Lopez J, Cedena T, et al. Recipient and donor thrombophilia and the risk of portal venous thrombosis and hepatic artery thrombosis in liver recipients. *BMC Gastroenterol*. 2011; 11:130. [PubMed: 22123067]

2. Iida T, Kaido T, Yagi S, et al. Hepatic arterial complications in adult living donor liver transplant recipients: a single-center experience of 673 cases. *Clinical transplantation*. 2014; 28:1025–30. [PubMed: 24974916]
3. Stine JG. Pre-transplant portal vein thrombosis is an independent risk factor for graft loss due to hepatic artery thrombosis in liver transplant recipients. *HPB*. 2015. <http://dx.doi.org/10.1016/j.jhpb.2015.10008> in press, epub Dec 9 2015
4. Cescon M, Zanello M, Grazi GL, et al. Impact of very advanced donor age on hepatic artery thrombosis after liver transplantation. *Transplantation*. 2011; 92:439–45. [PubMed: 21712754]
5. Warner P, Fusai G, Glantzounis GK, et al. Risk factors associated with early hepatic artery thrombosis after orthotopic liver transplantation - univariable and multivariable analysis. *Transplant international: official journal of the European Society for Organ Transplantation*. 2011; 24:401–8. [PubMed: 21210866]
6. Vivarelli M, Cucchetti A, La Barba G, et al. Ischemic arterial complications after liver transplantation in the adult: multivariate analysis of risk factors. *Archives of surgery (Chicago, Ill: 1960)*. 2004; 139:1069–74.
7. Oh CK, Pelletier SJ, Sawyer RG, et al. Uni- and multi-variate analysis of risk factors for early and late hepatic artery thrombosis after liver transplantation. *Transplantation*. 2001; 71:767–72. [PubMed: 11330540]
8. Dunn TB, Linden MA, Vercellotti GM, Gruessner RW. Factor V Leiden and hepatic artery thrombosis after liver transplantation. *Clinical transplantation*. 2006; 20:132–5. [PubMed: 16556168]
9. Mas VR, Fisher RA, Maluf DG, Wilkinson DS, Garrett CT, Ferreira-Gonzalez A. Hepatic artery thrombosis after liver transplantation and genetic factors: prothrombin G20210A polymorphism. *Transplantation*. 2003; 76:247–9. [PubMed: 12865818]
10. Dowman JK, Gunson BK, Mirza DF, Bramhall SR, Badminton MN, Newsome PN. Liver transplantation for acute intermittent porphyria is complicated by a high rate of hepatic artery thrombosis. *Liver Transpl*. 2012; 18:195–200. [PubMed: 21618697]
11. Graziadei IW, Wiesner RH, Marotta PJ, et al. Long-term results of patients undergoing liver transplantation for primary sclerosing cholangitis. *Hepatology*. 1999; 30:1121–7. [PubMed: 10534330]
12. Moon JJ, Barbeito R, Faradji RN, Gaynor JJ, Tzakis AG. Negative impact of new-onset diabetes mellitus on patient and graft survival after liver transplantation: Long-term follow up. *Transplantation*. 2006; 82:1625–8. [PubMed: 17198248]
13. Stine JG, Shah NL, Argo CK, Pelletier SJ, Caldwell SH, Northup PG. Increased risk of Portal Vein Thrombosis in Patients with Cirrhosis due to Non-Alcoholic Steatohepatitis (NASH). *Liver Transpl*. 2015
14. Wanless IR, Wong F, Blendis LM, Greig P, Heathcote EJ, Levy G. Hepatic and portal vein thrombosis in cirrhosis: possible role in development of parenchymal extinction and portal hypertension. *Hepatology (Baltimore, Md)*. 1995; 21:1238–47.
15. Stine JG, Shah PM, Cornella SL, et al. Portal vein thrombosis, mortality and hepatic decompensation in patients with cirrhosis: A meta-analysis. *World J Hepatol*. 2015; 7:2774–80. [PubMed: 26644821]
16. Sharma R, Kashyap R, Jain A, et al. Surgical complications following liver transplantation in patients with portal vein thrombosis--a single-center perspective. *Journal of gastrointestinal surgery: official journal of the Society for Surgery of the Alimentary Tract*. 2010; 14:520–7. [PubMed: 19960270]
17. Feng S, Goodrich NP, Bragg-Gresham JL, et al. Characteristics associated with liver graft failure: the concept of a donor risk index. *Am J Transplant*. 2006; 6:783–90. [PubMed: 16539636]
18. Akkina SK, Asrani SK, Peng Y, Stock P, Kim WR, Israni AK. Development of organ-specific donor risk indices. *Liver Transpl*. 2012; 18:395–404. [PubMed: 22287036]
19. Avolio AW, Siciliano M, Barbarino R, et al. Donor risk index and organ patient index as predictors of graft survival after liver transplantation. *Transplant Proc*. 2008; 40:1899–902. [PubMed: 18675083]

20. Bonney GK, Aldersley MA, Asthana S, et al. Donor risk index and MELD interactions in predicting long-term graft survival: a single-centre experience. *Transplantation*. 2009; 87:1858–63. [PubMed: 19543065]
21. Stewart ZA, Locke JE, Segev DL, et al. Increased risk of graft loss from hepatic artery thrombosis after liver transplantation with older donors. *Liver Transpl*. 2009; 15:1688–95. [PubMed: 19938120]
22. Mateo R, Cho Y, Singh G, et al. Risk factors for graft survival after liver transplantation from donation after cardiac death donors: an analysis of OPTN/UNOS data. *Am J Transplant*. 2006; 6:791–6. [PubMed: 16539637]
23. Zocco MA, Di Stasio E, De Cristofaro R, et al. Thrombotic risk factors in patients with liver cirrhosis: correlation with MELD scoring system and portal vein thrombosis development. *J Hepatol*. 2009; 51:682–9. [PubMed: 19464747]
24. Vaidya S, Dighe M, Kolokythas O, Dubinsky T. Liver transplantation: vascular complications. *Ultrasound quarterly*. 2007; 23:239–53. [PubMed: 18090835]
25. Wozney P, Zajko AB, Bron KM, Point S, Starzl TE. Vascular complications after liver transplantation: a 5-year experience. *AJR American journal of roentgenology*. 1986; 147:657–63. [PubMed: 3529892]
26. Bursac Z, Gauss CH, Williams DK, Hosmer DW. Purposeful selection of variables in logistic regression. *Source code for biology and medicine*. 2008; 3:17. [PubMed: 19087314]
27. Sun GW, Shook TL, Kay GL. Inappropriate use of bivariable analysis to screen risk factors for use in multivariable analysis. *Journal of clinical epidemiology*. 1996; 49:907–16. [PubMed: 8699212]
28. Tector AJ, Mangus RS, Chestovich P, et al. Use of extended criteria livers decreases wait time for liver transplantation without adversely impacting posttransplant survival. *Ann Surg*. 2006; 244:439–50. [PubMed: 16926570]
29. Wolf DC, Freni MA, Boccagni P, et al. Low-dose aspirin therapy is associated with few side effects but does not prevent hepatic artery thrombosis in liver transplant recipients. *Liver transplantation and surgery: official publication of the American Association for the Study of Liver Diseases and the International Liver Transplantation Society*. 1997; 3:598–603.
30. Shay R, Taber D, Pilch N, et al. Early aspirin therapy may reduce hepatic artery thrombosis in liver transplantation. *Transplant Proc*. 2013; 45:330–4. [PubMed: 23267805]
31. Vivarelli M, La Barba G, Cucchetti A, et al. Can antiplatelet prophylaxis reduce the incidence of hepatic artery thrombosis after liver transplantation? *Liver Transpl*. 2007; 13:651–4. [PubMed: 17457885]
32. Intagliata NM, Henry ZH, Maitland H, et al. Direct Oral Anticoagulants in Cirrhosis Patients Pose Similar Risks of Bleeding When Compared to Traditional Anticoagulation. *Dig Dis Sci*. 2016
33. Qi X, De Stefano V, Li H, Dai J, Guo X, Fan D. Anticoagulation for the treatment of portal vein thrombosis in liver cirrhosis: a systematic review and meta-analysis of observational studies. *European journal of internal medicine*. 2015; 26:23–9. [PubMed: 25566699]
34. Siegal DM, Curnutte JT, Connolly SJ, et al. Andexanet Alfa for the Reversal of Factor Xa Inhibitor Activity. *N Engl J Med*. 2015; 373:2413–24. [PubMed: 26559317]
35. Arshad F, Lisman T, Porte RJ. Hypercoagulability as a contributor to thrombotic complications in the liver transplant recipient. *Liver international: official journal of the International Association for the Study of the Liver*. 2013; 33:820–7. [PubMed: 23490221]
36. Villa E, Camma C, Marietta M, et al. Enoxaparin prevents portal vein thrombosis and liver decompensation in patients with advanced cirrhosis. *Gastroenterology*. 2012; 143:1253–60. e1–4. [PubMed: 22819864]
37. Cui SB, Shu RH, Yan SP, et al. Efficacy and safety of anticoagulation therapy with different doses of enoxaparin for portal vein thrombosis in cirrhotic patients with hepatitis B. *Eur J Gastroenterol Hepatol*. 2015
38. Seal JB, Bohorquez H, Reichman T, et al. Thrombolytic protocol minimizes ischemic-type biliary complications in liver transplantation from donation after circulatory death donors. *Liver Transpl*. 2015; 21:321–8. [PubMed: 25545787]

39. Burlage LC, Karangwa SA, Lisman T, Martins PN, Porte RJ. Thrombolytic protocol minimizes ischemic-type biliary complications in liver transplantation from donation after circulatory death donors. *Liver Transpl.* 2015; 21:1231–2. [PubMed: 26045355]
40. Gillespie BW, Merion RM, Ortiz-Rios E, et al. Database comparison of the adult-to-adult living donor liver transplantation cohort study (A2ALL) and the SRTR U.S. Transplant Registry. *Am J Transplant.* 2010; 10:1621–33. [PubMed: 20199501]
41. Stange BJ, Glanemann M, Nuessler NC, Settmacher U, Steinmuller T, Neuhaus P. Hepatic artery thrombosis after adult liver transplantation. *Liver Transpl.* 2003; 9:612–20. [PubMed: 12783404]
42. Melada E, Maggi U, Rossi G, et al. Back-table arterial reconstructions in liver transplantation: single-center experience. *Transplant Proc.* 2005; 37:2587–8. [PubMed: 16182752]
43. Silva MA, Jambulingam PS, Gunson BK, et al. Hepatic artery thrombosis following orthotopic liver transplantation: a 10-year experience from a single centre in the United Kingdom. *Liver Transpl.* 2006; 12:146–51. [PubMed: 16382467]
44. Ravaioli M, Zanella M, Grazi GL, et al. Portal vein thrombosis and liver transplantation: evolution during 10 years of experience at the University of Bologna. *Ann Surg.* 2011; 253:378–84. [PubMed: 21183851]
45. Hibi T, Nishida S, Levi DM, et al. When and why portal vein thrombosis matters in liver transplantation: a critical audit of 174 cases. *Ann Surg.* 2014; 259:760–6. [PubMed: 24299686]
46. Quintini C, Spaggiari M, Hashimoto K, et al. Safety and effectiveness of renoportal bypass in patients with complete portal vein thrombosis: an analysis of 10 patients. *Liver Transpl.* 2015; 21:344–52. [PubMed: 25420619]
47. Bhangui P, Lim C, Salloum C, et al. Caval inflow to the graft for liver transplantation in patients with diffuse portal vein thrombosis: a 12-year experience. *Ann Surg.* 2011; 254:1008–16. [PubMed: 21869678]

Table 1

Recipient baseline characteristics comparing those with portal vein thrombosis (PVT) to those without PVT

	Portal Vein Thrombosis (n=4,555)	No Portal Vein Thrombosis (n=55,849)	p-value
Recipient Characteristics			
Age at transplant, mean years	55.7 +/- 9.5	53.8 +/- 10.2	<0.001
Male gender, n (%)	3,191 (70.1)	37,350 (66.9)	<0.001
African American race, n (%)	285 (6.3)	5,572 (10.0)	<0.001
Diabetes, n (%)	1,376 (30.2)	12,740 (22.8)	<0.001
On dialysis at transplantation, n (%)	511 (11.2)	5,821 (10.4)	NS
BMI, mean kg/m ²	28.7 +/- 5.7	28.2 +/- 5.7	<0.001
Etiology of liver disease, n (%)			
Alcoholic liver disease	500 (11.0)	6,592 (11.8)	NS
Autoimmune disease	123 (2.7)	1,384 (2.5)	NS
Cholestatic disease	275 (6.0)	4,180 (7.5)	<0.001
Hepatitis B	75 (1.7)	1,201 (2.2)	0.023
Hepatitis C	1,151 (25.3)	16,847 (30.2)	<0.001
NASH	794 (17.4)	6,058 (10.9)	<0.001
Other	1,637 (35.9)	19,587 (35.0)	NS
Severity of Liver Disease			
MELD score at transplantation, mean	21.9 +/- 9.8	21.2 +/- 10.4	<0.001
HCC, n (%)	1,204 (26.4)	12,136 (21.7)	<0.001
Ascites grade > 2 at transplant, n (%)	1,465 (32.2)	15,533 (27.8)	<0.001
HE > 2 at transplant, n (%)	474 (10.4)	6,124 (11.0)	NS
Laboratory values			
Serum bilirubin, mg/dL, mean	7.74 +/- 10.72	7.79 +/- 10.52	NS
INR, mean	1.89 +/- 0.97	1.87 +/- 1.28	NS
Serum albumin, g/dL, mean	3.03 +/- 0.73	3.03 +/- 0.73	NS
Creatinine, g/dL, mean	1.54 +/- 1.16	1.55 +/- 1.38	NS
Serum sodium, mEq/L, mean	135.7 +/- 5.2	136.1 +/- 5.2	<0.001
Donor characteristics			
Age donor, mean years	41.7 +/- 16.9	41.5 +/- 16.8	NS
Male donor, n (%)	2,678 (58.8)	32,904 (58.9)	NS
African American donor, n (%)	803 (17.6)	9,120 (16.3)	0.023
Anoxic cause of death, n (%)	969 (21.3)	10,289 (18.4)	<0.001
Cerebrovascular attack as cause of death, n (%)	1,786 (39.3)	22,315 (40.0)	NS
Regional organ sharing, n (%)	924 (20.3)	11,704 (21.0)	NS
National organ sharing, n (%)	210 (4.6)	2,683 (4.8)	NS
CMV donor positivity n (%)	2,860 (65.5)	35,000 (65.5)	NS
DRI, mean	1.77 +/- 0.40	1.78 +/- 0.40	NS

	Portal Vein Thrombosis (n=4,555)	No Portal Vein Thrombosis (n=55,849)	p-value
High-risk donor (DRI > 1.7), n (%)	2,250 (49.4)	27,634 (49.5)	NS
Macrovesicular fat content of donor liver, mean percent	8.3 +/- 11.7	8.5 +/- 12.1	NS
Surgical characteristics			
Heparin use at cross clamp, n (%)	4,091 (89.8)	47,372 (84.8)	<0.001
DDAVP use, n (%)	66 (1.5)	396 (0.7)	<0.001
CIT, mean hours	6.95 +/- 3.34	6.81 +/- 3.43	0.015

BMI=Body mass index; CI=Confidence interval; CIT=Cold ischemia time;

CMV=Cytomegalovirus; DRI=Donor risk index; DDAVP=Desmopressin; HE=Hepatic encephalopathy; HCC=Hepatocellular carcinoma;
INR=International normalized ratio; NASH=Non-alcoholic steatohepatitis; NS=Not significant; PVT=Portal vein thrombosis

Table 2

Baseline characteristics of high-risk donors compared to low-risk donors

	High-risk Donor (n=29,884)	Low-risk Donor (n=30,520)	p-value
Recipient Characteristics			
Age at transplant, mean years	54.4 +/- 10.1	53.4 +/- 10.2	<0.001
Male gender, n (%)	19,291 (64.6)	21,250 (69.6)	<0.001
African American race, n (%)	2,681 (9.0)	3,176 (10.4)	<0.001
Diabetes, n (%)	7,151 (23.9)	6,965 (22.8)	0.001
On dialysis at transplantation, n (%)	2,476 (8.3)	3,856 (12.6)	<0.001
BMI, mean kg/m2	27.0 +/- 5.66	28.4 +/- 5.66	<0.001
Etiology of liver disease, n (%)			
Alcoholic liver disease	3,578 (12.0)	3,514 (11.5)	NS
Autoimmune disease	726 (2.4)	781 (2.6)	NS
Cholestatic disease	2,530 (8.5)	1,925 (6.3)	<0.001
Hepatitis B	648 (2.2)	628 (2.1)	NS
Hepatitis C	8,279 (27.7)	9,719 (31.8)	<0.001
NASH	3,584 (12.0)	3,268 (10.7)	<0.001
Other	10,539 (35.3)	10,685 (35.0)	NS
Severity of Liver Disease			
MELD score at transplantation, mean	20.3 +/- 10.0	22.1 +/- 10.6	<0.001
HCC, n (%)	6,688 (22.4)	6,652 (21.8)	NS
PVT, n (%)	2,250 (7.5)	2,305 (7.6)	NS
Ascites grade > 2 at transplant, n (%)	8,041 (26.9)	6,957 (29.4)	<0.001
HE > 2 at transplant, n (%)	3,065 (10.3)	3,534 (11.6)	<0.001
Laboratory values			
Serum bilirubin, mg/dL, mean	7.33 +/- 9.97	8.57 +/- 11.20	<0.001
INR, mean	1.83 +/- 1.23	1.92 +/- 1.28	<0.001
Serum albumin, g/dL, mean	3.04 +/- 0.72	3.01 +/- 0.73	<0.001
Creatinine, g/dL, mean	1.45 +/- 1.24	1.64 +/- 1.48	<0.001
Serum sodium, mEq/L, mean	136.1 +/- 5.20	136.1 +/- 5.20	NS
Donor characteristics			
Age donor, mean years	52.5 +/- 14.5	30.9 +/- 11.1	<0.001
Male donor, n (%)	14,043 (47.0)	21,539 (70.6)	<0.001
African American donor, n (%)	5,346 (17.9)	4,577 (15.0)	<0.001
Anoxic cause of death, n (%)	3,447 (11.5)	7,811 (25.6)	<0.001
Cerebrovascular attack as cause of death, n (%)	17,078 (57.2)	7,023 (23.0)	<0.001
Regional organ sharing, n (%)	7,744 (25.9)	4,884 (16.0)	<0.001
National organ sharing, n (%)	2,756 (9.2)	137 (0.5)	<0.001
CMV donor positivity n (%)	19,507 (71.0)	18,353 (60.6)	<0.001
Macrovesicular fat content of donor liver, mean percent	8.54 +/- 11.88	8.41 +/- 12.43	NS

	High-risk Donor (n=29,884)	Low-risk Donor (n=30,520)	p-value
Surgical characteristics			
Heparin use at cross clamp, n (%)	24,453 (81.8)	27,010 (88.5)	<0.001
DDAVP use, n (%)	5,526 (18.5)	7,321 (24.0)	<0.001
CIT, mean hours	7.18 +/- 4.08	6.48 +/- 2.58	<0.001

BMI=Body mass index; CI=Confidence interval; CIT=Cold ischemia time;

CMV=Cytomegalovirus; DRI=Donor risk index; DDAVP=Desmopressin; HE=Hepatic encephalopathy; HCC=Hepatocellular carcinoma; INR=International normalized ratio; NASH=Non-alcoholic steatohepatitis; NS=Not significant; PVT=Portal vein thrombosis

Table 3

Univariate and multivariable analysis of risk factors for graft loss due to hepatic artery thrombosis within 90 days of liver transplantation

	Univariate analysis (OR, 95% CI)	Multivariable analysis (OR, 95% CI)	p-values for multivariable model
Recipient Characteristics			
Age at transplant, mean years	0.98 (0.98–0.99)	0.98 (0.97–0.99)	0.001
Male gender	0.94 (0.79–1.11)		
African American race	1.05 (0.81–1.36)		
Diabetes	0.78 (0.64–0.95)	0.88 (0.68–1.14)	0.318
On dialysis at transplantation	0.79 (0.60–1.04)		
BMI, mean kg/m ²	1.01 (0.99–1.02)		
Etiology of liver disease			
Alcoholic liver disease	0.91 (0.70–1.17)		
Autoimmune disease	1.37 (0.85–2.12)		
Cholestatic disease	1.24 (0.94–1.64)		
Hepatitis B	1.07 (0.63–1.82)		
Hepatitis C	0.97 (0.82–1.16)		
NASH	1.14 (0.90–1.44)		
Severity of Liver Disease			
HCC	0.82 (0.67–1.01)		
Ascites grade > 2 at transplant	0.90 (0.85–1.07)		
HE > 2 at transplant	1.08 (0.85–1.39)		
Laboratory values			
Serum bilirubin, mg/dL	0.99 (0.98–1.00)		
INR	0.88 (0.79–0.97)	0.87 (0.76–0.98)	0.027
Serum albumin, g/dL	0.97 (0.87–1.08)		
Creatinine, g/dL	0.92 (0.86–0.98)	0.91 (0.83–0.99)	0.035
Serum sodium, mEq/L	0.99 (0.98–1.01)		
Donor characteristics			

	Univariate analysis (OR, 95% CI)	Multivariable analysis (OR, 95% CI)	p-values for multivariable model
Male donor	0.52 (0.45–0.61)	0.53 (0.44–0.64)	<0.001
High DRI	1.99 (1.69–2.35)	---	
Macrovesicular fat content of donor liver	1.00 (0.96–1.01)		
Surgical characteristics			
Heparin use at cross clamp	0.52 (0.43–0.63)	0.61 (0.47–0.79)	0.002
DDAVP use	0.90 (0.74–1.10)		
Thrombosis			
PVT and HRD[#]	3.84 (2.83–5.21)	3.56 (2.52–5.02)	<0.001
No PVT and HRD[#]	2.00 (1.67–2.39)	2.25 (1.49–3.42)	<0.001
PVT and SRD[#]	1.95 (1.32–2.89)	1.71 (1.36–2.15)	<0.001
PVT	1.93 (1.53–2.43)	---	

BMI=Body mass index; CI=Confidence interval; CIT=Cold ischemia time;

CMV=Cytomegalovirus; DRI=Donor risk index; DDAVP=Desmopressin; HE=Hepatic encephalopathy; HCC=Hepatocellular carcinoma; HRD=High-risk donor (DRI>1.7); INR=International normalized ratio; NASH=Non-alcoholic steatohepatitis; NS=Not significant; PVT=Portal vein thrombosis; SRD=Standard-risk donor (DRI 1.7)

[#] compared to reference group of no PVT and NRD

Final variables included in the logistic regression model included recipient age at transplantation, gender, BMI, African American race, diabetes, HCC, HCV, cholestatic liver disease, autoimmune hepatitis, encephalopathy (which was divided into those with encephalopathy score >2), ascites (similarly divided by score > 2), laboratories at transplantation (INR, bilirubin, creatinine, albumin, sodium), PVT], operative (heparin use at cross clamp, and donor factors (age, gender, HRD, DDAVP). Interaction terms for PVT and HRD, HCV and donor age, NASH and BMI and NASH and diabetes were included in the final model. Possible covariates for the PVT-HRD interaction variable included PVT with HRD, no PVT with HRD, as well as PVT with SRD.