In brief

Staff fail to treat mentally ill patients with respect: A survey of 416 patients who had stayed on mental health wards in England and Wales in the last two years showed that only one in five felt they were treated with respect and dignity by hospital staff, according to the mental health charity Mind. It also found that 23% of respondents had been in mixed sex accommodation. A summary of *Ward Watch* is at www.mind.org.uk

China passes its first AIDS related law: China's parliament, the National People's Congress, has passed a law that requires the government to provide funding for prevention and control of AIDS and bans the sale and purchase of human blood. In Henan province one million people were infected through blood selling schemes in the 1990s.

Long term care costs set to increase: The amount of money spent in the United Kingdom on providing long term care for an increasing population of old people will rise from the current £12bn (\$21bn; €18bn), or 1.4% of gross domestic product, to £53.9bn, or 1.8%, in 2051, says a report from the Joseph Rowntree Foundation. *Future Demand for Long-Term Care in the UK* is available at www.jrf.org.uk

US doctors sue health management organisations: 600 000 doctors have brought a class action suit against several management organisations, including Humana, Prudential, and United Healthcare, claiming that the organisations routinely put them through costly and time consuming exercises that made it unrewarding to challenge payments for medical procedures.

Long term conditions group launches manifesto: The Partnership on Long-term Conditions has launched a joint manifesto, "17 Million Reasons," in the House of Commons aimed at improving the lives of the 17 million British people with asthma, diabetes, arthritis, multiple sclerosis, or cancer. Members from the partnership have already met representatives from all political parties.

Poor lung function is five times commoner in teenagers exposed to high particulate concentrations

Scott Gottlieb *New York*

Current concentrations of air pollution can have chronic adverse effects on lung development in children, which lead to clinically important deficits in lung function by adulthood, a new study has shown (*New England Journal of Medicine* 2004; 357:1057-67).

Until now, whether exposure

to air pollution adversely affected the increase in lung function during the period of rapid lung development that occurs between the ages of 10 and 18 years was unknown.

In a prospective study, researchers recruited 1759 children (average 10 years old) from schools in 12 southern California

communities and measured lung function annually for eight years. About 10% of children were lost to follow up each year. The communities represented a wide range of ambient exposures to common pollutants, including ozone, acid vapour, nitrogen dioxide, and particulate matter.

The researchers—led by Dr W James Gauderman of the department of preventive medicine, University of Southern California, Los Angeles—measured the relationship between concentrations of air pollution and the forced expiratory volume in one second (FEV₁) and other spirometric measures. They found that, over the eight years, deficits in the growth of FEV₁ were associated with exposure to nitrogen dioxide (P=0.005), acid vapour (P=0.004), particulate matter (P=0.04), and elemental carbon (P=0.007).

The estimated proportion of 18 year old subjects with a low FEV₁ (defined as a ratio of observed to expected FEV₁ of less than 80%) was 4.9 times greater at the highest level of exposure to particulate matter with an aerodynamic diameter of less than 2.5 micrometres as at the lowest level of exposure (7.9% v 1.6%; P=0.002). □

Smoggy skies over Los Angeles could adversely affect the development of lung function during the teenage years

Thiomersal doesn't cause developmental disorders

Janice Hopkins Tanne *New York*

Two studies report that vaccines that contain thiomersal do not cause behavioural problems or developmental delay in young children. The studies, published in *Pediatrics*, are the University of Bristol's Avon longitudinal study of parents and children (*Pediatrics* 2004;114:577-83) and a review article by US authors (2004;114:793-804).

The Bristol study, also known as the "children of the 90s" study, is the largest of its kind in the world. It followed nearly 13 000 children from their births in 1991 and 1992 in southwest England. The children were born when the recommended vaccines contained thiomersal and when an accelerated vacci-

nation schedule was introduced.

Thiomersal, known as thimerosal in the United States, is a preservative that was used until recently in many children's vaccines. In the United States it is either not used at all in vaccines or used in very small amounts. Its use in vaccines in the United Kingdom will be discontinued from the end of this month.

Thiomersal contains about 50% ethylmercury, an organic compound that is metabolised into mercury. Concern about thiomersal arose because a related compound, methylmercury, was toxic in several environmental disasters. Low doses of methylmercury seem to have adverse developmental effects on children who are exposed before birth or in the first few months of life.

It was suggested that low doses of ethylmercury used as a preservative in vaccines might have similar detrimental effects, although ethylmercury is more quickly metabolised and excreted than methylmercury.

In the Bristol study, immuni-

sation records showed when the children received the vaccines. Researchers calculated the mercury exposure the children received at 3, 4, and 6 months. The children's cognitive and behavioural development between 6 and 91 months was assessed in standard international tests of social, emotional, hyperactivity, and motor skills.

The researchers found no convincing evidence that early exposure to the additive in vaccines had any deleterious effect on neurological or psychological outcomes.

In the review article researchers from the University of Colorado and the Centers for Disease Control and Prevention reviewed 10 epidemiological and two pharmacokinetic studies of ethylmercury, seeking evidence that the additive might cause autism or related disorders. They reported that the epidemiological studies did not show a link and that the pharmacokinetic studies made the link unlikely. □