Case report

Subungual osteochondroma: Nail sparing excision

Anurag Tiwari^{a,*}, Nidhi Agrawal^b, Tarun Verma^c, Hitesh Lal^d^a Department of Orthopaedics, Maulana Azad Medical College, New Delhi, India^b Department of Anesthesia, Vardhman Mahavir Medical College, New Delhi, India^c Department of Orthopaedics, Maulana Azad Medical College, New Delhi, India^d Department of Orthopaedics, Vardhman Mahavir Medical College, New Delhi, India

ARTICLE INFO

Article history:

Received 7 May 2016

Accepted 14 June 2016

Available online 23 June 2016

Keywords:

Subungual
Osteochondroma
Excision
Ring tourniquet
Nail

ABSTRACT

Subungual osteochondroma is a relatively uncommon benign bone tumor affecting mostly children and young adults, and is a major source of pain and nail deformity. Treatment consists of marginal excision and meticulous wound closure. In this report, we present three cases of subungual osteochondroma arising from the dorsal aspect of distal phalanx of the great toe, which were managed by marginal excision with preservation of nail apparatus under digital block anesthesia with the use of a ring tourniquet. We emphasize on the use of ring tourniquet made by glove's finger and the technique of preserving the nail apparatus to prevent nail deformity. At final follow-up, there is no evidence of recurrence or nail deformity with good functional and cosmetic result.

© 2016 Delhi Orthopedic Association. All rights reserved.

1. Introduction

Subungual osteochondroma is a rare, benign bony tumor that usually involves the phalanges of the toes or fingers, most commonly the distal phalanx of hallux.¹ It was first described by Dupuytren in 1817, as a bony outgrowth of the distal phalanx of the great toe that is now popularly known as “Dupuytren's exostosis”. We report on three cases of subungual osteochondroma. All the cases were originated from dorsal aspect of distal phalanx of the great toe and were excised under digital block anesthesia using ring tourniquet made of glove's finger with careful preservation of the nail apparatus. In two cases, we were able to keep the entire nail plate intact and the soft tissue defect thus created was covered by suturing the nail edge with the skin edge.

2. Case presentation

2.1. Case 1

A 14-year-old female presented with a mass on the tip of left great toe beneath the nail of six months duration. The mass was

painless and slowly progressive in size, to the extent that it has lifted the nail plate causing ugly nail deformity (Fig. 1). On examination, the swelling was approximately 10 mm × 10 mm × 5 mm in size, situated beneath the nail. The mass was nontender, firm in consistency and covered with dry, scaly skin with no restriction of movements of proximal and distal interphalangeal joints. There were no nail changes. Radiographs revealed a bony outgrowth off the dorsum of the distal phalanx of the great toe (Fig. 2). A provisional diagnosis of subungual osteochondroma was made and the mass was excised under digital block using ring tourniquet. The ring tourniquet was made by cutting a sterile glove's finger through its base and encircling this piece of latex around the base of the great toe and tying both the ends of glove finger with a hemostat as shown in Fig. 3. The mass was carefully separated from the nail plate and an osteotome was used to divide the base of the mass from the distal phalanx of the toe (Fig. 4). This technique created soft tissue defect of 1 cm × 1 cm area (Fig. 5), which was closed by suturing the nail edge with the skin (Fig. 6). At latest follow-up, there is no evidence of recurrence or nail deformity.

2.2. Case 2

An otherwise healthy female, 19 years of age, presented with a slowly growing mass under the right great toenail for the past two years. There was no history of trauma. On examination, there was a nontender, fixed, firm nodule on the dorsal aspect of

* Corresponding author at: Room No 8/9, Old Residents Doctors Hostel, MAMC, New Delhi 110002, India. Tel.: +91 9013211215.

E-mail address: dranurag.tiwari@gmail.com (A. Tiwari).

Pre Op

Fig. 1. Pre-operative clinical photograph of hallux of case 1 showing the subungual lesion.

Pre Op Radiograph

Fig. 2. Pre-operative anteroposterior and lateral radiograph of foot of case 1 showing bony outgrowth arising from the dorsal surface of distal phalanx of hallux.

the right great toe, beneath the nail plate. The lesion was about 5 mm × 5 mm in size and was not associated with nail plate deformity or onycholysis. Radiograph revealed a bony exophytic lesion arising from the distal phalanx of the hallux suggestive of subungual osteochondroma. 3-D Computed Tomography (CT) of the involved toe showed the lesion was arising from the dorsal surface of distal phalanx and was quite eccentric in location (Fig. 7). The patient underwent excision with careful preservation of the germinal matrix of nail bed. At latest follow-up, the patient is free of recurrence and nail deformity.

2.3. Case 3

An 18-year-old female presented with 5 months history of swelling beneath the right great toe nail. There was history of insignificant trauma 5 months back. Examination revealed a subungual nodule of size 10 mm × 5 mm. The lesion was non-tender, non-reducible and non-fluctuant on palpation. Radiograph

demonstrated the presence of exostosis arising from the distal phalanx of great toe. Excision was performed in a similar way as performed in cases 1 and 2.

In all the three cases, bloodless field was achieved using ring tourniquet, and adequate anesthesia was achieved with digital block. Histopathological examination of the lesion in all the cases confirmed the diagnosis of osteochondroma.

3. Discussion

Subungual osteochondroma is a relatively rare, benign osteocartilaginous tumor occurring mainly in second and third decades of life. It affects both sexes equally. They are most commonly seen in the distal phalanx of the great toe. However, other toes may be involved. Clinically, it appears as small, pink colored, hard, nodule that projects beyond the free edge of the nail. The overlying nail may become brittle and may be lifted or become detached.

Intra Op

Fig. 3. Intraoperative clinical photograph of case 1 showing the ring tourniquet made of glove's finger tied with hemostat.

Sketch Representation

Fig. 4. Sketch diagram lateral view of the great toe showing the subungual lesion and the incision used for its excision. Dotted line denotes the incision line.

Post Excision

Fig. 5. Clinical photograph of same patient after the excision of the lesion. Note the size of large soft tissue defect.

The exact etiology of subungual osteochondroma is unknown, although trauma, hereditary abnormality, chronic irritation, and long standing infection have been suggested as possible etiologies of subungual exostosis^{2,3}; however, chronic infection seems to be the result rather than the cause of this lesion.⁴

Some consider it to be a reactive metaplasia resulting from microtrauma^{1,5}; however, there is no conclusive evidence in the literature to support a particular pathogenesis.^{6,7}

Subungual osteochondroma is often misdiagnosed because of its rare incidence and similarity to a number of nail, soft tissue, and bony pathologies, like subungual verruca, onychomycosis, fibroma, pyogenic granuloma, keratoacanthoma, glomus tumor, myositis ossificans, squamous cell carcinoma and melanoma.⁸ Diagnosis is made without the need for biopsy through radiographs. Unlike these lesions, subungual osteochondroma contains an osseous component, the cortical and medullary component of which is continuous with the distal phalanx of the toe that is readily visualized on plain radiographs.⁹

Surgical excision is the appropriate management of this condition. Excision is usually performed under digital block anaesthesia. Although multiple specific surgical techniques and postoperative wound closure strategies have been reported,^{10–12,6,13} but no technique has significant evidence to show its superiority, as a result of a lack of comparative analyses. Suga and Mukouda⁶ suggested that if the excision creates a significant nail bed defect, you can either allow the defect to heal spontaneously or augment the wound with artificial skin allowing it to epithelialize. DaCabra et al.,¹⁰ suggested a method of wound management using a vacuum assisted closure (VAC) device to promote faster healing with good results. Complete excision of the lesion is required to prevent recurrence, and meticulous separation from underlying nail bed structures is the most important factor in preventing post-operative nail problems. If the lesion is located more proximally under the nail plate, the nail plate may need to be partially or totally avulsed to facilitate the entire removal of the lesion.

A recurrence rate of 5–11% has been reported. The recurrence rate is a result of incomplete surgical excision. However, no case of malignant transformation has been reported. We reported on three cases of subungual osteochondroma, managed by excision with nail preservation technique.

Post Suturing

Fig. 6. Clinical photograph of patient after the excision. The soft tissue defect was closed by suturing the nail edge with the skin.

3-D CT Scan

Fig. 7. 3-D CT Scanogram of the involved toe of case 2 showing the lesion arising from the dorsal surface of the distal phalanx. Note the eccentric location of the lesion.

Conflicts of interest

The authors have none to declare.

Acknowledgement

We thank Dr. Mihika Dube for making the beautiful illustration.

References

1. Mavrogenis A, Papagelopoulos P, Soucacos P. Skeletal osteochondromas revisited. *Orthopedics*. 2008;31:1018–1028.
2. Storlazzi C, Wozniak A, Panagopoulos I, et al. Rearrangement of the COL12A1 and COL4A5 genes in subungual exostosis: molecular cytogenetic delineation of the tumor specific translocation t(X; 6)(q1314; q22). *Int J Cancer*. 2006;118:1972–1976.
3. Dave S, Carounanidy U, Thappa DM, Jayanthi S. Subungual exostosis of the thumb. *Dermatol Online J*. 2004;10:15.
4. Singh R, Jain M, Goel R, Siwach R, Kalra R, Kaur K. Subungual exostosis of the great toe: a case report and tumor overview. *Foot Ankle Spec*. 2011;4:376–378.
5. Toyoda M, Morohashi M. An ultrastructural study of subungual exostosis, with special reference to intranuclear inclusions. *Med Electron Microsc*. 1998;31:142–146.
6. Suga H, Mukouda M. Subungual exostosis. *Ann Plast Surg*. 2005;55:272–275.
7. Grisafi PJ, Lombardi CM, Sciarrino AL, Rainer GF, Buffone WF. Three select subungual pathologies: subungual exostosis, subungual osteochondroma, and subungual hematoma. *Clin Podiatr Med Surg*. 1989;6:355–364.
8. Nowillo KS, Simpson RL. *Subungual Exostosis of the Finger With Nail Plate Induction*. New York, NY: Hand; 2009.
9. Guarneri C, Guarneri F, Risitano G, et al. Solitary asymptomatic nodule of the great toe. *Int J Dermatol*. 2005;44:245–247.
10. DaCampra MP, Gupta SK, FerrideBarros F. A novel management strategy for subungual exostosis. *BMJ Case Rep*. 2013;2013(August).
11. De Berker D, Langtry J. Treatment of subungual exostoses by elective day case surgery. *Br J Dermatol*. 1999;140:915–918.
12. Lokiec F, Ezra E, Krasin E, Keret D, Wientroub S. A simple and efficient surgical technique for subungual exostosis. *J Pediatr Orthop*. 2001;21:76.
13. Valinsky MS, Hettinger DF, Gennet PM. Subungual exostosis removal utilizing a minimal incision technique. *Curr Podiatr Med*. 1991;40:10–15.