Published in final edited form as:

Angew Chem Int Ed Engl. 2016 September 19; 55(39): 11882–11886. doi:10.1002/anie.201606381.

Novel Path to Aryl(isoquinoline)iodonium(III) Salts and Synthesis of Radiofluorinated Isoquinolines**

Zheliang Yuan^{#a}, Ran Cheng^{#b,c}, Dr. Pinhong Chen^a, Prof. Dr. Guosheng Liu^{*,a}, and Prof. Dr. Steven H. Liang^{*,c}

^a State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic Chemistry, Chinese Academy of Sciences 345 Lingling Road, Shanghai, 200032 (China)

^b School of Pharmaceutical Science and Technology, Tianjin University, Tianjin, 300072, China

^c Gordon Center for Medical Imaging & Division of Nuclear Medicine and Molecular Imaging, Massachusetts General Hospital and Harvard Medical School, 55 Fruit St., Boston, MA (USA)

[#] These authors contributed equally to this work.

Abstract

Iodonium compounds play a pivotal role in ¹⁸F-fluorination of radiopharmaceuticals containing non-activated arenes. However, preparation of these species is limited to oxidation conditions or exchange with organometallics that are prepared from aryl halides. Herein we describe a novel ‘one-pot’ process to assemble aryl(isoquinoline)iodonium salts in 40-94% yields from mesoionic carbene silver complex and Aryl-I-Py₂(OTf)₂. The method is general, practical, and compatible with well-functionalized molecules as well as useful for the preparation of a wide range of ¹⁸F-labeled isoquinolines resulting in up to 92% radiochemical conversion. As proof of concept, a fluorinated isoquinoline alkaloid, ¹⁸F-aspergillitine is prepared in 10% isolated radiochemical yield from the corresponding phenyl(aspergillitine)iodonium salt.

Keywords

Fluorine-18; Ag-mediated; diaryliodonium salts; [¹⁸F]isoquinolines; positron emission tomography

Fluorine-18 (¹⁸F, *t*_{1/2} = 109.7 min) is the most widely used nuclide for positron emission tomography (PET). New methodologies to incorporate [¹⁸F]fluoride into a variety of unprecedented targeted molecular probes has become a major focus of academic and industrial PET research programs.^[1] Attributed to the increasing prevalence of fluorine-containing pharmaceuticals and obstacles to the access of low specific activity electrophilic [¹⁸F]F₂, focused efforts have been made to design novel methods to radiolabel non-activated arenes using nucleophilic [¹⁸F]fluoride ion.^[2] Among these advances include transition metal-mediated reactions,^[3] and metal-free sulfonium^[4] & iodonium-based approaches.^[5] In particular, radiofluorination based on iodoniumylide precursors has demonstrated a wide

* gliu@mail.sioc.ac.cn, liang.steven@mgh.harvard.edu.

Supporting information for this article is given via a link at the end of the document.

range of compatibility in the labeling of well-functionalized molecules and radiopharmaceuticals, for instance, [^{18}F]FPPMP,^[5h] [^{18}F]FPPMP,^[5h] [^{18}F]-5-fluorouracil,^[5i] [^{18}F]-safinamide,^[5m] [^{18}F]-*meta*-fluorobenzylguanidine ([^{18}F]mFBG),^[5m] [^{18}F]fluoro-*meta*-tyrosine ([^{18}F]FMT)^[5m] and [^{18}F]3-fluoro-5-(2-pyridinylethynyl)benzonitrile ([^{18}F]FPEB),^[5j] as well as application of ^{18}F -azido arenes in bioconjugation reactions.^[5k,5l] Diaryliodonium salts have also shown to be useful in the production of PET radiotracers, including [^{18}F]TFB,^[5d] [^{18}F]flumazenil,^[6] [^{18}F]mFBG,^[7] [^{18}F]6-fluorodopamine,^[5g,8] [^{18}F]FPhe,^[5g] [^{18}F]DAA1106,^[5g] and [^{18}F]6-F-DOPA.^[5e,9] Despite the significant role of radiofluorination using iodonium methods in the PET tracer development, the preparation of these hypervalent iodine precursors^[10] necessitates oxidation of iodoarene in acid media^[11] or exchange with organometallic species,^[12] for example stannylated/boronated compounds, which are prepared from “Miyaura borylation” type cross-coupling reactions with the corresponding aryl halides (Scheme 1a). These common procedures have potential risks of contamination, if heavy metal reagents, for example tin, is utilized or show incompatibility with oxidation conditions in the presence of sensitive functionalities, including alkene and nitrogenous compounds. Therefore there is an unmet need for a robust and efficient method to prepare iodonium precursors for radiofluorination to fully realize the translational potential of these methods in the preparation of radiopharmaceuticals for molecular imaging and personalized medicine.

Isoquinoline is often found in biological natural products and utilized as a pharmacophore in drug discovery.^[13] For instance, the natural product berberine shows therapeutic potential in treatment of cancer and diabetes,^[14] and aspergillitine, isolated from *Aspergillus versicolor* in marine sponge, exhibited antibacterial activity against *Bacillus subtilis*.^[15] Fluorine incorporation into isoquinoline has been proven to reduce *in vivo* metabolism,^[16] particularly as a bioisostere of hydrogen to block oxidation at C-4 (the major metabolism), and thus serves as a valuable fluorine-containing component in pharmaceutical design,^[17] for example Glanatec®.^[17d] For the purpose of molecular imaging by PET, ^{18}F -isotopologues of fluoroisoquinoline would provide a unique and non-invasive way to track *in vivo* behavior of labeled drug candidates and their interactions with biological targets. However, there are only limited methods for the synthesis of fluoroisoquinolines with common theme focused on the electrophilic fluorinating agents, including Selectfluor^[18] and *N*-fluorobenzenesulfonylimide (NFSI).^[19] Applying these approaches for radiofluorination would require an electrophilic fluorinating agent, nearly all of which are generated from gaseous [^{18}F]F₂ and only available in a few PET centers worldwide with specialized apparatus, and also associated with serious shortcomings, *i.e.*, unselective fluorination and low specific activities. In order to obviate the need for electrophilic fluorinating reagents during ^{18}F -fluoroisoquinoline preparation, we report a new and efficient process for synthesis of aryl(isoquinoline)iodonium salts by sequential silver-catalyzed amination of alkynes and exchange with hypervalent iodine reagents (Scheme 1b). This approach represents the first example of diaryliodonium salts prepared from mesoionic carbene-metal [(mic)M] complex (**2** or **3**), which eradicates the use of aryl halides in oxidation or metal halide exchange conditions. Using concomitant isoprene emission as the driving force, the ‘one-pot’ cyclization provides high yields for a wide range of

aryl(isoquinoline)iodonium salts and demonstrates their usefulness in the (radio)fluorination of isoquinolines and synthesis of fluorinated natural product ^{18}F -fluoroaspergillitine.

Our mechanistic investigations^[19d] (Section 11 for plausible mechanism in supplementary information, SI) on the Ag-catalyzed amination of alkynes delineated an unusual mesoionic carbene silver [(mic)Ag] complex **3**, which makes the isoquinoline C-4 position accessible after the alkyne cyclization. We found that when silver complex **3** was treated with hypervalent iodine reagent $\text{PhIPy}'_2(\text{OTf})_2$ ($\text{Py}' = p\text{-methoxypyridine}$), an unprecedented I(III) adduct appeared in near quantitative yield, which was subsequently characterized as **4a** by X-ray crystallography (Scheme 2 and Section 14, SI). This serendipitous discovery inspired us to speculate that an apparent ‘umpolung’ would occur at C-4 if nucleophiles, for example fluoride ion, can be introduced to the newly formed aryl(isoquinoline)iodonium(III) species to furnish the synthesis of fluorinated isoquinoline instead of “ ^{18}F ” electrophiles. Therefore, we adopted this one-pot tandem protocol involving Ag-catalyzed amination of alkynes and exchange with I(III) reagent to create a new and efficient approach to prepare diaryliodonium precursors for the synthesis of [^{18}F]fluoroisoquinolines.

Initial efforts were focused on the exploration of a concise and general protocol to assemble diaryliodonium products (Table 1). The imine **1a** was readily synthesized in 90% yield via condensation reactions between the corresponding aldehyde and *tert*-butyl amine. Treatment of substrate **1a** by catalytic amount of AgNO_3 (20 mol%) at 0 °C for 20 minutes, followed by addition of $\text{PhIPy}'_2(\text{OTf})_2$, provided product **4a** in 80% yield. (entry 1). Efforts towards optimization of this one-pot method indicated that addition of Pyox ligand or base did not effectively increase the reaction yields (entries 2-3). Silver salts screening showed that AgNO_3 is most effective (entries 4-6). Increase of AgNO_3 loading (to 1 equiv.) improved the yield to 95% (entries 7-8). When other hypervalent iodine reagents, such as $\text{PhI}(\text{OAc})_2$ and $\text{PhI}(\text{OCOCF}_3)_2$, were employed, the desired product **4a** was also provided, yet in 30% yield using $\text{PhI}(\text{OAc})_2$, and 78% yield using $\text{PhI}(\text{OCOCF}_3)_2$ (entries 9-10). When we switched $\text{PhIPy}'_2(\text{OTf})_2$ to $\text{PhIPy}_2(\text{OTf})_2$, the reaction could avoid contamination with an unknown impurity from AgPy' complex, and improve the yield to 98% by ^{19}F -NMR spectroscopy. As a result, the final product **4a** was obtained in 88% isolated yield (entry 11). Finally, other transition metal catalysts, such as $\text{Cu}(\text{CH}_3\text{CN})_4\text{OTf}$, $\text{Pd}(\text{O}_2\text{CCF}_3)_2$, PtCl_2 and AuCl_3 which are used in the cyclization of alkyl-imine **1a** in the literatures,²⁰ were also investigated. However, none of desired products **4a** was detected. These observations reveal the unique reactivity of (MIC)Ag complex to hypervalent iodine reagent.

The scope and practicality of this tandem alkynyl-imine cyclization and hypervalent iodine exchange was investigated and the results are summarized in Table 2. Alkynyl-imines with a series of substituents (R) on the (hetero)aromatic ring, including electron-withdrawing fluorine, electron-donating methoxy and methyl groups, were compatible with current transformation to afford products **4a-f** in good to excellent yields of 51-90%. For the substituents (R') in the terminal alkynes, various functional groups like ester, cyclopropyl and alkenyl, were tolerated to give desired products **4g-j** in 40-94% yields. Furthermore, substrates with various aryl groups in terminal alkyne showed excellent compatibility in the reaction and provided **4k-q** with 60-89% yields, along with one thienopyridine salt **4r** of

42% yield. It is worth noting that the reaction of **1a** can be readily scaled up to gram-scale to obtain the desired phenyl(isoquinoline)iodonium salt **4a** in 90% yield.

(1)

We next studied the regioselectivity of fluorination reactions on phenyl(isoquinoline) iodonium salts **4** under nonradioactive conditions. The fluorination was carried out with KF (1.1 equiv.) and 18-crown-6 (0.4 equiv.) in DMF at 100 °C for 40 min with or without Cu(OTf)₂ catalyst. In the presence of copper(II) catalyst, fluorination of iodonium salt **4a** predominately occurred in the phenyl group to provide fluorobenzene in 73% yield determined by ¹⁹F-NMR spectroscopy. The desired positional selectivity, *i.e.*, fluorination at the C-4 of isoquinoline was achieved in the absence of metal catalyst and delivered the favorable fluorinated product **5a** in 70% NMR yield (equation 1). These observations were consistent with prior reports,^[5d, 21] and indicated the translation to radiofluorination and control of ¹⁸F-regioselectivity could be achieved from nucleophilic [¹⁸F]fluoride ion under transition metal free conditions. In addition, this method can be used to prepare non-radioactive standard compound in line with radiolabeling, and requires no ad hoc route for ¹⁹F-standard preparation.

A variety of phenyl(isoquinoline)iodonium salts were fluorinated with nucleophilic ¹⁹F or [¹⁸F]fluoride ion under transition metal free protocols (See conditions B in Table 3 and Section 7.3 in SI for optimal radiolabeling conditions). Fluorinated and radiofluorinated isoquinolines were obtained in 54-88% isolated chemical yields and 11-92% radiochemical conversions, respectively, which demonstrated the compatibility with diverse functionalities with alkyl, aryl, alkenyl, halide, cyclopropyl, and ester groups. The scope of this method was further extended to other fused pyridines. The results of fluorination (82%) and radiofluorination (77%) showed that this protocol was equally applicable to the synthesis of thienopyridine (Table 3, **5r** and **6r**). In particular, to verify the efficiency and practical use of this method, two radiochemical purification methods, *i.e.*, solid phase extraction and semi-preparative HPLC, were utilized to isolate and purify [¹⁸F]fluoroisoquinolines. Compounds **6j**, **6m**, **6p**, **6r** and **6a**, **6l** were isolated in 51-65% yields by SPE and 40-48% yields by HPLC, respectively. The specific activity of 3-*n*-butyl-4-[¹⁸F]fluoro-7-fluoroisoquinoline (**6a**) was determined to be 1.53 Ci/μmol and is suitable for the majority of in vivo PET imaging studies, including most low density biological targets.^[23] We then studied the regioselectivity of (radio)fluorination by measuring the ratio of desired fluoroisoquinoline **5a/6a** and byproduct 4-fluorobiphenyl (**8a/9a**) under both non-radioactive and ¹⁸F-labeling conditions. As shown in Equation 2, labeling precursor **7** was prepared from a new hypervalent reagent Ph-PhIPy₂(OTf)₂ in 77% yield. Both fluorination and radiofluorination demonstrated excellent and desired regioselectivity (>20:1) based on the results on iodonium

salt **7**.^[22] In addition to fluoride, other nucleophiles, such as acetate and azide, are also compatible to the nucleophilic substitution of phenyl(isoquinoline)iodonium salts. The related isoquinoline derivatives **10-16** were obtained in good to excellent yields (58-84%, See Section 5 in SI).

(2)

To demonstrate the utility of this method, we synthesized both fluorinated and radiofluorinated marine natural product aspergillitine by the corresponding phenyl(isoquinoline)iodonium salt derived from the tandem cyclization and iodine(III) exchange. The synthesis commenced with triflate **17**,^[15b] followed by Sonogashira coupling with propyne and condensation with *t*-butyl amine to generate alkyne **19** in two steps around 50% yield (Scheme 3). An efficient silver-mediated amination of alkyne **19** followed by rapid exchange with PhIPy₂(OTf)₂ furnished phenyl(aspergillitine) iodonium salt **20** in 80% yield. Fluorination using KF and 18-crown-6 gave fluoroaspergillitine (**21**) in 32% yield and the radiosynthesis of [¹⁸F]fluoroaspergillitine (**22**) was achieved in 26 ± 2% radiochemical conversions with 10 ± 1% isolated radiochemical yields (*n* = 3) by HPLC.

In summary, a novel method for the convenient access to [¹⁸F]isoquinolines has been developed, which is enabled by a sequential process involving a silver-mediated amination of alkynes and fluorination of phenyl(isoquinoline)-iodonium salts. The methodology has proven to apply to a broad scope of substrates and afford the desired isoquinolines and [¹⁸F]isoquinolines in satisfactory to excellent yields. As proof of concept, fluorinated natural product ¹⁸F-fluoroaspergillitine was prepared in 10% isolated radiochemical yield. The conceptual advantages of diaryliodonium salts preparation from mesoionic carbene silver intermediate, followed by iodonium exchange showcased in the simple setup and easy handling procedure, and excellent compatibility with a variety of functional groups, all of which warrant this method useful for further applications in [¹⁸F]fluoroisoquinoline and other related fluorinated heterocyclic aromatic syntheses.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

We are grateful for financial support from the National Basic Research Program of China (No. 973-2015CB856600), and the National Natural Science Foundation of China (No. 21225210, 21472219 and 21532009). R.C. is supported by China Scholarship Council (201506250036). S.H.L. is a recipient of NIH career development award (DA038000).

References

1. a Phelps ME. *Proc Natl Acad Sci.* 2000; 97:9226–9223. [PubMed: 10922074] b Ametamey SM, Honer M, Schubiger PA. *Chem. Rev.* 2008; 108:1501–1516. [PubMed: 18426240] c Miller PW, Long NJ, Vilar R, Gee AD. *Angew. Chem. Int. Ed.* 2008; 47:8998–9033.
2. a Cai L, Lu S, Pike VW. *Eur. J. Org. Chem.* 2008; 2008:2853–2873. b Brooks AF, Topczewski JJ, Ichiishi N, Sanford MS, Scott P. *Chem. Sci.* 2014; 5:4545–4553. [PubMed: 25379166] c Preshlock S, Tredwell M, Gouverneur V. *Chem. Rev.* 2016; 116:719–766. [PubMed: 26751274]
3. a Lee E, Kamlet AS, Powers DC, Neumann CN, Boursalian GB, Furuya T, Choi DC, Hooker JM, Ritter T. *Science.* 2011; 334:639–642. [PubMed: 22053044] b Lee E, Hooker JM, Ritter T. *J. Am. Chem. Soc.* 2012; 134:17456–17458. [PubMed: 23061667] c Tredwell M, Preshlock SM, Taylor NJ, Gruber S, Huiban M, Passchier J, Mercier J, Génicot C, Gouverneur V. *Angew. Chem.* 2014; 53:7751–7755. [PubMed: 24916101] d Mossine AV, Brooks AF, Makaravage KJ, Miller JM, Ichiishi N, Sanford MS, Scott PJ. *Org. Lett.* 2015; 17:5780–5783. [PubMed: 26568457]
4. a Mu L, Fischer CR, Holland JP, Becaud J, Schubiger PA, Schibli R, Ametamey SM, Graham K, Stellfeld T, Dinkelborg LM, Lehmann L. *Eur. J. Org. Chem.* 2012; 2012:889–892. b Sander K, Gendron T, Yiannaki E, Cybulska K, Kalber TL, Lythgoe MF, Arstad E. *Sci Rep.* 2015; 5. Article number: 9941.
5. For a recent review, see Preshlock S, Tredwell M, Gouverneur V. *Chem Rev.* 2016; 116:719–766. [PubMed: 26751274] , in particular, references 263–325 cited therein; For most recent iodonium-based radiofluorination reports (2013–present), see Chun J-H, Pike VW. *Org. Biomol. Chem.* 2013; 11:6300–6306. [PubMed: 23942997] Chun J-H, Telu S, Lu S, Pike VW. *Org. Biomol. Chem.* 2013; 11:5094–5099. [PubMed: 23804017] Richarz R, Krapf P, Zarrad F, Urusova EA, Neumaier B, Zlatopolskiy BD. *Org. Biomol. Chem.* 2014; 12:8094–8099. [PubMed: 25190038] Edwards R, Westwell AD, Daniels S, Wirth T. *Eur. J. Org. Chem.* 2015:625–630. Ichiishi N, Brooks AF, Topczewski JJ, Rodnick ME, Sanford MS, Scott PJH. *Org. Lett.* 2014; 16:3224–3227. [PubMed: 24890658] Zlatopolskiy BZ, Zischler J, Krapf P, Zarrad F, Urusova EA, Kordys E, Endepols H, Neumaier B. *Chem. - Eur. J.* 2015; 21:5972–5979. [PubMed: 25708748] Cardinale J, Ermert J, Humpert S, Coenen HH. *RSC Adv.* 2014; 4:17293–17299. Rotstein BH, Stephenson NA, Vasdev N, Liang SH. *Nat. Commun.* 2014; 5:4365. [PubMed: 25007318] Stephenson NA, Holland JP, Kassenbrock A, Yokell DL, Livni E, Liang SH, Vasdev N. *J. Nucl. Med.* 2015; 56:489–492. [PubMed: 25655630] Jacobson O, Weiss ID, Wang L, Wang Z, Yang X, Dewhurst A, Ma Y, Zhu G, Niu G, Kiesewetter DO, Vasdev N, Liang SH, Chen X. *J. Nucl. Med.* 2015; 56:1780–1785. [PubMed: 26315836] Wang L, Jacobson O, Avdic D, Rotstein BH, Weiss ID, Collier L, Chen X, Vasdev N, Liang SH. *Angew. Chem. Int. Ed.* 2015; 54:12777–12781. Rotstein BH, Wang L, Liu RY, Patteson J, Kwan EE, Vasdev N, Liang SH. *Chem. Sci.* 2016; 7:4407. [PubMed: 27540460]
6. Moon BS, Kil HS, Park JH, Kim JS, Park J, Chi DY, Lee BC, Kim SE. *Org. Biomol. Chem.* 2011; 9:8346–8355. [PubMed: 22057475]
7. Hu B, Vavere AL, Neumann KD, Shulkin BL, DiMagno SG, Snyder SE. *ACS Chem. Neurosci.* 2015; 6:1870–1879. [PubMed: 26313342]
8. Neumann KD, Qin L, Vavere AL, Shen B, Miao Z, Chin FT, Shulkin BL, Snyder SE, DiMagno SG. *J Labelled Comp Radiopharm.* 2016; 59:30–34. [PubMed: 26695865]
9. a Kuik WJ, Kema IP, Brouwers AH, Zijlma R, Neumann KD, Dierckx RA, DiMagno SG, Elsinga PH. *J. Nucl. Med.* 2015; 56:106–112. [PubMed: 25500826] b Qin L, Hu B, Neumann KD, Linstad EJ, McCauley K, Veness J, Kempinger JJ, DiMagno SG. *Eur. J. Org. Chem.* 2015:5919–5924.
10. Merritt EA, Olofsson B. *Angew. Chem. Int. Ed.* 2009; 48:9052–9070.
11. a Shah A, Pike VW, Widdowson DA. *J. Chem. Soc., Perkin Trans.* 1997; 1:2463–2466. b Zhdankin VV, Scheuller MC, Stang PJ. *Tetrahedron Lett.* 1993; 34:6853–6856. c Bielawski M, Olofsson B. *Chem. Commun.* 2007:2521–2523. d Bielawski M, Aili D, Olofsson B. *J. Org. Chem.* 2008; 73:4602–4607. [PubMed: 18505294] e Ye C, Twamley B, Shreeve J. n. M. *Org. Lett.* 2005; 7:3961–3964. [PubMed: 16119942] f Qin L, Hu B, Neumann KD, Linstad EJ, McCauley K, Veness J, Kempinger JJ, DiMagno SG. *European J Org Chem.* 2015:5919–5924.
12. a Carroll MA, Pike VW, Widdowson DA. *Tetrahedron Lett.* 2000; 41:5393–5396. b Stang PJ, Zhdankin VV, Tykwinski R, Zefirov NS. *Tetrahedron Lett.* 1991; 32:7497–7498. c Ochiai M, Toyonari M, Nagaoka T, Chen D-W, Kida M. *Tetrahedron Lett.* 1997; 38:6709–6712. d Koser GF,

- Wettach RH, Smith CS. *J. Org. Chem.* 1980; 45:1543–1544. e Pike VW, Butt F, Shah A, Widdowson DA. *J. Chem. Soc., Perkin Trans.* 1999; 1:245–248.
13. a Bentley, KW. *The Isoquinoline Alkaloids*. Hardwood Academic; Amsterdam: 1998. b Shamma, M. Vol. 25. Academic Press and Verlag Chemie; 2012. c Ibrahim SR, Mohamed GA. *Fitoterapia*. 2015; 106:194–225. [PubMed: 26388556]
14. a Ni WJ, Ding HH, Tang LQ. *Eur. J. Pharmacol.* 2015; 760:103–112. [PubMed: 25912800] b Ortiz LM, Lombardi P, Tillhon M, Scovassi AI. *Molecules*. 2014; 19:12349–12367. [PubMed: 25153862]
15. a Lin W, Brauers G, Ebel R, Wray V, Berg A, Sudarsono, Proksch P. *J. Nat. Prod.* 2003; 66:57–61. [PubMed: 12542346] b Simonetti SO, Larghi EL, Bracca AB, Kaufman TS. *Org. Biomol. Chem.* 2012; 10:4124–4134. [PubMed: 22531888]
16. a LaVoie EJ, Adams EA, Shigematsu A, Hoffman D. *Carcinogenesis*. 1983; 4:1169–1173. [PubMed: 6883639] b Boyd DR, Sharma ND, Dorrity MRJ, Hand MV, McMordie RAS, Malone JF, Porter HP, Dalton H, Chima J, Sheldrake GN. *J. Chem. Soc., Perkin Trans.* 1993; 1:1065–1071.
17. a Yamada, RS.; M.. 2010. Patent 201093789b Zeng, QY.; C. C.; Yao, G.; Wang, X.; Tadesse, S.; Jean, DJS., JR; Reichelt, A.; Liu, Q.; Hong, F-T.; Han, N.; Fotsch, C.; Davis, C.; Bourbeau, MP.; Ashton, KS.; Allen, JG. 2010. WO 201083246c Matsubara, KI.; A.; Oomura, A.; Kawasaki, K.; Yamada, R.; Seto, M. 2009. U.S. Patent 200948223 A200948221d Garnock-Jones KP. *Drugs*. 2014; 74:2211–2215. [PubMed: 25414122]
18. Si C, Myers AG. *Angew. Chem. Int. Ed.* 2011; 50:10409–10413.
19. a Xu T, Liu G. *Org. Lett.* 2012; 14:5416–5419. [PubMed: 23088282] b Xu T, Mu X, Peng H, Liu G. *Angew. Chem. Int. Ed.* 2011; 50:8176–8179. c Liu Q, Wu Y, Chen P, Liu G. *Org. Lett.* 2013; 15:6210–6213. [PubMed: 24261675] d Liu Q, Yuan Z, Wang H.-y. Li Y, Wu Y, Xu T, Leng X, Chen P, Guo Y.-l. Lin Z, Liu G. *ACS Catal.* 2015; 5:6732–6737.
20. a Subbarao KPV, Raveendra Reddy G, Muralikrishna A, Reddy KV. *J. Heterocyclic Chem.* 2014; 51:1045–1050. b Roesch KR, Larock RC. *J. Org. Chem.* 2002; 67:86–94. [PubMed: 11777443] c Malkov AV, Westwater M-M, Gutnov A, Ramirez-Lopez P, Friscourt F, Kadlcikova A, Hodacova J, Rankovic Z, Kotor M, Kocovsky P. *Tetrahedron*. 2008; 64:11335–11348. d Zhang M, Zhang H-J, Ruan W, Wen T-B. *Eur. J. Org. Chem.* 2015:5914–5918.
21. a Ichiishi N, Canty AJ, Yates BF, Sanford MS. *Org. Lett.* 2013; 15:5134–5137. [PubMed: 24063629] b Ichiishi N, Canty AJ, Yates BF, Sanford MS. *Organometallics*. 2014
22. It is possible that an alternative fluorination of aryne pathway, initiated from the elimination of phenyl(isoquinoline)iodonium **4**, could occur; however, such intermediate and related fluorinated isomer was not observed. We also compared the results of (radio)fluorination between iodonium salts **4a** and traditional precursors for nucleophilic aromatic substitution, including brominated (Table 5, **12**) /iodinated (**13**) /nitro (**14**) isoquinolines. While iodonium salt **4a** gave 90 yield in fluorination (**5a**) and 92 yield in radiofluorination (**6a**), respectively, no desired products **5a** and **6a** were observed from traditional precursors (See Section 12 and 13 in SI), indicating the unique fluorination reactivity of iodonium precursor **4**.
23. Lapi SE, Welch MJ. *Nucl. Med. Biol.* 2012; 39:601–608. [PubMed: 22261142]

(a) Traditional methods for the preparation of diaryliodonium salts (*previous work*) (b) New 'one-pot' protocol for diaryliodonium salts (*this work*) **Scheme 1.**

(a) Traditional preparation of diaryliodonium salts and (b) this work.

Scheme 2.

Hypothesis and formation of phenyl(isoquinoline)iodonium salt

[a][b] The same procedures in Table 4. [c] isolated yields by semi-preparative HPLC.

Scheme 3.

Synthesis of non-radioactive and ^{18}F -labeled fluoroaspergillitine

Table 1Preparative conditions for aryl(isoquinoline)iodonium salts.^a

Entry	catalyst (mol%)	I(III) reagent	yield (%) ^b
1	AgNO ₃ (20)	PhlPy' ₂ (OTf) ₂	80
2 ^c	AgNO ₃ (20)	PhlPy' ₂ (OTf) ₂	78
3 ^d	AgNO ₃ (20)	PhlPy' ₂ (OTf) ₂	27
4	AgNO ₂ (20)	PhlPy' ₂ (OTf) ₂	74
5	AgOTf (20)	PnlPy' ₂ (OTf) ₂	79
6	Ag ₂ CO ₃ (10)	PhlPy' ₂ (OTf) ₂	79
7	AgNO ₃ (50)	PhlPy' ₂ (OTf) ₂	86
8	AgNO ₃ (100)	PhlPy' ₂ (OTf) ₂	95
9	AgNO ₃ (100)	Phl(OAc) ₂	30
10	AgNO ₃ (100)	Phl(O ₂ CCF ₃) ₂	78
11	AgNO ₃ (100)	PhlPy ₂ (OTf) ₂	98 (88)

^a Reaction conditions: **1a** (0.1 mmol), I(III) reagent (0.12 mmol), reaction was initiated at 0 °C then r.t. for 3 h.^b ¹⁹F-NMR yields with CF₃-DMA (2,2,2-trifluoro-*N,N*-dimethylacetamide) as internal standard.^c 20 mol% Pyox.^d 1 equiv. of Li₂CO₃. PhlPy'(OTf)₂: Py' = 4-methoxypyridine, Pyox = (*S*)-4-isopropyl-2-(6-methylpyridin-2-yl)-4,5-dihydrooxazole

Table 2

Substrate scope for aryl(isoquinoline)iodonium salts^[a,b]

[a] Reactions in 0.5-1 mmol scale; [b] ¹H NMR yields with 2,2,2-trifluoro-*N,N*-dimethylacetamide as internal standard (isolated yield in parenthesis). [c] Upon completion, the mixture was heated at 50 °C for 4 h to remove *tert*-butyl group.

Table 3

Fluorination and radio fluorination of aryl(isoquinoline)iodonium salts

 5a 70% 6a 92% (48% ^d)	 5b 77% 6b 25%
 5c 65% 6c 72%	 5d 54% 6d 86%
 5e 61% 6e 65%	 5f 67% 6f 87%
 5g 61% 6g 52%	 5h 57% 6h 11%
 5i 78% 6i 17%	 5j 75% 6j 77% (51% ^c)
 5k 82% 6k 86%	 5l 87% 6l 85% (40% ^d)
 5m 88% 6m 76% (57% ^c)	 5n 83% 6n 80%
 5o 85% 6o 75%	
 5p 81% 6p 80% (65% ^c)	 5q 87% 6q 83%
 5r 82% 6r 77% (58% ^c)	

[a] Nonradioactive fluorination conditions (A): KF (1.1 equiv.), 18-crown-6 (0.4 equiv.), DMF, 100 °C, 40 min; [b] Radiofluorination conditions (B): precursor **4a** (7.2 μmol), [¹⁸F]tetraethylammonium fluoride ([¹⁸F]TEAF), DMSO, 100 °C, 10 min. Radiochemical conversions and product identify were determined by radioTLC and radioHPLC, respectively. For a detailed step-by-step radiofluorination procedure, see supplementary information; [c] isolated yield by solid phase extraction; [d] isolated yield by HPLC.