*Interactive case report***A 35 year old woman with diabetic nephropathy who wants a baby: case presentation**

Giorgina Barbara Piccoli, Elisabetta Mezza, Giorgio Grassi, Manuel Burdese, Tullia Todros

Chair of
Nephrology,
Department of
Internal Medicine,
University of Turin,
Corso Dogliotti,
14-10126 Turin,
Italy

Giorgina Barbara
Piccoli
nephrologist

Elisabetta Mezza
nephrologist

Manuel Burdese
nephrologist

Diabetic Care Unit,
Department of
Internal Medicine,
University of Turin
Giorgio Grassi
diabetologist

Department of
Obstetrics and
Gynaecology,
University of Turin
Tullia Todros
associate professor

Correspondence to:
G B Piccoli
gbpiccoli@
hotmail.com

BMJ 2004;329:674

Maria Tomasa (Tommy) is a 35 year old woman with diabetic nephropathy. She has been diabetic since the age of 15 and has developed several complications, including retinopathy (requiring laser therapy) and mild hypertension. She desperately wants a baby despite being advised against it by her diabetologist. Indeed, she changed where she was referred for diabetic care until she found a group of care givers mildly tolerant of a pregnancy.

Five years ago she had a spontaneous abortion in the first trimester, complicated by hypertension and nephrotic proteinuria and was referred to an outpatient unit dedicated to diabetic patients with renal diseases. She had a renal biopsy after the miscarriage, which showed nodular and diffuse nephropathy with initial sclerosis (five glomeruli, two with global ischaemic sclerosis, the others with signs of nodular and diffuse diabetic glomerulopathy).

Despite the presence of renal damage, we decided not to discourage her wish; none of us was enthusiastic about the idea, but the attitude was "let nature decide." She was put on a strict nephrological and diabetological follow up and a low protein diet (0.8 g/kg/day) and was instructed to stop taking angiotensin converting enzyme inhibitors in case of pregnancy. One year later

Patient's renal function

Variable	Value
Serum creatinine ($\mu\text{mol/l}$)	133
Creatinine clearance (ml/min)	57
Proteinuria (g/day)	2.6
Serum albumin (g/l)	38
Total protein (g/l)	68

Questions

1 What would you tell Tommy about the risks of assisted reproduction?

2 How would you search for evidence of the risks and what search strategy would you use?

3 How would you balance the risks against Tommy's determination to have treatment?

Please respond through bmj.com, remembering that Tommy is a real patient and that she and her doctors will read your response

she had a tubal pregnancy and needed salpingectomy. The surgeon suspected obstruction of the residual salpinx.

Over the next two years, Tommy's attempts to become pregnant continued to be unsuccessful, as expected in the case of a mechanical fertility problem. At the age of 35, she asked for assisted fertilisation. The table gives her renal data. Control of her diabetes was good (glycated haemoglobin 6.7-8%), but she needed a further cycle of laser therapy to stabilise her retinopathy.

Competing interests: None declared.

This is the first of a three part case report where we invite readers to take part in considering the diagnosis and management of a case using the rapid response feature on bmj.com. Next week we will report the case progression and in four weeks' time we will report the outcome and summarise the responses.

*A memorable patient***The gambler**

He was a poker player—one of Europe's finest—and he faced death. The stress, the smoke, and the stillness had wrecked his heart: a change in lifestyle was called for. His casino confrères were sanguine. "What makes this guy the best," said one, "is that he knows when to quit."

The truth was more complex. The difference between great poker players and good ones, he told me, is one of kind. Good players are skilled at risk assessment, bluff, and the reading and concealment of telltale clues, or "tells." Great players can do something more: they can make part of their game unpredictable even to themselves. "If you know how you're going to play," he said, "so do others."

He lived alone and had played almost every day for 40 years. Many of his casino friends had died. For some it was the heart that gave out. For others it was the brain. Often, players losing heavily would "hit the tilt button." Their mind, like a pinball

machine pushed too hard, would cease to function. That is part of poker's appeal: every player's hand is a potential gateway to ruin.

He had faced ruin himself, over a glass of water. Opponents had sensed when his hand was weak. Eventually he identified the tell: it was the sips that followed his inspection of the cards. Some players never identify their tells, and there is no one to help identify them. One man's sips can be another man's fortune.

Three weeks after leaving hospital, he had a fatal myocardial infarction. In the clinic, days before, he had told me about his return to poker. The risks, he realised, were high, but taking risks was what his life was about. He mentioned, however, that the return would be gradual and that he would avoid the stressful, big money games. I could not tell if he was bluffing.

David Cummins *consultant haematologist, Harefield Hospital, Middlesex*