UK heart surgeons are among the best in the world

Zosia Kmietowicz *London*

The most thorough investigation ever into individual surgeons' performance at the operating table has found that the United Kingdom's cardiac surgeons are among the best in the world, with more adults surviving surgery despite patients being sicker and older than elsewhere.

All 222 cardiac surgeons who

operate on adults in the United Kingdom met safety limits laid down by the Society of Cardiothoracic Surgeons, with none working at a mortality higher than 6% for coronary bypass surgery and many with much lower rates.

Of almost 25 000 bypass operations done in the United Kingdom each year, the average mortality for 2003 was 1.8%, slightly less than that reported by surgeons in the United States. Professor Bruce Keogh, president elect of the society and author of the report, however, acknowledged that American surgeons deal with a more complex case mix, with many more obese and diabetic patients than UK surgeons, which accounted

for the difference in outcomes.

The report is the fifth collection of outcome measures on adult cardiac surgery since 1996 and is based on the analysis of 210 000 patients who have recently had heart surgery. What started out as a 12 page pamphlet has grown, however, into a 340 page document, mainly due to demands from the public and the profession to measure performance more rigorously after the Bristol children's heart surgery scandal in the late 1990s (*BMJ* 2001;323:181).

Then it took a whistleblower to report on surgeons at the Royal Infirmary who were operating on babies and children despite higher mortality than similar units elsewhere. Now the society has taken the lead and can identify any surgeon whose results are outside the norm and investigate the reasons why, said Professor Keogh.

The society has stopped short of publishing individual surgeon's death rates, however, because it says it cannot yet take account of each patient's personal risk during a procedure. Instead, the society has published the number of bypass operations each surgeon has done in a year, whether surgeons meet the society's safety criteria (they all do), and the

hospital's average survival rate.

Professor Keogh said that presenting results that were not risk adjusted would be grossly unjust and might lead to surgeons wanting to avoid high risk cases to improve their performance.

Currently, more experienced surgeons tend to take on more complex cases, and the types of procedures vary between hospitals. Although some surgeons deal only with elective patients, others take on a large proportion of emergency or urgent cases which carry greater risks.

To do more robust analyses on outcomes for each surgeon, the society needs to collect more complete data sets using a system called EuroSCORE (European system for cardiac operative risk evaluation), which looks at 15 variables for each patient, including age, ejection fraction, blood pressure, diabetes status, and weight. This information is missing for 30% of patients currently undergoing cardiac surgery in the United Kingdom, according to the report. □

Copies of the *Fifth National Adult Cardiac Surgical Database Report* are available for £69 including postage and packing from Dr Peter K H Walton (tel 01491 411288; email publishing@e-dendrite.com).

Food agency launches campaign to reduce salt intake

Lynn Eaton *London*

The Food Standards Agency, the United Kingdom's food safety watchdog, has launched a campaign to encourage people to reduce the amount of salt in their diet.

It recommends that individuals eat no more than 6 g of salt a day, a figure that is based on advice from the Committee on Medical Aspects of Food and Nutrition Policy in 1994 and endorsed last year by the Scientific Advisory Committee on Nutrition in its report *Salt and Health*.

However, the agency says men are eating on average 11 g of salt a day while women eat an average of 8.1 g a day.

A high salt intake can increase blood pressure, says the agency, citing the 2001 annual report of

the chief medical officer of the Department of Health. And it cites figures from the National Institute for Clinical Excellence that the cost to the NHS of prescriptions for reducing high blood pressure is around £840m (\$1510m; €1230m), nearly 15% of the total annual cost of all primary care drugs.

John Krebs, chairman of the agency, said, "We want to encourage everyone to see how they can reduce the salt they eat. By cutting back on adding salt to food at home and when eating out, as well as looking at what's on the labels of processed foods, it's possible for people to make positive health choices."

But he acknowledged that the biggest battle was with processed foods, including bread

and cereals, which are estimated to account for 75% of the population's salt intake.

"The food industry is about two thirds of the way to reaching our target of a 1 g reduction in processed foods by the end of 2005," said Sir John. "However, to reach the ambitious target of 6 g per day by 2010 will require further action by both consumers and industry if we are to reduce the human and health costs of eating too much salt."

Graham MacGregor, professor of cardiovascular medicine at St George's Hospital and a long time campaigner against high salt content in foods, welcomed the move but felt it put too much of the onus on the consumer, rather than putting pressure on the food industry through legislation.

He dismissed the Salt Manufacturers' Association claim that it was solely overeating and drinking too much alcohol that caused high blood pressure, rather than high levels of salt, as "complete rubbish."

The difficulty of achieving a reduction in blood pressure by advising people to reduce their salt intake was highlighted, however, by a systematic review published in the *BMJ* two years ago.

The review, whose lead author was Lee Hooper, a research fellow at the University Dental Hospital of Manchester, concluded: "Intensive interventions, unsuited to primary care or population prevention programmes, provide only small reductions in blood pressure and sodium excretion, and effects on deaths and cardiovascular events are unclear" (*BMJ* 2002;325:628).

Ms Hooper told the *BMJ* that one of the reasons that such interventions had not proved effective was that a high proportion of the salt that people consumed was in processed food. Unless something was done to reduce salt in processed foods, people would find it difficult to cut down, she said. □