years before, it would not have been renewed, and seemingly without disapproval at Somerset House. The case serves to show how needful it is, for many reasons, that a searching inquiry should be made into all the circumstances connected with the disastrous prevalence of yellow fever in our ships of war, of recent years, on the West Indian and other stations. The welfare of the seaman and the interests of the public alike call for it. In the official investigation which, it is understood, has been instituted by the War Office respecting the late calamitous outbreak at Bermuda, when both services sustained such heavy losses, the points we have specially alluded to well deserve the most careful examination.

REVIEW III.

1. *Entozoa: an Introduction to the Study of Helminthology, with reference more particularly to the Internal Parasites of Man.* By J. SPENCER COBBOLD, M.D., F.R.S., Lecturer on Comparative Anatomy at the Middlesex Hospital.—London, 1864. pp. 480.
2. *On Poisoning by Diseased Pork: being an Essay on Trichinosis or Flesh-worm Disease; its Prevention and Cure.* By JULIUS ALTHAUS, M.D., M.R.C.P. Lond., Physician to the Royal Infirmary for Diseases of the Chest.—London, 1864. pp. 34.
3. *On Domestic Animals in Health and Disease.* By JOHN GAMGEE, Principal of the New Veterinary College, Edinburgh.—Edinburgh, 1862-4.
4. *The Edinburgh Veterinary Review and Annals of Comparative Pathology.*—Edinburgh. Vols. I.—VI. 1859-64.

THERE are three classes of persons who are especially concerned in the subject-matter of Dr. Cobbold's book—the zoologist, the political economist, and the physician, with whom we may place the sanitarian. Of course, whatever concerns the two latter classes of inquirers concerns also the general public. Without any further preface, then, we will proceed, taking Dr. Cobbold's excellent work as the basis of our remarks, to discuss the subject of entozoa from the standing-place from which each of these three classes of persons would view it; not following, therefore, the precise order adopted by the author, but one which for our purpose is more convenient. But, first, let us pay a tribute of respect to another class of observers in whose libraries this book must of necessity find a place. We mean the gentlemen who are engaged in the special practice of veterinary medicine. The cow-leech of our younger days, ignorant, quackish, and conceited, is fast being replaced in this country, especially in the estimation of the higher grade of stock-owners, by a race of men some of whom, for scientific culture and general attainments, are not surpassed by the best educated

members of our own profession. The names of the late Mr. Youatt, of Prof. Simonds, and Prof. Gamgee are familiar, not only to graziers and veterinarians, but also to those amongst ourselves whose range of reading is not circumscribed by the strict boundaries of human pathology. To such as these (for we only mention them as samples of an estimable order of scientific men) we, as physicians and sanitarians, and the public at large, owe a debt of gratitude which has hitherto received but little public recognition. It is not for us at the present time to point out in what particulars their researches have tended towards the advancement of the science which it is our business to cultivate; we simply desire, for ourselves, to acknowledge our obligations to the learned veterinarians both of this country and of the Continent, in the expression of which we have placed on our list with Dr. Cobbold's work the title of a recent work of Prof. Gamgee, and of one of the journals which circulate among the members of the veterinary profession, and which, we are happy to say, find a place upon the table of the public libraries to which we are all accustomed to resort.

The first part of Dr. Cobbold's work is devoted to systematic helminthology, and he there gives a general account of the habits, structure, development, affinities, distribution, and classification of the entozoa and their allies. In his second part he enters more fully upon the consideration of those amongst them which take up their abode in the body of man. But, first of all, he defends the practice of those who, like himself, in place of distributing these parasitic animals amongst the several classes of invertebrata into which their structure would severally locate them, group them all together in one class of "Entozoa"—regarding them, in fact, as "a distinct *fauna*, destined to occupy an equally peculiar territory." This he does on the ground of convenience, and also of the remarkable character which they exhibit of migrating at different phases of their existence, not only from one class of animal or one individual of a class to another, but even sometimes to localities which would lead and which have led observers to overlook their relation to the entozoa altogether: "None of the entozoal parasites continue in one stay; all have a tendency to roam; migration is the very soul of their prosperity; change of residence the *sine quâ non* of their existence; whilst a blockade in the interior, prolonged beyond the proper period, terminates only in cretification and death." (p. 4.) It is not always possible to trace distinctly the wanderings which every species undertakes, but where observation fails a surmise may be made from what is known of the habits of allied species.

The following scheme exhibits the classification of entozoa adopted by Dr. Cobbold. We have, as will be seen by reference to his work, united here several tables into one, and have included in ours the several species which take up their residence in the body of man:

CLASS HELMINTHA.

Subclass I.—Sterelmintha.

Orders.	Families.	Genera and species infesting man.
1. Turbellaria	{ 1. Planariidæ. 2. Nemertidæ. 1. Monostomidæ.	<i>Fasciola hepatica</i> , Linnæus. <i>Distoma lanceolatum</i> , Mehlis. „ <i>ophthalmobium</i> , Diesing. „ <i>crassum</i> , Busk. „ <i>heterophyes</i> , Siebold. <i>Bilharzia hæmatobia</i> , Cobbold.
2. Trematoda	{ 2. Distomidæ . . . 3. Tristomidæ. 4. Polystomidæ . 5. Gyrodactylidæ.	{ <i>Tetrastoma renale</i> , Delle Chiaje. <i>Hexathyridium pinguiicola</i> , Treutler. „ <i>renarum</i> , Treutler.

Subclass II.—Cælelmintha.

	1. Anguillulidæ.	
	2. Gordiidæ . . .	{ <i>Dracunculus medinensis</i> , Cobbold. „ <i>Loa</i> , Cobbold.
	3. Oxyuridæ . . .	<i>Oxyuris vermicularis</i> , Bremser. <i>Trichocephalus dispar</i> , Rudolphi.
	4. Filaridæ . . .	<i>Filaria lentis</i> , Diesing. „ <i>trachealis</i> , Cobbold. <i>Trichina spiralis</i> , Owen.
3. Nematoda . .	5. Cheiracanthidæ.	
	6. Ascaridæ . . .	<i>Ascaris lumbricoides</i> , Linnæus. „ <i>mystax</i> , Rudolphi.
	7. Strongylidæ . .	<i>Strongylus bronchialis</i> , Cobbold. <i>Eustrongylus gigas</i> , Diesing. <i>Sclerostoma duodenale</i> , Cobbold.
	8. Cucullanidæ.	

Subclass III.—Anenterelmintha.

4. Acanthocephala	Echinorhynchidæ.	
		<i>Tænia solium</i> , Linnæus. „ <i>mediocanellata</i> , Küchenmeister. „ <i>acanthotrias</i> , Weinland. „ <i>flavopuncta</i> , Weinland. „ <i>nana</i> , Von Siebold. „ <i>elliptica</i> , Batsch. „ <i>marginata</i> , Batsch. „ <i>echinococcus</i> , Von Siebold.
5. Cestoda . .	1. Tæniadæ . . .	
	2. Bothriocephalidæ	<i>Bothriocephalus latus</i> , Bremser. „ <i>cordatus</i> , Leuckart.
	3. Tetrarhynchidæ.	

From this scheme it appears that there are thirty-one distinct animals which, in some phase or other of their existence, infest the body of man, and that these animals are furnished out of three orders of worms and belong to nine families. In our review of this part of the subject we shall confine our attention almost exclusively to these species, families, and orders.

I. The *Trematoda*, or flukes, during the several phases of their development, perform migrations from parasitic to non-parasitic abodes. The family, *Distomidae*, is distinguished by the presence of two suckers, one connected with the mouth, the other being usually on the ventral surface. In the *Distoma conjunctum*, which Dr. Cobbold takes as the type of the family, the mouth communicates through the medium of an oval œsophageal bulb, with two long digestive canals terminating in cæcal ends. There is in the same individual both a male and a female generative system, ovary and testis, each having a separate outlet at the "reproductive papilla," situated near the ventral sucker. There is further an excretory, or "water-vascular system," in the form of two straight tubes, which, passing downwards, coalesce in the median line, where they combine to form a broad sigmoidal duct. The most interesting part about these animals is the method of their development. It has been supposed that the adult animal was self-impregnating, but Dr. Cobbold tells us that he has seen an actual sexual congress between two individuals of the species *D. conjunctum*. The changes undergone by the *Distomidae* from the ovum to the mature sexual animal vary considerably with altering conditions of season, weather, &c., but the following appears to be a sufficiently accurate outline of their mode of development:—*a.* The egg, which may or may not be provided with a lid, first exhibits the usual process of yolk segmentation, and there is then formed within it a finely ciliated embryo, which escapes free either prior or subsequently to its arrival upon or within the bodies of certain mollusks. *b.* The embryo, having become free, develops into a "*sporocyst*" or germ-sac, or into organized "*rediae*," or nurses, provided with head, mouth, œsophageal bulb, and cæcal stomach. Within the creature at this stage are developed a progeny of *cercariæ* or larvæ, each presenting two suckers and a dark forked line indicating the first appearance of the diastomic digestive system. *c.* These cercarial larvæ escape from the *rediae* or *sporocysts*, and are then free, swimming, tailed animals. In this condition they again become parasitic in or upon some mollusk or aquatic insect, where they assume a pupa condition, becoming encysted, and losing their tails. *d.* In this condition, by the accidental swallowing of its "host," the distoma becomes transferred to the intestine of some higher animal, being then provided with cephalic hooks, and once in its final resting-place its reproductive organs become developed, and it assumes the characters of an adult distome. In the *Fasciola hepatica*, or common liver fluke, all the tissues of the body are contractile, but its locomotion is aided by the two suckers, the lower one acting as an anchor. In this animal the

primary digestive tubes are beautifully branched, and contain biliary matters, epithelium scales, and blood corpuscles, on which it feeds, although Dr. Cobbold thinks it probable that it also derives nutriment by absorption through its skin. The generative organs are very complicated, and the male system is said to terminate in an intermittent organ, which is commonly found protruding in the dead fluke. The eggs when placed in water do not, according to Cobbold, acquire an imperfect embryonic development under a period of six weeks. It has, when escaped, a ciliated covering, and, like other animalcules, displays great activity in water, describing gyrations and circles, but when the ciliated covering drops off its mode of progression is "creeping." Little appears known of the higher larval conditions of the fluke, so that its progress to the mature form is matter rather of inference from what happens in other allied species than of observation. The following is Dr. Cobbold's summary of what is known and what is probable as to the phases, migrations, and habits of this animal:

"1. The *Fasciola hepatica*, or sexually mature fluke, is especially prevalent in sheep during the spring of the year, at which time it constantly escapes from the alimentary canal of its *host*, and is thus transferred to open pasture grounds. 2. It has been shown by dissections that the liver of a single sheep may, at any given time, harbour several hundred specimens of the fluke, and it is certain that every mature entozoon will contain many thousands of minute eggs. 3. The escaped flukes do not exhibit powers of locomotion sufficient to prove them capable of an extended migration, but their movements may subserve the purpose of concealing them within the grass or soft soil where they have fallen. Their habit of coiling upon themselves probably facilitates the expulsion of their eggs. 4. The eggs can only escape from the oviduct of the entozoon one at a time, but there is no doubt whatever that prodigiously large numbers of loose ova are expelled the infested sheep in the same manner as the flukes themselves. 5. By the dispersing agencies of winds, rains, insects, feet of cattle, dogs, rabbits, and other animals, and even by man himself, the eggs are carried in various directions, not a few of them ultimately finding their way into pools, ponds, ditches, canals, and running streams. 6. The freed eggs, at the time of their maturity, contain ciliated embryos, capable of active progression when brought in contact with dew on blades of grass, rain drops, pools of water, ponds, and lakes. The prolonged action of moisture without, aided by vigorous movements of the perfected embryo within, serves to loosen the lid-like end of the egg-shell, by the opening of which the animalcule is set free. 7. The ciliated embryo, which is furnished with a solitary X-shaped eye, after a longer or shorter period of activity, loses its ciliated covering, and becomes comparatively inert. It alters its form, and probably soon afterwards gains access to the body of a fresh-water mollusc, or, possibly, into the tissues of a land snail. 8. Once within the viscera or substance of its so-called *intermediate host*, the now-ciliated larva probably becomes transformed into a large sac, and develops new larvæ within its interior. These sac-like larvæ are called 'nurses' or 'sporocysts,' or when rather highly organized, 'redia.' 9. The contained nurse-progeny or higher trematode larvæ, are probably furnished with tails as in other flukes. When fully developed they constitute the well-known *Cercariæ*. 10. The *Cercariæ* have a tendency to migrate from the bodies of their molluscan hosts, and they are quite capable of an independent existence. During these wanderings in the water, they are occasionally brought in contact with the human

body, and in a few instances appear to have succeeded in penetrating the skin. 11. It is not certain whether the Cercariæ are taken into the bodies of quadrupeds when the latter are drinking water or eating solid food, but it is probable that they are passively transferred in either way. It is not unlikely that they are often swallowed while still resident with the bodies of their molluscan hosts. 12. From the digestive organs of sheep or cattle the Cercariæ make their way into the liver, in which new situation they probably part with their tails and become encysted. This constitutes the so-called *pupa* stage. 13. The pupa, thus encysted for many weeks or even months, attains a higher organization, at last becoming converted into the sexually mature *Fasciola hepatica*. It gains access to the liver ducts, then passes into the common biliary outlet, and from thence is transferred into the intestinal canal, being finally expelled its vertebrate host in the manner previously described." (p. 181.)

The *Distoma lanceolatum* is smaller than the *Fasciola hep.*, measuring rather more than $\frac{1}{3}$ inch in length, and about $1\frac{1}{2}$ line in breadth. Its most common habitat is the liver of the ox, but it has also been met with in the sheep, red and fallow deer, rabbit, hare, and cat. In this animal the contents of the ova attain a tolerably high degree of development before the eggs quit the oviduct of the parent; and hence it is not considered improbable that the subsequent larval metamorphoses are accomplished more rapidly than in the *Fasciola*. The free ciliated embryo is globe-shaped, with a central boring spine. The *Distoma ophthalmobium* is suspected by Cobbold and Weinland to be identical with the *Monostoma lentis* of Von Nordmann; and both, Cobbold thinks, may be referable to the species last described, *D. lanceolatum*. *D. Crassum* measures from $1\frac{1}{2}$ to 3 inches, with a breadth of $\frac{5}{8}$ inch, and it is characterized by its uniform and considerable thickness. The most minute of the human trematodes, however, is the *D. heterophyes*, which measures only $\frac{3}{4}$ line in length, and $\frac{1}{4}$ of a line in breadth; it has an oblong, pyriform outline, the body is compressed throughout, and the surface is covered with numerous minute spines. In the *Bilharzia hæmatobia*, the male and female reproductive organs occur in separate individuals. The male is a cylindrical vermiform worm $\frac{1}{2}$ inch, or rather more, in length; while the female is filiform, longer, and much narrower than the male, being about $\frac{4}{5}$ inch long. It is the *D. hæmatobium* of Dr. Bilharz, of Cairo. It infests in the mature condition no animal lower than man and monkeys. Griesinger conjectures that the young of the animal exist in the waters of the Nile, in its fishes, and even in bread, grain, and fruit.

The *Nematoda*, so named from the threadlike form of their body, are usually called *round worms*. They bear, as a whole, some resemblance in external form to the earthworm. Most of them are only parasitic during a portion of their lifetime, although this period of their non-parasitic existence is comparatively short. The family of *Gordiiæ* is characterized by an extremely elongated body, with a mouth and intestinal canal, the latter ending in a cæcum, and not by an anus. At present the female only of the *Dracunculus medinensis* is known. It measures from 1 to 12 feet in length, and about $\frac{1}{10}$ in.

in thickness. Its mode of reproduction, like the Gordiidæ generally, is viviparous, the numerous young, invested by the uterine walls, almost filling up the entire cavity of the body. The head has a rather flatly convex outline, and near its centre four papillæ or rudimentary tentacles project at equal distances from each other, and from the oral aperture. The digestive apparatus consists of a cylindrical tube extending to the concavity of the tail, where no anus is perceptible. Bastian, who has fully described this worm, was unable to discover any vagina or vulva, and as this coincides with the observations of others, it is clear that the young can only escape by the rupture of the maternal body either before or after its decomposition, and escape from the "host." We may again quote from Dr. Cobbold his summary of the life history of this worm :

"1. The guinea worm, as commonly known, is the adult female condition of a nematode parasite, forming a distinct generic type, for which the title of *Dracunculus* ought still to be retained. 2. The female *D.* is parasitic only during the final stages of its life-period, after it has taken up its residence in the subcutaneous and intermuscular cellular tissue of man, dogs, and horses. 3. The adult female reproduces viviparously; and in consequence of an enormous development of the internal reproductive organs, contains myriads of embryos in all stages of development. 4. The embryos, when set free, are capable of sustaining an independent existence; this vitality is so tenacious that, after a more or less complete desiccation, they will revive on the application of moisture. 5. The proper habitat of the free active embryos may be either the fresh water of natural and stagnant pools, the damp mould and mud of low-lying, marshy districts, or especially the soft, ochreous, argillaceous, clayey soil forming the bottom of wells, tanks, and other artificial reservoirs. 6. The male sexually mature *Dracunculus* is at present unknown; in all probability it is very much smaller than the female, non-parasitic, as far as animal bearers are concerned, and a *constant* inhabitant of fresh water, mud, or moist earth. 7. In all likelihood, the male *Dracunculus* bears a close resemblance to *Urobates palustris*, or to some other of the numerous allied microscopic forms of *Filaria*, which abound in the natural and artificial water reservoirs above mentioned. 8. The young *Dracunculi* probably acquire sexual maturity shortly after their escape from the parent; the sexes associating in muddy waters during the monsoon, after which period the males, in all likelihood, rapidly perish, the females being left to watch their opportunity for further development by migration into the human body. 9. The impregnated females do not gain access to their bearers by the mouth or stomach of the latter; on the contrary, there is every reason to believe that they enter the body in a direct manner, probably by penetrating the sudorific ducts of the skin. 10. Within the tissues of the bearer they grow with considerable rapidity, and in about twelve months, more or less, give rise to a formidable entozootic disease (*dracontiasis*) which, after prolonged monsoons, is not unfrequently severely endemic." (p. 387.)

Dr. Cobbold regards the *D. Loa* as a distinct worm from the *Filaria oculi* of Nordmann and others. The term "*Loa*" is the native name of the worm, which has a wide geographical distribution.

The only representative of the family of the *Oxyuride* which we need notice is the *Oxyuris vermicularis*, the common threadworm which is such an annoyance to young children. One of its most remarkable anatomical characters is its possession of a strong œsophageal

bulb, with tooth-like crushing organs; below this is the stomach, which merges into an intestinal canal, the latter terminating in an anus. The most important of its characters is, of course, its mode of development, but of this little is known. Dr. Cobbold, however, asserts in opposition to Küchenmeister, that in fresh specimens of impregnated oxyuridæ he has almost invariably discovered well-formed tadpole-shaped embryos. Nothing definite is known of the migration of the larvæ :

"I have introduced the eggs containing embryos into various animals, but have not yet succeeded in rearing oxyurides. I have also introduced them into the pulpy parenchyma of apples and pears, but I have not been able to satisfy myself that any of the young nematodes, which I subsequently procured by thousands in one or two of the pears, were the result of these experiments." (p. 369.)

Dr. Cobbold says that the genus *Filaria* has hitherto served as a sort of refuge for the destitute. With a view to assist in an improved classification of these nematodes, he says :

"Provisionally I shall include in the above family title [*Filaridæ*] all those nematodes presenting the following characters: body filiform, smooth, or finely striated transversely, furnished with a simple round or triangular mouth, generally surrounded by a variable number of papillæ; head continuous with the body; anus terminal, or nearly so; tail of the male obtuse, bluntly pointed or slightly expanded; penis consisting of a long spiculum, which is often accompanied by a short accessory piece." (p. 68.)

Passing over the structural peculiarities of the three other members of this family, which infest man, we shall limit our notice here to the most important of the four, *Trichina spiralis*—a worm to which public attention has been much directed during the last two years, and the acquaintance with which by medical men in this country has been greatly promoted by the papers published by Dr. Althaus. And here, for the first time, we come upon an animal which gains access to the human frame through the eating of the flesh of animals which are infested with it. The following is Dr. Cobbold's description of the general and specific characters of the worm in the mature state :

"An extremely minute nematode helminthe; the male, in its fully developed and sexually mature condition measuring only $\frac{1}{18}$ th of an inch, whilst the perfectly developed female reaches a length of about $\frac{1}{8}$ "; body rounded and filiform; usually slightly bent upon itself, rather thicker behind than in front, especially in the males; head narrow, finely pointed, unarmed, with a simple central minute oral aperture; posterior extremity of the male furnished with a bi-lobed caudal appendage, the cloacal or anal aperture being situated between these divergent appendages; penis consisting of a single spicula cleft above, so as to assume a V-shaped outline; female shorter than the male, bluntly rounded posteriorly, with the genital outlet placed far forward at about the end of the first fifth of the long diameter of the body; eggs measuring $\frac{1}{1270}$ " from pole to pole; mode of reproduction viviparous." (p. 335.)

In its larval condition it is found in the form of a spirally coiled worm in the interior of small globular, oval or lemon-shaped cysts, which appear as minute specks scarcely visible to the naked eye. Dr.

Cobbold considers with Leuckart that the cysts are not requisite. They are, in fact, an abnormal condition which may be absent. The larvæ may exist in one "bearer" to the number of at least twenty millions. Pigs and men are most commonly infested with these creatures, but they have been found in dogs, cats, certain birds, and the horse, ox, sheep, and other ruminants, as well as being obtained from rabbits, rats, mice, and guinea-pigs, which have been fed with trichinous flesh. When such flesh is eaten, what happens is this: the flesh is digested, and thus the larvæ are set free; from the torpid state in which they lay they awake up and begin to move about, lose their spiral figure, and become stretched; at the same time they grow rapidly, and the generative organs are developed. Larval trichinæ have attained sexual maturity within less than forty-eight hours after introduction into their new "bearer." Most females contain from three hundred to five hundred ova. In six days the female parasites will contain perfectly developed and free embryos in their interior; and when these have acquired their full size they pass out at the vaginal opening. Soon after birth the larval trichinæ begin their wanderings, penetrating directly through the intestinal walls and peritoneum, and then, without entering the blood-vessels, rapidly working their way through the tissues in all directions until they arrive at the several voluntary muscles of their "host." They seldom stop in any non-muscular organ. We have, however, ourselves seen what we believed to be larval trichinæ in thousands in the lungs of a cat, and Cobbold appears to have seen the same. Virchow has seen them in the heart. Having selected their permanent resting-place, they become encysted by alteration of the sarcolemma. The same cyst may contain from one to three trichinæ. After a few months' residence in this situation the cysts become calcified. The vitality of these worms varies with their condition. The adult worms, if their bearer die, do not survive him above six hours; but the larvæ will live for a long time even in flesh which has already undergone putrefaction. In this way, as has been suggested by Davaine, "the *débris* of an animal devoured by carnivora may become fatal to rodents, or a carcass near a marsh or rivulet may communicate the parasites to the ruminants which drink the water, or to pigs." We shall conclude what we have to say upon this part of the subject by the following summary, which Dr. Cobbold derives from Leuckart:

"1. *Trichina spiralis* is the juvenile condition of a little round worm, hitherto unknown, to which, however, the generic names of *Trichina* must remain attached. 2. The sexually mature trichina inhabits the intestinal canal of numerous warm-blooded animals, especially mammalia (also of man), and constantly in great numbers. 3. At the second day after their introduction, the intestinal trichinæ attain their full sexual maturity. 4. The eggs of the female trichina are developed within the uterus of the mother, into minute filaria-like embryos, which from the sixth day are born without their egg-shells. 5. The new-born young soon after commence their wandering. They penetrate the walls of the intestine, and pass directly through the abdominal cavity into the muscles of their bearers, where, if the conditions are otherwise favourable, they are developed into the form hitherto known. 6. The direction in which

they proceed is in the course of the intermuscular connective tissue. 7. The majority of the wandering embryos remain in those sheathed muscular groups which are nearest to the cavity of the body (abdomen and thorax), especially in those which are smaller, and most supplied with connective tissue. 8. The embryos penetrate into the interior of the separate muscular bundles, and here already, after fourteen days, acquire the size and organization of the well-known *Trichina spiralis*. 9. Soon after the intrusion of the parasite, the infested muscular fibre loses its original structure. The fibrillæ collapse into a finely granular substance, whilst the muscular corpuscles change into oval nucleated cells. 10. The infested muscular bundle retains its original sheathing up to the time of the complete development of the young trichinæ, but afterwards its sarcolemma thickens, and begins to shrivel at the extremities. 11. The spot inhabited by the rolled up parasites is converted into a spindle-shaped widening, and within this space, under the thickened sarcolemma, the formation of the well-known lemon-shaped or globular cysts commences by a peripheric hardening and calcification. 12. The migration and development of the embryos also take place after the transportation of impregnated trichinæ into the intestines of a new host. 13. The further development of the muscle trichinæ into sexually mature animals is altogether independent of the formation of the calcareous shell, and occurs as soon as the former have reached their completion. The male and female individuals are already recognisable in their larval state." (p. 342.)

Of the family of Ascaridæ, the *A. lumbricoides* requires no description. We may merely say that it discharges its ova by the vaginal outlet, and that they pass off from the bowels of the host *per anum*. Probably they complete their development in open waters, where it seems that an interval of some six months must elapse before the embryo is formed. It is not known what wanderings it now undergoes, or in what form it gains access to its future host, but it is probably taken in with river or pond water. The *A. mystax* of the cat is also liable to infest the human body. It is a moderate sized nematode, characterized more especially by the presence of conspicuous alaform appendages, one on either side of the head. The male may reach $2\frac{1}{2}$ inches, and the female about 4 inches in length. It is unnecessary that we should enter at all at length into the structure or development of this worm, which probably gains access to the stomach through the same media as the *A. lumbricoides*.

We pass now over the family of *Strongylidæ* to the consideration of the important order of cestode worms, or tape-worms, of the life history of which our knowledge is much more extensive and accurate. The following extract will enable the reader the better to follow the descriptions we are about to give :

"The tapeworm may be looked upon as a creature compounded of a series of partly dissimilar animal forms, closely linked in single file; and the welfare of this peculiarly associated colony is, in a great measure, dependent on its connexion with the so-called head. . . . Every cestode passes through several distinct phases during its life-history. In the ordinary colonial or tapeworm condition it has been termed the *strobila* (Van Beneden.) The separate joints of which the strobile is composed are denominated *proglottides*, or zooids. The anterior segment forms the *head*, and remains barren, those of the neck and front part of the body being sexually immature during the process of strobile formation. The mature proglottides at the caudal end are capable of realising

an independent existence, and the eggs which they contain develop the six-hooked embryos, or *proscolices* (Van Beneden), in their interior. These latter become metamorphosed into *scolices* or nurses, representing the well-known cysticercal state, which, in its sterile or aborted condition, forms the common *hydatid*." (p. 104.)

The family of the *Tæniadæ*, or true tapeworms, have a small distinct head, with four single oval or round suckorial disks, and a rostellum or proboscis which is retractile, and frequently armed with a crown of hooks. The reproductive orifices are mostly at or near the margins of the joints, which are bisexual. The family of *Bothriocephalidæ* have heads furnished with two or four opposed bothria, or fossettes, often armed with hooks. The mature proglottides vary much in size; but, in the more typical forms, the segments are broad and narrow, having their reproductive outlets side by side at or near the centre of each joint. With this introduction, then, we shall proceed to the several cestode worms found in man, chiefly, however, confining our attention to those points in their life history and development, which are of the highest practical importance. Although Dujardin, Eschricht, and Von Siebold had already shown that the cystic worms, hydatids, acephalocysts, &c., were only phases in the development of cestodes, yet the merit of demonstrating this experimentally lies with Küchenmeister. The only point in the ordinary anatomy of the *Tænia solium* to which we shall refer is one to which Dr. Cobbold particularly calls attention—we mean the water-vascular system, which is still commonly regarded as a digestive system. It consists of two main channels, one passing down on either side of the worm, both being connected by a transverse vessel at one end of every joint. The male reproductive organs in each joint consist of testicular sacs, a vas deferens, and penis; and the female organs of an ovarium, branched uterus, seminal reservoirs, and a vaginal canal. The *scolex*, or larval condition of this *tænia* constitutes the *Cysticercus cellulosæ* which is so commonly found as the "measle" in the pig, and which presents in pork the appearance of an elliptical or reniform hydatid, of the average diameter of $\frac{1}{3}$ ". If the measle be dissected, it is found that the vesicular portion merely consists of the bladder-like caudal extremity of the larva, within which the head, neck, and vermiform body lie retracted and coiled up. The development of the *tænia* is thus described by our author:

"After the proglottis (which, it must be borne in mind, is furnished with male and female reproductive organs), has undergone impregnation by contact with another proglottis, there results from this the formation of eggs within it, which eggs, while still in the body of the parent, develop into embryos, the latter still retaining the egg coverings. At this time the proglottis is about to undergo a passive migration; and having detached itself from the strobila, it is soon expelled from the bowel of the host, thus finding its way into some cesspool, or it may be, into the open fields. The proglottides move about for a time, but the growth of the multitudes of embryos within their interior causes the proglottides sooner or later to burst, and the embryos thus become dispersed; some are then conveyed down drains and sewers, others are lodged by the roadsides in ditches and waste places, while great quantities are scat-

tered far and wide by winds or insects in every conceivable direction. Each embryo is furnished with a special boring apparatus, having at its anterior end three pairs of hooks; the entire group of embryos of any single proglottis is consequently called 'the six-hooked brood.' After a while—by accident, as it were—a pig coming in the way of these embryos or of the proglottides is liable to swallow them with matters taken in as food. The embryos, immediately on being transferred to the digestive canal of the pig, escape the egg-shells and bore their way through the living tissues of the animal, and having lodged themselves in the fatty parts of the flesh, they there rest to await their further transformation, or destiny. The animal thus infested becomes measled, its flesh constituting the so-called measled pork. In this situation the embryos drop their hooks or boring apparatus, and become transformed into the *cysticercus cellulosæ*. A portion of this measled meat being eaten by ourselves, either in a raw or imperfectly cooked condition, transfers the cysticercus to our own alimentary canal, in which locality the cysticercus attaches itself to the wall of the human intestine, and having secured a good anchorage, begins to grow at the lower or caudal extremity, producing numerous joints or buds to form the strobila, or tapeworm colony." (p. 220.)

The *T. mediocanellata* may be distinguished from the *T. solium* by the greater size and thickness of the segments, the absence of rostellum and hooks, and the blackish appearance of the head in consequence of much pigment granules around the suckers. It derives its name from the presence of a main transverse channel of the water vascular system running through the free space between the four sucking discs, and sending branches to and around the suckers. The "cysticerci," or "measles," which constitute the larval condition of this worm, are met with in the muscles and internal organs of cattle.

Referring to Dr. Cobbold's book for what is known of the other tæniæ, we proceed to the consideration of the last one on the list, the *T. echinococcus*, the importance of which to us as medical men cannot well be exaggerated. The tapeworm in its mature state is very small, seldom exceeding one-fourth of an inch in length, and developing but four segments. In this, its strobila condition, it is only found in the dog and wolf. Dr. Cobbold agrees with Leuckart in maintaining, in opposition to Küchenmeister, that the several varieties of hydatids to which the human body is subject proceed from only one species of tapeworm. Into this controversy, however, we must not enter, but go on to describe, as nearly as we can in Dr. Cobbold's own words, the development of this remarkable and interesting worm, and especially the formation of the hydatids, which prove so fatal to man. It is indeed difficult to do this without reference to the admirable engravings with which Dr. Cobbold illustrates this and the other subjects in his book. We may start from the time when the six-hooked embryo which is met with as in other tæniæ has bored its way to the liver, or has been carried in the circulating stream to other organs. When arrived at its resting-place it becomes transformed into the acephalocyst, or hydatid. And here it is to be observed, that it differs from the other larval tæniæ in retaining a more or less globular figure, in place of presenting the distinctions of head, neck, and body. Although the mature worm is confined to the dog and wolf, the hydatid is met with, not in man only, but in the ruminants generally, the horse, ass, &c. When

a hydatid is removed from its cyst and dissected, it is found to consist of two membranes, an *ecto-cyst* and an *endo-cyst*.

"The first indication (of what is subsequently to become an echinococcus head, or a group of heads), consists of a slight papilliform elevation at the inner surface of the endocyst. This prominence—which does not hitherto appear to have been recognised in its very earliest stage—after a short while displays in its interior a vacuole-like cavity, the latter being occupied, however, with a clear limpid fluid. Its margin becomes more and more clearly defined, until the cavity is by and by seen to be lined with a distinct cuticular membrane. The papilla increasing in size, becomes at first elongated or oval, eventually, scoliceform, or even, perhaps, a true echinococcus head. The scolex-like development has now—possibly, it may be, not in every case—has now, I repeat, to sacrifice itself by developing in its interior a brood of scolices or echinococcus heads. In other words, it becomes gradually transformed into the so-called brood-capsules of Leuckart and other authors. All observers who have microscopically examined fresh echinococci must have seen these capsules. Portions of the essential vital endocyst here forming the outer wall of the brood capsule now thicken in the same way that the original maternal endocyst had done, and these thickened portions in their turn become true scolices, or in some cases, scoliceoid formations. By a process of inversion—precisely similar to that which has been so often described and figured in the case of *Cœnurus*—the heads are withdrawn, as it were, into the cavity of the brood capsules; and thus the latter display the bird's-nest-like appearance. Before proceeding further, I may here remark that many modifications of this process have been witnessed as regards individual scolices, some of which abort, whilst others, possibly, become new or secondary brood capsules. All those numerous instances where scolices have been seen free in the cavity of the capsules or in the great general cavity of the maternal hydatid, or in the cavities of the (subsequently to be described) daughter and granddaughter hydatids, are simply examples of morbid phenomena. In this condition the scolices are generally quite dead." (p. 264.)

The proliferation of the hydatid, by which "daughter" and "granddaughter" (secondary and tertiary) hydatids become formed in its interior, is held by our author as well as by Naumyn, whom he quotes upon this subject, to be brought about by a differentiation of the bud developments "which, instead of becoming brood-capsules, become daughter and granddaughter vesicles, constantly developing in their interior secondary and tertiary brood capsules and scolices, but sometimes, it would appear, developing neither the one nor the other." (p. 269.)

Dr. Cobbold thus sums up his view of the process:

"With Huxley, I still regard the endocyst as forming solitary scolices. Some of the scolices become differentiated to form brood capsules, a portion of whose individual echinococcus heads may in their turn become secondary brood capsules, whilst others fail to become either scolices or secondary brood capsules. It also accords more with our knowledge of the general plan of development to believe that the daughter and granddaughter hydatids are likewise peculiarly modified scolices; in which case it is obvious that they must commence by a process of budding from the endocyst, as in point of fact Naumyn states they do. It does not seem probable that the secondary and tertiary hydatids should arise from independent granular points within the laminae of the non-vital ectocyst. For the acceptance of my view it is not

even necessary to perceive a thorough scolecoform character in the primary buddings destined to become daughter vesicles; it is sufficient if the original process of differentiation clearly resembles the papilliform commencement of the ordinary scolex head." (p. 270.)

Of the *Bothriocephalus latus*, Dr. Cobbold says:

"At present our knowledge of the history of its development is incomplete; but sufficient, at least, has been accomplished to enable us to set aside the erroneous notions advanced by Vogt and others as to its mode of introduction into the human body."

The 6-hooked embryo is ciliated at first, and swims about like the volvox. After it loses its ciliated covering, the embryo probably takes up its abode in some intermediate host, but what particular kind of host is undetermined, Dr. Cobbold thinks probably some kind of fish of the salmon or trout family.

II. Let us now consider the subject very briefly as respects those animals which we use as food, and in its relation to diseased meat.

1. *Measles in the Pig*.—The extent to which this disease prevails is stated by Mr. Gamgee to be much greater in Ireland than in Britain. The Irish have an adage that "every pig has its measles," and Mr. Gamgee has been informed by a Wiltshire bacon factor that, whereas not one pig reared in England or Scotland is measled, Irish pigs in some years suffer to the extent of six, seven, or eight per cent. It is most prevalent in those parts of Ireland where pigs are reared in small lots by poor people.

"I found," he says, "that measles was very rife in some parts of Cork, in Limerick, Tipperary, and Queen's County. . . . It is certain, however, that those pigs suffer most from measles that live in common with human beings; that are allowed to roam about at will, and to eat human excrement around the cottages in the roadside, &c. A very few people affected with tapeworm discharge joints enough to contaminate an immense number of pigs."

The fact of a pig being measled is no bar to its sale. Measled pigs are brought to market with healthy pigs, and the only difference made is that, if seriously measled, a reduction is made in the price. "Measly bales of ham and bacon are regular articles of trade, and though sold but little in Ireland (where the disease is known), are largely exported for Glasgow and other large towns in England and Scotland."

Such bacon, we are told, is sold at about three shillings per flitch less than sound bacon. Mr. Gamgee says that some retailers order measly flitches specially, and all that Waterford can supply go to London. That pigs reared in large towns should suffer from a variety of parasitic diseases cannot be matter of surprise, from the mode of feeding commonly adopted. Although pigs first introduced into town piggeries often refuse at first the food offered them, yet in time they learn to feed like their brethren. Mr. Gamgee tells us how that is in Edinburgh, and we have reason to believe that similar practices are pursued in London. It is here, at any rate, a common practice to

establish piggeries in neighbourhoods where offal and entrails of sheep and cattle can be procured readily.

"It is a common practice to feed pigs on the flesh of horses and of other diseased animals, or on the entrails of cattle and sheep that cannot be utilised so profitably in any other way. This animal food is supplied, as a rule, without being cooked, and only mixed with the vegetable matter and fæces contained in the alimentary canal of the herbivorous quadrupeds. . . . I have referred to the gross mismanagement of slaughter-houses. Not an insignificant part of that mismanagement in Edinburgh is permitting entrails, diseased organs, and sometimes condemned carcases, to be carted away by farmers, who throw the whole, raw, to their pigs. It is evident that no more certain means could be devised to spread parasitic diseases in such animals, and to render them unfit for human food; and I think it will be admitted that animals should not be fed so that they may merely increase rapidly in bulk, to be slaughtered as human food, when such system of feeding renders them diseased.

"I have made careful inquiries as to the health of carnivorous pigs, and find as follows: That sows fed on flesh give birth to pigs which cannot be reared, but die shortly after being born; that if any young pigs are admitted to the flesh diet they also soon die; that pigs of two or three months old seem to thrive on flesh, though the mortality amongst them is sometimes considerable; that pigs fed on flesh have a peculiarly soft, diffuent fat, emit a strong odour from their bodies, and after death putrify more rapidly than others. I also find that when any pigs die on the establishments above noticed they are devoured by their companions, and frequently boiled for that purpose." (*Loc. cit.*, p. 281.)

Possibly some of those instances in which pork has been found to produce disturbance of the digestive system of those who have eaten it, but nevertheless where no morbid condition has been apparent in the meat, may find an explanation in this unnatural and disgusting method of feeding.

It is unnecessary to enter at any length into the physical appearances presented by measles pork. A few cysticerci may escape observation, but when the flesh is badly measles it presents on section a disgusting spawn-like appearance from the confluence of the cysticerci. "The presence of many cysticerci in the flesh," says Mr. Gamgee, "leads to an open condition of the texture favourable to the imbibition of fluids, and for this reason measles pigs are easily pickled."

2. We have no data at present on which to form even a surmise as to the degree of prevalence of *Trichina disease* amongst our pigs. Probably it is a comparatively rare disease in this country. All that Dr. Cobbold can say about it is contained in the following passages:

"Persons qualified to judge affirm that the trichina appears in pigs of all races, and that since cooking does not always kill the worm, it is necessary, whenever a pig is killed, to examine the meat through a microscope. . . . The notion that particular breeds of swine are more liable to be infested than others is scarcely likely to prove correct, especially when it is considered how many other kinds of domesticated animals are liable to the same parasite-invasion." (p. 344.)

"The signs of the disease in the animal are said to be loss of appetite, a hoarse voice, quietness, and aversion to all kinds of movement or running, the latter function, when attempted, being attended with dragging of the extremities." (p. 245.)

But yet an instance is quoted where a pig showed none of these symptoms, being considered "*all right*;" yet the unfortunate butcher who killed the pig has since died from eating its flesh. Trichinous pigs moderately affected may even look fat and plump.

It is evident, from what has been said, that the detection of trichinous pork can only be accomplished by a microscopical examination. The meat at first sight *may* present no appearances calculated to raise a suspicion of its poisonous character. The mode of conducting such an examination is fully detailed by Dr. Althaus :

"I have already mentioned that the cysts are visible to the naked eye as whitish, round, or ovoid specks, with which the surface of the muscle is sprinkled. If these are touched with a drop of acetic acid, or better still with diluted hydrochloric acid, the lime is dissolved and the white colouring disappears. This experiment is, however, not perfectly reliable, if larger pieces of flesh are examined; for the acid then produces a deposit from the muscular juice, whereby the whole surface becomes indistinct and turbid. It is therefore the best plan to cut off a very small piece of flesh with a fine pair of scissors, to tear this asunder with needles, and to free the cysts as much as possible from the flesh. If this is done with the glass resting on a dark object, the cysts may be clearly distinguished as whitish grains, and the dissolving power of the acids becomes quite apparent. If the spots retain their colour, it is probable that small pieces of fat, nervous fibres, or similar formations, are present. But as pieces of fat may be connected with the cysts, a negative result is not so decisive as a positive one; and it is therefore in doubtful cases always better to use the microscope in order to decide the point.

"It does not matter very much from which muscles the pieces to be examined are taken. Even if there are only a few trichinæ present, they generally exist in all the muscles of the body, excepting the heart. The heart of pigs may therefore be eaten with impunity. Trichinæ are, however, more numerous in the tendinous extremities of the muscles, which is probably due to the circumstance that a large number of trichinæ penetrate as far as possible in the muscular tissue, and only stop their progress if certain impediments (as tendons) offer. . . .

"*Trichinæ which have not yet become encysted can only be recognised by means of the microscope.* A thin layer of flesh should be cut out with a sharp scalpel, spread over a glass plate, and moistened with a drop of water. If it is then covered with a thin sheet of glass, we may distinguish trichinæ, if there are any, with a magnifying power of 50. Their intimate structure, however, can only be recognised with a power of 300." (p. 17.)

3. "*Sturdy*," "*staggers*," "*gid*," "*turnsick*," are synonymes under which the disease produced by the *cœnurus cerebri* is known when it occurs in sheep. It prevails wherever sheep are kept. The hydatid which occasions it is the larval condition of the *tenia cœnurus* of the dog. This has been fully and satisfactorily established not only by the experimental researches of German helminthologists, but also by those of our countryman, Mr. Gamgee. This relation is also shown by the circumstances under which the disease is practically observed to arise. Mr. Gamgee says that it occurs

"more especially in the autumn and winter months. I find, however, that in some districts there is a greater prevalence of sturdy in summer. This occurs when, during the hot months, sheep are kept on unenclosed pastures on

hills where they must constantly be 'herded,' whereas during the winter the flock is transferred to the enclosed fields, and dogs are more or less removed from them. Sturdy will always be found to prevail on farms with open pastures, where flocks constantly need the guardianship of shepherds and dogs; or on enclosed farms where sheep are fed on turnips, confined daily within limited space, *with one or more dogs amongst them.*" (*Loc. cit.*, p. 236.)

To lessen or eradicate "sturdy," therefore, amongst our flocks, attention must evidently be directed to the dogs which go amongst them, and this both in withholding them from the food, through the medium of which the larvæ may be introduced, and also in subjecting the animals from time to time to such treatment as shall ensure the evacuation of any *tæniæ* which they may harbour, confining them at such times within boundaries so that the *fæces* passed may be collected and destroyed by fire.

4. "*Rot*" in sheep and cattle is a disease produced by the presence of "flukes" in the liver of the animals. It is more prevalent in some seasons than in others, and especially in those years which, like 1860, are characterized by excessive and continued rains; in fact, that year will long be recollected by agriculturists for its destructive effects. Mr. Simonds states that he is acquainted with instances where six hundred or seven hundred sheep in a flock were lost from this cause alone. One person he mentions who buys eight hundred Welsh ewes annually, who had not more than forty or fifty which escaped; and he states that in some parishes in Devonshire five-sixths of the sheep perished, or were sold for a few shillings each for slaughtering. In 1862 Mr. Gamgee says that the mortality was in many parts still higher than in 1860, and that he is quite certain that not less than five hundred thousand sheep suffered in that year from "*rot*" in the United Kingdom, reducing them in value two-thirds and more, and leading to a loss of several hundred thousand pounds to the country at large. "Scores of cases," says Dr. Cobbold, "are on record where our English farmers have individually lost three, four, five, six, seven, and even eight hundred sheep in a single season; and many agriculturists have thus become completely ruined."

As respects the causes of this prevalence of the disease it may be said that it occurs most frequently not only in wet seasons, but upon low lands and marshy pastures, and that it develops most readily from the month of June to the month of October. An exception to the influence of marshy ground is met with, however, in the case of those flocks which feed in the salt-water marshes of our eastern shores.

A "rotten" sheep may be readily known. At first he appears to lay on fat; and we have been informed that it has been customary with some graziers to place their sheep intended for the market on damp land with a view to the production of this primary result, selling them to the butcher before the subsequent ill results follow. When the disease, however, becomes fully developed, it is to be known by the ordinary symptoms of marked *spanæmia*; the skin becomes less unctuous than is natural, and is seen to be scaly where uncovered by

hair, the fleece drops off in patches, the vessels of the conjunctiva are seen to be turgid with pale or yellowish-coloured blood, and, later still, may be scarcely capable of recognition; pressure over the loins causes the animal to wince, and gives a yielding sensation very unlike to the firm resistance the hand meets with in a sound sheep; dropsical swellings succeed, with listlessness, almost stupor, and the sheep dies in a state of hectic.

The meat of a "rotten" sheep may be known by the wasted, flabby, watery condition of all the tissues, and the absence of the normal red colour. Of course, its recognition is aided materially by the examination of the liver when the organs have not been removed from the carcass. Mr. Cobbold states also that it is by no means unfrequent to find one, or even several other kinds of entozoa existing with the fasciola in the same sheep, the most common form being that of the larval echinococcus.

As regards the prevention of the disease all that can be said is, that "moisture being essential to the growth and development of the fluke-larvæ, it is clear that sheep cannot be infested so long as they remain on high and dry grounds; and even in low pastures they can scarcely take the disease so long as they are folded, and fed on hay, turnips, and fodder procured from drier situations." The practice of mixing salt with their food seems to have been attended with good results both in a preventive and curative point of view.

III. As our object in the present article is to direct attention to the sanitary rather than the medical aspect of the subject of "entozoa," as they affect ourselves, we shall, with a single exception, confine our final remarks within the limits which such a scheme would appear to prescribe. We will take the parasites which infest man in the order in which Dr. Cobbold treats of them.

Fasciola hepatica, no doubt, occasionally infests man. Dr. Cobbold has faith in fifteen of the recorded instances. Leuckart believes only in five. Professor Partridge has found it in the gall-bladder, and Mr. Fox of Topsham, and Mr. Harris of Liverpool, have found the fluke located the former beneath the skin behind the ear, the latter under the scalp. The cercariæ, in the latter instance, probably gained access by directly penetrating the skin. Professor Simonds states that, on more than one occasion, he has known persons taken seriously ill when engaged in opening many rotten sheep at a time; and one case in which a country butcher was seized in the evening after being thus occupied with choleraic symptoms, which proved fatal in two days. The flesh of rotten beasts is necessarily lean, and hence less likely to be digestible, as well as being wanting in nutritive qualities.

Three instances are on record of the discovery of the *Distoma lanceolatum* in the human body; one of these is especially instructive. It is that of a young girl accustomed to look after sheep fed on a pasture enclosed by woods, and traversed by two water dykes, and also supplied by ten little stagnant pools which harboured numerous amphibia and molluscs. The child often quenched her thirst from the

half putrid water, and probably also partook of the watercresses growing in the ditches. She was emaciated, the abdomen enlarged. After death the gall-bladder was found to contain forty-seven of these distomæ, and eight calculi.

Distoma ophthalmobium has been found in the human eye. How they have come there is a question difficult to solve. Gescheidt found four in the eye of a child five months old, born with lenticular cataract. They were situated between the lens and the capsule, and could be recognised as so many dark spots on the surface of the lens.

Fourteen specimens of *D. crassum* were discovered by Mr. Busk in the duodenum of a Lascar. No other case has occurred to Dr. Cobbold's knowledge. The museums of the Royal College of Surgeons, and of the Middlesex Hospital, contain some of these specimens.

The *D. heterophyes* was discovered in 1851 by Dr. Bilharz, of Cairo, in the small intestine of a boy; and on a second occasion he collected several hundred specimens under very similar circumstances.

The *Bilharzia hæmatobia* is also a parasite prevalent on the borders of the Nile, and has also been found in South Africa and the Mauritius. At the Cape of Good Hope it gives rise to an endemic disease. Its favourite habitat is the portal system of blood-vessels and those of the bladder. Dr. Bilharz believes that half the grown-up people in Egypt are infested with it. The symptoms it gives rise to consist of disturbance of the uro-poietic functions; diarrhœa and hæmaturia occur in the advanced stage of the disease, often associated with anæmia, colic pains and prostration of the vital powers. The discovery of the ova of the worm in the urine and other excreta is conclusive of the nature of the illness. The disease is said to be more virulent in the summer months, probably from the cercæan larvæ being most abundant in the spring.

The *Tetrastoma renale* was found once by Lucarelli in the urine of an old woman with symptoms of calculus. Five specimens were collected, but no more were found after death.

Only one instance, also, is on record of the *Hexathyridium pinguiicola* being found—namely, Treutler's case—he found it in a tumour about the size of a nut, which was connected with the left ovary.

Four instances are recorded of the discovery of the *H. venarum*; in one instance it was found in the blood which flowed from a ruptured anterior tibial vein; in two instances in the sputa brought up in hæmoptysis; and in a fourth instance in some venous blood. Dr. Cobbold regards these instances as authentic.

The *Tænia solium* may infest our bodies both in the adult and in the larva form. With respect to the symptoms referable to the abdomen in the former case we need say nothing, but would especially direct attention to the importance of keeping this parasite in mind when searching out the cause of some of the more severe forms of nervous disturbance. The *Cysticercus cellulosæ* is of comparatively rare occurrence in the human brain; nevertheless, Griesinger alone has collected fifty or sixty such cases, two of which came under his own observation. The symptoms observed have, however, not been

sufficiently characteristic for diagnostic purposes; epilepsy, mental disturbance, signs of cerebral irritation, or torpor, either singly or conjoined, have been those observed. The origin of human tapeworm in the use of imperfectly cooked meat need be no further dwelt upon except to quote from Dr. Cobbold some less obvious modes in which the larvæ may gain access to the digestive canal. In badly meat,ed pork,

"there is danger even in the cutting up of the flesh. The knife employed by the butcher—it is sometimes even incautiously placed in his own mouth—is indiscriminately used to cut up any other meats at hand, and not unfrequently the vesicles are transferred from meat to meat, and from meat to mouth. This is one reason why so many butchers suffer from tapeworm. Cooks also, in like manner, will cut bread, cheese, and other foods with the same knives which they have just previously been employing to slice or chop up pork which may have contained a few or even many cysticerci; and thus, again, the mode of infection is accounted for." (p. 229.)

Dr. Cobbold also warns helminthologists and medical men against the danger which they run, in handling fresh tapeworms, of the eggs becoming concealed under the nails, or in the clothing; they may subsequently drop, or be brushed into our food, and being swallowed, become converted within us into cysticerci.

"In like manner our neighbours, who complacently devour choice salads made from the stores of the market gardener, also run a certain amount of risk not only as regards this entozoon, but, indeed, as respects several others. The appropriate vegetables may be manured with night soil containing myriads of tapeworm eggs, or they may be watered with fluid filth, and other abominations, into which these eggs have been cast by various possible ways. In such cases one or more tapeworm ova may be transferred into our digestive organs, unless the vegetables are carefully cleaned before they appear on the table. In the same way one perceives how fallen fruits, all sorts of edible plants, as well as pond, canal, and even river water, procured from the neighbourhood of human habitations, are liable to harbour the embryos capable of gaining an entrance to our bodies. It thus becomes evident also how one individual suffering from tapeworm may infect a whole neighbourhood, rendering the swine measly, these animals, in their turn, spreading the disease far and wide. Such a person may even prove dangerous to the life of his friends and neighbours, and especially also to members of the same household. A case of this kind has come to my knowledge, and it may be explained as follows: one or more free proglottides, or joints with their eggs, had escaped from the canal either at night, voluntarily on their part, or during the day passively, in some such way, at all events, as secured the liberation of scores, possibly thousands of ova. One of these escaped eggs had somehow or other come in contact with human food, or possibly had been blown into the mouth, or even, perhaps, had been carried on the legs of some fly into the milk or sugar on the table; in some such manner, at least, the egg found its way to the stomach, and from thence it got into the circulation, ultimately making its passage to the brain, where its presence gave rise to convulsions and death. I offer this as a probable explanation in the case of a young woman whose sister suffered from tapeworm; but I need not say that, out of the sixty or seventy deaths recorded from the ascertained presence of this 'worm in the brain,' it is more than likely that a similar train of reasoning would afford a true solution of the so-called mystery as to how these entozoa become our brain guests." (p. 230.)

A similar idea seems to have occurred to Dr. Ogle, and although he adduces no case in which cysticeri have been traced in the human body to such an origin (a most difficult thing to do, by the way), yet his warning cannot be regarded as misplaced. He says:

"In almost all cases there is reason to think that a tænia does not exist very long within the intestine without portions or joints being passed, not only by stool—i.e., with the fæces, but also by their spontaneously working their way through the anus and 'skedaddling' (to use an expressive Scotch phrase) from the person, adhering to the garments or falling on the ground. Now I have known cases in which there is reason to think that these fragments, which soon desiccate, shrivel, and become reduced to a small, brownish-coloured mass, may have been picked up by children, often crawling infants, who are ever ready to pounce upon and put in their mouth every crumb or crumb-like substance which they see, and then have been introduced into the digestive organs. I have no certainty of this, but I think it is a point worthy of consideration and attention, and one to be mentioned to parents in cases where the tænia is known to exist in any members of a family."¹

In the number of the same periodical for April 4th, Dr. Ogle further remarks:

"Will not the ovum of the tænia, under exceptional circumstances, hatch in the human bowel, and the resultant embryo pass into the hydatid, cysticercus, the larval, or scolex form in the human body, as it does in other of the vertebrata?"

Another possibility may be suggested—namely, that of the bearer of a tapeworm infecting himself with cysticeri. The wonder is, that considering the frequency of tapeworm, the cysticercal form of the disease in man is not more commonly observed. That it is observed in occasional instances, however, suggests, as we have said, the possibility of self-infection. Dr. Ogle informs us, that a tænia was once brought to him which had apparently undergone a process of digestion in the intestine. Had this process been sufficiently complete to have liberated the ova, there appears to have been no obstacle then to the infection of the system extensively with the larval disease. We have ourselves had tænia brought us with joints partially softened, but have been accustomed to regard this as the effect of maceration in water.

Some important statistics are quoted by Dr. Cobbold from Herr Wauruch, of Vienna—the more valuable because culled at a time long antecedent to our recent helminthological discoveries—from which it appears that out of 3864 persons treated at an hospital in that city, 206 were affected with tænia, and that the persons most affected were cooks, butchers, and other persons concerned with animal provisions. Professor Gamgee calculates that for every measly pig in the United Kingdom there is at least one human being with tapeworms, but supposing the proportion to be only one for every other measly pig, we should have 17,275 individuals thus affected, or 1 in every 2000 people. In confirmation of this, Dr. Haldane states that he has found a tapeworm three if not four times in the bodies of 2000 persons which he has dissected.

The *Tænia mediocanellata* is believed by Dr. Cobbold to equal in prevalence the *T. solium*. That we obtain the larvæ of this

¹ British Med. Journal, March 14th, 1863, p. 267.

worm from imperfectly-cooked veal and beef may be accepted as established.

"Drs. Barclay, W. T. Gairdner, and many others in this country, have published cases where their patients were disgustingly addicted to the habit of eating raw flesh; but of all the striking facts which have come to light on this score, none will bear comparison with those recorded by Kaschin. This professional observer, speaking of the Burätes, or Cossacks of the Baikal region, says: 'In Irkutsk, the Burätes stationed there as Cossacks, and who, in part, have been away from their homes for many years, are, almost all of them, infested with tapeworms; some individuals harbouring as many as fifteen specimens at a time. In 130 post-mortem examinations, only two bodies were found entirely free from these parasites, and among 500 other persons treated in hospital, the existence of these entozoa was demonstrated in all. Certainly, the presence of these parasites cannot cause surprise, when the habits of their bearers are duly considered. The Burätes are herdsmen, and feed almost exclusively upon flesh, especially that of calves, sheep, camels, and horses; they neither clean the meat of these animals properly, nor cook it completely; and, moreover, they eat from tables which immediately before have served for the purpose of cutting up the flesh. The tables are just as little familiarized with the application of water as are the dishes and the people themselves; in other words, they are never cleaned. Fat, liver, and kidneys are eaten quite raw, and diseased animals are as much relished as half-rotten carcases. Moreover, these Burätes are so voracious that any two of them can demolish a one-year old lamb at a single meal.' (!) Kaschin supposes these tapeworms to be referable to *T. solium*, but in this, as Leuckart points out, he is evidently mistaken, for not only are the ruminants the principal source of their food, but they comparatively rarely partake of pork." (p. 241.)

The following is Dr. Brinton's reply to a question put him by the Chairman of the Special Committee on the Cattle Importation, &c. Bill:

"Do you believe that the habit of consuming raw meat exists to any extent amongst the population?—I do. Tapeworm is not uncommon, and, speaking roughly, I should say that 60 per cent. of the cases of tapeworm I have seen own to eating raw meat, and I suspect that of the remaining 40 per cent. a good many do it and do not own to it." (p. 58.)

As regards the frequency of this worm in our own country, Dr. Cobbold states that half the worms at the Middlesex Hospital belong to this species, and quotes Dr. Aitken, to the effect that amongst the soldiers at Fort Pitt it is more abundant than the *T. solium*. He believes that while the latter chiefly infests the poor who feed much on pork, the *T. mediocanellata* is met with mostly among the rich.

The *Tænia acanthrotrias*, *T. flavopuncta*, *T. nana*, *T. elliptica*, and *T. marginata*, are of very rare occurrence; respecting the last also, it may be stated that only one authentic and indisputable instance of its larva, the *Cysticercus tenuicollis*, occurring in the human body, is on record. We may, therefore, pass on to a much more important tænia, the larva of which is far more extensively distributed—the *T. echinococcus*.

Referring to Dr. Cobbold's book for the relative frequency with which *hydatids* infest particular organs, we shall confine ourselves here to the statistics of the disease, and the precautions which science appears to indicate as the only method of diminishing its frequency. The echinococcus disease is endemic in Iceland, where for every 100 inhabitants there are 1100 head of horned cattle, and every peasant

has on an average six dogs. It is estimated that, at any one time, there are in the island 10,000 persons affected with echinococci, who are mostly treated by quacks, with whom such "remedies" as dogs' urine and fresh dogs' excrement play a conspicuous part. The disease is much more common in the interior than on the coast, and among women than among men; the latter circumstance being probably due to their more constantly drinking the water about their dwellings to which the dogs have access, and to their more implicit obedience to the dictates of the "quacks." Dr. Cobbold attributes the rarity of the disease amongst sailors to their being debarred from drinking water from localities where dogs abound, and its comparative rarity among our upper classes to the custom of drinking beer, where the poorer classes are under the necessity of drinking water. With respect to the advice to be offered as to the checking of the prevalence of these larval parasites he quotes with commendation the following remarks by Leuckart:

"In order to escape the danger of infection the dog must be watched, not only within the house, but whilst he is outside of it. He must not be allowed to visit either slaughter-houses or knackeries, and care must be taken that neither offal nor hydatids found in such places are accessible to him. In this matter the sanitary inspector has many important functions to perform. The carelessness with which these offals have hitherto been disposed of, or even purposely given to the dog, must no longer be permitted, if the welfare of the digestive organs of mankind are (*sic*) to be considered." (p. 286.)

The precautions to be adopted in short are to ensure the destruction of all echinococcus vesicles, and the expulsion and annihilation of the *tænia* from the dog by timely physicking, as suggested by Dr. Leared, and burning of the excreta. Dr. Cobbold adds:

"I again invite attention to the importance of observing this rule: *All the entozoa which are not preserved for scientific investigation or experiment should be thoroughly destroyed by fire, when practicable, and under no circumstances whatever should they be thrown aside as harmless refuse.* In the case of the *Tænia echinococcus*, the greatest difficulty likely to be experienced lies in the fact of the extreme smallness of this tapeworm. As an additional security, therefore, I would recommend that boiling-hot water be occasionally thrown over the floors of all kennels where dogs are kept, for in this way not only the escaped tapeworms, but also the little free embryos themselves, would be effectually destroyed." (p. 288.)

The *Bothriocephalus latus* is most frequent in West Switzerland and the adjoining French districts, and is common in the north-western and northern provinces of Russia, Sweden, and Poland. It appears to be most prevalent where the *T. solium* is rare. Dr. Cobbold expresses his strong persuasion that people infect themselves with this parasite by eating imperfectly cooked fish. The *B. cordatus* has only been found once in the human subject.

Warned by the length to which this article has already extended, we must pass over the other members of the Trematode group, to concentrate attention upon three of the most important of them.

The first of these is the *Trichina spiralis*. Although when we review the past literature of the profession, and although when each of us who has practised for many years looks back to his own experi-

ence and regards it by the light of modern investigation, we must hesitate to pronounce *trichinosis* to be a new disease, yet it is one of so recent recognition as to demand at our hands something more than a few cursory remarks. Its history, since the *trichina spiralis* was regarded as a mere curiosity of the dissecting-room, only dates back for a period of five years, since it was in January, 1860, that Zencker found the larval trichinæ as the cause of death in a young servant girl treated in the Dresden Hospital, and who exhibited symptoms similar to those which we shall presently describe. In that year Dr. Zencker recognised an epidemic of the disease in Dresden and its neighbourhood, and showed the existence of trichinæ in a ham and in pork sausages which the affected persons had eaten. Since then, numerous instances of the disease have been recognised in Germany, enough indeed almost to create a pork-panic; and especially in the kingdom of Saxony, and the Prussian provinces of Saxony, as in Magdeburg, Leipzig, Eisleben, &c. In the spring of 1862 thirty cases occurred at Plauen, in Saxony; and altogether there appear to have occurred in the place 80 cases, of which one was fatal. In Magdeburg, during 1858-62, 300 fell ill with the disease, and two died. In Calbe, near Magdeburg, during the summer of 1862, out of 38 persons attacked eight died; and in Blankenburg, out of 60 persons two died. From the end of October to the middle of December, 1863, about 158 persons were affected with the disease in Hettstädt, near Eisleben, and of these twenty-eight died. It can scarcely be imagined that the cases of disease recorded as happening in these places are all that have occurred. But what it is of chief moment to dwell upon is, that the mode in which the outbreaks of trichinosis take place is often calculated to raise a suspicion of foul play on the part of persons engaged in the preparation of the food partaken of. An instance or two of this may not be out of place. They will be found in Dr. Althaus's pamphlet. One such instance is described by Dr. Tüngel as occurring on board a merchant vessel trading between Valparaiso and Hamburg:

"On leaving the former place the cook of the ship bought a pig, which was slaughtered, and part of which (about 30 pounds) was eaten by the crew; the rest of the animal was salted. Most of the crew fell ill, and two of them died. In a boy who died, and where an autopsy was made, numerous live trichinæ were found in the muscles. A piece of the salted pork was sent for examination to Professor Virchow of Berlin, who discovered in it a number of trichinæ, all of which were dead." (p. 10.)

Another:

"In 1863 Professor Langenbeck, of Berlin, excised a tumour which had grown on the neck of a man. During the operation he noticed that the muscles which were laid bare contained a number of encysted trichinæ. The patient being questioned whether he had not at one time or another been affected with a remarkable illness, related the following history:—In 1845, a committee of eight gentlemen, being engaged in the inspection of schools in Saxony, dined together at an inn, and partook, amongst other dishes, of ham and sausages. They all, with the exception of one, who merely drank a glass of claret, fell ill, and four died. Suspicion fell upon the meal and the host. The wine from which they had drunk was analysed; and although no poison was found the host continued to be suspected, and was at last obliged to emigrate." (p. 7.)

The symptoms produced in man by trichina infection will depend for severity upon the number of parasites swallowed in the poisonous meal. Dr. Althaus distinguishes three stages in the distemper. The *first* lasts from four to eight days—i.e., until the first of the progeny are born. In this stage the symptoms are not distinctive, *malaise*, symptoms of indigestion, vomiting, and diarrhœa. The *second* lasts from the commencement of migration until the trichinæ have fully established themselves by becoming encysted in the muscles. Briefly, the symptoms in this, the most important stage, are—fever, sometimes high, the dryness of skin being succeeded by profuse perspirations, œdema of the face, with inflamed conjunctivæ, photophobia, and pain in moving the eyes, especially on looking upwards; scanty, high-coloured urine, diarrhœa, and tenderness of the abdomen; rigidity, swelling, and pain in the moving of the muscles, first of the neck and back, and proceeding to the arms and thighs, and next the forearms and legs; dyspnœa and hiccough arise in consequence of invasion of the respiratory muscles, and hoarseness and loss of voice in consequence of invasion of the laryngeal muscles; when the muscles of mastication and deglutition are invaded, these processes as well as that of speaking becomes impossible, and the patient lies on his back with his extremities slightly flexed, unable to move or speak. At last, with increase of the fever, there occur fainting fits and delirium, meteorism, hæmoptysis, lobular pneumonia, and pleural effusion, and finally death by exhaustion. A fatal issue may take place four or five days from the commencement of the illness, but it usually occurs during the fourth or fifth week. In other cases, however, recovery sets in about this period. This is the *third* stage, the symptoms all gradually lessening in severity, but stiffness and weakness of the muscles may remain for some time, and baldness and desquamation of the skin may succeed. In some insidious cases the characteristic symptoms are so slight that the patients are not even laid up, yet death may ensue in three or four weeks by pneumonia or peritonitis. Children, from their tendency to get rid of offending matters by vomiting and diarrhœa, are less apt to suffer than adults. The only absolute certainty attainable in the diagnosis of trichinosis lies in the discovery of the trichinæ in the evacuations or in the muscular tissues. To obtain a portion of the latter for microscopic examination, Middeldorpf's harpoon may be used, or an incision may be made near the lower insertion of the deltoid muscle, and a minute piece of the flesh snipped off with scissors.

For full information as to the treatment of this affection we must refer our readers to Dr. Althaus' pamphlet; suffice it for us to say that an early emetic and calomel purge are not only rationally indicated, but seem practically to be efficacious in dislodging some at least of the trichinæ introduced in a meal, and the subsequent administration of sweet oil has been suggested. A desideratum is a substance which shall act as a poison to the trichinæ and not to man. Benzine has been suggested, and although poisonous both to trichinæ and man, yet has been given by Dr. Mosler in doses sufficient to destroy the trichinæ while within the intestine without damaging the person who has taken it. The grand object, however, to be attained is the pre-

vention of the disease. *Thorough* cooking must kill the trichinæ in pork; but how rarely is pork, especially in the form of sausages, thoroughly cooked! More particularly, how prone are the poor to eat this and other meats more or less underdone! *Thorough* salting, again, seems to kill the trichinæ; but then it is to be kept in mind that many persons object to bacon and ham that have been thoroughly salted, and instances of infection by these preserved meats have been recorded, probably because bacon and ham are not saleable at a high price if too salt. It is a question of importance whether the State should not place itself between the pig-feeder and the public consumer, and demand from the former a guarantee that the meat supplied is not of a nature to originate disease.

The only other worm which we shall notice here is the guinea-worm, the *Dracunculus medinensis*, and we shall say very little about it. The "time-honoured" method of extracting the worm is as satisfactory as it is ancient; but now, perhaps, above all, the rule should be observed of not throwing away the worm to become the source of infection to others, but of consuming it entirely by fire. Prophylactically speaking, also, persons who travel in the districts infested by the larvæ of this worm should carefully protect the feet and legs during the rainy season, and after washing or bathing should take the precaution of drying their bodies most thoroughly by means (if possible) of warm or hot towels.

We must now conclude our notice of Dr. Cobbold's book; but before doing so we must express the obligation we are under not only to Dr. Cobbold, but to his publisher and printer, for presenting the profession with so really handsome a volume. People who regard books only from a strictly utilitarian point of view, may be content with the miserable blotting-paper and flimsy covering of the works which issue from the continental press; but good printing on good paper, attractive and artistic coloured illustrations, and an elegant and substantial binding, go far, in our observation, to popularize any subject, and especially such a subject as Dr. Cobbold discusses, in itself to some minds a somewhat repulsive one. In this instance also we can commend all concerned in the preparation of the work, inasmuch as the contents of the book are eminently worthy of the extrinsic attractions which have been thrown around it. The sixty pages of bibliography which complete the book are alone worth the purchase-money to anyone who would make himself familiar with the literature of the entozoa.

REVIEW IV.

1. *On the Two Forms or Dimorphic Condition in the species of Primula, and on their Sexual Relations.* By CHARLES DARWIN, M.A., F.R.S. ('Journal of Linnean Society,' vol. vi., Botany. 1862. p. 77.)
2. *On the Existence of Two Forms, and on their reciprocal Sexual Relation, in several species of the genus Linum.* By CHARLES DARWIN. ('Journ. of Linnean Soc., Botany, vol. vii. 1864. p. 94.)