ART. XII.

Principles of Physiology, General and Comparative. By WILLIAM B. CARPENTER, M.D. F.R.S. F.G.S., Examiner in Physiology and Comparative Anatomy in the University of London, &c. &c. Third Edition.—*London*, 1851. 8vo, pp. 1098. With 321 Wood-Engravings.

THE general character of the treatise, of which the goodly volume before us constitutes a new and greatly-enlarged edition, is sufficiently well known, we presume, to our readers, to have rendered it unnecessary for us to have occupied much space in directing their attention to it, had it been *merely* a new and enlarged edition. But it comes before us in some degree with the claim of a new work; as will be apparent from the following extract from the author's preface.

"The second edition of his '*Principles of General and Comparative Physiology*' having been for some time out of print, the author commenced, about two years ago, the task of preparing for the re-appearance of that work; in the expectation that, although much addition and modification would be requisite, in order to embody in it the most important of those discoveries in which the Physiological researches of recent times have been so fertile, yet that it would be found possible to interweave these with the original matter, in such a mode as to preserve the unity of the whole. On proceeding with his task, however, he soon perceived that it would be impossible to do it the justice he desired, without putting aside his former work altogether, and entirely reconstructing the treatise; only employing the materials of the original, where they might chance to be appropriate to his purpose. And thus it happens that the present work may be considered as essentially a new one; since, out of the 1080 pages of which it consists, not above 150, or less than one-seventh, belong to the previous edition. The author hopes that this fact may be admitted as evidence of his desire to spare no pains, in order to render his treatise a faithful exposition of the present state of physiological science; and that it will be deemed a sufficient justification of the slight change, by which he considers that the title has been made to represent more appropriately the present character of the work." (p. vii.)

We feel it right, therefore, to give our readers some opportunity of judging for themselves as to its scope and character.

The author further informs us, in his Preface, that "notwithstanding this almost complete renewal of the *materials* of his Treatise, he has found no important modification to be requisite in its *plan*;" for, in fact, he had framed his physiological system as if with an anticipation of subsequent discoveries. The reader of the first edition, published in 1839, will be at no loss to discern that the leading idea of the work was that of reducing every form of organic structure, and every functional change, to its simplest expression; and that the lowest vegetable organism, consisting of a single cell, was continually recurred-to as the starting point from which to trace out the essential character of each physiological action, through its successive complications in the higher forms of vegetable and animal life. In no other treatise, that we are aware of, had this been even hinted at; and we have high authority for saying, that the introduction of this method of study, and the elevation of the comparative physiology of Plants to the rank of an integral portion of general physiology, made this publication the commencement of a new era in the science. Certain it is, that much of what has since been done to extend

and consolidate it, has been in the path thus opened up; so that, again to use the author's language, "the 'cell-doctrine' of Schwann and Schleiden, perfected and extended by the labours of the numerous observers, who have subsequently prosecuted the same line of research, has fallen (so to speak) into its appropriate place, and now serves as a complete confirmation of the principles which the author had previously propounded." (Preface.)

It is obvious from the list given by the author in his Preface, of the doctrines which he regards as more particularly his own, that what he has himself done for the advancement of Physiological Science has been chiefly in the way of reasoning upon the facts collected by others. Such, for example, was his application of the doctrines of Reflex action to the Nervous System of Invertebrated Animals, contained in his Prize Thesis for 1839; the validity of which is now universally recognised, we believe, by Physiologists, although it is not always attributed to its original author. Such, also, is his application of the "Correlation" idea to the Vital Forces, and to their relation with the Physical, of which we gave a sketch in our last number. Several other doctrines, first advanced by him (some of which we shall presently notice), are, in like manner, based upon the observations of others; but are not the less new in themselves, or less likely to take their due rank in Physiological Science. One important original investigation, however, has been entirely worked out by himself, namely, the inquiry into the Organic Structure of the Shells of the Mollusca, Echinodermata, and Crustacea, which was published in the Reports of the British Association for 1844 and 1847.

The Treatise is divided into Two Books, of nearly equal extent; the first being the "General," and the second the "Special and Comparative" Physiology. The following are the titles of the Chapters included in the First Book:—I. On the Nature and Objects of the Science of Physiology. II. Of the General Characters of Organised Structures. III. Of the Nature and Conditions of Vital Phenomena. IV. Of the Component Structures of Organised Fabrics. V. Of the Distinctive Characteristics of the Vegetable and Animal Kingdoms. VI. General View of the Vegetable Kingdom. VII. General View of the Animal Kingdom. VIII. On the General Plan of Organic Structure and Development.

We do not intend to give any regular analyses of these chapters; but shall make a few selections from some of them, that may serve to show the mode in which their subjects are treated. As the first two chapters do not afford us any passages capable of such isolation, and as we have already brought under the notice of our readers the views which our author has recently enunciated respecting the "Nature and Conditions of Vital Phenomena," we shall pass on to the fourth, from which we shall, in the first place, extract a statement, that will show how much closer is the alliance between the Plant and the Animal, in regard to the essential composition of their primary tissues, than has been until recently supposed. In describing the vegetable cell, the author remarks:

"Although we have hitherto spoken of the *cell-wall* as a simple membrane, yet it is now well known to be made up, in most if not in all instances, of two layers of very different composition and properties. The inner of these layers, which has received the name of 'primordial utricle,' appears to be the one first formed, and most essential to the existence of the cell; it is extremely thin and delicate, so

that it escapes attention so long as it remains in contact with the external layer, and is only brought into view when circumstances occasion its separation from this; it seems to consist of an azotised compound, probably of an albuminous nature; and it appears to participate actively in the vital operations of the cell. The external layer, on the other hand, though commonly regarded as the proper 'cell-wall,' seems to be generated on the external surface of the primordial utricle, after the latter has completely enclosed the cavity and its contents, so that it cannot be regarded as essential to the cell; it is usually thick and strong in comparison with the other, but it may possess various degrees of consolidation, from mere mucus to a firm and tenacious substance; it is composed of cellulose, a substance nearly identical with starch; and it does not appear to take any active share in the vital operations of the cell, its principal office being to protect, locate, and isolate the matter it contains. This external layer may consist of many laminae, the results of successive deposits from the surface of the primordial utricle; but it still usually remains readily permeable to fluid, although no pores can be distinguished in it, even under the highest magnifying power." (pp. 88-9.)

Hence it appears that the *real* cell-wall of the Plant has the same albuminous composition as that of the Animal; and that the envelope of cellulose by which it is generally invested has no more to do with the cell itself, than has the envelope of chondrine which encloses and connects the cartilage-cells of the latter, or than that of horny matter which constitutes the firm intercellular substance that gives tenacity to the otherwise delicate cellular residue left after the decalcification of certain shelly structures.

On the minute structure of the Vegetable and Animal tissues, we do not find any new observations of importance; and it is obvious that the limits imposed upon the author have prevented him from entering into many details that find their appropriate place in treatises of less comprehensiveness. We would direct the attention of our readers, however, to some interesting observations, too long for extract, which present themselves under the head of "Transformation of Tissues;" wherein a parallel is sketched out between the histological metamorphosis observed in ascending the scale of organisation, and that which is witnessed in the successive stages of embryonic development, this parallel being, for various reasons, most complete in the vegetable kingdom. We here find a doctrine explicitly stated, with regard to the real character of the lower tribes of Infusorial Animalcules, which has been hinted at by many preceding writers, but which has not been, so far as we know, so clearly expressed before:

"Many of the simpler forms of (so called) Animalcules must be probably regarded as nothing else than simple cells, which are endowed by their ciliary appendages with the power of free locomotion, but which contain no variety of organs, and multiply themselves (like other independent cells) by spontaneous fission. Now such beings cannot be properly compared to any of the higher Animals, in the fully-developed state of the latter; but they do present a very close correspondence to their earliest embryonic condition. For the life of the highest commences in a single cell, in which not the least vestige of its future organs or tissues can be traced; and this cell speedily multiplies itself by a process of binary subdivision, until a homogeneous mass of cells is produced, bearing a very close resemblance to some of those composite clusters of Animalcules, which are produced in like manner by the repeated bi-partition of a single individual." (p. 177.)

The parallel was first drawn, we believe, by Dr. Barry; but, among those who have since noticed it, these Animalcules have been still

described as possessing sensibility, and as executing their movements under the guidance of an intelligent will, or, at any rate, of a consciously-exerted instinct. We are satisfied that such an assumption is altogether illogical; and that, so long as it is the fashion to interpret the actions of the simplest organisms by those of the most complex, instead of using the former to enable us rightly to comprehend the latter, we shall have no sound physiology. This part of the subject is more fully discussed, however, in the ensuing chapters, in which the author applies himself to the determination of the distinctive characters of the Animal and Vegetable kingdoms; and, as the question is one of much interest, and has recently been the subject of much debate, we shall take this as a sample of the mode in which the higher departments of general physiology are discussed in the present work. We may premise, however, that some naturalists of our time assert that no real distinction between the two kingdoms exists in nature; others, that there must be an intermediate kingdom for the reception of a group of organisms that strictly belongs to neither; whilst there are others who yet sustain the antiquated notion, that there are beings which are Vegetables in one part of their lives, and Animals in the other.

Dr. Carpenter first inquires into our ideas of a *typical* Plant and a *typical* Animal; and he comes to the conclusion that the existence of consciousness and the power of spontaneous motion must be regarded as the chief attributes, which must be superadded in our "idea" of an Animal to those which, being involved also in our "idea" of a Plant, are therefore the characteristics of organised beings in general; so that, if we always possessed the means of determining where consciousness and spontaneity *do* and *do not* exist, we should have comparatively little difficulty in drawing a definite line of demarcation between the two. "But," he continues,—

"We have no other means of judging of the presence or absence of these psychical endowments, than by studying the *movements* of the beings whose character we seek to determine; and it is frequently impossible to say, with anything like certainty, from what source these movements proceed. This difficulty is increased by the fact, that the most remarkable movements exhibited by undoubted Plants are seen in a group whose lowest forms present the nearest approximation to the simplest types of the Animal kingdom; and hence it has happened that beings have been invested with the attributes of Animality, whose subsequent history has clearly indicated their Vegetable nature. On the other hand, there are beings to which, from the absence of sensible movements, we should be disposed to refuse the endowment of consciousness; although the general characters of their structure would lead us to suppose their place to be rather in the Animal than in the Vegetable Kingdom. Such beings, however, present a strong resemblance to the embryonic or transitory forms of higher types of organisation; and as the (presumed) absence of consciousness in the early embryo does not lead us to exclude it from the Animal Kingdom, so it should not warrant the exclusion of a class of beings, the *tendency* of whose development is the same with that of undoubted animals, although the higher stages of that development are never attained. Here, as elsewhere, it happens that in the separation and circumscription of really 'natural' groups, we are obliged to leave a margin around each, for the reception of forms to which the definitions that we frame to express their most characteristic features are not applicable; and in the assignment of such 'aberrant' forms to one or the other of these groups, we must be rather guided by their general resemblance to those which display its typical cha-

racters, than by the degree in which they may themselves possess them. Thus, when we meet with beings possessing the general structure of Plants, and corresponding with them in the mode of their development, we are justified in referring them to the Vegetable Kingdom, although their movements may bear a strong resemblance to those of Animals. On the other hand, when we encounter beings possessing the general structure of Animals, and corresponding with them in the mode of their development, we seem justified in referring them to the Animal Kingdom, even if we never witness any such movements in them as can be fairly considered indicative of consciousness." (pp. 182-3.)

According to this view, then, although the presence of consciousness is the most important feature of the Animal kingdom, taken as a whole, yet it is not that on which the most satisfactory diagnosis can be founded; and we must look elsewhere for characters of more ready applicability. These may be found, in the Author's opinion, in the chemical composition of the doubtful fabric, and in the mode in which it acquires its food; provided that due discrimination be employed.

"If its cell-walls and fibres be found to consist solely of albuminous or gelatinous matter, the presumption is strongly in favour of its Animal nature. If, on the other hand, the tissues should be chiefly composed of cellulose, the presumption is equally strong in favour of its Vegetable character. In the application of such a test, however, great discrimination is required; for, as already pointed out, albuminous matter is not merely stored up in the cell-cavities of Plants, but even forms the most essential part of their cell-walls; whilst, on the other hand, it is quite certain that cellulose (contrary to what was formerly maintained) may be an integral part of the bodies of certain animals. A distinction is to be drawn between the cases in which, as in the *Corallines* and *Nullipores*, it forms part of the proper cell-walls; and those in which it is merely deposited within or around the cells, as happens in certain Compound Tunicata whose food consists largely of starchy matters, the cellulose appearing to be stored up in certain parts of their fabric, just as fatty matter is in other instances. Thus, although the mere presence of starchy substances in the fabric is not of itself a sufficient proof of the Vegetable nature of a doubtful being, it may be held to be so when there is no distinct proof to the contrary, and when the starch is found as an integral part of the solid fabric, and is generated by the being itself, instead of being a mere deposit derived from external sources.—This leads us to consider, in the *second* place, the inferences derivable from the mode in which the nutritive materials are taken in. For if the aliment of a being whose other characters leave its place doubtful, be *formed* by it at the expense of carbonic acid, water, and ammonia,—as is indicated especially by its decomposition of carbonic acid, and its liberation of oxygen under the influence of light,—there is a strong presumption of its Vegetable nature: whilst, on the other hand, if it be dependent for its support upon organic compounds previously formed by other beings, and especially if these be received into internal cavities for the purpose of digestion, the presumption is equally strong in favour of its Animal character. Here, too, discrimination is needed; for Plants may seem to live at the expense of matters previously organised (as is the case with the whole group of Fungi), when they really only thrive by the decay of these, taking in the carbonic acid and ammonia which are thus set free; whilst, on the other hand, there are Animals which appear to subsist on inorganic matters, such as sand or earth, although they are really supported by the small quantity of organic substances which these may include." (p. 191.)

Now it is quite true that many *reputed* Animalcules possess the power of decomposing carbonic acid under the influence of light; but it is also certain that the tendency of microscopic research has been recently to refer these, on other grounds, to the vegetable kingdom. The most active movement, effected by the agency of cilia, cannot be admitted as any evidence of animality, since we know that the 'zoospores' of many undoubted

Algæ are equally active, their propulsion being effected by the very same instruments; and when no other proof than this can be given that a certain organism is an 'animalcule,' it seems to us obvious that agreement with the vegetable kingdom in a fundamental condition of its organic existence, ought at once to determine its true place to be on that side of the boundary.—It is curious to observe how strongly the current of opinion is now taking this direction. The authority of Ehrenberg, who withdrew from the domain of Botany a very large number of the doubtful forms, to find a place among his "Infusionsthierchen," for a long time bore down every other; so great was the weight attached to his character as a most industrious and accurate observer, and as a sagacious and discriminating reasoner. But gradually that authority has been undermined by the labours of the vast number of observers who have devoted themselves to this field of inquiry; and it now becomes apparent that the whole work has to be done over again, in order to complete what Ehrenberg has left undone, to correct much of what he was supposed to have done, and thus to afford data for satisfactory generalisation on points which he is found to have determined under the influence of strong prejudice, and, not unfrequently, in the very teeth of evidence. The consequence has already been, the transfer of several important groups (including most of the so-called, "fossil-animalcules") back to the Vegetable kingdom, from which he removed them; and the probable allocation in the same division of a large proportion of those so called Polygastrica, which are distinguished by the greenness of their colour, notwithstanding that some of these have hitherto ranked among the most indubitable "Animalcules." What will our readers say, for example, if the *volvox globator*, so familiar to every microscopist, should prove to be a plant? Yet such may now be considered as all but demonstrated. We find the following notice on this subject in a subsequent chapter of Dr. Carpenter's book.

"It is not by any means certain that the *Volvocine* are not to be transferred to the Vegetable Kingdom. Their green colour leads to the suspicion that they decompose carbonic acid; and the stomachs described by Ehrenberg in the component Monads, are not more distinct than the stomachs which he has represented as existing in several other beings, whose Vegetable nature is now generally admitted. It is considered by Braun, who has paid much attention to the development of the inferior Algæ, that the *Volvocine* are of the same type with certain 'Zoospores,' which become composite by fissiparous multiplication." (p. 251.)

We have recently seen a series of preparations by Mr. Williamson of Manchester, who has paid much attention to the development of the *Volvox*, which leave not the slightest doubt in our minds that the entire hollow sphere is originally composed of cells, formed by the multiplication of a single one; and that these cells are of the same essential character with those of the inferior Algæ generally; it being a part of their nature to secrete a great thickness of pellucid mucilaginous matter around the primordial utricle, and thus the green utricles, which are originally in close approximation, become separated from each other by its intervention, still remaining connected, however, by certain thread-like prolongations, which Ehrenberg has erroneously described as vessels.

The chapters which contain the general view of the Vegetable and Animal Kingdoms occupy a far larger proportion of the Treatise than they did, in the previous editions, together making, in fact, not less than 365 pages. Had he been able to refer to any concise outline that suited

his purpose, the author would probably have forborne to enlarge his work by the allotment of so much space to what must be considered rather as Natural History than as Physiology; but to those who desire to become acquainted with those types of organic structure, which are of special interest to the Physiologist, but which are insufficiently or not at all described in elementary treatises, without going through the vast mass of (to him) useless matter contained in the larger systematic works on Botany, Zoology, and Comparative Anatomy, these outline views, filled-in where most requisite for the purpose intended, seem almost indispensable. Novelty is not here aimed at by the author; a faithful sketch being all that he aims at presenting. And we shall therefore be extremely brief in our notice of this portion of his work.

We find the Animal kingdom subdivided into *five*, not *four*, sub-kingdoms; the fifth being termed *Protozoa*,—a designation which has been of late much employed by German writers, but which has been very variously applied. Dr. Carpenter limits them to those in which the very simplest type of form and structure presents itself, but which must yet be regarded as deserving a place in the animal kingdom, on account of their chemical composition, and their mode of obtaining their aliment, and also on account of their resemblance to the embryonic condition of higher animals; and he thus gives them a corresponding rank with those equally simple vegetable organisms, to which the name *Protophyta* has been given. The best idea of the characters of this group may be derived, he considers, from an examination of the *Amœba*, commonly known as the ‘Proteus,’ on account of the unlimited variety of forms which it may assume, but superseded in that designation by the well-known perennibranchiate Batrachian. This creature is a not uncommon inhabitant of fresh waters and stagnant infusions; and the following account of it, given by the author in great part from his own observations, contains so many points of interest, that we shall place it in full before our readers:

“At first sight, this creature seems to be nothing else than a minute particle of gelatinous matter; but its spontaneous changes of shape soon give evidence of its possession of vitality; and it becomes evident, on further examination, that it may be considered as a *simple cell*, of unusual dimensions (being sometimes, in the largest species, as much as 1-70th of an inch in diameter), analogous to that which constitutes each individual of the simplest Protophytes. Like these, it lives for and by itself, and it imbibes its nutriment through its cell-wall, having no oral orifice; but it possesses a motor power with which they are not endowed; and this power is obviously related to the character of the food on which it is to be supported. For, whilst *they* can obtain, wherever they may be situated, the materials required for their development, this Animal cell, being dependent upon matter previously assimilated, and not endowed with any means of drawing such matter towards itself, must go (as it were) in search of it. This movement is accomplished by the continual changes of form that take place in its body, the typical figure of which may be considered globular, but which may assume almost any shape whatever. The change of form seems due rather to actions taking place in the interior of the cell, than to any irritability of the cell-wall; for if the movements of an *Amœba* be attentively watched, the extension of the gelatinous body in any particular direction (so as to form one of the digitate prolongations) will be seen to be preceded by the setting of a current of the moving molecules within the cell, in that direction; to which current the protrusion of the cell-wall is really due. A continuation of the same current distends the prolongation, and the whole mass of the body is gradually carried onwards (so to speak) into it, so that its place in the field

of the microscope is slowly changed. After a short time, the particles in the interior of the cell are again seen to move in a definite course; a new prolongation is thrown out, in the same or some other direction; and the body is again absorbed into it. Thus its locomotion is effected by the agency of the currents which traverse the interior of the cell, and which are obviously connected (as in the cells of *Chara* and other plants) with the organic processes concerned in its growth and maintenance; and it must consequently be regarded as entirely independent of any spontaneous control on the part of the individual, and cannot even be allowed to afford the least indication of consciousness. The nutrition of the *Amœba* is derived from animal and vegetable substances; but these are not received into an internal cavity (as affirmed by Ehrenberg), their enclosure within the gelatinous body being rather apparent than real. When the creature, in the course of its progress, meets with a particle capable of affording it nutriment, its body spreads itself over or around this, so as even to invest it completely; and the particle thus seems to be included in a digestive cavity, while it is really in contact with the external surface only, through which its nutritive elements are absorbed into the cavity of the proteus-cell. It is interesting to see such a creature thus manifesting the peculiar *nisus* of animal development; *making*, as it were, a stomach for itself, by wrapping its body round the alimentary matter, which it is not able to receive into its interior." (pp. 244-5.)

This account closely corresponds with that which Mr. Carter has given of the *sponge-proteus*, from his observation of the fresh-water sponges of Bombay. The fact that the change of form of the cell appears to be consequent upon the direction of currents in its interior, is one of great physiological interest, in reference to the production of the stellate, caudate, and other prolongations of cells, which are so frequent in the pigmentary, nervous, and other tissues.—Further, we have in this animal, as Dr. Carpenter remarks, just such a manifestation of animal *tendencies*, rather than of animal *powers*, as might be expected in beings that constitute the transition-group between the two kingdoms; and these tendencies are still further carried out in the three higher groups, which he comprehends under the general designation Protozoa,—namely, the true *Animalcules*, the *Rhizopods*, (corresponding with the group commonly designated as *Foraminifera*,) and the *Sponges*.

"Thus in the (so-called) *Polygastric Animalcules*, we seem to find the highest development which animal cells can attain as individuals; their nourishment being directly received into their interior by an orifice in the cell-wall; and the movements of the cilia with which they are beset, being adapted not merely for the propulsion of the body through the water, but also for the introduction of food through this orifice. But this is a type of structure, which does not seem capable of any further development; for, although we meet with forms among these *Animalcules*, which seem like sketches of fabrics much more elevated in the scale, we do not meet with any which present a decided approximation towards higher groups. So in the *Rhizopods*, we have composite fabrics, presenting a considerable regularity of form, which seem to have their origin in the gemmiparous multiplication of beings, closely allied to the *Amœba* in their general structure and mode of life, each individual still living for and by itself. In the *Sponges* and their allies, on the other hand, we meet with vast aggregations of proteiform cells, united into composite fabrics; of which the several parts, though almost precisely similar to each other, have yet a certain degree of interdependence; and through these we are led towards the true *Zoophytes*, in which the individuality of the component cells and fibres is merged (as it were) in that of those composite structures which are termed Polypes." (pp. 245-6.)

The section on *Entozoa* contains several facts and doctrines which will

probably be new to most of our readers ; we shall pass it over, however, on the present occasion, as we intend to take an early opportunity of reviewing in detail some of the more interesting amongst recent contributions to our knowledge on this subject, which differ in this from what had been previously brought together,—viz. that the history of their *development* is now made the object of most attentive research, and, when fully made out, has not merely removed many difficulties and cleared up many obscurities, but has revealed a new order of facts, of the most remarkable character.

A large proportion of this chapter is occupied with a general account of the organisation of the different classes of vertebrated animals ; and in this is included an outline of the “Vertebral Theory” of Professor Owen, a description and figure of whose “Archetype Skeleton” is given in the introductory account of the vertebrata generally ; whilst under the several heads of Fishes, Reptiles, and Birds, the chief modifications which this archetype undergo are pointed out. In treating of the cranium of the Mammalia, however, a different plan is followed ; each bone being analysed, so to speak, and reduced to its proper elements in the vertebral system, which are generally found to be marked in the embryonic condition of the bone, by so many distinct centres of ossification. The whole of this exposition, as we are informed in the Preface, has had the advantage of Professor Owen’s revision ; and, consequently, it may be recommended with confidence to those who desire to gain a general acquaintance with the system, as presenting an authorised outline of it.

In the last chapter of the First Book, “On the General Plan of Organic Structure and Development,” we find an exposition of the much-canvassed doctrine of *Unity of Composition*, in a form which, we believe, is perfectly consistent with the known facts of anatomical science, and which, even in its present imperfect development, enables these to be far more readily apprehended than they could otherwise be. In assigning to any particular being its place in the organised creation, our author remarks, we always proceed *from the general to the special* ; for each group, in a perfect classification, should have its “archetype” or ideal model ; and in proportion as we pass from the larger to the smaller groups, from those most comprehensive to those most restricted, should we find the general characters of the archetype presenting modifications of successively increasing degrees of speciality, until we arrive at those distinctions which are characteristic of the most restricted groups. All scientific classification, in fact, aims at the determination of such archetypes ; and it is in helping us to the knowledge of what are the most general or fundamental characters of groups, and what are the more special and (so to speak) accidental, that the study of development seems likely to exert its most important influence upon zoological science. That the real history of the development of the higher organisms is *not* expressed by saying that they pass through forms or conditions which are permanent in the lower, was shown by Dr. Carpenter in his former edition. The true law is here, too, that of progress *from the general to the special*, as first propounded by Von Bär, and more fully set forth by Dr. Martin Barry ; and this is fully elucidated by the general remarks on the subject contained in this chapter, and by the more detailed account of the evolution of the several organs and systems given under various heads in the Special Physiology.

The doctrine of the 'progressive development' of the organised creation peopling this earth, during successive geological epochs, has been, under one form or another, pretty generally held, from the time when a scientific knowledge of Palæontology began to prevail, until a comparatively recent period; for however few there may have been, who gave credence to the hypothesis of the actual transmutation of the lower species of one epoch into the higher species of another, yet still the fact of a *succession* of organic life is indisputable, as is also that of the *general tendency* towards a more elevated set of types as we come down in geological time from the remoter epochs towards the more recent. But when this general doctrine is carried into detail, it is immediately met by such a vast number of difficulties, that many palæontologists have of late altogether refused to acknowledge its validity; and the fact of the succession has of late stood by itself, no law or general expression of the phenomena having been proposed. By Prof. Agassiz and others, indeed, it has been maintained that the general plan of development of terrestrial organic life corresponds with that of the embryonic development of the higher organisms; but they have not been guided by a correct notion of what that plan is. Dr. Carpenter suggests that the law of progress from the more general to the more special, will here also be found applicable; for, as he remarks, "many indications present themselves that the types of each principal group first introduced were *not* the lowest, but that they presented *in combination* those characters which are found to be *separately* distributed, and more distinctly manifested, among groups that subsequently made their appearance." As an example of this law, he specially cites the very peculiar distribution *in time* of the several orders of the class Echinodermata. This class seems to have been first represented, during the Palæozoic epoch, by an order that combined some of the most remarkable characters of all the orders which have since shown themselves; and this order, *Cystidea*, must be regarded as superior in general organisation to the Crinoidea, notwithstanding that the greatest multiplication of the latter occurred at a later period. Moreover, such of the other orders as showed themselves during the Palæozoic period, exhibited a decided approximation to the cystidean type; and it was not until the secondary period, when the *Cystidea* disappeared altogether, that the most characteristic types of Crinoidea, Asterozoa, Ophiurida, and Echinida presented themselves; these orders dividing among them (so to speak) the characters possessed in combination by the *Cystidea*, and carrying them out separately as the distinctive peculiarities of their respective types.—In the extraordinary multiplication of the Tetrabranchiate or Nautiloid Cephalopods, too, during the Palæozoic period, our author finds another example of the same general plan; for these seem to have occupied the place of two other groups which came into consequence as it dwindled away, namely, the carnivorous Gasteropods and the higher Cephalopods.—The class of Fishes presents many indications of the same kind; and it serves to indicate the difference between the view of Prof. Agassiz and that of Dr. Carpenter, that the former continually represents the 'homocercal' tail, of which the earliest fishes were almost invariably possessed, as being an embryonic character, because it is found at an early period of the development of existing fishes, whose tail subsequently becomes 'heterocercal;' whilst by Dr. Carpenter, the homocercal tail is considered as the

“most general” character in the class of Fishes, because it is common to them all in some stage of their development; whilst the heterocercal is peculiar to certain sections of the fishes of later epochs.—It is in the Reptilian class, however, that we find the most curious exemplifications of this principle; for there is every reason to believe that this class, at one time, occupied the place, and performed the duties, of the whole series of air-breathing Vertebrates, including groups which represented Fishes, Birds, and Mammals respectively, and thus having a *more general* character than the class at present exhibits. These groups, which can in no degree be regarded as paralleled by any embryonic conditions of existing reptiles, subsequently gave place to those *more special* forms, which carry out most exclusively the reptilian type. And when, as Dr. Carpenter remarks, we look at the earliest forms of reptilian life of which we have any cognizance, we find them to present very remarkable combinations of the characters which are now distributed among different groups. Thus, at the very commencement of the secondary period, we meet with the extraordinary Labyrinthodon, which, whilst essentially a gigantic Batrachian, was in many respects formed after the Crocodilian model. So, again, the same formation contains the remains of Reptiles, which, while essentially Saurian in their general structure, had the horny mandibles, and probably many other characters, of the Turtle tribe.—It seems to us that there is a sufficiently strong *prima facie* case in favour of this doctrine, to warrant its being systematically put on its trial. The one in place of which it is proposed, cannot stand a rigorous criticism for a moment; and it will be a strong indication of unity of design, if the palæontologist should be able to trace, throughout the long succession of geological epochs, and the repeated metamorphoses which have occurred in the types of organic life which prevailed during each, one harmonious plan, and that plan the same as the embryologist discerns in watching the development of any single organism.

The Second Book commences with a “General View of the Functions of Animated Beings, and of their Mutual Relations;” and the author then proceeds with each function in detail, according to the following plan. Under the head of “General Considerations,” the essential nature of the functional change is considered, and the conditions on which it is dependent are set forth. The function is then traced, in its various stages of complication, from the lowest to the highest forms of the Vegetable kingdom, and its gradual *specialisation* pointed out; and a similar specialisation is traced in the history of the embryonic development of the higher forms of plants. The same course is then followed in regard to Animals; the function being first examined under its simplest conditions, and then traced upwards through the ascending scale of Animal existence, so as to show its gradual limitation to a certain set of organs, which, whilst becoming more and more distinct from the remainder of the organism, and more and more devoted to that purpose only, perform it with more and more of energy and completeness; and the progress of the evolution of these organs in the embryonic state of the higher types, being then exhibited, by way of showing the essential conformity in *plan* between the two series of phases. Such having been the scheme on which this portion of the work was originally constructed, we do not think it requisite now

to give any detailed exemplification of it; since, although it is much more fully and elaborately carried out in the present edition, the readers of the former will here find no essential departure from the original plan. We may, however, notice a few of the more interesting facts, which we now, for the first time, encounter.

It was originally laid down by our author, that the development of the special circulating apparatus is proportional,—not so much to the limitation of the respiratory surface to a particular part of the organism, as stated by Cuvier and others,—as to the degree of restriction of the digestive cavity and of the power of absorption, whereby the parts imbibing aliment are removed from those requiring supplies. In Zoophytes, we find that no part of the organism is sufficiently far removed from the parietes of the digestive cavity, to prevent it from obtaining the requisite amount of alimentary matter, by imbibition through them; and in the Acalephæ and Trematode Entozoa, we frequently find the digestive cavity extended through the body in a very remarkable manner, so as to supersede the necessity of any special circulating apparatus. On the other hand, in the Echinodermata among the Radiated classes, and in all but the lowest Mollusca and Articulata, we find the digestive cavity no longer channelled out among the tissues, but constituting a distinct sac or canal, with which a large part of the body has no close proximity; and a special absorbing and circulating apparatus, of greater or less complexity, is interposed, to take up the nutrient materials from the walls of the digestive cavity, and to convey them to the parts which are destined to appropriate them. Now the very simplest condition of this apparatus would be that in which the general visceral cavity, instead of a distinct system of vessels, is employed for the purpose; and such has been observed in a group of animals, whose general type of structure would give them a comparatively high place in the scale, although, in the evolution of their apparatus of organic life, they present a remarkable degradation. We refer to the tribe of *Pycnogonidæ*, which, in regard to their habits of life, seem to be, among Crustaceans, something what the Sloths are among Mammals. They are found lying inertly upon sea-weeds, with their long, sprawling limbs spread out around the body; and when they change their position, it is not by walking, or springing, or swimming, but by fastening one of their hooked feet to some fixed point, and then drawing the body towards it by the flexure of the limb, just as the Sloth will do when placed upon the ground. Like the Sloths, again, these little animals seem naturally to live in situations where they can imbibe their food with the least possible trouble; for their mouth is suctorial, and it seems probable that their aliment consists of the gelatinous mucus covering the fronds of the sea-weeds on which they live. The mode in which the alimentary matter is conveyed to the tissues, constitutes the most curious feature in the organisation of this animal; and it is thus described by Dr. Carpenter, from his own observations, which coincide with those of M. de Quatrefages. We should mention, however, that the extension of the digestive cavity into the limbs was first noticed by M. Milne-Edwards.

“The mouth leads to a very narrow œsophagus, which passes backwards to the central stomach, situated in the middle of the thorax; and from the posterior part of this cavity, a narrow intestine passes off, to terminate at the posterior extremity of the body. From this central stomach, however, five pairs of cæcal prolongations

radiate, one pair entering the feet-jaws, and the other four penetrating the thoracic legs, and passing along them as far as the last joint but one; when these cæca are distended with fluid, they present a series of contractions and dilatations, which correspond with the articulations of the limbs. That these curious organs are veritable extensions of the stomach into the limbs, and not mere secreting cæca, is proved by the curious peristaltic action which the transparency of the body permits to be observed in them; for they all contract and dilate alternately, so as to produce a continual flow of the contents of this curious digestive cavity from one part into another. Their membranous walls are covered, however, with a layer of brownish-yellow granulations; and these, which are scarcely perceptible on the wall of the central stomach, and not at all upon that of the intestine, may probably be regarded as a diffused and rudimentary condition of the liver. In the fluid which is transmitted from one portion of the digestive cavity to another, minute granular particles can be seen, which are probably of a faecal nature. The digestive sac is freely suspended in the general solid cavity of the body, by occasional fibrous bands; and nothing intervenes between its external surface and the muscular lining of the crustaceous envelope. Thus a wide *lacuna* is left around it, occupying the greater part of the thoracic cavity, and also the hollow of the legs; and this lacuna is filled with a transparent liquid, in which are seen floating a number of minute transparent corpuscles, of irregular size. This fluid is kept in continual motion, not only by the general movements of the body and limbs, but by the more special and constant action of the different parts of the digestive sac; for, when the cæcum in any one of the legs undergoes dilatation, a part of the circumambient liquid will be pressed out from the cavity of the limb, either into the thorax or into some other limb whose cæcum is contracting; and when, in its turn, the first cæcum contracts, the space around it will be again filled with liquid, forced into it by the dilatation of the central stomach or of some other cæcum. There is no other circulating apparatus whatever; and it cannot be doubted that the liquid which fills the great cavity of the body must be regarded as the nutrient material derived from the digestive organs, that is, blood; and that its movement from one part of the body or limbs to another is to be regarded as a great lacunar circulation, no part of it being as yet circumscribed within distinct vessels, and no special organ of impulsion being developed. There is no special organ of respiration; but the aeration of the circulating fluid must be accomplished by exposure to the surrounding medium through the crustaceous envelope, which does not possess great density in any of these animals." (pp. 402-3.)

This condition of the circulating system is not so exceptional as it may at first appear; for the visceral cavity seems to be the sole channel for the conveyance of the nutritious fluid from the digestive sac to the tissues, both among the Bryozoa (which now rank as Mollusks) and the Rotifera; and even among the higher Mollusca and Articulata, it usually forms part of the circuit. The curious discovery that in Invertebrated animals, with probably no exception, the circulation of nutritious fluids is partly *lacunar*, the blood escaping from the vessels to meander through the interstitial spaces, and employing, even in animals as high as Cephalopods, the peritoneal and pericardiac cavities as great venous sinuses, is chiefly due to the researches of Professor Milne-Edwards. Generally speaking, the arterial circulation is truly *vascular*; but there is no proper system of systemic veins, the passage of the blood from the tissues to the respiratory organs taking place entirely through lacunar channels, and it being only whilst returning to the heart after passing through the gills, that it is again conveyed by proper vessels. As justly observed by Professor Milne-Edwards, there is no part of the circulating apparatus in the Vertebrata which can be properly compared with the lacunar portion of that of the Invertebrata, except the commencement of the lymphatic system; for that superfluous

fluid, which, having transuded the walls of the capillary blood-vessels, is not appropriated by the surrounding tissues, appears to find its way through their interstices into the peripheral portions of the lymphatic system, whose origin has not yet been clearly made out. It seems almost impossible to contemplate the peculiar arrangement which has been described as prevalent in the Invertebrata, without perceiving that the branchial circulation must be maintained by some other force than the heart's action, which can no longer be imagined to communicate a *vis a tergo* to the blood, when this is meandering through the lacunæ of the tissues. In the Cephalopoda and higher Crustacea, as well as in certain among the Annelida, we find distinct contractile cavities (apparently of a propulsive character) at the base of the gills; but where these are wanting, there would seem to be no other sustaining power than that for which Professor Alison has always contended, as arising out of the changes to which the blood is rendered subservient in passing through the capillaries themselves.

Some of the most interesting novelties in the Second Part will be found in the Chapter on *Generation and Development*; and of one or two of these we may stop to take a brief notice. The general view of the antagonistic relation between the act of *sexual* Reproduction, or true Generation, and reproduction or multiplication by gemmation, is that which has been already expressed in our own pages; and it has received full confirmation from the researches of Mr. Huxley, who has had ample opportunity of studying the wonderful aberrant forms of the Acalephan class with which tropical seas abound, and who has succeeded, with the aid of the microscope, in unravelling their structure, and in making out their real character, to an extent that is really surprising. In fact, Mr. Huxley had arrived at precisely the same conclusions, when studying these creatures on the coast of New Guinea, that Dr. Carpenter had formed from the analysis and comparison of the facts previously known to him. He considers the *Medusæ*, whatever their degree of independent activity and self-nutrition, as nothing else than the detached generative apparatus of some polypoid organism, which, without the medusa-bud, cannot be regarded as a complete individual; and thus he holds, with Dr. Carpenter, that the polype-stock and free medusa, instead of being "alternate generations," constitute in reality but a single generation. He considers, in fact, that a metamorphosis will be required in our ideas, and, perhaps, also in our language, in consequence of our having formed our ideas of *individuality* entirely upon the higher organisms, in which the whole series of parts evolved as the product of a single act of generation, is so mutually dependent as to constitute but one individual. Among the lower tribes, this is the exception rather than the rule; for the same germ may be developed by gemmation into a great number of parts, which, being able to live independently of each other, commonly take rank as distinct individuals. These may be all equal and similar one to another, or one may be, so to speak, the complement of the other; and of both kinds of gemmation do we see examples among many of the lower tribes. For the generative apparatus to be separate as a distinct gemma, and to be made more or less self-supporting, is really a very common occurrence; and when looked at in this point of view, even the most aberrant and apparently complex forms among the Acalephæ are resolved into simplicity. They are really Zoophytic in their character, being composed of a repetition of a number of similar polype-like organs developed

from each other by gemmation ; and what shows that we must not regard each of these parts, but rather the entire stock, as the real *individual*, is that the generative medusa-buds are not developed from the polypoid bodies, but from the common stem which bears them. Mr. Huxley has proposed the term *Zooids* for these separable segments ; and we believe that it might be adopted with advantage.

This gemmiparous power seems greater, as pointed out by Mr. Paget, in the inverse proportion to the grade of development which the organism has already attained ; and consequently, in the higher tribes, we should expect that the reproductive nusus, at an early period, might be in some degree comparable to that which normally shows itself at a later period in the lower. Thus we find, in General Physiology, the probable explanation of certain Teratological phenomena, which have hitherto been a source of much perplexity. For, as Dr. Carpenter remarks :

“By the knowledge of these facts and principles, we seem justified in the surmise, that the occurrence of supernumerary or multiple parts is not always due (as usually supposed) to the ‘fusion’ of two germs, but that it may result from the subdivision of one ; for if it be supposed that this subdivision has taken place when the developmental process has advanced no further than in a Hydra or a Planaria, it seems by no means impossible that each part might, as in those creatures, advance in its development up to the attainment of its complete form.” (p. 873.)

This view has much to recommend and support it. The separation of the primitive germ-mass into two or more parts, does not seem a very uncommon phenomenon among the lower animals ; and even where the complete organism is not multiplied by this process of fission, imperfect but temporarily-independent bodies are sometimes formed. The following are the very curious observations made by Nordmann on the embryonic development of *Tergipes*, one of the Nudibranchiate Gasteropoda.

“It is not unfrequently seen that some of the cells of the vitelline mass detach themselves from the principal cluster, become clothed with long cilia, and continue to move about actively within the egg, until the escape of the embryo. It is even affirmed by Nordmann, that they increase by partial subdivision, and that thus from a single detached cell may be produced a cluster having a very definite form, and furnished with long cilia, so as very strongly to resemble a parasitic animal. It has not been shown, however, that these bodies ever advance to a higher condition, or are capable of generating their kind ; and the correct view is probably to regard them (with Vogt) simply as portions of the embryonic mass, exactly resembling those that form the ciliated lobes, which, being detached from the rest, preserve their vitality for an unusually long time,—such vitality, however, not being different in kind from that of an ordinary ciliated epithelium-cell, though greater in degree.” (p. 949.)

Now Professor Simpson exhibited to the Physiological Section of the British Association, at its meeting in Edinburgh last year, two living subjects, and several preserved specimens, exhibiting an obvious attempt at the reparation of limbs, after their spontaneous amputation *in utero* ; thus fully supporting the view, that the gemmiparous power exists, even in the human subject, during the earlier period of development ; and justifying the surmise, that the primitive germ-mass might undergo such a complete subdivision as to produce two connected bodies, or that it might bud forth more than its accustomed number of appendages. It has always seemed to us a most extravagant supposition, that the least excess (for example, a double thumb,) must be due to the partial incorporation or

fusion of a second germ ; and if a second thumb or a sixth finger should bud forth from the hands, why should not a second arm with all its appendages bud forth from the shoulder, as in the case which has been recently noticed in the public prints ; or why should not a second head be developed by spontaneous fission of the germ-mass at its cephalic extremity, or an additional trunk and lower extremities proceed from a single head by fission at the posterior part of the primitive trace, the anterior remaining single ? These monstrosities are usually explained on a principle for which we have no warrant in any other part of the animal kingdom ; whilst for their production by the abnormal occurrence of that process in man and the higher animals, which is normal in the lower, we seem to have a strong *à priori* probability. And to the whole class of "monstrosities by inclusion," such a view is obviously more applicable than any other ; it being very easy to understand that the abnormal growth may be the result of an imperfect gemmation, commenced at a very early period ; and this not less readily in the male subject than in the female. The occurrence of cysts containing hair, teeth, bones, &c., is really not very unfrequent ; and their origin seems to us to find a ready solution in the view here propounded.

The inquiry into the history of the *Development* of Animals may be regarded as only now being brought into the prominence it deserves ; and with respect to that of the lower Invertebrata, it may be said that we, as yet, know next to nothing. Unfortunately, it is only by a prolonged residence near the seacoast, and by patient and attentive watchfulness for many months or even years, that the history of even a single species can be fully made out. Among the recent contributions of this kind, one of the most curious is the account given by Professor Müller of the early condition of a species of *Star-fish*. The body first developed from the embryonic mass is a larva, of which little remains in the permanent structure ; and the star-fish is subsequently developed out of a mass of unconverted yolk at its anterior extremity. The larva is of elongated form, and has seven arm-like appendages on either side, which are fringed with cilia, and its bilobed tail is furnished with the same appendages ; by these it moves actively through the water, whilst as yet the mass at the extremity does not present the least indication of what it is permanently to become. Moreover, this larva possesses not only a mouth, but an intestinal canal and anal orifice of its own ; and these have no relation to the mouth which the star-fish is to possess, this being afterwards formed by the thinning-away of the integuments at the part most remote from the attachment of the larva. The young star-fish, when sufficiently matured to maintain its own existence, is separated from the larva (which from its shape has received the name of *Bipinnaria*) by the forcible contractions of the connecting pedicle ; and the larva continues active for some days after its detachment. Strange as this process may seem, it is not so great a departure from the plan of development with which we are familiar in the higher animals, as it at first sight appears ; for, as Dr. Carpenter remarks,—

"We here find the portion of the yolk-mass that first undergoes organisation, converted into a structure which is destined only to possess a transient existence, and which disappears entirely by the time that the development of the principal part of the yolk-mass has advanced so far, that it begins to assume the characters

of the permanent organism. This, however, is what takes place in the higher Vertebrata; for the structures first developed in the egg of the Bird hold nearly the same relation to the rudimentary chick, that the 'Pluteus' bears to the incipient Echinus or Ophiura, or the 'Bipinnaria' to the incipient Star-fish. The only essential difference consists in this,—that the development of these temporary structures proceeds so much further in the latter case, as to give them more the character of distinct individuals; and they are endowed with self-moving powers, whereby they are dispersed through the water in this stage of their existence, so as to prevent that accumulation in particular localities which would otherwise result from the comparatively sluggish habits of these animals in their adult condition." (p. 939.)

In fact, in the Mammalia, we may regard the foetal portion of the placental apparatus as nearly corresponding, homologically, with the bipinnarian larva of the star-fish; and the separation of the former at birth may be considered as representing the detachment of the latter. In each case we find that the portion of the germinal mass which first undergoes organisation, is adapted to provide nutriment for that which is to be more permanent, rather than to be itself converted into it; and the early condition of the embryo is scarcely more unlike what it is afterwards to become, in the one case, than it is in the other.

We do not think it necessary to detain our readers with any account of our author's views on the offices of the Nervous system and organs of sense, which constitute the concluding portion of the work, since they have been already placed before the readers of this journal and of one of its predecessors; and we have only to add, in reference to the external characters of this volume, that it is one of the handsomest, best printed, and best illustrated, that has ever issued from the medical press. A large part of its 321 wood-engravings have been executed by Mr. W. Bagg, whose well-known reputation is a guarantee for their artistic excellence; and the author informs us that he has been careful to have them taken direct, where practicable, from the original monographs of the authors who have specially treated of the several subjects. To all these, reference is made in a separate list; and there are besides, numerous references in the foot-notes, chiefly to the sources of those new facts and doctrines which cannot be yet said to have become part of the common stock of physiological science. The author has followed the plan originally adopted by him, of numbering the paragraphs, and of making frequent reference, by means of these numbers, from one part of the volume to another. And when we add that there is a copious index, we think we have said enough to show that he has spared no pains to make the work available to those for whose use it is intended. We shall end, as we began, with an extract from his Preface, which we think, in justice to him, ought to be placed before our readers.

"The Author thinks it but due to himself to state, that as the almost constant occupation of his time in other indispensable avocations leaves him much less opportunity than he would desire for the prosecution of those original researches in which he feels the greatest satisfaction, so he trusts that the same plea may in some degree avail him in regard to any deficiencies which may be found in the present work. The whole of his disposable time has been devoted to it, for nearly two years, not merely without the anticipation of the slightest pecuniary reward, but under the certain loss involved in the relinquishment of other literary engagements of a remunerative character; and the sale of the entire edition will not do more than remunerate his liberal Publisher for the very large outlay which has been incurred in the production of the work." (*Preface*, p. x.)