more clear, namely, that in proportion as one attains such a perfection of technique as to enable one to open the abdomen without fear of thereby doing injury, so one must discard rough and ready methods like injections or inflations and go straight to the injured spot by means of abdominal section. And this not in intussusception alone.

TUBERCULOSIS IN THE ZOOLOGICAL GARDENS.

DR. WOODS HUTCHINSON,¹ Professor of Comparative Pathology, University of Buffalo, places on record a series of observations made by him in the post-mortem room of the London Zoological Gardens during the winter and spring of 1898-99, the object being to ascertain whether any light could be thrown upon the predisposing causes of tuberculosis from a study of the incidence of the disease upon the various classes of animals there confined. The three great predisposing factors which were specially considered were race, food, and housing, and the death records of the gardens for the past five years were considered in conjunction with Dr. Hutchinson's own observations. Some very curious points came out, but whenever an attempt was made to arrive at a wide generalisation some puzzling exception always cropped up, sufficient to show that we have not yet quite got to the heart of the matter. Take the mammals. The highest order, the primates, is by far the most liable to tuberculosis, while next to it comes the lowest division, the marsupials. Ungulates and carnivora, which are usually ranked as on about the same level of specialisation, are widely separated, the former having six times the death-rate of the latter. Within the class of ungulates itself again the most striking contrasts occur, the odd-toed animals—even horses and asses under domestication—being almost completely exempt, while the even-toed ones are extremely liable. Hence a classification according to race tells us nothing. When we come to food and food habits we seem at first on surer ground, the figures pointing strongly towards a greatly increased susceptibility to the disease among vegetable feeders, and a much reduced liability among meat eaters, both mammals and birds. This corresponds closely with the results of veterinary experience, according to which cattle, rabbits, and guinea-pigs are highly susceptible; pigs much less so, and dogs, cats and rats, almost immune. Here, again, however, there are serious exceptions to the apparent rule; the horse, the ass—indeed all the odd-toed ungulates—the goat and the sheep, all pure vegetable feeders, having marked resisting powers and very seldom becoming infected. Evidently although food-habits coincide more nearly with the distribution of tubercle than do racial divisions, yet there is some other factor involved. On the question of housing and artificial heat it was impossible to say much, for on the one hand the animals shut up in warmed houses are mostly tropical animals which would probably die more quickly still out of doors, while on the other hand it is to be noted that some of the older buildings and sheds are worse ventilated and lighted than the modern-heated houses; so that even out-door animals may be under conditions as far from ideal as indoor ones. Certain observations among the monkeys, however, go to enforce very strongly modern ideas regarding the advantages of air-space and light. Coming back then to the apparent exceptions to the general rule as to the immunity of the carnivora, Dr. Hutchinson inquires whether it is the

nature of the food, or the habits of life involved in obtaining a carnivorous diet, which is at the root of the matter. That a carnivorous method of life in the majority of instances demands a higher degree of activity, speed, and endurance than a herbivorous one needs little demonstration, and he concludes that where from any cause this higher degree of vitality and dash has been acquired by herbivora then immunity from tuberculosis follows. As then the degree of this vigour and activity may be roughly indicated by the size of the heart relative to the body weight, Dr. Hutchinson has arranged his animals according to this ratio, and certainly there is much in his tables to confirm the view suggested. As confirmation of this view it is to be remembered that size and vigour of the heart have been held by more than one authority to be an important element in the predisposition to tuberculosis among men. Taken altogether, the conclusion arrived at from this preliminary survey of the question of the distribution of tubercle is that it is a matter of vigour, endurance, and resisting power, rather than of race, food, or exposure to infection, and that as these are usually higher in flesh-eaters than in vegetable feeders, the former possess a marked relative immunity.

¹ New York Med. Rec., Aug. 25.

TOTAL EXTIRPATION OF THE PROSTATE.

THE lawyers are apt to say that the essence of a crime lies in the intent rather than in the act committed, and it is possible that if the question now being discussed with a certain amount of acrimony as to priority of invention of Mr. Freyer's "new operation" for the extirpation of the prostate were considered in the same way, much of the difference would vanish. As we described in a recent article, Mr. Freyer would treat certain cases of enlarged prostate by enucleation of the gland. He maintains that the prostate is from the first a double gland, and that the obstruction to the outflow of urine is due to the urethra being compressed between its two halves which, although they grow together, are in origin distinct. Having reached that point he advises that an operation should be done, the object and intent of which is the enucleation of these two halves, knowing that if that be done the urethra will be set at liberty. With the urethra itself he endeavours not to interfere. This is his "new operation." No sooner, however, did Mr. Freyer claim that this operation was "new" than he was assailed from many quarters. It was pointed out, with great justice, that the prostate had been enucleated many times before, and with striking success. Although the late Mr. McGill in his earlier cases attacked the obstructing portion of the prostate, it is clear that later both he and those who followed him went much further and enucleated considerable portions of the gland, perhaps the whole of it. Hence they are probably right in maintaining that what Mr. Freyer does they had done, and that in that sense the mere enucleation of the prostate is not a new thing. Moreover, it seems evident that Mr. Freyer much under estimated the success which some operators had met with in their operative proceedings. A man must be very sure of his ground before he sets up a claim for novelty nowadays. Still novelty may lie in the intent as well as in the thing done, and from a careful perusal of all the letters which have been published on the subject, while we cannot but regret the tone in which the discussion has been carried on, we come to the conclusion that if we take a lesson from the lawyers and consider the intent of the opera-

tion instead of concentrating our attention upon the actual details of the proceedings, Mr. Freyer is right in speaking of it as a "new operation." The others aim at removing the obstruction, and, although they often remove a good deal more, if they have removed the obstruction they have done what they set out to do. Mr. Freyer on the other hand, putting on one side the question what particular part of the prostate may at that particular time be doing the mischief, seems to say, "If I can but extirpate this prostate the man will be cured"; so his aim is to extirpate the prostate, and we cannot but feel that looking at the matter in this light his operation is new.

THE TREATMENT OF RAYNAUD'S DISEASE.

IN lecturing a little time ago at the London Hospital on the subject of that group of phenomena which is commonly spoken of as Raynaud's disease, Mr. Jonathan Hutchinson first pointed out the many causes by which interference with the circulation of the extremities might be so interfered with as to produce conditions which might be described, according to their variations, as local syncope, local asphyxia, or symmetrical gangrene of the extremities. But he insisted upon the fact that, in order to constitute a typical resemblance to Raynaud's phenomena, the patient ought to be affected in a paroxysmal manner, that is to say that the symptoms ought to come and go. In regard to treatment, he said that if it seemed that the nervous system was very susceptible in a reflex manner to the influence of temperature, the chief method

of treatment would be to protect the individual from cold, and in young children the main measure was to insist upon warm clothing, and if possible to take care that they lived in a warm climate. In his private practice, Mr. Hutchinson said, he always tried to send a patient who was liable to Raynaud's phenomena to winter in Egypt—not to a tropical but to a warm climate—and he could speak in clear terms as to the benefits derived by such patients from habitually spending the cold part of the year out of England. If that could not be done, there should be every precaution in the shape of warm clothing, to whatever extent might be necessary, to protect them from the injurious influence of external temperature. Then, if the case were one of scleriosis or morphea, and the patient were an adult who could not go away from England during cold parts of the year—and this remark applied to all forms of gangrene of the extremities—the very best remedy for preventing undue susceptibility in regard to reflex influences and for maintaining the circulation in the extremities was opium. If purgatives be given in small quantities, such as from three to five drops of laudanum three or four times a day, with tonics, and careful attention to the liver, and these agree, other remedies may be useful, such as the moderate use of stimulants or the employment of galvanism; but, says Mr. Hutchinson, "my faith is in the use of tonics, the maintenance of the external temperature, and the exhibition of small doses of opium for a long time; it may be even extending to years."

PROGRESS IN OTOLGY.

(Continued from page 399.)

A correct knowledge of the course which the air takes in passing through the nose has important bearings upon aural surgery, as without it we are apt to misdirect our efforts for the relief of nasal insufficiency and the establishment of efficient ventilation of the middle ear. In other words, if we recognise the inferior meatus as the main inspiratory air-path, and the middle and upper meati as having a very subsidiary, if any, importance in the direction taken by the current, we may suffer obstructions in these regions to go untreated which all the while are creating manifest aural phenomena. The downward aspect of the anterior nares in itself suggests the course taken by the inspired air in following its comparatively straight line to the entrance of the middle meatus, yet the inferior is the so-called respiratory tract, to reach which the air must take a strong curve backwards at the limen vestibuli. Zarniko proved by experiment a considerable time ago, the upward course of the inspired air, and later observers, amongst them C. A. Parker,⁶ have demonstrated that it takes a still higher one than even he believed. Chevalier Jackson⁷ directs attention to the influence of certain abnormalities of the septum (usually taken little notice of) upon the inspiratory blast as it descends to reach the posterior choanae on its way to the nasopharynx. These are the small pads or swellings often seen upon the septum posteriorly, close to its free border. By the aid of experiments on the cadaver, Jackson has endeavoured to prove that these swellings deflect the course of the atmosphere in a lateral direction on to the Eustachian cushions and tube mouths. He adduces clinical evidence in support of his contentions and reports seeing deposits of black material, over the Eustachian region in aural patients, the subjects of these growths.

He records that the aural trouble disappeared when the growths were extirpated by the curette. Jackson recommends this practice in every case in which these swellings occur, as a prophylaxis against deafness and tinnitus. This advice is scarcely likely to be followed unless the growths attain dimensions sufficient to cause manifest nasal obstruction.

Remedies for chronic non-suppurative otitis media, the bane of otologists, still come, and "practical cures" are still reported. At the last Otological Congress, Turnbull passed in review the measures which he considered have stood the test of the last 35 years. Foremost amongst these the Politzer bag, after which, Delstanché's rarefacteur, and one or two vibratory methods, were almost the only ones mentioned. Perhaps the bougie and intratympanic injections might be included with these as occasionally rendering valuable aid. Yet it would be a mistake to adopt a hypercritical or cynical attitude towards new remedies enthusiastically eulogised by their advocates lest we stifle research by so doing, and postpone the day, apparently still so far distant, when this inveterate disease shall be placed on the list of curable infirmities. Unfortunately a more accurate knowledge of the natural history and pathology of chronic dry deafness is still a desideratum, and although a clear dividing line between the catarrhal and sclerotic forms cannot be drawn, a full recognition of the two distinctive types is of considerable importance in practice. Amongst the therapeutic agents recently brought forward and awaiting the test of fuller experience is thermotherapy, and there can be no questioning the honesty which Messrs. Lermoyez and Mahu⁷ have exhibited in their account of failures and successes with its employ-