Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/mjafi

Case Report

A rare musculo-septal aponeurotic tunnel in the arm with anomalous arterial pattern

Aparna Muraleedharan^a, Raveendranath Veeramani^{b,*}, Parkash Chand^c^a Assistant Professor (Anatomy), Pondicherry Institute of Medical Sciences, Ganapathichettikulam, Kalapet, Puducherry 605014, India^b Assistant Professor (Anatomy), Jawaharlal Institute of Postgraduate Medical Education & Research (JIPMER), Dhanvantri Nagar, Puducherry 605006, India^c Professor & Head (Anatomy), Jawaharlal Institute of Postgraduate Medical Education & Research (JIPMER), Dhanvantri Nagar, Puducherry 605006, India

ARTICLE INFO

Article history:

Received 11 May 2016

Accepted 17 August 2016

Available online 17 October 2016

Keywords:

Axillary artery

Biceps brachii muscle

Brachial artery

Radial artery

Superficial brachioradial artery

called as bicipital aponeurosis (BBA) or lacertus fibrosus strengthens the deep fascia of the cubital fossa.¹ Supernumerary heads are the most commonly reported variations in this muscle.² The insertion of all the accessory heads was reported to be to the radial tuberosity. There is paucity of literature regarding the variations in the insertion of this muscle.^{3,4}

The median nerve (MN) and the brachial artery (BA) that form the main neurovascular structures of the arm descend close to the medial aspect of the BBM and the brachialis muscles and are also closely related to the medial intermuscular septum (MIS) of arm.¹ The proximity to the musculoaponeurotic structures makes them susceptible to varying degrees of entrapment during anatomical variations in the region. Thus knowledge of the variant insertion of BBM is significant as it can give rise to MN and/or BA entrapment.

Axillary artery (AA) continues as the BA at the lower border of teres major muscle. BA continues distally in the anterior compartment of arm till it reaches the cubital fossa. During its course in the arm it is crossed anteriorly by MN which also descends to the cubital fossa medial to BA. At the level of neck of radius the artery divides into its terminal branches – radial (RA) and ulnar arteries (UAs). RA usually begins at the level of the neck of the radius and terminates by forming the deep palmar arch in the hand by anastomosing with the deep branch of the UA.¹ Variations in the arterial pattern of upper limb have also been reported by several investigators. A brachioradial artery is the most commonly reported variation in the arterial pattern of upper limb with an incidence close to 20%. The incidence of superficial radial artery has been found

Introduction

A number of variations have been previously reported in the muscles, nerves and arteries of the upper limb. The presence of any such variations in the arm and forearm is vulnerable and prone for injury during diagnostic procedures, surgeries and accidents. The biceps brachii muscle (BBM) originates by two heads. Both the tendons unite in the arm, cross the elbow joint and attaches as a common tendon to the posterior aspect of radial tuberosity. A triangular expansion from the tendon

* Corresponding author.

E-mail address: dr_raveendra@rediffmail.com (R. Veeramani).<http://dx.doi.org/10.1016/j.mjafi.2016.08.006>

0377-1237/© 2016 Published by Elsevier B.V. on behalf of Director General, Armed Forces Medical Services.

to be as low as 1%. A superficial brachioradial artery (SBRA) is a very rare variation with an incidence lower than 0.26%.⁵

Here we describe the case of a rare musculoseptal aponeurotic tunnel due to a variant insertion of tendon of BBM which has not been reported previously in literature with coexistent entrapment of MN and BA and the presence of SBRA. Such multiple variations suggest an embryological anomaly. The knowledge of these types of variations is important in diagnostic purposes like cardiac catheterization, arterial grafting and other angiographic procedures.

Case report

Routine dissection of the left upper limb of a 50-year-old male cadaver preserved in 10% formalin was performed. The branches of brachial plexus and AA and the muscles were dissected carefully and the fascial planes including the medial and lateral intermuscular septa were delineated.

There was a well-developed triangular aponeurosis of length 9 cm in the lower part of arm. It extended from the medial aspect of bicipital tendon (BT) to the MIS. It had a well-defined sharp crescentic free margin of 4 cm (Fig. 1). This aponeurosis formed the roof of a musculo-septal aponeurotic tunnel that extended up to the cubital fossa. MN and BA were found to pass through this tunnel with the MN coursing medial to the BA (Fig. 1). Beyond the tunnel the MN passed between the two heads of the pronator teres muscle. Neither the MN nor BA was found to give any branches inside the tunnel after they were opened by dissection. The floor of the tunnel was formed by the MIS with the origin of superficial flexors of forearm. The BBA was found to be thinner and lesser developed than usual.

There was an abnormal medial branch from AA given off 3 cm proximal to the lower border of teres major muscle and 5 cm proximal to the MN formation (Fig. 2). This branch

coursed superficially throughout the arm giving only cutaneous branches to the arm and muscular branches to BBM. At the elbow it crossed from medial to lateral aspect by passing superficial to the tunnel and continued superficially through the forearm as RA and terminated by forming the deep palmar arch along with deep branch of UA. Hence the artery was named as SBRA as per the terminology by Rodríguez-Niedenführ et al.⁵ The larger continuation of AA coursed deep to the MN and gave profunda brachii, superior and inferior ulnar collateral arteries. Beyond the elbow it gave off a recurrent artery which anastomosed with the radial collateral artery (anterior descending branch of profunda brachii). At the level of neck of radius, it divided into UA and a common trunk which later gave off the median artery (MA) and interosseous arteries. The MA was found to pass through the MN splitting the nerve (Fig. 2, inset). The nerve fibers reunited after the artery passed through. The UA coursed normally and terminated in the hand by forming the superficial palmar arch along with a superficial branch of SBRA. There was no communication between SBRA and BA.

The other neurovascular and musculo-fascial structures were found to be normal. The right upper limb was also found to be normal.

Discussion

BBM is a highly variable muscle.⁶ Several authors have reported variations in the number and pattern of fusion of its heads. BBM has been found to have additional heads in as high as 20% of the population.² The present case reports the presence of a musculo-septal aponeurotic tunnel due to a variant insertion of BT with associated entrapment of MN and BA which has not been reported in the past and is rare. Awareness of such a variation helps the surgeon in treating the

Fig. 1 – Lower third of left arm and cubital fossa showing the musculo-septal aponeurotic tunnel. *** Sharp crescentic free margin of the abnormal expansion from bicipital tendon, BBM – biceps brachii muscle, BT – bicipital tendon, BBA – bicipital aponeurosis, BR – brachialis muscle, BA – brachial artery, SBRA – superficial brachioradial artery, MN – median nerve, MIS – medial intermuscular septum, MCV – median cubital vein, MACV – median antecubital vein, MCNF – medial cutaneous nerve of forearm, CV – cephalic vein, IUCA – inferior ulnar collateral artery.

Fig. 2 – Left axilla showing the branches of brachial plexus and abnormal branching pattern of axillary artery into superficial brachioradial and brachial artery 5 cm proximal to formation of median nerve. Inset shows the persistent median artery which passes through median nerve after cutting and reflecting insertion of pronator teres medially. AXA – axillary artery, AXV – axillary vein, ATA – acromiothoracic artery, P.maj – pectoralis major, P.min – pectoralis minor, LHBB – long head of biceps brachii, SHBB – short head of biceps brachii, CB – coracobrachialis, LC – lateral cord of brachial plexus, MCN – musculocutaneous nerve, MN – median nerve, MCNA – medial cutaneous nerve of arm, MCNF – medial cutaneous nerve of forearm, UN – ulnar nerve, SBRA – superficial brachioradial artery, BA – brachial artery, BT – bicipital tendon, FDS – flexor digitorum superficialis, MA – median artery, UA – ulnar artery, CIA – common interosseous artery, PTr – pronator teres reflected.

vascular and nerve entrapment symptoms by simple release of these surrounding compressing structures. Hence, possible anatomical variations in the fibrous and/or muscular structures which can result in entrapment of the neuro-vascular structures most importantly MN and BA require special attention.

Entrapment neuropathies are common. The MN can get compressed at several locations anywhere from the arm to the carpal tunnel owing to its long course.^{7,8} In the present case, it has occurred at the cubital fossa deep to the abnormal aponeurotic tunnel. Generally, such a longstanding entrapment would result in weakness and wasting of the distal musculature which was not seen in this case. The close proximity of the BA to the MN results in entrapment of the BA also in most of the compressions in the arm as is seen in the present case. This result is ischemia of the upper limb leading to the formation of collaterals or manifesting as pain, pallor, paraesthesia of the distal limb.

The morphology of upper limb arterial variations has been studied and classified in detail by several researchers.⁵ Superficial BA, accessory BA, superficial brachioulnar, superficial UA, brachioulnar artery, brachioradial artery, superficial radial artery, SBRA, brachiointerosseous artery, superficial brachio-median, superficial brachioulnoradial artery, absence of the UA and absence of the radial artery are some of the variations in upper limb arterial pattern.⁵ SBRA is a variation in the arterial pattern of the upper limb with prevalence reported as low as 0.26% and is defined as a RA arising high in the arm which courses superficially associated with a normal BA that usually branches into ulnar and interosseous arteries in the cubital fossa.⁵ Even in such cases there would be communications

between SBRA and BA or any deep arteries in the arm. Hence, this case is unique in that there was no communication between SBRA and BA anywhere in the arm, cubital fossa or forearm.

Variations in the vascular patterns are usually the result of anomalous vasculogenesis.⁹ The reason for variations in the arteries could be attributed to the persistence or failure of regression of the embryonic blood vessels.⁹ In the present case, the variant arterial pattern could be due to the collateral formation as a result of compression of BA in the abnormal tunnel during development. The ischemia and hypoxia following compression stimulate the local release of vascular endothelial growth factor (VEGF) which in turn would have stimulated the sprouting of the collateral vessel from the AA which has persisted as SBRA. This view is supported by the fact that the SBRA coursed above the tunnel so as to ensure adequate blood supply to the distal limb.

The clinical significance of upper limb blood vessels is well known. RA is used for coronary artery bypass grafting by many surgeons in addition to great saphenous vein or internal mammary artery (RA). RA is a choice for coronary bypass grafts by some surgeons as there is usually no age limit for its use whereas IMA is usually used in less than 65 years. The biological characteristics of the RA graft like adequate luminal size and very less or no chance of atherosclerosis, introduction of new antispastic management are also reasons for preferring RA over IMA. But higher wall thickness, less elastic and more muscular tunica media, higher incidence of spasm, and non-availability of antispastic management are well known contraindications for its use.¹⁰ Most cases result in reduced hand perfusion following the procedure which has always been a matter of concern though the exact reasons are not

known.¹¹ There is a higher probability of dangerous ischemia of fingers especially in such a situation as seen in our case where the BA too cannot compensate as it is compressed in the lower third of arm.

Conclusions

There was a well-developed triangular aponeurosis in the lower part of arm which extended from the medial aspect of BT to the MIS. The above aponeurosis formed the roof of a tunnel through which the MN and BA passed through. There was associated multiple variations in the arterial pattern of upper limb all of which could be correlated embryologically. The knowledge of these types of variations is important in cardiac catheterization, arterial grafting and angiographic procedures.

Conflicts of interest

The authors have none to declare.

REFERENCES

1. Standring S. *Gray's Anatomy The Anatomical Basis of Clinical Practice, Expert Consult – Online and Print*. 40th ed. Edinburgh: Churchill Livingstone; 2008.
2. Cucca YY, McLay SV, Okamoto T, Ecker J, McMenamin PG. The biceps brachii muscle and its distal insertion: observations of surgical and evolutionary relevance. *Surg Radiol Anat*. 2010;32(4):371–375.
3. Cho CH, Song KS, Choi JJ, et al. Insertional anatomy and clinical relevance of the distal biceps tendon. *Knee Surg Sports Traumatol Arthrosc*. 2011;19(11):1930–1935.
4. Fogg QA, Hess BR, Rodgers KG, Ashwood N. Distal biceps brachii tendon anatomy revisited from a surgical perspective. *Clin Anat*. 2009;22(3):346–351.
5. Rodríguez-Niedenführ M, Vázquez T, Nearn L, Ferreira B, Parkin I, Sañudo JR. Variations of the arterial pattern in the upper limb revisited: a morphological and statistical study, with a review of the literature. *J Anat*. 2001;199:547–566.
6. Macalister A. *Additional Observations on Muscular Anomalies in Human Anatomy (Third Series) with A Catalogue of the Principal Muscular Variations Hitherto Published*. Dublin: Academy; 1872.
7. Bilecenoglu B, Uz A, Karalezli N. Possible anatomic structures causing entrapment neuropathies of the median nerve: an anatomic study. *Acta Orthop Belg*. 2005;71:169–176.
8. Wertsch JJ, Melvin J. Median nerve anatomy and entrapment syndromes: a review. *Arch Phys Med Rehabil*. 1982;63:623–627.
9. Rodríguez-Niedenführ M, Vázquez T, Parkin IG, Sanudo JR. Arterial patterns of the human upper limb: update of anatomical variations and embryological development. *Eur J Anat*. 2014;7:21–28.
10. He GW. Arterial grafts for coronary artery bypass grafting: biological characteristics, functional classification, and clinical choice. *Ann Thorac Surg*. 1999;67:277–284.
11. Hayes EC, L'Ecuyer KM. A standard of care for radial artery grafting. *Am J Crit Care Off Publ Am Assoc Crit-Care Nurses*. 1998;7:429–435.