Published in final edited form as:

Psychol Med. 2017 January ; 47(1): 1–17. doi:10.1017/S0033291716002075.

The Association Between Cognitive Function and Subsequent Depression: A Systematic Review and Meta-Analysis

Matthew A. Scult, MA^a, Athelia R. Paulli^a, Emily S. Mazure, MSI, AHIP^b, Terrie E. Moffitt, PhD^a, Ahmad R. Hariri, PhD^{a,*}, and Timothy J. Strauman, PhD^{a,*}

^aDepartment of Psychology & Neuroscience, Duke University, Durham, NC, 27708, USA

^bDuke University Medical Center Library & Archives, Duke University Medical Center, Durham, NC, 27710, USA

Abstract

Despite a growing interest in understanding the cognitive deficits associated with major depressive disorder (MDD), it is largely unknown whether such deficits exist before disorder onset or how they might influence the severity of subsequent illness. The purpose of the present study was to conduct a systematic review and meta-analysis of longitudinal datasets to determine whether cognitive function acts as a predictor of later MDD diagnosis or change in depression symptoms. Eligible studies included longitudinal designs with baseline measures of cognitive functioning, and later unipolar MDD diagnosis or symptom assessment. The systematic review identified 29 publications, representing 34 unique samples, and 121,749 participants, that met the inclusion/exclusion criteria. Quantitative meta-analysis demonstrated that higher cognitive function was associated with decreased levels of subsequent depression ($r=-0.088$; 95% CI: $-0.121, -0.054$; $p<0.001$). However, sensitivity analyses revealed that this association is likely driven by concurrent depression symptoms at the time of cognitive assessment. Our review and meta-analysis indicate that the association between lower cognitive function and later depression is confounded by the presence of contemporaneous depression symptoms at the time of cognitive assessment. Thus, cognitive deficits predicting MDD likely represent deleterious effects of subclinical depression symptoms on performance rather than premorbid risk factors for disorder.

Introduction

A growing literature has found that depression is associated with impaired cognitive functioning (Snyder 2013). For example, broad deficits in neuropsychological functioning have been demonstrated in a number of reviews and meta-analyses on depression (Christensen *et al.* 1997; Rogers *et al.* 2004; Snyder 2013; Rock *et al.* 2014), and cognitive deficits have been found to correlate with symptom severity (McDermott & Ebmeier 2009). Theoretical models have been proposed suggesting that cognitive dysfunction could either

Correspondence should be addressed to matthew.scult@duke.edu. Department of Psychology & Neuroscience, Duke University, Durham, NC, 27708, USA. Phone: 919-660-5757.

*Co-equal authorship

Conflicts of Interest

None

be a risk factor for later depression or that cognitive function may be impaired as a result of depression (Stern 2003; Barnett *et al.* 2006). In support of the latter hypothesis, two recent meta-analyses concluded that the deficits in cognitive function persist after remission from depression (Bora *et al.* 2013; Rock *et al.* 2014). However, evidence showing that deficits in cognitive function persist after remission from depression, which are primarily derived from cross-sectional studies, does not rule out the possibility that poor cognitive function preceded the onset of depression. The present review and meta-analysis sought to comprehensively summarize the state of the research on cognitive function and its association with future depressive symptoms using data from longitudinal studies.

While the clinical diagnosis of depression is relatively codified according to DSM and ICD criteria, and depression symptoms are readily measured with a number of validated instruments (Lewinsohn *et al.* 2000; Cuijpers *et al.* 2004), a similar canonical approach to measuring cognitive function is not present. Thus, it is important to specify a working definition of cognitive function before examining its possible association with depression. Empirical work has reinforced the concept of a single construct of cognitive function through high correlations in performance among disparate cognitive tasks (Spearman 1904). This has led to the derivation of a single factor representing general cognitive function or ability known as “*g*” (Johnson *et al.* 2004). The “*g*” factor is commonly determined by administering a wide range of tasks and using factor analysis to determine the shared variance across these tasks (Ree & Earles 1991). Even in cases when “*g*” is not calculated as such, there are other customary ways of capturing this general factor by combining across cognitive tasks, e.g., by creating an index of executive functions (Miyake *et al.*, 2000) or measuring the intelligence quotient (Gottfredson 1997).

Although executive functions and intelligence are often considered independently, evidence has suggested that they may best be characterized by an integrated framework based on a common neural network (Barbey *et al.* 2012), supporting the idea that they reflect a shared general factor of cognitive ability. In light of these findings and the observation that individuals with depression commonly exhibit deficits across a range of cognitive tasks, our review used the broadest measure of cognitive function that was available in a given study. When a higher-order construct was not available, results from individual measures were used. We further restricted analyses to prospective cohort studies with depression assessment before age 65 in order to minimize potential influences of age-related cognitive decline.

We follow our literature review with a quantitative meta-analysis to (1) determine whether cognitive function predicts later depression, (2) evaluate whether differences in the measurement of depression (categorical vs. continuous) influence observed associations, and (3) examine whether sex or age of participants may moderate effects, and (4) whether effects are confounded by depression symptoms at baseline.

Methods

We followed the Meta-Analyses of Observational Studies in Epidemiology (MOOSE) guidelines (Stroup *et al.* 2000) (Supplemental Table 1) and the literature search strategies suggested by Atkinson *et al.* (Atkinson *et al.* 2014).

Search Process

Systematic searches were conducted in February 2015, and updated in December 2015 in three electronic databases: PubMed, EMBASE and PsycInfo. Searches were conducted by a research librarian, EM, and overseen by MS. The search syntax was adapted for each database and designed to capture the participants, predictors, comparisons and outcomes described below, according to the PICOS framework (Moher & Liberati 2009). The Boolean operator “OR” was used within categories and the operator “AND” was used between categories. The complete search strategy is described in Supplemental Materials: Search Strategy. Hand searches were conducted of the reference lists of included articles.

Eligibility Criteria

The review included English language studies of empirical studies investigating the longitudinal association of cognitive function (IQ, neuropsychological tests, executive functions) with unipolar depression diagnosis or symptoms. Detailed inclusion/exclusion criteria were as follows:

Participant Characteristics—Eligible studies included children, adolescents and adults under the age of 65. This age range was specified because of the possible impact of cognitive decline in older populations. Articles examining a highly specific population (e.g., with a particular medical diagnosis) were excluded so as to not limit the generalizability of results. Studies that required MDD diagnosis, hospitalization or admission to a treatment center as part of the inclusion criteria at baseline were also excluded from the review so as to permit conclusions about depression onset. This was to ensure that a majority of study participants did not have depression at baseline. However, cohort studies were included even if they did not measure depression at baseline, assuming that these samples would have depression prevalence rates that were comparable to population norms. Subgroup analyses were subsequently performed on the studies that specifically excluded for any history of depression. If there were several papers about one cohort, the paper with larger sample size or better study quality was included.

Predictor Type—Eligible studies included a measure of cognitive, executive or neuropsychological function. The study included a range of cognitive predictors, given that previous studies have found broad deficits in cognitive function in depression (Christensen *et al.* 1997; Rogers *et al.* 2004; Snyder 2013; Rock *et al.* 2014). When more than one cognitive predictor was reported as an outcome in a study, the broadest measure of cognitive functioning was used. If no measure captured broad cognitive functioning, the measure that was most closely aligned with constructs on a standard IQ test (working memory, processing speed, verbal comprehension, or perceptual reasoning) was used in the meta-analysis.

Studies were excluded if cognitive function was assessed via educational achievement, given that these measures can be more easily influenced by learning disabilities than by general cognitive functioning (Siegel 1999). Self-report descriptive measures of cognitive functioning were also excluded in favor of more objective measures. Lastly, since this study sought to investigate general information processing abilities rather than cognitive

attributional style, studies with cognitive attribution measures as the sole cognitive measure were excluded.

Comparisons—Search terms used to indicate the possible association between cognitive function and depression included: *risk, premorbid, prodromal, onset, predict* and *association* (for a full list, see search terms in the Supplemental Materials). When data were reported from multiple time-points, the analysis accounting for the longest time between assessments was used.

Outcomes—Eligible studies reported depression diagnosis or symptoms determined by investigator or self-report. Presence of comorbidity was permitted given that depression has been found to be highly comorbid with other disorders (Melartin *et al.* 2002; Kessler *et al.* 2003). Articles not adequately specifying how depression was assessed or articles that did not uniquely measure depression (ie. only had an index of “mental disorders” or the category of “mood disorders” lumped together (Gale *et al.* 2010)), were excluded if no unique measure of depression could be obtained from study authors. This is because, although comorbidity was permitted, these could have represented instances of pure anxiety disorders with few or no depressive symptoms or instances of bipolar disorder.

Study Design—Eligible study search terms included: *cohort studies, longitudinal, epidemiological, prospective, retrospective, follow-up* or *case-control* studies. Studies not following a longitudinal design were excluded during full-text review. Additionally, studies where cognitive function was assessed only concurrently with or after depression were excluded, given that the purpose of the review was to investigate whether cognitive function predicted later depression.

Studies were included regardless of whether they were reported in peer-review journals in order to account for potentially biased reporting of results. Review articles were included in the search terms to find additional relevant references, but were not included in the meta-analysis. Baseline and protocol studies without follow-up data were excluded.

Study Selection

A flow diagram of the process of study selection shows the overall procedures (Figure 1). Records identified from the search processes were combined in EndNote software version X7 and duplicates were removed. Titles and abstracts were reviewed independently by two reviewers (MS & AP) according to the eligibility criteria described above, and marked as either potentially eligible or not eligible. If either author marked an article as potentially eligible, it was included in the full-text review for closer examination by MS and AP, who used an electronic form to indicate if an article should be included or excluded in the final selection. When an article was excluded, the reason for exclusion was selected from a hierarchical list of exclusion criteria presented in Figure 1. Both MS and AP were initially blinded to the other’s decisions. After both MS and AP completed their reviews, discrepancies about whether a study should be included or excluded were resolved via discussion and consensus amongst MS, AP, TM, AH, and TS.

Data Extraction

Study information was extracted independently by MS and AP using an electronic form created for this study. Information was collected and coded based on report characteristics, participant characteristics, study setting, predictors, outcome measures and quality assessment (Cooper 2010). Any discrepancies (approximately 3% of data points) were resolved via consensus.

For studies with categorical depression diagnoses, effect sizes were generally presented as odds ratios. For these studies, the odds ratio plus the upper and lower bound of the confidence interval were recorded. Two of the studies (Zammit *et al.* 2004; Gale *et al.* 2008) reported odds ratios greater than one to indicate that lower IQ was associated with greater risk of depression. The other studies reported odds ratios in the reverse, leading to odds ratios less than one for the same direction of the effect. Therefore the odds ratios from the first two studies were recalculated by dividing (1/odds ratio). Another study (McCord & Ensminger 1997) reported results from chi-squared tests. Additional studies reported mean premorbid IQ scores rather than odds ratios, in which case the means, sample sizes and/or p-values and number of tails of the t-test were utilized to calculate the effect size.

For studies in which depression was measured as a continuous variable, effect sizes were presented as correlation coefficients. The direction of the effect (positive or negative) and the sample size were also extracted to be used in the meta-analysis. Two studies reported only beta weights (Simons *et al.* 2009; Rawal & Rice 2012) and we were not able to obtain zero-order correlations from the authors. Given that previous reports have suggested that beta weights can be used as an estimate of correlation coefficients in meta-analysis (Peterson & Brown 2005), the beta weights were used instead, and analyses were run both including and excluding these data points.

When insufficient information was presented in the article, authors were contacted requesting further detail. An additional 6 studies were included at this point. One study (Hatch *et al.* 2007) included effect sizes for the female half of the cohort, but did not provide statistics for the males, but did indicate a null effect. Only women from this study were included in the meta-analyses, but imputing a score of 0 for the males yielded similar overall results.

Follow-up analyses directly compared studies that reported zero-order correlations between IQ at Time 1, Depression at Time 1 and Depression at Time 2 (additional details described in Supplemental Methods).

Statistical Methods

Data were analyzed using Comprehensive Meta-Analysis software (Version 3.0, Biostat Inc, Englewood, NJ, USA). A random-effects model was used for expected heterogeneity across studies. A random-effects model also allows for broader generalization of results to the population at large. Heterogeneity of effects was calculated using the Q statistic, where a significant p-value indicates that true effect varies across studies. The I^2 statistic was used to calculate the ratio of the true heterogeneity to total variation, acting as an index of signal-to-

noise ratio. When I^2 is large, it has been recommended to consider subgroup analysis or meta-regression to account for the variance (Borenstein *et al.* 2011).

Subgroup analyses were conducted using a mixed-effects analysis in which a random effects model was used to combine studies within each subgroup, and a fixed effect model was used to combine subgroups and yield the overall effect (the study-to-study variance is not assumed to be the same for all subgroups). Subgroup analyses were conducted comparing type of outcome (continuous vs. categorical), broad vs. specific cognitive measures, and outcomes that were adjusted vs. unadjusted for baseline depression symptoms.

Meta-regression was performed using the following variables on unadjusted analyses: age at baseline, age at follow-up, time between assessments, IQ, percent white, year of cognitive assessment, year of follow-up assessment, percent female, and study quality score.

Study Quality and Publication Bias

Study quality was evaluated independently by MS and AP using a checklist adapted from Luppino & Wit (Luppino *et al.* 2010). The checklist contained 12 items and studies were rated with a “+” if it met criteria for that item, a “-” if it did not meet criteria for an item, and a “?” if unclear. “+” were coded as a score of 1 and “-/?” were coded as a score of 0. Total scores and percentages were calculated. Articles with total scores of 0–3 were considered to be of low quality, scores of 4–7 were considered a medium quality, and scores of 8–12 were considered to be of high quality. Any discrepancies (approximately 3% of data points) were resolved via consensus.

Publication bias was assessed using a funnel plot, which presents study size on the vertical axis (as standard error, in this case) as a function of effect size on the x-axis. When no publication bias exists, the studies should be distributed symmetrically about the combined effect size (Borenstein *et al.* 2011). If evidence of publication bias was observed, Duval and Tweedie’s Trim and Fill Method was used (Duval & Tweedie 2000), which recalculates effects based on imputing data from studies that are likely to be missing. Lastly, the Fail-Safe N was calculated (Cooper 1979), which indicates the number of missing studies that would need to exist to nullify the observed effect.

Results

A flow diagram of the search process is shown in Figure 1. The search identified 2,773 records through PubMed, 2,158 through EMBASE and 1,345 through PsycInfo, for a total of 6,276 records. Duplicate records (1,709) were removed in EndNote, resulting in 4,567 unique records. After title and abstracts were reviewed, 329 records were identified as potentially eligible for the meta-analysis. These 329 records were reviewed in full and 300 articles were excluded hierarchically if they were: a review, baseline or protocol paper (n=6), if the sample had a mean age over 65 at the depression assessment (n=41), if the sample had a specific medical illness (n=23), if the study was not a longitudinal design (n=19), if MDD diagnosis, hospitalization or admission to treatment center was part of the inclusion criteria of the study (n=18), if cognitive function was assessed only concurrently with or after depression (n=36), if cognitive function was assessed by educational achievement or self-

report ($n=26$), if cognitive function was assessed by attribution style rather than neuropsychological measure ($n=7$), if there was no measure of cognitive function ($n=18$) or no uniquely measured depression outcome ($n=24$). Furthermore, if articles reported on the same sample as another article with higher quality or more information, the duplicate articles were excluded ($n=8$). Lastly, a number of studies appeared to measure both cognitive function and depression, but did not report enough information to be included in the meta-analysis and did not respond to requests for additional information ($n=74$, approximately 59 unique samples). In total, 29 publications were included in the meta-analysis, comprising 34 samples (one study included two separate birth cohorts and another four reported data separately for men and women).

Study Characteristics

The characteristics of included samples are detailed in Table 1. Sample sizes ranged from 43 to 50,053 with a median of 339.5. The most common types of samples were school samples, (9/34), community samples (8/34), and birth cohorts (7/34). The samples were majority white, with two samples having exclusively black participants. Additionally, 8 samples were men only, 6 samples were women only, and the rest reported combined effect sizes. Mean age of participants at baseline ranged from 3.5 to 59 years with a median of 12 years. Most samples did not report whether comorbid diagnoses were present (23/34).

Of the 74 papers excluded for having insufficient information (representing 60 unique samples), the median number of subjects was not statistically different between those studies and the studies included in the meta-analysis ($n=288$ and $n=339.5$, respectively; $p=0.19$), nor did the average age of study participants at baseline statistically differ (median across studies = 11 vs. 12, respectively; $p=0.57$). The types of cognitive measures used were also similar across the excluded and included studies, with the most common measure being IQ (28/60 and 20/29), followed by Verbal Comprehension (8/60 and 5/29).

Meta-Analysis

The overall analysis yielded a significant association between cognitive function and subsequent depression outcomes ($r=-0.088$; 95% CI: $-0.121, -0.054$; $p<0.001$; Figure 2). Results were similar when excluding the two studies that used beta-weights ($r=-0.071$; 95% CI: $-0.100, -0.042$; $p<0.001$) or when using episodic memory in place of processing speed in a study reporting both outcomes ($r=-0.085$; 95% CI: $-0.118, -0.051$; $p<0.001$). Follow-up analyses were conducted in the sample of studies reporting educational attainment and self-report outcomes (Supplemental Table 2). Heterogeneity across samples was significant ($Q=801.54$, $p<0.001$), suggesting substantial variation across the individual samples. The ratio of the true heterogeneity to total variation was large ($I^2=95.88\%$). I^2 is on a relative scale and values close to 100 indicate that most of the observed variance is real rather than spurious and suggests that subgroup analyses or meta-regression may help to explain this variability (Borenstein *et al.* 2011).

Subgroup Analyses and Meta-Regression

A significant difference was found between categorical and continuous effect sizes ($p<0.001$). The studies assessing continuous outcomes were found to have a larger effect ($r=$

−0.121; 95% CI: −0.168, −0.073; $p < 0.001$; $n = 22$) than studies assessing categorical outcomes ($r = -0.035$; 95% CI: −0.054, −0.017; $p < 0.001$; $n = 12$). No differences were observed between studies utilizing broad vs. specific cognitive measures ($p = 0.57$).

Outcomes were also compared adjusting for baseline depression symptoms at the time of cognitive assessment. Nine studies provided enough detail to calculate the partial correlation coefficients adjusting for baseline depression symptoms. Using the partial correlation coefficients led to a null result ($r = -0.032$; 95% CI: −0.078, 0.014; $p = 0.169$; $n = 9$; Figure 3). The null result does not seem to be due to a lack of power because using the unadjusted correlation coefficients for the same nine studies still led to a significant result ($r = -0.120$; 95% CI: −0.169, −0.070; $p < 0.001$; $n = 9$; Figure 3). As a follow-up, the difference in effect sizes between the adjusted and unadjusted values for these 9 studies were tested ($r = 0.082$; 95% CI: 0.069, 0.095; $p < 0.001$; $n = 9$), suggesting that the effects are significantly different when directly comparing the values that do and do not control for baseline depression symptoms.

Expanding the subgroup analyses to include studies that controlled for baseline depression plus other covariates (Zammit *et al.* 2004; Simons *et al.* 2009; Rawal & Rice 2012), and studies that explicitly excluded for a history of depression diagnosis (Gale *et al.* 2008; Vinberg *et al.* 2013; Pappmeyer *et al.* 2015), did not change the null result ($r = -0.027$; 95% CI: −0.058, 0.005; $p = 0.097$; $n = 15$).

None of the meta-regression analyses were significant, indicating that the between-study variance was not attributable to age at baseline, IQ, percent of sample that was female, or percent of the sample that was white age at follow-up, time between assessments, year of cognitive assessment, year of follow-up assessment, or study quality score.

Study Quality and Publication Bias

Multiple samples within a study were combined for the purposes of quality assessment, given that studies within the same publication were found to employ the same methodology. Of the 29 publications reviewed, 12 were rated as high quality and 17 were rated as medium quality. Details are provided in Supplemental Table 1. Comparable results to the overall findings were obtained when just using high quality studies ($r = -0.085$; 95% CI: −0.141, −0.029; $p = 0.003$).

Inspection of the funnel plot (Figure 4) found no evidence for publication bias to support the hypothesis. In fact, imputation of negative correlations suggested that additional studies favoring the hypothesis may be missing. While the calculated effect size is −0.088 (95% CI: −0.121, −0.054), using Duval and Tweedie's Trim and Fill Method, the imputed effect size estimate is larger at −0.137 (95% CI: −0.176, −0.099; imputed studies $n = 11$). Lastly, the Fail-Safe N procedure found that 3,583 null studies would need to be located for the combined 2-tailed p -value to exceed 0.05. This suggests that significant results are not likely to be confounded by publication bias and may even underestimate true effect sizes.

Discussion

The present systematic review and meta-analysis of 29 longitudinal publications, including 121,749 participants, revealed that after accounting for baseline depression symptoms, variability in cognitive function did not predict subsequent depression. Consistent with prior reports, there was a significant contemporaneous association between higher depression symptoms and lower cognitive function. These patterns have implications for understanding the association between cognitive function and depression and may be important for advancing etiologic and treatment research.

First, our findings reinforce models positing that cognitive function tracks with depression severity (McDermott & Ebmeier 2009). One possible explanation for this contemporaneous association is that depression symptoms interfere with the capacity to complete cognitive assessments, possibly through the general lack of motivation that is one hallmark of the disorder. While this explanation represents an experimental confound for studies focused on assessment of cognitive function, it likewise represents an opportunity for studies to focus more comprehensively on the assessment of both depression and mood disorder symptoms, since performance on cognitive tests may be a sensitive measure of motivational state that could augment self-reported symptoms. Alternatively, the co-occurrence of cognitive deficits and depression symptoms may reflect a shared genetic etiology (Hagenaars *et al.* 2016) and dysfunction of neural circuits supporting both cognitive and emotional processes (Scult *et al.* in press). In particular, dysfunction of prefrontal and striatal circuits have been associated with both depression and cognitive function (Keedwell *et al.* 2005; Aarts *et al.* 2011).

Second, our findings suggest that prior studies of links between cognitive function and depression that did not assess baseline symptoms may have overestimated the potential protective role of higher cognitive function. While the importance of accounting for subclinical symptoms is not novel, the implementation of such accounting is often neglected in psychiatric epidemiology, given the current categorical, threshold-based diagnostic system. In our meta-analysis, larger effects were observed for continuous as compared to categorical outcomes, which aligns with growing emphasis on dimensional measures of psychopathology (Cuthbert & Kozak 2013).

Third, the associations between cognitive function and either current or later depression were not moderated by data quality or participant age, sex, and race. However, there are a number of additional potential moderators that could not be addressed in the current review and meta-analysis. For example, the number and type of comorbid conditions with depression were often not reported in the included studies, but could represent an important moderator of the observed effects. Modeling the effects of comorbid illness is an important avenue for future research.

One limitation of the current literature is that only a handful of studies reviewed explicitly excluded for depression diagnosis at baseline (Gale *et al.* 2008; Vinberg *et al.* 2013; Pappmeyer *et al.* 2015); however, consistent with the finding that controlling for baseline depression symptoms led to a null association between cognition and subsequent depression, there was no effect observed when including these studies in the subgroup analyses, further

suggesting that the association between cognitive function and depression is likely contemporaneous. It will be important to follow-up these analyses with individual studies that include rigorous assessment of baseline depression symptoms and other potentially confounding factors to further clarify this relationship.

An open question that remains for future research is whether deficits in cognitive function are likely to predict the development of more severe or recurrent depression later in life. Only one of the included studies (Koenen *et al.* 2009), specifically investigated recurrent depression and found that lower IQ was associated with greater persistence of disorder.

A final consideration is that while our review and meta-analysis focused on general cognitive function mainly assessed via IQ, there may be specific subtypes of cognitive function that are more predictive of later depression. For example, a lack of cognitive flexibility may more closely match with the ruminative style characteristic of depression, and therefore tasks specifically measuring this component of cognitive function might have some predictive utility. While our review did not find differences between broad and specific measures of cognitive function, there were not enough studies identified to allow for precise consideration of differences between individual cognitive domains.

A limitation of the meta-analysis is that while the present study followed standard guidelines (Stroup *et al.* 2000; Atkinson *et al.* 2014), decisions about study selection, data extraction, and quality assessment, include a number of decision points that necessarily involve a level of subjectivity. Reliability was ascertained by using independent raters, however, different inclusion/exclusion criteria or a different approach to the evaluation of those criteria, could result in a different set of studies being included. Ultimately, the quality of the meta-analysis is dependent upon the underlying studies that are analyzed. Furthermore, although studies with insufficient information to be included in the meta-analysis were not found to differ based on basic study data from those studies included in the meta-analysis, nonetheless it is unknown how if at all data from these studies could change the overall results.

Acknowledging these limitations, our present review and meta-analysis found that while an association is evident between cognitive function and later depression, general cognitive function does not appear to be a risk factor for depression, but rather is more likely related to performance decrements associated with concurrent depressive symptoms. Our results suggest that low cognitive function is therefore probably not a causal factor of depression and that clinical practice may benefit more from a focus on how decreased cognitive function in the depressed state is likely to influence treatment outcomes (Gyurak *et al.* 2015). Our findings have important implications for better understanding depression and for the design of future studies in highlighting the need to control for subthreshold depression symptoms when investigating risk factors for mental illness as well as when assessing cognitive function more generally.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial Support

M.A.S. is supported by an NSF Graduate Research Fellowship. A.R.H. is supported by NIH grants RO1DA033369 & RO1AG049789. T.E.M. is supported by RO1AG049789.

We would like to thank Harris Cooper PhD at Duke University, who provided invaluable insight into conducting the meta-analysis. We thank all of the authors of studies who provided additional information for this meta-analysis.

References

- Aarts E, van Holstein M, Cools R. Striatal Dopamine and the Interface between Motivation and Cognition. *Frontiers in psychology*. 2011; 2:163. [PubMed: 21808629]
- Atkinson KM, Koenka AC, Sanchez CE, Moshontz H, Cooper H. Reporting standards for literature searches and report inclusion criteria: making research syntheses more transparent and easy to replicate. *Research Synthesis Methods*. 2014
- Baer LH, Tabri N, Blair M, Bye D, Li KZH, Pushkar D. Longitudinal associations of need for cognition, cognitive activity, and depressive symptomatology with cognitive function in recent retirees. *The journals of gerontology. Series B, Psychological sciences and social sciences*. 2013; 68:655–664.
- Barbey AK, Colom R, Solomon J, Krueger F, Forbes C, Grafman J. An integrative architecture for general intelligence and executive function revealed by lesion mapping. *Brain : a journal of neurology*. 2012; 135:1154–1164. [PubMed: 22396393]
- Barnett JH, Salmond CH, Jones PB, Sahakian BJ. Cognitive reserve in neuropsychiatry. *Psychological medicine*. 2006; 36:1053–1064. [PubMed: 16854246]
- Beaujean AA, Parker S, Qiu X. The relationship between cognitive ability and depression: a longitudinal data analysis. *Social psychiatry and psychiatric epidemiology*. 2013; 48:1983–1992. [PubMed: 23474611]
- Belsky DW, Caspi A, Arseneault L, Bleidorn W, Fonagy P, Goodman M, Houts R, Moffitt TE. Etiological features of borderline personality related characteristics in a birth cohort of 12-year-old children. *Development and psychopathology*. 2012; 24:251–265. [PubMed: 22293008]
- Betts, KS.; Williams, GM.; Najman, JM.; Alati, R. *Journal of Psychiatric Research*. Vol. 72. Elsevier Ltd; 2016. Predicting spectrums of adult mania, psychosis and depression by prospectively ascertained childhood neurodevelopment; p. 22-29.
- Bora E, Harrison BJ, Yücel M, Pantelis C. Cognitive impairment in euthymic major depressive disorder: a meta-analysis. *Psychological medicine*. 2013; 43:2017–2026. [PubMed: 23098294]
- Borenstein, M.; Hedges, LV.; Higgins, JPT.; Rothstein, HR. *Introduction to meta-analysis*. John Wiley & Sons; 2011.
- Canals J, Domènech-Llaberia E, Fernández-Ballart J, Martí-Henneberg C. Predictors of depression at eighteen. A 7-year follow-up study in a Spanish nonclinical population. *European child & adolescent psychiatry*. 2002; 11:226–233. [PubMed: 12469240]
- Christensen H, Griffiths K, Mackinnon A. A quantitative review of cognitive deficits in depression and Alzheimer-type dementia Selection of Studies. 1997:631–651.
- Connolly, SL.; Wagner, Ca; Shapero, BG.; Pendergast, LL.; Abramson, LY.; Alloy, LB. *Journal of behavior therapy and experimental psychiatry*. Vol. 45. Elsevier Ltd; 2014. Rumination prospectively predicts executive functioning impairments in adolescents; p. 46-56.
- Cooper HM. Statistically combining independent studies: A meta-analysis of sex differences in conformity research. *Journal of Personality and Social Psychology*. 1979; 37:131–146.
- Cooper, HM. *Research Synthesis and Meta-Analysis: A Step-by-Step Approach*. 4th. Washington (DC): U.S: SAGE Publications; 2010.
- Cuijpers P, de Graaf R, van Dorsselaer S. Minor depression: risk profiles, functional disability, health care use and risk of developing major depression. *Journal of affective disorders*. 2004; 79:71–79. [PubMed: 15023482]

- Cuthbert BN, Kozak MJ. Constructing constructs for psychopathology: the NIMH research domain criteria. *Journal of abnormal psychology*. 2013; 122:928–937. [PubMed: 24016027]
- Der, G.; Batty, GD.; Deary, IJ. Intelligence. Vol. 37. Elsevier Inc; 2009. The association between IQ in adolescence and a range of health outcomes at 40 in the 1979 US National Longitudinal Study of Youth; p. 573-580.
- Dubow EF, Boxer P, Huesmann LR. Childhood and adolescent predictors of early and middle adulthood alcohol use and problem drinking: the Columbia County Longitudinal Study. *Addiction*. 2008; 103:36–47. [PubMed: 18426539]
- Duval S, Tweedie R. Trim and Fill: A Simple Funnel-Plot-Based Method of Testing and Adjusting for Publication Bias in Meta-Analysis. *Biometrics*. 2000; 56:455–463. [PubMed: 10877304]
- Evans LD, Kouros CD, Samanez-Larkin S, Garber J. Concurrent and Short-Term Prospective Relations Among Neurocognitive Functioning, Coping, and Depressive Symptoms in Youth. *Journal of clinical child and adolescent psychology : the official journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division*. 2015; 53:1–15.
- Franz, CE.; Lyons, MJ.; O'Brien, R.; Panizzon, MS.; Kim, K.; Bhat, R.; Grant, MD.; Toomey, R.; Eisen, S.; Xian, H.; Kremen, WS. The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry. Vol. 19. American Association for Geriatric Psychiatry; 2011. A 35-year longitudinal assessment of cognition and midlife depression symptoms: the Vietnam Era Twin Study of Aging; p. 559-570.
- Gale CR, Batty GD, Tynelius P, Deary IJ, Rasmussen F. Intelligence in Early Adulthood and Subsequent Hospitalization for Mental Disorders. *Epidemiology*. 2010; 21:70–77. [PubMed: 19907333]
- Gale CR, Deary IJ, Boyle SH, Barefoot J, Mortensen LH, Batty GD. Cognitive ability in early adulthood and risk of 5 specific psychiatric disorders in middle age: the Vietnam experience study. *Archives of general psychiatry*. 2008; 65:1410–1418. [PubMed: 19047528]
- Gale, CR.; Hatch, SL.; Batty, GD.; Deary, IJ. Intelligence. Vol. 37. Elsevier Inc; 2009. Intelligence in childhood and risk of psychological distress in adulthood: The 1958 National Child Development Survey and the 1970 British Cohort Study.; p. 592-599.
- Gjerde PF. Alternative Pathways to Chronic Depressive Symptoms in Young Adults: Gender Differences in Developmental Trajectories. *Child Development*. 1995; 66:1277–1300. [PubMed: 7555216]
- Gottfredson LS. Mainstream science on intelligence: An editorial with 52 signatories, history, and bibliography. *Intelligence*. 1997; 24:13–23.
- Guyer, AE.; Choate, VR.; Grimm, KJ.; Pine, DS.; Keenan, K. Journal of the American Academy of Child and Adolescent Psychiatry. Vol. 50. Elsevier Inc; 2011. Emerging depression is associated with face memory deficits in adolescent girls; p. 180-190.
- Gyurak, A.; Patenaude, B.; Korgaonkar, MS.; Grieve, SM.; Williams, LM.; Etkin, A. Biological psychiatry. Vol. 79. Elsevier; 2015. Frontoparietal Activation During Response Inhibition Predicts Remission to Antidepressants in Patients with Major Depression; p. 1-8.
- Hagenaars, SP.; Harris, SE.; Davies, G.; Hill, WD.; Liewald, DCM.; Ritchie, SJ.; Marioni, RE.; Fawns-Ritchie, C.; Cullen, B.; Malik, R.; METASTROKE Consortium IC for BPG, SpiroMeta Consortium, CHARGE Consortium Pulmonary Group CCA and LG. Worrall, BB.; Sudlow, CLM.; Wardlaw, JM.; Gallacher, J.; Pell, J.; McIntosh, AM.; Smith, DJ.; Gale, CR.; Deary, IJ. Molecular psychiatry. Nature Publishing Group; 2016. Shared genetic aetiology between cognitive functions and physical and mental health in UK Biobank (N=112 151) and 24 GWAS consortia; p. 031120
- Hatch, SL.; Jones, PB.; Kuh, D.; Hardy, R.; Wadsworth, MEJ.; Richards, M. Social science & medicine (1982). Vol. 64. Elsevier Ltd; 2007. Childhood cognitive ability and adult mental health in the British 1946 birth cohort; p. 2285-2296.
- Hipwell AE, Sapotichne B, Klostermann S, Battista D, Keenan K. Autobiographical memory as a predictor of depression vulnerability in girls. *Journal of clinical child and adolescent psychology : the official journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division* 53. 2011; 40:254–265.
- Horowitz, JL.; Garber, J. Journal of the American Academy of Child and Adolescent Psychiatry. Vol. 42. The American Academy of Child and Adolescent Psychiatry; 2003. Relation of intelligence

and religiosity to depressive disorders in offspring of depressed and nondepressed mothers; p. 578-586.

- Johnson W, Bouchard TJ, Krueger RF, McGue M, Gottesman II. Just one g consistent results from three test batteries. *Intelligence*. 2004; 32:95–107.
- Keedwell PA, Andrew C, Williams SCR, Brammer MJ, Phillips ML. The neural correlates of anhedonia in major depressive disorder. *Biological Psychiatry*. 2005; 58:843–853. [PubMed: 16043128]
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, Rush AJ, Walters EE, Wang PS. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003; 289:3095–3105. [PubMed: 12813115]
- Koenen KC, Moffitt TE, Roberts AL, Martin LT, Kubzansky L, Harrington H, Poulton R, Caspi A. Childhood IQ and adult mental disorders: a test of the cognitive reserve hypothesis. *The American journal of psychiatry*. 2009; 166:50–57. [PubMed: 19047325]
- Lewinsohn PM, Solomon A, Seeley JR, Zeiss A. Clinical implications of 'subthreshold' depressive symptoms. *Journal of Abnormal Psychology*. 2000; 109:345–351. [PubMed: 10895574]
- Luppino FS, de Wit LM, Bouvy PF, Stijnen T, Cuijpers P, Penninx BWJH, Zitman FG. Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Archives of general psychiatry*. 2010; 67:220–229. [PubMed: 20194822]
- Mccord J, Ensminger ME. Multiple risks and comorbidity in an African-American population. *Criminal Behaviour and Mental Health*. 1997; 7:339–352.
- McDermott, LM.; Ebmeier, KP. *Journal of Affective Disorders*. Vol. 119. Elsevier B.V; 2009. A meta-analysis of depression severity and cognitive function; p. 1-8.
- Melartin TK, Rytälä HJ, Leskelä US, Lestelä-Mielonen PS, Sokero TP, Isometsä ET. Current comorbidity of psychiatric disorders among DSM-IV major depressive disorder patients in psychiatric care in the Vantaa Depression Study. *The Journal of clinical psychiatry*. 2002; 63:126–134. [PubMed: 11874213]
- Meyer SE, Carlson Ga, Wiggs Ea, Martinez PE, Ronsaville DS, Klimes–Dougan B, Gold PW, Radke–Yarrow M. A prospective study of the association among impaired executive functioning, childhood attentional problems, and the development of bipolar disorder. *Development and Psychopathology*. 2004; 16:461–476. [PubMed: 15487606]
- Miyake A, Friedman NP, Emerson MJ, Witzki aH, Howerter A, Wager TD. The unity and diversity of executive functions and their contributions to complex 'Frontal Lobe' tasks: a latent variable analysis. *Cognitive psychology*. 2000; 41:49–100. [PubMed: 10945922]
- Moher D, Liberati A. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Annals of internal Medicine*. 2009; 151:264–269. [PubMed: 19622511]
- Papmeyer M, Sussmann JE, Hall J, McKirdy J, Peel a, Macdonald a, Lawrie SM, Whalley HC, McIntosh aM. Neurocognition in individuals at high familial risk of mood disorders with or without subsequent onset of depression. *Psychological Medicine*. 2015:1–11.
- Peterson RA, Brown SP. On the Use of Beta Coefficients in Meta-Analysis. *Journal of Applied Psychology*. 2005; 90:175–181. [PubMed: 15641898]
- Pine DS, Cohen P, Brook J, Coplan JD. Psychiatric symptoms in adolescence as predictors of obesity in early adulthood: a longitudinal study. *American Journal of Public Health*. 1997; 87:1303–1310. [PubMed: 9279265]
- Quinn ME, Joormann J. Control When It Counts: Change in Executive Control Under Stress Predicts Depression Symptoms. *Emotion*. 2015; Junio 22:1–10.
- Rawal, A.; Rice, F. *Journal of the American Academy of Child and Adolescent Psychiatry*. Vol. 51. Elsevier Inc; 2012. Examining overgeneral autobiographical memory as a risk factor for adolescent depression; p. 518-527.
- Ree MJ, Earles JA. The stability of g across different methods of estimation. *Intelligence*. 1991; 15:271–278.
- Rock PL, Roiser JP, Riedel WJ, Blackwell AD. Cognitive impairment in depression: a systematic review and meta-analysis. *Psychological Medicine*. 2014; 44:2029–2040. [PubMed: 24168753]

- Rogers MA, Kasai K, Koji M, Fukuda R, Iwanami A, Nakagome K, Fukuda M, Kato N. Executive and prefrontal dysfunction in unipolar depression: a review of neuropsychological and imaging evidence. *Neuroscience research*. 2004; 50:1–11. [PubMed: 15288493]
- Sculd MA, Knodt AR, Swartz JR, Brigidi BD, Hariri AR. Thinking and Feeling: Individual Differences in Habitual Emotion Regulation and Stress-Related Mood Are Associated With Prefrontal Executive Control. *Clinical Psychological Science*. in press.
- Sharp, C.; Petersen, N.; Goodyer, I. *Journal of the American Academy of Child and Adolescent Psychiatry*. Vol. 47. The American Academy of Child and Adolescent Psychiatry; 2008. Emotional reactivity and the emergence of conduct problems and emotional symptoms in 7- to 11-year-olds: a 1-year follow-up study; p. 565-573.
- Siegel, LS. *Journal of Learning Disabilities*. Vol. 32. Boston University; 1999. Issues in the Definition and Diagnosis of Learning Disabilities: A Perspective on Guckenberger v; p. 304-319.
- Simons CJP, Jacobs N, Derom C, Thiery E, Jolles J, van Os J, Krabbendam L. Cognition as predictor of current and follow-up depressive symptoms in the general population. *Acta psychiatrica Scandinavica*. 2009; 120:45–52. [PubMed: 19133876]
- Slykerman RF, Thompson J, Waldie K, Murphy R, Wall C, Mitchell EA. Maternal stress during pregnancy is associated with moderate to severe depression in 11-year-old children. *Acta Paediatrica*. 2015; 104:68–74. [PubMed: 25174703]
- Snyder HR. Major depressive disorder is associated with broad impairments on neuropsychological measures of executive function: A meta-analysis and review. *Psychological Bulletin*. 2013; 139:81–132. [PubMed: 22642228]
- Sørensen, HJ.; Sæbye, D.; Urfer-Parnas, A.; Mortensen, EL.; Parnas, J. *Journal of affective disorders*. Vol. 136. Elsevier B.V; 2012. Premorbid intelligence and educational level in bipolar and unipolar disorders: a Danish draft board study; p. 1188-1191.
- Spearman, C. *The American Journal of Psychology*. Vol. 15. JSTOR; 1904. 'General Intelligence,' Objectively Determined and Measured; p. 201
- Stern Y. The Concept of Cognitive Reserve: A Catalyst for Research. *Journal of Clinical and Experimental Neuropsychology (Neuropsychology, Development and Cognition: Section A)*. 2003; 25:589–593.
- Stroup D, Berlin J, Morton S, Olkin I, Williamson GD, Rennie D, Moher D, Becker BJ, Sipe TA, Thacker SB. Meta-analysis of Observational Studies in Epidemiology: A Proposal for Reporting. *JAMA*. 2000; 283:2008. [PubMed: 10789670]
- Vinberg M, Miskowiak KW, Kessing LV. Impairment of executive function and attention predicts onset of affective disorder in healthy high-risk twins. *The Journal of clinical psychiatry*. 2013; 74:e747–e753. [PubMed: 24021515]
- Zammit S, Allebeck P, David AS, Dalman C, Hemmingsson T, Lundberg I, Lewis G. A longitudinal study of premorbid IQ Score and risk of developing schizophrenia, bipolar disorder, severe depression, and other nonaffective psychoses. *Archives of general psychiatry*. 2004; 61:354–360. [PubMed: 15066893]

Figure 1. Flow Diagram of the Search Process

Meta Analysis

Figure 2. Meta-Analysis of the Association Between Cognitive Function and Subsequent Depression

A forest plot for all studies that investigated associations between cognition and later depression. Results are reported as correlation coefficients denoted by squares, and 95% confidence intervals indicated by lines (effect sizes are converted to correlation coefficients if reported as odds ratios). Meta-analysis results are displayed as the diamond. The overall analysis found a significant effect of cognition on depression ($r = -0.088$; 95% CI: $-0.121, -0.054$; $p < 0.001$).

Meta Analysis

Figure 3. Meta-analysis Comparing Values Adjusted for Baseline Depression Symptoms vs. Unadjusted

A Forest plot for analysis comparing results from a subset of the same set of studies reported in Figure 2 when using unadjusted values compared to using values that are adjusted for baseline depression. Effects are significant for unadjusted values ($r=-0.120$; 95% CI: $-0.169, -0.070$; $p<0.001$), but not for adjusted values ($r=-0.032$; 95% CI: $-0.078, 0.014$; $p=0.169$). Results are reported as correlation coefficients denoted by squares, and 95% confidence intervals indicated by lines. Meta-analysis results are displayed as diamonds.

Figure 4. Trim and Fill Funnel Plots for All Samples

Fisher's Z, a measure of effect size, is plotted on the x-axis and the standard error is plotted on the y-axis. Larger studies appear toward the top of the graph and smaller studies towards the bottom. In the absence of publication bias, the studies are symmetrically distributed around the mean. Actual studies are shown in open circles and imputed studies would be shown in black circles. No additional studies would be expected in the opposite direction (to the right) of the observed effect, but an additional 11 studies are imputed to the left of the mean effect size. While the calculated effect size is -0.088 (95% CI: $-0.121, -0.054$), using Duval and Tweedie's Trim and Fill Method, the imputed effect size estimate is larger at -0.137 (95% CI: $-0.176, -0.099$; imputed studies $n=11$).

Table 1

Characteristics of included studies. Sample size is the number of subjects included in reported analyses of interest. NR = not reported. ~ indicates only a range was reported.

Study	Name of Cohort	Type of Cohort	Sample Size	Comorbidity	How Depression Assessed	Mean IQ at Baseline	Name of Cognitive Measure	Type of Cognitive Measure	Name of Depression Outcome	Type of Depression Outcome	Covariates Included in Meta-Analyses	Percent Female	Race/Ethnicity	Continent	Mean Age at Baseline	Time Between Assessments (years)	Year of Initial Assessment
Bauer et al. 2013	Concordia Longitudinal Retirement Project	Retirees	333	NR	Study Assessment	NR	Montreal Cognitive Assessment	Verbal Comprehension, Perceptual Reasoning, Working Memory	Center for Epidemiological Studies Depression Scale (CES-D)	Continuous	Depression symptoms at Time 1	NR (both male and female)	NR	North America	59.06	4	~2005
Beautrais et al. 2013	National Longitudinal Study of Adolescent Health	Community Sample	14,322	NR	Study Assessment	100.61	Add Health Picture Vocabulary Test (AHPVT) – an abridged version of the PPVT-R	Verbal Comprehension	Created for this study	Continuous	Depression symptoms at Time 1	50	76.31% White; 16.74% Black; 4.22% Asian; 2.5% American Indian; 11.83% Hispanic; 0.05% Mixed Race	North America	15.98	7	1994
Belsky et al. 2012	Environmental Risk Longitudinal Twin Study (E-Risk)	Birth Cohort	2,123	conduct disorder, anxiety, borderline personality disorder	Study Assessment	100	Wechsler Preschool and Primary Scale of Intelligence (WPPSI)	General Intelligence, Verbal Comprehension, Perceptual Reasoning	Children's Depression Inventory (CDI)	Continuous	None	51	~90% White	Europe	5	7	1999
Betts et al. 2016	Mater University Study of Pregnancy (MUSP)	Pre-Birth Cohort	1,934	Psychosis, Mania	Study Assessment	NR	Peabody Picture Vocabulary Test-Revised (PPVT-R)	Verbal Comprehension	Composite International Diagnostic Interview (CIDI-Auto)	Continuous (latent factor)	None	NR	NR	Australia	5	16	1989
Casals et al. 2002 (females)	NR	Community Sample	99	NR	Study Assessment	NR	Academic aptitude test (AAT)	General Intelligence, Verbal Comprehension, Perceptual Reasoning	Schedules for Clinical Assessment in Neuropsychiatry	Categorical	None	100	NR	Europe	~11	6	NR
Casals et al. 2002 (males)	NR	Community Sample	100	NR	Study Assessment	NR	Academic aptitude test (AAT)	General Intelligence, Verbal Comprehension, Perceptual Reasoning	Schedules for Clinical Assessment in Neuropsychiatry	Categorical	None	0	NR	Europe	~12	6	NR
Connelly et al. 2014	Temple University Adolescent Cohort and Enrichment (ACE) Project	Community Sample	200	NR	Study Assessment	NR	Digit Span from Wechsler Intelligence Scales for Children (WISC)	Working Memory	Children's Depression Inventory (CDI)	Continuous	Depression symptoms at Time 1	56.5	45.2% White; 51.3 % Black	North America	12.41	1	NR
Der et al. 2009	US National Longitudinal Survey of Youth 1979	Community Sample	7,458	Noted a range of health conditions	Study Assessment	NR	Armed Services Vocational Aptitude Battery (ASVAB)/Armed Forces Qualification Test (AFQT)	General Intelligence, Verbal Comprehension	Center for Epidemiological Studies Depression Scale (CES-D)	Continuous	Depression symptoms at Time 1	52	31% Black; 19% Hispanic	North America	17.9	~22	1979
Dubow et al. 2008 (females)	Columbia County Longitudinal Study (CCLS)	School Children Grade 3	215	NR	Study Assessment	~104	California Short-Form Test of Mental Maturity	General Intelligence	Minnesota Multiphasic Personality Inventory (MMPI) Scale 2	Continuous	None	100	90% White; 3% Black; <1% Asian, <1% Hispanic	North America	8	11	1960
Dubow et al. 2008 (males)	Columbia County Longitudinal Study (CCLS)	School Children Grade 3	211	NR	Study Assessment	~104	California Short-Form Test of Mental Maturity	General Intelligence	Minnesota Multiphasic Personality Inventory (MMPI) Scale 2	Continuous	None	0	90% White; 3% Black; <1% Asian, <1% Hispanic	North America	8	11	1960
Evans et al. 2015	NR	School Children Grades 5-9	192	Some with ADHD	Study Assessment	111	Wechsler Abbreviated Scale of Intelligence (WASI)	General Intelligence, Verbal Comprehension, Perceptual Reasoning, Working Memory, Set-Shifting	Children's Depression Inventory (CDI)	Continuous	Depression symptoms at Time 1	52.1	71.4% White; 18.2% Black; 2.6% Asian; 3.6% Hispanic; 4.2% Mixed Race	North America	12.36	0.3	~2000s
Frantz et al. 2011	Vietnam Era Twin Study of Aging	Military Cohort	1,231	NR	Study Assessment	NR	Armed Forces Qualification Test (AFQT Form 7A)	General Intelligence, Perceptual Reasoning, Working Memory, Processing Speed, Set-Shifting, Inhibition	Center for Epidemiological Studies Depression Scale (CES-D)	Continuous	None	0	86% White	North America	20	35	~1970
Gale et al. 2008	Vietnam Experience Study (VES)	Military Cohort	3,258	Mix	Study Assessment	100	General technical section of the Army Classification Battery	General Intelligence, Verbal Comprehension, Arithmetic reasoning	Diagnostic Interview Schedule (DIS)	Categorical	Socioeconomic Status, Ethnicity, Place of service	0	80.8% White; 12.7 % Black	North America	20.4	~17	~1964
Gale et al. 2009 (1958 Cohort)	1958 National Child Development Survey	Birth Cohort	6,369	NR	Study Assessment	102.8	General ability test, devised by the National Foundation for Educational Research in England and Wales	General Intelligence	Rutter's Malaise Inventory	Continuous	None	NR (both male and female)	NR	Europe	11	22	1969

Study	Name of Cohort	Type of Cohort	Sample Size	Comorbidity	How Depression Assessed	Mean IQ at Baseline	Name of Cognitive Measure	Type of Cognitive Measure	Name of Depression Outcome	Type of Depression Outcome	Covariates Included in Meta-Analysis	Percent Female	Race/Ethnicity	Continent	Mean Age at Baseline	Time Between Assessments (years)	Year of Initial Assessment
Gale et al. 2002 (1970 Cohort)	1970 British Cohort Study	Birth Cohort	6,074	NR	Study Assessment	101.8	British ability Scale	General Intelligence	Rutter's Malaise Inventory	Continuous	None	NR (both male and female)	NR	Europe	10	20	1980
Gjerdet et al. 1995 (females)	NR	Recruited from a nursery school	51	NR	Study Assessment	118.7	Wechsler Adult Intelligence Scale (WAIS)	General Intelligence	General Behavior Inventory (GBI) Depression scale	Continuous	None	100	~66% White; ~25% Black; ~0.05% Asian	North America	18	5	1983
Gjerdet et al. 1995 (males)	NR	Recruited from a nursery school	45	NR	Study Assessment	111.12	Wechsler Adult Intelligence Scale (WAIS)	General Intelligence	General Behavior Inventory (GBI) Depression scale	Continuous	None	0	~66% White; Black; ~0.05% Asian	North America	18	5	1983
Hatch et al. 2007	1946 British Cohort Study	Birth Cohort	957	NR	Study Assessment	NR	National Foundation for Educational Research Test	General Intelligence, Verbal Comprehension, Perceptual Reasoning	General Health Questionnaire (GHQ-28)- severe depression subscale	Continuous	None	51	NR	Europe	8	45	1954
Hipwell et al. 2011	Pittsburgh Girls Study (PGS)	Community Sample	195	NR	Study Assessment	99.54	Wechsler Intelligence Scale for Children (WISC) III-R	Verbal Comprehension	Schedule for Affective Disorders and Schizophrenia - Age Children - Present and Lifetime Version (K-SADS-PL)	Continuous	Depression symptoms at Time 1	100	70% Black or multiracial	North America	11.54	1	1998
Horowitz et al. 2003	NR	Elementary school cohort	196	NR	Study Assessment	104.61	Wechsler Intelligence Scale for Children (WISC) - R	General Intelligence, Verbal Comprehension, Perceptual Reasoning	Schedule for Affective Disorders and Schizophrenia - Age Children - Present and Lifetime Version (K-SADS-PL)	Categorical	Depression symptoms at Time 1	54.2	89% White, 14.7% Black, 3.3% Other	North America	11.86	6	~1980
Koenen et al. 2009	Dunedin Cohort	Birth Cohort	730	None	Study Assessment	NR	Wechsler Intelligence Scale for Children (WISC)	General Intelligence, Verbal Comprehension, Perceptual Reasoning, Working Memory, Processing Speed	Diagnostic Interview Schedule for DSM-IV	Categorical	Sex, Socioeconomic Status, Physical Health, childhood maltreatment	48	"Primarily White"	New Zealand	9	23	1981
McCord & Ensminger (females)	Woodlawn Cohort	Elementary school cohort	346	Some with alcoholism	Study Assessment	NR, (13% of sample had scores over 110)	Unspecified IQ	General Intelligence	Composite International Interview Schedule (according to DSM-III-R)	Categorical	None	100	100% Black	North America	~6	27	1966
McCord & Ensminger 1997 (males)	Woodlawn Cohort	Elementary school cohort	313	Some with alcoholism	Study Assessment	NR, (11% of sample had scores over 110)	Unspecified IQ	General Intelligence	Composite International Interview Schedule (according to DSM-III-R)	Categorical	None	0	100% Black	North America	~6	27	1966
Meyer et al. 2004	NR	Risk Sample	86	NR	Study Assessment	~118	WISC	General Intelligence, Verbal Comprehension, Perceptual Reasoning	SCID	Categorical	None	63	99% White, 9% Black, 1% Hispanic	North America	11	13	1989
Papmeyer et al. 2015	Scottish Bipolar Family Study	Risk Sample	111	NR	Study Assessment	Controls: 108 Patients: 107	National Adult Reading Test (NART)	Verbal Comprehension	Structured Clinical Interview for DSM Disorders (SCID) symptoms	Categorical	None	52	NR	Europe	21	2	NR
Pine et al. 1997	NR	Community Sample	644	NR	Study Assessment	100.7	Unspecified "IQ"	General Intelligence	Based on Diagnostic Interview Schedule for Children	Continuous	Depression symptoms at Time 1	52	91% White	North America	13.8	9	1983
Quim & Joormann 2015	NR	Undergraduate Students	43	NR	Study Assessment	NR	n-back	Working Memory	Beck Depression Inventory II (BDI-II)	Continuous	Depression symptoms at Time 1	63	58% White, 30% Hispanic or Latino, 19% Asian, 2% African American, 2% American Indian, 2% Native Hawaiian, 2% Indian	North America	19	0.14	NR
Rawal & Rize 2012	Early Prediction of Adolescent Depression study	Risk Sample	230	None	Study Assessment	97.46	Wechsler Intelligence Scale for Children (WISC)	General Intelligence, Verbal Comprehension, Perceptual Reasoning, Working Memory, Processing Speed	Child and Adolescent Psychiatric Assessment (CAPA)	Both	Depression symptoms at Time 1, Age, Overgenerality to	59.6	NR	Europe	13.71	1	NR
Sharp et al. 2008	The Child Behaviour Study	Community Sample	439	NR	Study Assessment	105	Wechsler Intelligence Scale for Children (WISC)	General Intelligence, Verbal Comprehension, Perceptual Reasoning	The Mood and Feelings Questionnaire	Continuous	Depression symptoms at Time 1	52	97% White, 0.5% Black, 2.5% Asian	Europe	9.4	1	NR
Simons et al. 2009	East Flanders Prospective Twin Survey	Twin Registry	444	NR	Study Assessment	NR	Principal component from Stroop, Concept Shifting Test, and Letter Digit Substitution	Processing Speed, Set Shifting, Inhibition	Structured Clinical Interview for DSM Disorders (SCID)	Continuous	Depression symptoms at Time 1	100	NR	Europe	28	2	NR