Published in final edited form as:

Ann Surg Oncol. 2017 January ; 24(1): 202–210. doi:10.1245/s10434-016-5565-9.

Surgical strategy and outcomes in duodenal gastrointestinal stromal tumor

Ser Yee Lee, MD, MSc, MMed, FRCS(Ed)^{1,2,3}, Brian K. P. Goh, MBBS, MSc, MMed, FRCS(Ed)^{2,3}, Eran Sadot, MD¹, Rahul Rajeev, MBBS¹, Vinod P. Balachandran, MD¹, Mithat Gönen, PhD⁴, T. Peter Kingham, MD¹, Peter J. Allen, MD¹, Michael I. D'Angelica, MD¹, William R. Jarnagin, MD¹, Daniel Coit, MD¹, Wai Keong Wong, MBBS, FRCS^{3,5}, Hock Soo Ong, MBBS, FRCS^{3,5}, Alexander Y. F. Chung, MBBS, FRCS^{2,3}, and Ronald P. DeMatteo, MD¹

¹Department of Surgery, Memorial Sloan Kettering Cancer Center, Singapore General Hospital, Singapore

²Department of Hepatopancreatobiliary and Transplant Surgery, Singapore General Hospital, Singapore

³Duke - National University of Singapore (NUS) Medical School, Singapore

⁴Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, Singapore General Hospital, Singapore

⁵Department of Upper Gastrointestinal & Bariatric Surgery, Singapore General Hospital, Singapore

Abstract

Introduction—The surgical management of duodenal gastrointestinal stromal tumors (DGIST) is poorly characterized. Limited resection may be technically feasible and oncologically safe, but anatomical considerations may compromise the resection margins due to the proximity of critical structures, thereby necessitating more extensive resections such as pancreaticoduodenectomy.

Methods—Patients undergoing surgery for DGIST at 2 institutions from 1994–2014 were identified. Clinicopathologic and survival data were analyzed to compare outcomes in patients treated with limited or radical resection.

Results—Sixty patients underwent surgery for DGIST. Pancreaticoduodenectomy was performed in 38% while the rest underwent limited resections. The most common type of limited resection was wedge resection and primary closure (49%) followed by segmental resection with an end-to-end or side-to-side duodenojejunostomy (27%). The pancreaticoduodenectomy group tended to have larger tumors with the majority located in D2/3 (87%) and at the mesenteric border (91%). The pancreaticoduodenectomy group also had significantly greater intraoperative blood loss, longer operative time, longer hospital stay, and higher 90-day morbidity and readmission rates. The 5- year relapse-free survival (RpFS), recurrence-free survival (RFS), and overall survival (OS)

for the pancreaticoduodenectomy vs. limited resection were 81% vs. 56% ($p = 0.05$), 64% vs. 53% ($p = 0.5$), and 76% vs. 72% ($p = 0.6$), respectively. A surgical algorithm based on the location and size of the tumor is proposed.

Conclusion—Limited resection of DGIST is safe, but may be associated with lower 5-year relapse-free survival. Pancreaticoduodenectomy is recommended for selected patients with DGIST when an R0 resection cannot be performed without removing the ampulla or part of the pancreas.

INTRODUCTION

Gastrointestinal stromal tumors (GISTs) are the most common mesenchymal tumors of the gastrointestinal tract and arise from the interstitial cells of Cajal.¹ GISTs are characterized by overexpression of the KIT (CD117) tyrosine kinase receptor protein and usually harbor a gain-of-function mutation in either KIT or platelet-derived growth factor receptor alpha (PDGFR α).^{2, 3} Historically, nearly 50% of patients were deceased by 5 years after surgery, but with the development of the tyrosine kinase inhibitor imatinib mesylate (Gleevec, Novartis Pharmaceuticals), outcomes in advanced and metastatic GIST have dramatically improved.^{1, 4} The standard of care for primary localized GISTs is surgical resection. Segmental resections usually suffice since GISTs do not typically infiltrate adjacent structures, lack submucosal spread, and rarely metastasize to lymph nodes.

Although GISTs occur in the duodenum only 1–5% of the time, they account for 30% of primary duodenal neoplasms.⁵ The surgical management of duodenal GISTs (DGIST), including the optimal procedure and techniques of reconstruction, is not well defined due to the complexity of their anatomical location.^{6–10} In particular, there is ongoing debate whether radical surgery such as pancreaticoduodenectomy or a limited resection (LR) is the most appropriate procedure for DGIST.¹¹ Specifically, LR may be technically feasible and oncologically sound, but anatomical considerations may render it more challenging due to the proximity of critical structures, including the papilla, pancreas, and biliary and pancreatic ducts. Radical resection provides good oncological control and is warranted in some patients. Although pancreaticoduodenectomy is generally safe and associated with low mortality in specialized centers, its associated morbidity remains.^{11, 12} Here, we analyze the 20-year experience of the surgical management of DGIST of 2 referral centers, including technical aspects as well as short- and long-term outcomes.

METHODS

Patients with DGIST who underwent surgery with curative intent from 1994–2014 at Memorial Sloan Kettering Cancer Center (MSK) and the Singapore General Hospital (SGH) were identified. Medical records were reviewed for clinicopathological variables, comorbidities, medical treatment (e.g., imatinib mesylate or sunitinib), tumor characteristics, and operative data (e.g., type of resection and reconstruction, blood loss, transfusion, operative time). Pathological data (tumor location, size, margins, and mitoses per 50 high-power fields [HPF]) and outcomes such as hospital stay, readmissions, adjuvant therapy, recurrence, and survival were also analyzed. Approval for the study was obtained from each center's institutional review board.

The preoperative assessment included a complete history and physical examination, laboratory investigations, and in many cases endoscopy with endoscopic ultrasonography (EUS) and biopsy. Routine imaging consisted of triphasic computed tomography (CT) or magnetic resonance imaging (MRI) of the abdomen and pelvis. Patients with DGIST were discussed at a weekly multidisciplinary disease management conference, which included gastrointestinal and hepatopancreatobiliary surgeons, medical oncologists, radiologists, gastroenterologists, and pathologists. The decision for neoadjuvant therapy was made by the primary surgeon and consensus at a multidisciplinary conference. The rationale for neoadjuvant therapy was an attempt to downstage the disease to avoid a radical resection if possible. The strategy of neoadjuvant treatment was initiated in 2002. The response to neoadjuvant treatment was assessed with Response Evaluation Criteria in Solid Tumors (RECIST version 1.1) and Choi response criteria with CT at least every 3 months.^{13, 14} Adjuvant imatinib was implemented as part of a clinical trial in 2002 (MSKCC).¹⁵ Decisions on adjuvant therapy were made by the primary surgeon and the medical oncologist with a multidisciplinary tumor board. In general, tumors that were regarded as high Miettinen risk or those with R1 or close margins were considered for adjuvant therapy (SGH).

Decision for a radical resection (i.e. pancreaticoduodenectomy) or a local resection was made primarily by the attending surgeon based on a variety of factors, including tumor size, location (D1/2/3/4), and especially the proximity to the mesenteric border. Technical and oncological considerations included the proximity to the ampulla or pancreas, adequacy of margins, and feasibility of bowel reconstruction. Generally, if the tumor was large, located in D2 or D2/3, or near the mesenteric border, pancreaticoduodenectomy was chosen.

Pancreaticoduodenectomy was performed at MSK using a previously reported technique.¹⁶ The technique of pancreaticoduodenectomy and LR for GIST at SGH has also been previously described.^{9, 10} Pathological specimens were reviewed by a sarcoma or gastrointestinal pathologist. The diagnosis of GIST was based on standard criteria - consistent histological features and immunohistochemistry that included CD117.¹⁷ Locoregional recurrence was defined as the recurrence at or near the initial site of the primary tumor or resection bed. Distant recurrence was defined as peritoneal, liver, or extra-abdominal disease.

Relapse-free survival (RpFS) was calculated from the date of surgery to the first recurrence. Recurrence-free survival (RFS) and overall survival (OS) were calculated from the date of surgery to the first recurrence or death or until the time of death, respectively. Patients who did not experience the event of interest by the end of the study were censored at the time of last follow-up. Survival rates were estimated using the Kaplan-Meier method and stratified log-rank test. Analyses were performed using statistical software (SPSS software, version 21). Continuous variables were compared using the Student *t* test or Mann-Whitney test, as appropriate for the type of distribution. Categorical variables were compared using χ^2 or the Fisher exact test depending on the number of observations. Two-tailed statistical analyses were used to calculate *p* values, and *p* < 0.05 was considered significant.

RESULTS

Patient and tumor characteristics

A total of 60 patients with DGIST was included (MSK $n = 37$; SGH $n = 23$). Males comprised 52% ($n = 31$) of the group, and the median age was 58 (interquartile range [IQR]: 50–69 years). The majority ($n = 37$, 62%) underwent some type of LR while the remaining patients ($n = 23$, 38%) had a pancreaticoduodenectomy (Table 1). The most common tumor site was D2 or D2/D3 junction ($n = 38$, 63%), followed by D3 or D3/D4 junction ($n = 13$, 22%). Tumor locations are summarized in Fig. 1a.

Neoadjuvant therapy

Six (10%) patients received neoadjuvant imatinib (400mg/day), one of whom also received sunitinib (50mg/day) as second-line therapy. The median duration of neoadjuvant treatment was 6.5 months (IQR: 2–18 months), with resulting stable disease in 4 patients (67%) and partial response in 2 patients (23%) according to the Response Evaluation Criteria in Solid Tumors (RECIST version 1.1) and Choi response criteria.^{13, 14} Of these 6 patients, 3 required pancreaticoduodenectomy, 3 received LR and all achieved an R0 resection. In the 2 cases of partial response, 1 patient underwent a pancreaticoduodenectomy and 1 patient underwent LR.

Surgical techniques and complications

The most common type of LR was wedge resection and primary closure (49%), followed by a segmental resection with an end-to-end or side-to-side duodenojejunostomy (27%). The different types of LR and reconstruction are summarized and illustrated in Table 2 and Fig. 1b. Pancreaticoduodenectomy at MSKCC was performed with a pancreaticojejunostomy, while pancreaticogastrostomy was used at SGH.

There was greater intraoperative blood loss for pancreaticoduodenectomy (1000 vs. 200 ml, $p = 0.001$), although perioperative transfusions were not different. The median operating time was longer in the pancreaticoduodenectomy group vs. LR group (317 vs. 140 minutes, $p < 0.001$). Overall 90-day morbidity and Grade 3 or worse complications were higher in the pancreaticoduodenectomy group than in the LR group (70% vs. 24%, $p = 0.001$; 43% vs. 5%, $p = 0.001$, respectively). There was no 90-day postoperative mortality in either group. After pancreaticoduodenectomy, the median duration of hospital stay was longer (11 vs. 8 days, $p = 0.003$) with a significantly higher 30- and 90-day readmission rate than in the LR group (30-day: 39% vs. 3%, $p < 0.001$; 90-day: 39% vs. 11%, $p = 0.02$) (See Table 1 and Table 2).

Pathologic findings

The tumors tended to be larger in the pancreaticoduodenectomy group (median 80 vs. 45 mm, $p = 0.1$). The majority of patients who underwent pancreaticoduodenectomy had tumors located in D2 or D2/D3 junction (87% vs. 49%) and involving the mesenteric/pancreatic aspect (91% vs. 5%, $p < 0.001$). All patients in the pancreaticoduodenectomy group had an R0 resection compared to 84% in the LR group ($p = 0.1$). According to the Miettinen risk classification, 10 (43%) GISTs in the pancreaticoduodenectomy group were

considered at high risk of recurrence compared with 14 (38%) in the LR group ($p = 0.8$) (see Table 1).¹⁸

Adjuvant treatment and survival

Adjuvant therapy (imatinib mesylate, 400mg/day) was initiated in 15 patients (25%) with a median treatment length of 18.5 months (IQR: 10.5–38.2 months); 6 were in the pancreaticoduodenectomy group and 9 in the LR group (26% vs. 25%, $p = 0.9$). There was no difference in the median duration of adjuvant therapy between the 2 groups (pancreaticoduodenectomy: 18.5 months [IQR: 10.5–43 months] vs. LR: 18.5 months [IQR: 5–41 months]; $p = 0.8$).

Overall, 15 patients (25%) had experienced recurrence at a median of 72 months (RpFS). Fourteen (93%) recurrences were distant (all in the liver) and 1 (7%) was locoregional (peritoneal nodule near previous resection bed). There were no significant differences in the recurrence pattern between the pancreaticoduodenectomy and LR groups ($p = 0.46$), although the numbers were small. For the entire population, after a median follow up of 38 months (IQR: 21–72 months), the median RpFS, RFS, and OS were 72, 69, and 95 months, respectively. Compared to LR, pancreaticoduodenectomy had a trend to improved median RpFS (not reached vs. 60 months, $p = 0.05$), but similar median RFS ($p = 0.5$) and median OS ($p = 0.6$). The 5-year RpFS, RFS, and OS for the pancreaticoduodenectomy group versus the LR group were 81% vs. 56%, 64% vs. 53%, and 76% vs. 72%, respectively (see Table 1 and Fig. 2).

DISCUSSION

Duodenum GISTs are rare and surgically challenging due to the complex anatomy and intricate relationship of critical structures in a small area.¹⁹ About a third of small-bowel GISTs are located in the duodenum.^{11, 20, 21} In a large study on DGIST by Miettinen et al., the most frequent site is the second portion of the duodenum, followed by the third, fourth and first portion, similar to our findings (see Fig. 1a).²²

Surgical resection is the treatment of choice, with resection with negative margins and avoidance of tumor rupture the critical goals. The optimal procedure for DGIST remains controversial and should be tailored to the tumor location and size and patient's fitness. There are only a handful of retrospective studies directly comparing perioperative and long-term outcomes of radical resections, such as pancreaticoduodenectomy, to LR in the management of DGISTs.¹¹ However, available data suggest that in selected patients, LR is safe and oncologically comparable to radical resection.^{10, 11, 23, 24} A recent meta-analysis compared 162 patients who underwent LR and 98 patients who underwent pancreaticoduodenectomy. Tumors in patients who underwent pancreaticoduodenectomy were more likely to be larger (> 5 cm; 76% vs. 37%, OR: 5.5), located at D2 (81% vs. 29%, OR: 10), have a high mitotic count ($> 5/50$ HPF; 34% vs. 19%, OR: 2.2), and be classified as high risk (60% vs. 32%, OR: 3.2) compared to tumors in patients who underwent LR. This study also found that LR was associated with a lower postoperative morbidity rate than pancreaticoduodenectomy (20.7% vs. 48.3%, RR: 0.43) and not associated with an increased rate of local recurrence. Limited resection had better disease-free survival compared to

pancreaticoduodenectomy (HR: 2.07) and a lower rate of distant metastasis (9% vs. 26%, OR: 0.28).¹¹ In contrast, we did not find a statistical difference in RFS or OS between local resection and pancreaticoduodenectomy in our data, although there was a trend towards improved RpFS ($p = 0.05$) in favor of radical resection (see Table 1 and Fig. 2). Referral bias, difference in the follow-up period between the 2 groups, limited sample size and patient selection likely account for disparate findings.

Surgical techniques

The main concern regarding LR for GISTs is the risk of involved margins and the accompanying theoretical increased risk of local recurrence, especially for large tumors in the pancreaticoduodenal complex due to the proximity of critical structures.^{11, 25–27} Achieving wide margins in the duodenum is not always possible; margins of fewer than 5 mm after resection of duodenal tumors are not uncommon.⁹ We found that only 84% of LR were R0 resections versus 100% in the pancreaticoduodenectomy group. This may be relevant as there was a trend favoring RpFS for the pancreaticoduodenectomy group ($p = 0.05$), despite the pancreaticoduodenectomy group having larger tumors. However, this did not translate to a difference in RFS or OS between the 2 groups. These similar outcomes may be related to the use of effective adjuvant therapy such as imatinib, especially in patients with high-risk tumors.

Different techniques of LR and reconstruction for DGIST have been described according to the site and the size of the tumor.^{9, 10, 12, 23, 24} The various procedures and types of reconstruction are illustrated in Fig. 1b. A surgical algorithm based on the site and size of the tumor is outlined in Fig. 3. Currently, there is no established consensus or algorithm guiding the type of resection for DGIST. For tumors in D1/D2 proximal to the ampulla, distal gastrectomy with a loop or Roux-en-Y gastrojejunostomy is an option (Fig. 1bi). Wedge resection with primary closure can be performed for small tumors, especially those on the anti-mesenteric border regardless of location, if preservation of the ampulla is possible and the remnant bowel lumen is adequate, (Fig. 1bii).^{9, 22} For larger tumors where the resultant defect cannot be approximated without tension or compromising the bowel lumen, a Roux-en-Y duodenojejunostomy can be performed (see Fig. 1bii).^{9, 10} Segmental duodenectomy with side-to-end or end-to-end duodenojejunostomy for larger tumors located distal to ampulla at D3/4 has also been described (Fig. 1biii).⁹

For tumors in D2/D3, located at the mesenteric or medial aspect involving the ampulla or the pancreas, major resection via a pancreaticoduodenectomy or a pancreas-sparing duodenectomy may be necessary.^{28, 29} In our combined experience, the majority ($n = 38$, 63%) of the patients have tumors at D2/3, of which approximately half ($n=18$, 47%) are amenable to LR. The remaining 20 patients (53%) with tumors located at D2/3 had involvement of the mesenteric border, thus requiring a pancreaticoduodenectomy (Fig. 1biv). Overall, we achieved an overall negative margin rate of 90% (see Table 1). In this study, patients who underwent pancreaticoduodenectomy tended to have larger tumors and tumors located at D2/3 as compared to patients who underwent LR. This tendency was not statistically significant.

Role of neoadjuvant therapy

Neoadjuvant imatinib has been utilized when an extended procedure is necessary, such as a pancreaticoduodenectomy for DGIST or abdominoperineal resection for rectal GIST is required. There is uncertainty whether residual microscopic disease at the resection margin will affect outcome, particularly if the patient is treated with postoperative imatinib.³⁰ Re-resection is recommended when the site of remnant microscopic disease can be identified and when the oncological benefits of re-resection outweigh surgical morbidity. Prospective, non-randomized studies have been conducted to evaluate preoperative imatinib for treatment of locally advanced GIST.^{31–33} In these studies, most GISTs became smaller and less vascular with 2–6 months of preoperative imatinib; these results might facilitate a LR and allow organ-sparing surgery.³⁴ In our series, only 6 (10%) patients received neoadjuvant imatinib, one of whom went to sunitinib as second line therapy, with 67% having stable disease and 23% with a partial response. Three members of the neoadjuvant group (50%) required pancreaticoduodenectomy and 3 (50%) received LR. Of the 2 patients with partial responses, one required a pancreaticoduodenectomy and the other a LR; all 6 had an R0 resection. We recommend neoadjuvant therapy to be used selectively in locally advanced GISTs that are not amenable to LR or preservation of the pancreas, although clinical judgment is necessary to estimate the likelihood of conversion to a LR.

This study's retrospective, non-randomized design, with its inherent biases in areas such as patient selection, is its biggest limitation. The details of the decision making in each case were not well captured due to the retrospective nature of the study (e.g., the precise reasons why LR was chosen over a radical resection or why neoadjuvant or adjuvant therapy was administered). In general, this was primarily based on the attending surgeon's assessment and consensus at a multidisciplinary disease management conference. Due to the limited numbers and the retrospective nature of the study, we are not able to meaningfully study important factors such as Eastern Cooperative Oncology Group (ECOG) performance status, Miettinen risk, and the role of imatinib therapy (neoadjuvant or adjuvant) on outcome. However, the proportion of patients in each group who received pre or post-operative therapy was similar. The ASA class (as a surrogate of ECOG performance status) and the proportion of patients in each Miettinen risk group was also not significantly different (Table 1). Taking that information into consideration, the pattern of recurrence (local vs. distant), RFS, and OS were not significantly different in the 2 groups. The disease spectrum and management between the different institutions may also vary, but as DGIST is uncommon, the culmination of the experience of different institutions contributes to the knowledge base and clarifies the strategy of DGIST. There are limited data in the literature to unequivocally establish LR as the procedure of choice for amenable tumors, as the studies and the descriptive nature of the available data have inherent selection and publication biases as well, although LR seems to be appropriate in carefully selected patients.¹¹

CONCLUSION

While limited resection may result in lower relapse-free survival, in terms of RFS and OS it is a safe and oncologically comparable procedure for selected patients with DGIST when compared to more radical resection, probably due to this tumor type's low propensity for

lymphatic metastasis and limited submucosal spread. Limited resection has the advantage of avoiding radical surgery and its potential complications if the tumor is in an anatomically advantageous location. Our proposed surgical strategy can help guide the procedure of choice for DGIST.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Supported by National Cancer Institute grants CA102613 (RPD) and Cancer Center Support Grant P30 CA008748.

References

1. Joensuu H, DeMatteo RP. The management of gastrointestinal stromal tumors: a model for targeted and multidisciplinary therapy of malignancy. *Annu Rev Med.* 2012; 63:247–258. [PubMed: 22017446]
2. Corless CL, Fletcher JA, Heinrich MC. Biology of gastrointestinal stromal tumors. *J Clin Oncol.* 2004; 22(18):3813–3825. [PubMed: 15365079]
3. Heinrich MC, Rubin BP, Longley BJ, et al. Biology and genetic aspects of gastrointestinal stromal tumors: KIT activation and cytogenetic alterations. *Hum Pathol.* 2002; 33(5):484–495. [PubMed: 12094373]
4. DeMatteo RP, Lewis JJ, Leung D, et al. Two hundred gastrointestinal stromal tumors: recurrence patterns and prognostic factors for survival. *Ann Surg.* 2000; 231(1):51–58. [PubMed: 10636102]
5. Schottenfeld D, Beebe-Dimmer JL, Vigneau FD. The epidemiology and pathogenesis of neoplasia in the small intestine. *Ann Epidemiol.* 2009; 19(1):58–69. [PubMed: 19064190]
6. Zhou B, Zhang M, Wu J, et al. Pancreaticoduodenectomy versus local resection in the treatment of gastrointestinal stromal tumors of the duodenum. *World J Surg Oncol.* 2013; 11(1):196. [PubMed: 23945012]
7. Bourgouin S, Hornez E, Guiramand J, et al. Duodenal gastrointestinal stromal tumors (GISTs): arguments for conservative surgery. *J Gastrointest Surg.* 2013; 17(3):482–487. [PubMed: 23229887]
8. Tien YW, Lee CY, Huang CC, et al. Surgery for gastrointestinal stromal tumors of the duodenum. *Ann Surg Oncol.* 2010; 17(1):109–114. [PubMed: 19841981]
9. Goh BK, Chow PK, Kesavan S, et al. Outcome after surgical treatment of suspected gastrointestinal stromal tumors involving the duodenum: is limited resection appropriate? *J Surg Oncol.* 2008; 97(5):388–391. [PubMed: 18163461]
10. Goh BK, Chow PK, Ong HS, et al. Gastrointestinal stromal tumor involving the second and third portion of the duodenum: treatment by partial duodenectomy and Roux-en-Y duodenojejunostomy. *J Surg Oncol.* 2005; 91(4):273–275. [PubMed: 16121353]
11. Chok AY, Koh YX, Ow MY, et al. A systematic review and meta-analysis comparing pancreaticoduodenectomy versus limited resection for duodenal gastrointestinal stromal tumors. *Ann Surg Oncol.* 2014; 21(11):3429–3438. [PubMed: 24854490]
12. El-Gendi A, El-Gendi S, El-Gendi M. Feasibility and oncological outcomes of limited duodenal resection in patients with primary nonmetastatic duodenal GIST. *J Gastrointest Surg.* 2012; 16(12):2197–2202. [PubMed: 23007283]
13. Choi H, Charnsangavej C, Faria SC, et al. Correlation of computed tomography and positron emission tomography in patients with metastatic gastrointestinal stromal tumor treated at a single institution with imatinib mesylate: proposal of new computed tomography response criteria. *J Clin Oncol.* 2007; 25(13):1753–1759. [PubMed: 17470865]
14. Eisenhauer EA, Therasse P, Bogaerts J, et al. New response evaluation criteria in solid tumours: revised RECIST guideline (version 1.1). *Eur J Cancer.* 2009; 45(2):228–247. [PubMed: 19097774]

15. Dematteo RP, Ballman KV, Antonescu CR, et al. Adjuvant imatinib mesylate after resection of localised, primary gastrointestinal stromal tumour: a randomised, double-blind, placebo-controlled trial. *Lancet*. 2009; 373(9669):1097–1104. [PubMed: 19303137]
16. Fischer M, Matsuo K, Gonen M, et al. Relationship between intraoperative fluid administration and perioperative outcome after pancreaticoduodenectomy: results of a prospective randomized trial of acute normovolemic hemodilution compared with standard intraoperative management. *Ann Surg*. 2010; 252(6):952–958. [PubMed: 21107104]
17. Fletcher CD, Berman JJ, Corless C, et al. Diagnosis of gastrointestinal stromal tumors: A consensus approach. *Hum Pathol*. 2002; 33(5):459–465. [PubMed: 12094370]
18. Miettinen M, Lasota J. Gastrointestinal stromal tumors: review on morphology, molecular pathology, prognosis, and differential diagnosis. *Arch Pathol Lab Med*. 2006; 130(10):1466–1478. [PubMed: 17090188]
19. Uchida H, Sasaki A, Iwaki K, et al. An extramural gastrointestinal stromal tumor of the duodenum mimicking a pancreatic head tumor. *J Hepatobiliary Pancreat Surg*. 2005; 12(4):324–327. [PubMed: 16133702]
20. Johnston FM, Kneuert PJ, Cameron JL, et al. Presentation and management of gastrointestinal stromal tumors of the duodenum: a multi-institutional analysis. *Ann Surg Oncol*. 2012; 19(11):3351–3360. [PubMed: 22878613]
21. Colombo C, Ronellenfitch U, Yuxin Z, et al. Clinical, pathological and surgical characteristics of duodenal gastrointestinal stromal tumor and their influence on survival: a multi-center study. *Ann Surg Oncol*. 2012; 19(11):3361–3367. [PubMed: 22843188]
22. Miettinen M, Kopczynski J, Makhlof HR, et al. Gastrointestinal stromal tumors, intramural leiomyomas, and leiomyosarcomas in the duodenum: a clinicopathologic, immunohistochemical, and molecular genetic study of 167 cases. *Am J Surg Pathol*. 2003; 27(5):625–641. [PubMed: 12717247]
23. Zhang Q, Shou CH, Yu JR, et al. Prognostic characteristics of duodenal gastrointestinal stromal tumours. *Br J Surg*. 2015; 102(8):959–964. [PubMed: 25980461]
24. Duffaud F, Meeus P, Bachet JB, et al. Conservative surgery vs. duodeneopancreatectomy in primary duodenal gastrointestinal stromal tumors (GIST): a retrospective review of 114 patients from the French sarcoma group (FSG). *Eur J Surg Oncol*. 2014; 40(10):1369–1375. [PubMed: 24994075]
25. Koh YX, Chok AY, Zheng HL, et al. A systematic review and meta-analysis comparing laparoscopic versus open gastric resections for gastrointestinal stromal tumors of the stomach. *Ann Surg Oncol*. 2013; 20(11):3549–3560. [PubMed: 23793362]
26. Karakousis GC, Singer S, Zheng J, et al. Laparoscopic versus open gastric resections for primary gastrointestinal stromal tumors (GISTs): a size-matched comparison. *Ann Surg Oncol*. 2011; 18(6):1599–1605. [PubMed: 21207158]
27. Cavallaro G, Polistena A, D'Ermo G, et al. Duodenal gastrointestinal stromal tumors: review on clinical and surgical aspects. *Int J Surg*. 2012; 10(9):463–465. [PubMed: 22939976]
28. Yamashita S, Sakamoto Y, Saiura A, et al. Pancreas-sparing duodenectomy for gastrointestinal stromal tumor. *Am J Surg*. 2014; 207(4):578–583. [PubMed: 24119884]
29. Winfield RD, Hochwald SN, Vogel SB, et al. Presentation and management of gastrointestinal stromal tumors of the duodenum. *Am Surg*. 2006; 72(8):719–722. discussion 722–3. [PubMed: 16913316]
30. McCarter MD, Antonescu CR, Ballman KV, et al. Microscopically positive margins for primary gastrointestinal stromal tumors: analysis of risk factors and tumor recurrence. *J Am Coll Surg*. 2012; 215(1):53–59. discussion 59–60. [PubMed: 22726733]
31. Wang D, Zhang Q, Blanke CD, et al. Phase II trial of neoadjuvant/adjuvant imatinib mesylate for advanced primary and metastatic/recurrent operable gastrointestinal stromal tumors: long-term follow-up results of Radiation Therapy Oncology Group 0132. *Ann Surg Oncol*. 2012; 19(4):1074–1080. [PubMed: 22203182]
32. Rutkowski P, Gronchi A, Hohenberger P, et al. Neoadjuvant imatinib in locally advanced gastrointestinal stromal tumors (GIST): the EORTC STBSG experience. *Ann Surg Oncol*. 2013; 20(9):2937–2943. [PubMed: 23760587]

33. Hohenberger P, Eisenberg B. Role of surgery combined with kinase inhibition in the management of gastrointestinal stromal tumor (GIST). *Ann Surg Oncol*. 2010; 17(10):2585–2600. [PubMed: 20407930]
34. Eisenberg BL, Trent JC. Adjuvant and neoadjuvant imatinib therapy: current role in the management of gastrointestinal stromal tumors. *Int J Cancer*. 2011; 129(11):2533–2542. [PubMed: 21671474]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Synopsis

A surgical strategy for duodenal gastrointestinal stromal tumors is proposed based on a 20-year experience from 2 tertiary institutions. Limited resection is safe, but may be associated with lower relapse-free survival. Pancreaticoduodenectomy may be necessary to obtain negative margins.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Wedge resection of D2 GIST followed by primary closure if amenable, if not - with a side to side Roux-en-Y duodenojejunostomy. **iii:** Segmental resection of a D2 or D2/3 GIST with an end-to-end or side-to-side duodenojejunostomy. **iv:** Resection of a mesenteric based GIST via pancreaticoduodenectomy with either a pancreaticojejunostomy or a pancreaticogastrostomy loop reconstruction.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 2. Outcomes after surgery

A. Relapse-free survival. **B.** Recurrence-free survival. **C.** Overall survival. *1 patient in the limited resection group was excluded from the survival analysis due to missing follow-up data.

Figure 3. Algorithm for surgical management of duodenal GISTs

----->: Possible option; CT: Computed tomography; MRI: Magnetic resonance imaging; EUS: Endoscopic ultrasonography; FNA: Fine needle aspiration; D1/2/3/4: First, second, third, fourth portion of the duodenum; DJ: duodenojejunostomy

Table 1

Radical versus limited resection.

Variable	Total	Radical Resection	Limited Resection	p§
Demographics				
Number of patients	60	23(38%)	37 (62%)	NA
Male gender, n (%)	31 (52%)	15 (65%)	16 (43%)	0.1
Median age yrs, (range)	58 (50–69)	57 (49–69)	59 (50–70)	0.3
Median ASA	2 (2–2)	2 (1–2)	2 (2–2)	0.5
Median follow up, months (range)	38 (21–72)	65 (22–84)	26 (17–51)	0.03
Operative details				
Estimated Blood loss, mls, range	400 (100–1025)	1000 (500–1560)	200 (100–475)	0.001
Blood transfusion, units (range)	0 (0–1)	0 (0–1)	0 (0–0)	0.9
Operative time, mins, median (range)	235 (131–306)	317 (253–350)	140 (110–236)	<0.001
Post-operative outcomes				
90 day morbidity	25 (42%)	16 (70%)	9 (24%)	0.001
90 day mortality	0	0	0	NA
Grade 3 complications	12(20%)	10 (43%)	2 (5%)	0.001
Length of hospital stay	9 (7–14)	11 (8–19)	8 (7–10)	0.003
Patients requiring ICU stay	3 (5%)	1 (4%)	2 (5%)	1
30 day readmission	10(17%)	9 (39%)	1 (3%)	<0.001
90 day readmission	13(22%)	9 (39%)	4 (11%)	0.02
Tumor characteristics				
Median tumor size mm (range)	52 (35–88)	80 (40–95)	45 (30–72)	0.1
Site of tumor				0.08
- D1 or D1/D2 junction	7 (12%)	1 (4%)	6 (16%)	
- D2 or D2,3 junction	38 (63%)	20 (87%)	18 (49%)	
- D3 or D3,4 junction	13 (22%)	2 (9%)	11 (30%)	
- D4 or D4/Jejunum junction	2 (3%)	0	2 (5%)	
Site of tumor at mesenteric border	23 (38%)	21 (91%)	2 (5%)	<0.001
Mitotic figures/50 HPF	3 (1–10)	4 (1–15)	2.5 (1–7.25)	0.5
Miettinen risk ¹⁸ - very low / low risk	24(40%)	8 (35%)	16 (43%)	0.8
- intermediate risk	12 (20%)	5 (22%)	7 (19%)	
- high risk	24 (40%)	10 (43%)	14 (38%)	
Resection status				
- R0	54 (90%)	23 (100%)	31 (84%)	0.1
- R1	6 (10%)	0	6 (16%)	
Adjuvant therapy e.g. imatinib mesylate	15 (25%)	6 (26%)	9 (25%)	0.9
Survival Outcome				
No. of patients with recurrence	15 (25%)	5 (22%)	10 (27%)	0.7
Type of recurrence	1 (7%)	0	1 (10%)	0.46
- Locoregional	14 (93%)	5 (100%)	9 (90%)	

Variable	Total	Radical Resection	Limited Resection	p [§]
- Distant				
RpFS, months; median (95%CI)	72 (51-NR)	Not reached	60 (40–80)	0.050
5Y-RpFS, % (95%CI)	69% (52–85)	81% (62–99)	56% (29–83)	
RFS, months; median (95%CI)	69 (56–82)	72 (31–NR)	60 (40–80)	0.5
5Y-RFS, % (95%CI)	60% (44–76)	64% (43–86)	53% (28–79)	
OS months; median(95%CI)	95 (49–141)	Not reached	86 (56–116)	0.6
5Y-OS, % (95%CI)	75% (61–89)	76% (57–94)	72% (49–94)	

Continuous variables presented as: median; (interquartile range). Categorical variables presented as: number of cases (percentage). ASA: American Society of Anesthesiologist; HPF: High power field. ICU: Intensive Care unit. RpFS: Relapse-free survival. RFS: recurrence-free survival. OS: Overall survival. NR: Not reached.

[§]The univariate analysis is performed for the radical resection and limited resection groups only and not the whole cohort.

Table 2

Type of limited resection and reconstruction and complications

Limited Resection - Procedure		n (%)	
Wedge resection and primary closure		18 (49%)	
Segmental resection end-to-end duodenojejunostomy or Side-to-side Roux-en-Y duodenojejunostomy		10 (27%)	
Wedge resection with side-to-side Roux-en-Y duodenojejunostomy		6 (16%)	
Distal gastrectomy with loop or Roux-en-Y gastrojejunostomy		3 (8%)	
Type of complications (any grade)	Whole population n (%)	Radical Resection n (%)	Limited Resection n (%)
- Pancreatic fistula	3 (5%)	2 (9%)	1 (3%)
- Anastomotic bowel leak	4 (7%)	3 (13%)	1 (3%)
- Bile Leak	2 (3%)	2 (9%)	0
- Pancreatitis	2 (3%)	2 (9%)	0
- Intra-abdominal collection, abscess or infection	14 (23%)	12 (52%)	2 (5%)
- Intra-abdominal bleeding or hematoma	2 (3%)	2 (9%)	0
- Paralytic ileus, delayed gastric emptying	6 (8%)	3 (13%)	3 (8%)
- Ascites	5 (8%)	3 (13%)	2 (5%)
- Pleural effusion/ Pneumonia	11(19%)	7 (30%)	4 (11%)
- Wound infection/ Line sepsis/ Urinary tract infection	6 (10%)	3 (13%)	3 (8%)
- Acute renal failure	2 (3%)	1 (4%)	1 (3%)
- Acute coronary syndrome e.g. myocardial infarction	1 (2%)	0	1 (3%)