projecting over the edge of the table. The index finger of the left hand guides the largest curette which will pass between the Eustachian prominences into the nasopharynx and up to the septum. A few sweeps are made downwards, and the remaining fragments cleared out with Bronner's curette and the sterilised finger-nail. After the operation the patient should be kept in bed for 24 hours on cold fluids.

Light food should be ordered for a few days, and exposure to cold winds avoided. No douche or spray should be prescribed.

¹ West London Med. Jour., April.

THE PRIMARY SEAT OF INFECTION IN TUBERCULOSIS.

MUCH controversy has raged round the question of how the tubercle bacillus commonly gains entrance into the human body. The two chief theories on the subject hold respectively that the usual portal is the respiratory system and the alimentary system, while a third party considers that the lymphoid tissue of the naso-pharynx is commonly the primary nidus. Though it is quite true that any one of these localities may be the first to become infected, the important question is still outstanding as to which is the most frequent primary source of trouble. Drs. J. O. Symes and Theodore Fisher¹ have made an investigation into the matter, analysing the post-mortem reports of 500 fatal cases of tuberculosis in the hospitals of Bristol. As they point out there are several sources of fallacy to be avoided in an examination such as they have carried out. Thus the discovery of disease in a certain part of the body does not necessarily imply that the cause of the disease reached that spot by the nearest route. There is a want of resistance in certain organs and the infecting agent may pass through more resistant parts of the body to invade these weaker organs. Thus they have found acute tuberculosis limited to the peritoneum when the only old focus in the body that could be discovered was a caseous bronchial gland; and they further point out that, as with the mesenteric glands, there is no doubt that the bronchial glands can become infected without the presence of any visible lesion of the lungs or bronchi. Again, if the matter were sufficiently simple to be settled by the evidence afforded by the situation of the oldest foci of disease in the body, there is still the difficulty in many cases of deciding which are the oldest foci. Cases of tuberculosis in children in which the disease is entirely limited to one cavity of the body are considerably in the minority, and the liability to fallacy of personal judgment cannot but influence the decision as to which cavity was first attacked.

In drawing their conclusions the authors have been careful to classify all those cases as doubtful where there would be the least possibility of difference of opinion, although where there are intestinal ulcers, and a few caseous mesenteric glands of apparently recent origin, should there be old pulmonary lesions such as a cavity and perhaps also large masses of caseous bronchial and mediastinal glands, they have generally considered it justifiable to classify the abdominal disease as produced by swallowed sputum infected with tubercle bacilli, and therefore secondary to the disease in the thorax. In 102 cases of tuberculosis in subjects under 12 years of age the site of the original focus was as follows:—definitely abdominal, 11·7 per cent.; definitely thoracic, 55·8

per cent.; doubtful whether abdominal or thoracic 23·5 per cent.; other situations, *e.g.* bones, skin, etc., 8·8 per cent. That is to say that the proportion of abdominal to thoracic cases was 1 : 4·7. Subjects under two years of age showed, however, a much higher proportion of abdominal cases. Curiously enough the proportion of definitely abdominal to definitely thoracic cases appeared to be almost as great in the second twelve years of life as in the first twelve years, though after the age of 24 the proportion of abdominal cases greatly decreased. It is probable, as the authors point out, that their percentages cannot be taken to represent the exact relative frequency of abdominal infection, as cases of abdominal tuberculosis are more likely to gain admittance to hospitals than are those where the lungs are principally affected, and the ratio of the former to the latter is therefore probably not so high as their statistics would at first sight seem to indicate.

¹ Brit. Med. Jour., April 16.

COMPLICATIONS AND SEQUELS OF PROSTATECTOMY.

IN view of the growing inclination that appears to exist towards recommending patients suffering from enlarged prostate to undergo radical treatment without waiting for the more serious symptoms to appear, it is well to consider what are the drawbacks to the operation. It is clear from the situation of the organ that its removal is not and never can be a trivial affair. Although the immediate mortality in careful hands is small, yet it is far from negligible, and, moreover, in a certain percentage of those who recover complications and sequelæ are almost certain to arise and detract from the success of the operation. In discussing this aspect of the matter Dr. J. E. Moore¹ points out that the most frequent cause of death following radical treatment is uræmia, consequently the most important preliminary examinations are those of the kidneys. Unfortunately, the enlarged prostate and inflammation of the bladder render it difficult, or impossible, to ascertain the condition of the kidneys; but the surgeon is justified in operating so long as there is no positive evidence of renal complications. The next greatest danger is sepsis, and in the presence of an infected bladder it is an ever-present one. The patients in whom the operation is required usually have not so much resistance against wound infection as younger men. All that can be done to counteract this is to thoroughly cleanse the wound, and to provide free drainage. The complication of cystitis is often at the present time the most positive indication for operation, and the only satisfactory treatment for it is drainage. Stone is a frequent complication of enlarged prostate and should always be sought for during prostatectomy. Cases are on record in which a calculus has been overlooked, and as a consequence the operation has been a failure. Such a mistake is by no means unlikely in perinaeal prostatectomy unless a deliberate search is made for stone. The incision, when too extensive, may cause various complications. In the suprapubic operation it may injure the peritoneum, and in the perinaeal one important muscles, blood-vessels, and nerves may be damaged. Ordinarily the peritoneum is easy to avoid, but with a contracted bladder the danger is great. Hæmorrhage is not a common complication;

when it does occur it is best controlled by ligature or forcipressure, as packing causes severe pain and may produce sloughing.

Too little effort is made, Dr. Moore believes, to protect the urethra, especially in operating by the perinaeal route. It simplifies the operation wonderfully to sacrifice the whole prostatic urethra, as is frequently done; and it is marvellous how quickly nature compensates for its loss; but it is not reasonable to suppose that traumatic stricture will not occur later, and a man with a traumatic stricture at the neck of his bladder may be in a worse condition than one with an enlarged prostate. The most difficult problem in the perinaeal operation is to avoid injuring the rectum. There are few, if any, operators who have never met with this accident. It is not a disastrous event when it does happen, though it is

¹ *Annals of Surgery*, April 1904.

always a serious complication. There is another drawback to prostatectomy as usually performed, namely, the sacrifice of the seminal ducts. An occasional sequel of removal of the prostate is dribbling of urine owing to injury of muscles or nerves at the neck of the bladder, but the condition can be avoided by care and gentle operating. A perinaeal fistula calling for a plastic operation is another occasional sequel, and may be caused by too protracted draining. It is rarely necessary to keep a tube in the bladder for more than a week. Epididymitis is a quite frequent consequence of the operation, and may become so troublesome as to require removal of the epididymis. Dr. Moore's opinion is that the operation of prostatectomy is altogether too grave a one to be resorted to as a routine treatment for every enlarged prostate.

PROGRESS IN BACTERIOLOGY.

Malaria.—Mary E. Rowley¹ states that in æstivo-autumnal malaria in addition to the ring shaped bodies, crescents, and ovoids, there are to be found two other varieties of parasites. One is an intra-corporal, elongated, sausage-shaped parasite with openings at the extremities, scattered pigment which is sometimes grouped at the end, and with dotted chromatin parallel to the long axis. The other a form of parasite intermediate between the crescent and the foregoing; curved, with the chromatin in dots at one end, and the pigment at the periphery. She inclines to the belief that these sausage-shaped bodies are developed from ring forms and are on their way to form crescents. Leishman² has recently demonstrated that important diagnostic points between the various types of malaria are brought out by Romanowsky's stain, especially if deep staining be employed. Thus, staining of moderate intensity in cases of tertian fever brought out Schüffner's dots in the red cells invaded by the parasite and this change was characteristic of a tertian invasion. Quartan fever did not affect the red cells in any such way. Malignant cases, he thought, might possibly be due to invasion by two varieties of parasites, one exciting malignant quotidian fever and the other exciting malignant tertian fever. Certainly the results obtained by staining were contradictory, for whilst it was often possible to demonstrate the dots in the red cells infected by young malignant rhizoids, and deep-stained capsules surrounding the mature gametes or crescents, yet in other cases no such changes could be detected although the same staining methods were employed. Leishman calls attention to forms which seem to indicate that crescents may be derived from a circular intra-corporal parasite similar to the gametes of benign tertian and quartan malaria.

Rheumatoid Arthritis.—Poynton and Paine³ have brought forward evidence to show that the intravenous injection of a diplococcus obtained from a joint affected by rheumatoid arthritis may excite in rabbits not only a multiple arthritis but a monarticular osteo-arthritis. In a later paper⁴ they state that the organism is indistinguishable from the diplococcus rheumaticus, and that from a second case they were able to excite in rabbits lesions which were both rheumatic and rheumatoid in character. Hale

White⁵ has put on record an acute case of osteo-arthritis in which a diplococcus was isolated from the knee-joint, but he does not regard it as identical with the diplococcus rheumaticus. Von Dungern and Schneider cultivated a diplococcus from a case of rheumatoid arthritis, and cultures inoculated into rabbits produced typical joint lesions; but, as Hale White points out, their results are discounted by the fact that the case had recently suffered from rheumatic fever. In White's case in addition to the diplococcus which was found in cultures made from a scraping of an affected knee-joint, there was also found in an enlarged mesenteric gland a diplobacillus, but neither of these organisms were pathogenic to animals. The diplococcus did not form chains, grew well on the ordinary media, produced acid but not gas, did not coagulate milk, and was a facultative anaerobe. It stained by Gram's method.

Acute Rheumatism.—Meyer,⁶ whose previous work on the tonsillitis of acute rheumatism confirmed the theory of a specific diplococcus as described by Triboulet, has made a further investigation into this matter. He failed in many attempts to isolate the diplococcus from the blood or joint fluid, but was successful in obtaining it from swabbings of the tonsils of 26 cases of rheumatic angina. Inoculated into rabbits the diplo-streptococcus gave rise to conditions of arthritis and endocarditis. There are, undoubtedly, great difficulties in isolating the organism from the blood, joint fluids, or urine. McCrae⁷ failed to obtain it in 270 consecutive cases investigated at the Johns Hopkins Hospital, and Phillipp⁸ failed in 30 cases at Prague. Sahli found a coccus morphologically resembling the staphylococcus citreus. He regarded the joint lesions as being due to multiple local infection by the bacteria. Schmidt⁹ has tried Menzer's anti-rheumatic serum in three cases of acute rheumatic fever, eight of subacute and four of chronic rheumatism. He injected 15 to 20 c.cms. in the neighbourhood of the inflamed joint, and was of opinion that benefit was obtained especially in subacute cases which did not yield to ordinary remedies. The reaction with redness, swelling, and erythema he believes to be beneficial. Poynton¹⁰ summarises Menzer's serum treatment of rheumatic fever as follows: The serum which is anti-bacterial is obtained from cultures