doses in order to get a high percentage in the urine. Two cases have recently come under our observation, in which hæmaturia followed 10-grain doses of urotropin, and in one of these instances the hæmorrhage was profuse. Dr. W. Coleman,¹ as the result of an extended search into the literature of the subject, mentions the following poisonous manifestations due to the drug. (1) Irritation of the stomach as shown by impairment of digestion and slight malnutrition. (2) Diarrhœa and slight abdominal pain. (3) Irritation of skin, with a rash resembling that of measles. (4) Headache and tinnitus aurium. (5) Albuminuria. (6) Irritation of the bladder, shown by strangury. (7) Hæmaturia and hæmoglobinuria. To lessen the chance of producing poisonous symptoms the drug should be freely diluted, five grains being given in half a pint of water; and if fixed rules are in any case desirable in therapeutics this is a harmless and useful one to follow, for it insures against producing an excessive percentage of urotropin in the patient's urine. Personally we have never ordered a patient more than ten-grain doses and we have had reason to be satisfied with the results, although M. Lubowski² characterises such a quantity as "remarkably small" and as the "lowest effective dose."

¹ Med. News, 1903, p. 894. ² Therapist, Aug. 15.

THE DISTRIBUTION OF FAT AND STRYCHNINE IN NUX VOMICA SEEDS.

In a paper read before the British Pharmaceutical Conference at Sheffield, Messrs. H. Wippell Gadd and Sydney C. Gadd called attention to some experimental work which they had carried out with a view to obviate the difficulties which are experienced in the manufacture of the official liquid extract of nux vomica on account of the fat which the seeds contain.

After steaming a quantity of nux vomica seeds to soften them, they passed them through a Carter's disintegrator. The centrifugal action of the disintegrator separated the light hairs from the heavier particles which make up the bulk of the seeds, and it was thought that it might be advantageous to reject these hairs, as they probably contained more fat than did the remainder of the seeds.

Before doing so, samples of the hairs (which amounted to about 5 per cent. of the whole) and of the residue of the powder were taken and examined by the following methods: (1) A portion of each

sample was freed from fat by prolonged treatment in a Soxhlet tube with boiling ether, with the following results:—

Percentage of fat found in hairs	7
Percentage of fat found in remainder of the seeds	2
Percentage of fat in whole seeds (by calculation)	2.25

Another portion of each sample was examined for strychnine by Bird's process¹ with the following results:—

Percentage of strychnine in hairs	0.5
Percentage of strychnine in remainder of seeds	1.5
Percentage of strychnine in whole seeds (by calculation)	1.4

From these results it is obvious that by rejecting the hairs a considerable portion of the fat would be removed, whilst the loss of strychnine would be

inmaterial. A liquid extract was therefore made from the seeds from which the hairs had been removed. The liquid extract was made by the official process, but the percolation was continued until the product ceased to give a precipitate with the usual alkaloidal reagents. The weaker percolates were concentrated, and the finished product adjusted to the strength of 1.5 per cent. The amount of fat in the liquid extract was then estimated by the following method: The extract was evaporated almost to dryness, and the residue ground with powdered glass. The fat was extracted by treating the mixture with boiling ether in a Soxhlet tube, with the following result:—

Percentage of fat in liquid extract 0.62

It was therefore evident that the amount of alcohol (70 per cent), which was sufficient to extract the whole of the strychnine present in the seeds (freed from hairs), only removed one-third of the fat. The liquid extract made thus was of excellent quality and free from any objectionable amount of fat. Further experiments showed that the skins of the seeds contain very little fat, and therefore there is no need to reject the skins as well as the hairs.

Summing up, the authors state that (1) the hairs of nux vomica seeds contain proportionately much more fat and less strychnine than do the inner portion of the seeds, and it follows that (2) the hairs can be rejected with advantage before making the liquid extract.

¹ Pharmaceutical Jour., 1900, vol. ii., p. 574.

RHEUMATISM IN CHILDHOOD.

WHILE everyone is agreed that the manifestations of rheumatism in children are not limited to chorea and arthritis, there is considerable difference of opinion as to the number of ailments which ought rightly to be designated rheumatic. Dr. James Burnet¹ takes rather a comprehensive view of the subject. He thinks that pharyngitis and tonsillitis should be more frequently regarded as rheumatic than they are at present; and he states that in some cases where sore throat is the chief or sole complaint careful light percussion of the heart shows it to be somewhat dilated, while judicious questioning may elicit the complaint of general soreness all over the body. Many of these cases yield rapidly to anti-rheumatic treatment, and the patient should be kept in bed until all doubts about the nature of the case have been dispelled. He also thinks that pneumonia may sometimes be a manifestation of rheumatism, and refers to a case where a child developed chorea shortly after an attack of pneumonia. Psoriasis also may have a direct connection with rheumatism for it most commonly develops in rheumatic subjects and is benefited by anti-rheumatic remedies. Appendicitis, he thinks, is undoubtedly a rheumatic affection in not a few cases, and it is moreover one which is frequently unrecognised. Sometimes the symptoms are clearly attributable to the appendix, but in others there are merely recurrent attacks of abdominal pain relieved by simple treatment. Pain in the side or in the upper part of the chest is a common complaint in childhood and is often quickly relieved by anti-rheumatic remedies. Lastly, children of rheumatic heredity do not always present any definite manifestation of the disease.