TUMOURS OF THE BREAST.

MR. ERNEST SHAW,¹ in a paper read before the Abernethian Society, gives a classification of breast tumours in which he combines the practical division, into innocent and malignant, with the more exact method based on histological characteristics. He deals seriatim with the various tissues entering into the formation of the mammary gland, and describes the different growths which may originate in them. The epithelial cells may grow (a) outwards or (b) inwards. In the former case the new growth may imitate closely the normal gland characters and a fibroadenoma result, or disorderly growth of cells may break through the basement membrane and invade adjacent structures, forming a carcinoma.

In case (b) an orderly growth is a duct papilloma; a disorderly invasion is a duct carcinoma.

Fibroma and sarcoma grow from the fibrous tissue of the breast; lipomata, angiomata, and perhaps neuromata may occur in connection with the fatty, vascular, and nervous elements. Secondary carcinomatous and sarcomatous nodules are sometimes found in the breast. Combinations of separate tumour formations are occasionally met with. One breast contained a sarcoma, a fibroadenoma and a nearly pure fibroma.

Mr. Shaw draws careful distinctions between the different forms of cysts with intracystic growths which, in the breast, cause so much confusion because of inaccurate descriptions and uncertain nomenclature. A cyst formed from a dilated duct may contain an adenomatous mass, the gland tubes of which have dilated; the intervening fibrous tissue broken down, and the broken ends projecting freely into the enlarged glandular spaces thus formed, resemble the processes of the next form, duct papilloma.

Duct papilloma is an epithelial ingrowth into the lumen of a duct. Cystic tumours occur near the nipple, they contain a little bloodstained fluid and a soft reddish mass attached to one wall and composed of delicate processes of loose fibrous tissue, and blood-vessels, the whole covered by a layer of columnar cells. Bleeding from the nipple and a tendency to malignant degeneration characterise these two growths. Duct carcinoma consists of a distinct papillary ingrowth within small cysts, and an infiltrating columnar-celled downgrowth; perhaps a malignant stage of duct papilloma.

Intracystic adenofibroma consists of gland tissue in the form of cystic spaces and masses of connective tissue polypoid in form, and either cellular and soft or fibrous and hard projecting into these cysts. This growth may rapidly progress and intracystic adenocarcinoma result. In one part of such a tumour there will be cysts with fibrous or myxomatous intracystic growths, in another part the cysts have disappeared, and a solid softish mass mottled with extravasated blood is found.

Occasionally there is a smooth thick-walled cyst into the cavity of which projects a small firm irregular nodule of carcinoma. It is not rare to find carcinoma beginning in the outer part of the wall of an apparently simple mammary cyst. The correct surgical treatment of cysts occurring in the breast, judging by pathological experience, is excision rather than incision and drainage.

¹ St. Bart's Hosp. Jour., June 1904.

VERONAL: A NEW HYPNOTIC.¹

VERONAL is a colourless crystalline substance, sparingly soluble in cold water, readily soluble in hot water. It is most conveniently given in tablets, which can be dissolved in hot milk. The doses range from 4 to 40 grains, but in most cases a dose of 7½ grains is sufficient to produce sleep within an hour, and lasting some seven or eight hours with, perhaps, an occasional interruption. The smaller doses were found to act admirably in weakly women suffering from advanced phthisis or from heart disease. When there is actual physical discomfort 15 grains may be needed to produce sleep, and in cases with severe pain, such, for example, as acute rheumatism or sciatica, veronal, like most of its fellows, cannot be successfully substituted for morphine. In the active mental excitement of delirium tremens, general paralysis, or epilepsy as much as 50 grains have been ordered with good effect, the patients not only being promptly calmed, but continuing in a more or less sleepy condition the whole of the following day. In some few instances a slight antipyrin-like rash has been observed, but with this exception veronal appears to be free from unpleasant after-effects. It has no action on the great vital centres, nor does it disturb the blood-pressure. It appears, also, that it does not create any "want" in the system, as when taken repeatedly the dose has rather to be diminished than increased, and there seems to be little or no tendency to the establishment of a veronal habit.

¹ Alfred C. Jordan, M.D. Brit. Med. Jour., March 5, 1904.

THE DETERIORATION OF VISION DURING SCHOOL LIFE.

IN A paper read at the International Congress of School Hygiene at Nuremberg, Dr. Ettie Sayer¹ reported that the result of an investigation into the condition of the eyesight of children attending the elementary schools in London showed that 10 per cent. of the boys and 11 per cent. of the girls had a visual acuity of $\frac{6}{18}$ or less. These figures refer to children from 8 to 14 years of age. When the tests were applied to younger children, that is, from 6 to 8 years of age, it was found that 81 per cent. could read $\frac{6}{6}$ with each eye separately, and only in 3.5 per cent. were there defects sufficiently serious to reduce the vision as low as $\frac{6}{18}$. That is, stated broadly, the serious defects among children of 6 years of age are only 3 per cent., whereas at 11 years the proportion has reached 11 per cent., after which the figures are somewhat less. To avoid this deterioration of vision during school life, Dr. Sayer suggests that all children should be tested at the outset, and that where vision is below the normal standard atropine should be employed and the refraction estimated. When there is excessive hypermetropia *plus* glasses should be worn until this is outgrown. Commencing myopia raises the question whether the child should be allowed to enter on any form of near work, and in any event a special curriculum with a minimum of near work should be ordered. All young children should be preserved from small type and small writing. As the law demands compulsory education, it ought to provide means of protection

for the many thousands of children whose eyesight and general health are prejudiced by the effort of accommodation involved in learning lessons for many hours every day of their school lives.

Another contribution bearing upon the effect of prolonged near work on vision comes from the anthropometrical laboratory of the University of Aberdeen.² Dr. G. Stoddart and Dr. C. H. Usher have conducted a series of observations on the eyes of 400 students and have tabulated the results. From these it appears that as regards acuteness of vision the Aberdeen students compare favourably with the students of certain foreign universities, and that in particular the proportion of myopes is much smaller. Thus in Aberdeen these numbered 18·7 per cent., as compared with 28 per cent. in some American universities, and 59 per cent. at the University of Breslau. An analysis of the observations also shows that myopes are more frequent in town-bred than in country-bred, students. Contrary to the general creed the average size of the pupil in the hypermetropes was found to be larger than in the myopes. Light-coloured eyes were found to have a slightly larger pupil than dark or grey-coloured eyes. Acuteness of vision did not appear to have any definite relationship either to size of pupil or colour of iris. Nearly 3 per cent. of the students tested were found to be colour-blind.

¹ Brit. Med. Journ., June 18th, 1904. ² Lancet, June 4th, 1904.

THE TREATMENT OF INEBRIETY BY ATROPINE.

DR. C. A. MCBRIDE¹ reports a number of cases of inebriety treated by atropine and expresses the conviction that a complete recovery with absence of all craving for stimulants may generally be expected after a six weeks' course of treatment. Long periods of detention are in the majority of cases wholly unnecessary. Dr. McBride, in addition to atropine, uses strychnine and a fluid extract of cinchona bark. He insists upon the need for the personal supervision of the medical attendant, and allowing for peculiarities of individual cases, advises:—1. Atropine sulphate (hypodermically) $\frac{1}{200}$ gr. thrice daily, increased during the first week until the throat is dry and the pupils are affected; in most cases $\frac{1}{10}$ gr. is found to be the maximum dose. 2. Strychnine nitrate given at the same time as the atropine, and also hypodermically, $\frac{1}{60}$ gr. increased to $\frac{1}{20}$ gr. 3. Fluid extract of red cinchona every three hours by the mouth. Warm baths, regulated exercises, and other measures affecting the general health are auxiliary influences of importance. Close personal attention is necessary on the part of the physician, and the treatment is hardly likely to be efficiently practised unless the patient is under the discipline of a sanatorium or other suitable institution.

¹ Brit. Med. Jour., April 30, 1904.

PROGRESS IN MEDICINE AND SURGERY.

DISEASES OF THE HEART AND CIRCULATION.

(Concluded from page 264.)

Left Auricle and Pulmonary Veins.—In an early stage of development the left auricle consists of auricular canal and auricle proper, and the pulmonary veins enter the auricle by a single trunk. In the course of development the heart sinks backwards between the lungs and the lungs grow forward round the heart, and the pulmonary veins come to open separately into the auricle. They are surrounded by circular fibres, acting as sphincters, derived from the musculature of the original vestibule. The difference between the symptoms of mitral stenosis and regurgitation is due to the fact that in stenosis the highest pressure within the auricle occurs during systole of that chamber, *i.e.* when the pulmonary orifices are closed, whereas in mitral incompetence the highest pressure takes place when the walls of the auricle are relaxed in diastole and the pulmonary veins stand wide open so that the ventricular wave of pressure can spread right within the venous system of the lung. In systole of the left auricle the wave of contraction spreads from the orifices of the pulmonary veins to the apex of the appendix. The negative pressure in the auricle during its diastole, is due, as on the right side, to the systole of the ventricle during which its posterior base moves downwards.

The Bulbus Arteriosus is surrounded, during early fetal life, by a considerable musculature. This bulbus separates into pulmonary artery and aorta,

and whilst the musculature disappears from the aorta it persists as loops which encircle the termination of the infundibulum of the right ventricle. These fibres are of importance normally in rendering the pulmonary valves competent and probably play a part in causing the systolic bruit of anæmia. For in anæmia the infundibulum becomes dilated and the muscular fibres form a functional stricture round the base of the pulmonary artery.

Irregularity of the Heart.—Mackenzie² has published a large number of tracings tending to show that the condition known as persistent irregularity of the heart or delirium cordis is due to the rhythm of the heart proceeding from the ventricles and not, as normally, from the great veins as they debouch into the auricles. The evidence is as follows: Sphygmographic tracings were taken of the jugular vein and the radial artery. On inspection of the former these two waves are normally seen, one due to the auricular systole and the second to the ventricular systole. Synchronous with the latter the arterial tracing shows a wave—the radial pulse. Now in cases of irregularity it is found that there occurs on the radial tracing a small wave synchronous with the auricular wave in the jugular vein, followed by a long pause, whereas in the jugular tracing there is no indication of the normal ventricular wave. Evidently, the normal peristaltic wave of contraction beginning in the auricles finds the heart muscle refractory on arriving at the ventricles, owing to the fact that a ventricular contraction has taken place before its time, *i.e.* the ventricle initiates its own contraction. The same