Published in final edited form as:

Tetrahedron Lett. 2016 November 16; 57(46): 5047–5049. doi:10.1016/j.tetlet.2016.10.009.

Ochrocephalamine A, a new quinolizidine alkaloid from *Oxytropis ochrocephala* Bunge

Li-Na Liu^{1,3,||}, Jian-Qiang Ran^{1,||}, Li-Jun Li¹, Yu Zhao⁴, Masuo Goto⁴, Susan L. Morris-Natschke⁴, Kuo-Hsiung Lee^{4,5,*}, Bao-Yu Zhao^{2,*}, and Cheng-Jian Tan^{1,*}

¹School of Ethnic Medicine, Guizhou Minzu University, Guiyang 550025, China

²College of Veterinary Medicine, Northwest A&F University, Yangling, Shaanxi, 712100, China

³Department of Biotechnology, Guizhou Medical University, Guiyang 550025, China

⁴Natural Products Research Laboratories, UNC Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, NC 27599, United States

⁵Chinese Medicine Research and Development Center, China Medical University and Hospital, Taichung 401, Taiwan

Abstract

One dimeric matrine-type alkaloid, ochrocephalamine A (**1**), was isolated from the poisonous plant *Oxytropis ochrocephala* Bunge. Its structure was elucidated by spectroscopic data and single-crystal X-ray diffraction. The insecticidal and cytotoxic activities of **1** were evaluated.

Graphical Abstract

*Corresponding Author, Phone: (+86) 851-83610313. tcj1229@163.com, zhaobaoyu12005@163.com, khlee@unc.edu.

||These authors contributed equally to this work

Supplementary Material

NMR spectra, bioactivity assay, and the X-ray crystallographic data (CIF) of **1** are available free of charge via the Internet at <http://>

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Poisonous plant; *Oxytropis ochrocephala* Bunge, *Oxytropis ochrocephala*; Alkaloid; Quinolizidine alkaloid

The plant *Oxytropis ochrocephala* Bunge is a common poisonous plant found in the western grasslands of China^{1–2}. After ingesting the plant, livestock gradually develop a chronic neurological disease, characterized by a staggering gait and muscular incoordination, giving rise to the vivid name ‘locoweed’ for this poisonous plant³. Previous phytochemical investigations on *O. ochrocephala* provided quinolizidine and indolizidine alkaloids, as well as flavonoids and saponins^{4–10}.

Quinolizidine alkaloids are important natural products exhibiting a broad array of pharmacological activities, such as antineoplastic, antibacterial, antiviral, and insecticidal properties¹¹. Recently, we reported the synthesis of sophoridine derivatives as potential antitumor agents¹². As part of our ongoing research program to identify bioactive constituents, particularly from this alkaloid class, we obtained a new quinolizidine alkaloid, ochrocephalamine A (**1**), from *O. ochrocephala*. Herein, we describe the isolation, structural elucidation and bioactivity of this compound.

The whole plant (20.0 kg) of *O. ochrocephala* was percolated three times with 95% EtOH to give a crude extract (3.0 kg). The extract was concentrated to dryness under reduced pressure, followed by partitioning between CH₂Cl₂ and 2% HCl. The aqueous phase was then adjusted to pH 11 with 3% NaOH and extracted with CH₂Cl₂ to give crude alkaloids (100 g). The crude alkaloids were subjected to a silica gel column chromatography eluted with CH₂Cl₂/MeOH (1:0 to 0:1) to obtain fractions A, B and C. Fraction B was chromatographed on silica gel [CH₂Cl₂/MeOH (10:1)] and then RP-18 (30% MeOH) columns to yield ochrocephalamine A (**1**, 20 mg).

Ochrocephalamine A (**1**)¹³ was obtained as a light-yellow solid (CH₂Cl₂), and its molecular formula was established as C₃₀H₄₂N₄O₂ by HR-EI-MS (*m/z* 490.3296 [M]⁺, calcd 490.3308), requiring 12 degrees of unsaturation. The ¹H NMR data of **1** (Table 1) showed two groups of signals characteristic for a matrine-type alkaloid at δ_{H} 4.29 (1H, dd, *J* = 12.8, 4.0 Hz) and 3.11 (1H, *brt*, *J* = 12.8 Hz) and at 4.09 (1H, dd, *J* = 12.8, 7.2 Hz) and 3.80 (1H, *brt*, *J* = 12.8 Hz)¹⁴. In addition, two coupled olefinic protons [δ_{H} 7.30 (1H, d, *J* = 7.6 Hz), 6.51 (1H, d, *J* = 7.6 Hz)] were distinguished by NMR analysis. The ¹³C-NMR data (Table 1) accounted for 30 carbon signals, classified as four sp² quaternary carbons including two lactam-CO signals at δ_{C} 170.9 and 164.6, two sp² and eight sp³ methines, and 16 methylenes. Six sp² carbons accounted for 4 out of the 12 degrees of unsaturation, and the remaining 8 degrees required that **1** is an octacyclic system.

Comprehensive analysis of the 1D NMR data suggested that **1** contains two matrine-type alkaloid moieties, sophoramine and matrine¹⁵. Considering the octacyclic ring system of **1**, this compound was likely a dimeric alkaloid of sophoramine and matrine connected through a C-C bond. Accordingly, the methine of sophoramine at C-14 (δ_{C} 116.4) and methylene of matrine at C-13 (δ_{C} 19.2)¹⁵ were replaced by the quaternary carbon (δ_{C} 130.1) and methine

(δ_C 30.0), respectively, in **1**. The HMBC correlations from H-13 to C-15, C-11 and C-13', H₂-14' to C-15', C-13' and C-14, H-13' to C-15' and C-14', and H₂-12' to C-7', C-11', C-13' and C-14 confirmed the C-14–C-13' connection. Thus, the gross planar structure of ochrocephalamine A was established as shown (Figure 2a).

Determinations of the stereo configurations of **1** were particularly challenging. The ROESY correlations of H-17' α to H-5', H-6' to H-7' indicated H-5', H-6' and H-7' were α configuration¹⁰. H-11 α was assigned in β -configuration on the basis of the mutual ROESY between H-17' β and H-11'¹⁰. The relative configuration of H-13' could not be determined by a ROESY experiment due to unobserved correlations between H-11' and H-13' as well as H-11' and H-14'. Additionally, the relative configuration of H-5, H-6 and H-7 also can't be determined by ROESY experiment due to unobserved correlations between one of them with another confirmed proton configuration. Thus, we attempted to prepare crystals of **1** for X-ray determination. Eventually, a single crystal of this compound was obtained from EtOH. The connection between C-14 and C-13' was confirmed by X-ray crystallographic diffraction with CuK α radiation and the relative configurations of H-5, H-6, H-7 and H-13' were α -configuration (Figure 2b)^{13,16}. The absolute configurations of **1** were also determined by the refined Hooft parameter value 0.01(6) for 1892 Bijovet pairs with a probability of 1.000 as shown in Figure 2b¹⁶.

From a biogenetic point of view, matrine could react with known sophoridine through a Michael addition reaction to give the intermediate B. Aromatization of this intermediate would then yield **1**.

We evaluated the contact toxicity of **1** against *Spodoptera litura* third-instar larvae by the micro-drip method. One μ L of **1** (3 mg/mL in EtOH) was dropped on the pronotum of larvae. After five days, 50% of larvae were killed. In cytotoxicity assays, compound **1** was not active against A549, KB, KB-VIN, and MDA-MB-231 cell lines (ED₅₀ > 20 μ M).

O. ochrocephala is widely distributed in the grasslands of eastern and central Asia, Australia, and western North and South America, where it poses an extreme risk to grazing animals, resulting in a chronic neurological disease^{1,3}. The indolizidine alkaloid swainsonine is claimed as the constituent poisonous to livestock³. In view of the plant itself, poisonous secondary metabolites can improve the plant's chances of survival and reproduction by deterring animals, such as sheep, cattle, and horses. In addition, quinolizidine alkaloids^{8–9, 17}, such as **1**, play an important role in defense against insects. Biogenetically, indolizidine and quinolizidine alkaloids originate from the same precursor L-lysine¹⁸. The use of one biogenetic pathway to produce two different types of alkaloids involved in plant defense against both livestock and insects, respectively, is a smart and efficient strategy to combat the survival challenge.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The work was supported financially by the Special Scientific Research Fund of Agriculture Public Welfare Industry (Grant No. 201203062), the Chinese National Natural Science Foundation (31360084, 31660103), and Guizhou High-level Innovative Talent to Chengjian Tan. Partial support was supplied by NIH Grant CA177584 from the National Cancer Institute awarded to Kuo-Hsiung Lee. We are grateful to Dr. X.-N. Li for help with the X-ray diffraction analysis.

References

1. Lu H, Wang SS, Zhou QW, Zhao YN, Zhao BY, Rangeland J. 2012; 34:329–339.
2. Academia Sinica. Flora of China. Vol. 42. Science Press; Beijing: 1998. p. 21
3. Molyneux RJ, James LF. Science. 1982; 216:190–191. [PubMed: 6801763]
4. Cao GR, Li SJ, Duan DX, Zhao XW. J Northwest Sci-Tech Univ Agri For. 1989; 17:1–7.
5. Meng XZ, Hu XQ, Zhang RM, You YT. Chin Trad Herb Drug. 1994; 25:61–64.
6. Li X, Zhang SD, Jin HZ, Dong F, Shan L, Zhang WD. Nat Prod Res. 2013; 27:554–557. [PubMed: 22494026]
7. Sun RQ, Jia ZJ, Cheng DL, Zhu ZQ. Planta Med. 1992; 58:211–213. [PubMed: 1529034]
8. Tan C-J, Liu L-N, Bi-sihaletu H, Tang H-M, Zhao B-Y. Prog Vet Med. 2015; 35:82–85.
9. Tan C-J, Liu L-N, Tang H-M, Shi B-Y, Ran J-Q, Zhao B-Y. Nar Pro Res Dev. 2015; 27:1365–1367.
10. Tan CJ, Yi P, Goto M, Morris-Natschke SL, Liu LN, Lee KH, Zhao BY. Helv Chim Acta. 2016; 99:225–227.
11. Wang LS, You YJ, Wang SQ, Liu X, Liu BM, Wang JN, Lin X, Chen MS, Liang G, Yang H. Bioorg Med Chem Lett. 2012; 22:4100–4102. [PubMed: 22578453]
12. Tan C-J, Zhao Yu, Goto M, Hsieh KY, Yang X-M, Morris-Natschke SL, Liu L-N, Zhao B-Y, Lee KH. Bioorg Med Chem Lett. 2016; 26:1495–1497. [PubMed: 26865176]
13. Ochrocephalamine A (**1**): yellow crystals (EtOH); mp 159–161 °C; ESIMS m/z 491 $[M + H]^+$; HREIMS at 490.3296 $[M]^+$ (calcd. 490.3308, $C_{30}H_{42}N_4O_2$); $[\alpha]_D^{24} +32.2$ (c 0.48, MeOH); UV (MeOH) λ_{max} (log ϵ) 202 nm (4.2), 311 nm (3.9), 373 nm (2.3); IR (KBr) ν_{max} 2932, 2860, 2805, 2764, 2745, 1639, 1597, and 1554 cm^{-1} ; 1H and ^{13}C NMR data (Table 1). Crystal data for **1**: $C_{30}H_{42}N_4O_2 \cdot 6(H_2O)$, $M = 598.77$, monoclinic, $a = 13.0528(3)$ Å, $b = 8.3014(2)$ Å, $c = 14.8105(4)$ Å, $\alpha = 90.00^\circ$, $\beta = 101.6050(10)^\circ$, $\gamma = 90.00^\circ$, $V = 1572.01(7)$ Å³, $T = 100(2)$ K, space group $P2_1$, $Z = 2$, $\mu(CuK\alpha) = 0.745$ mm⁻¹, 13217 reflections measured, 4874 independent reflections ($R_{int} = 0.0386$). The final R_I values were 0.0521 ($I > 2\sigma(I)$). The final $wR(F^2)$ values were 0.1565 ($I > 2\sigma(I)$). The final R_I values were 0.0525 (all data). The final $wR(F^2)$ values were 0.1572 (all data). The goodness of fit on F^2 was 1.082. Flack parameter = $-0.3(2)$. The Hooft parameter is 0.01(6) for 1892 Bijvoet pairs. Deposited to Cambridge Crystallographic Data Center with No. CCDC 1449099.
14. Liu X-J, Cao M-A, Li W-H, Shen C-S, Yan S-Q, Yuan C-S. Fitoterapia. 2010; 81:524–527. [PubMed: 20079811]
15. Galasso V, Asaro F, Berti F, Pergolese B, Kovac B, Pichierri F. Chem Phys. 2006; 330:457–468.
16. Hooft RWW, Straver LH, Spek AL. J Appl Crystallogr. 2008; 41:96–103. [PubMed: 19461838]
17. Sanchez MDC, Altares P, Pedrosa MM, Burbano C, Cuadrado C, Goyoaga C, Muzquiz M, Jimenez-Martinez C, Davila-Ortiz G. Food Chemistry. 2005; 90:347–355.
18. Wu, L-J.; Lou, H-X.; Zhou, J. The Chemistry of Natural Products. 6. People's Medical Publishing House; Beijing: 2012. p. 379–380.

- Ochrocephalamine A (**1**) was isolated from *Oxytropis ochrocephala* Bunge.
- The poisonous plant *O. ochrocephala* poses an extreme risk to grazing animals.
- Compound **1** is a dimeric matrine-type alkaloid.
- Its structure was solved from spectral data and single-crystal X-ray diffraction.
- Compound **1** showed contact toxicity against *Spodoptera litura* third-instar larvae.

Figure 1.
The structure of ochrocephalamine A (1)

Figure 2.
a) key HMBC correlations of **1**, b) X-ray crystal structure of **1**.

Scheme 1.
Biogenetic pathway proposed of ochrocephalamine A (**1**)

Table 1
 ^1H (400 MHz) and ^{13}C NMR (100 MHz) Data of Ochrocephalamine A (1) in CD_3OD

No	δ_{H} (mult, J)	δ_{C}	no.	δ_{H} (mult, J)	δ_{C}
2	2.82 (m) ^a 2.07 (m) ^a	57.8	2'	2.82 (m) ^a 2.07 (m) ^a	58.3
3	1.56–1.65 (m) ^a	22.2	3'	1.56–1.65 (m) ^a	21.6
4	1.79 (m) ^a	27.8	4'	1.67–1.71 (m) ^a	28.7
5	2.25 (m)	32.9	5'	1.80 (m)	37.5
6	2.33 (t, 4)	61.5	6'	2.29 (<i>brs</i>)	65.6
7	3.03 (<i>brs</i>)	39.5	7'	1.89 (m)	42.6
8	2.15 (m)	27.5	8'	2.44–2.50(m)	28.4
9	1.56–1.65 (m) ^a	22.2	9'	1.56–1.65 (m) ^a	21.3
10	2.82 (m) ^a 2.07 (m) ^a	57.6	10'	2.82 (m) ^a 2.07 (m) ^a	58.2
11		147.7	11'	3.94 (m)	53.8
12	6.51 (d, 7.6)	106.6	12'	2.14 (m)	29.8
13	7.30 (d, 7.6)	135.7	13'	3.38 (m)	30.0
14		130.1	14'	2.62(dd, 16.0, 4.0) 2.46(dd, 16.0, 8.0)	37.5
15		164.6	15'		170.9
17	4.09 (dd, 12.8, 7.2) 3.80 (<i>brt</i> , 12.8)	45.5	17'	4.29 (dd, 12.8, 4.0) 3.11 (<i>brt</i> , 12.8)	43.7

^a overlapped.