Figure 5 The wound is closed

Figure 6 Skin thinning

DISCUSSION

The technique described here has three advantages over the conventional technique: (i) improved cosmetic effect (thanks to avoidance of the abnormal hair growth that occurs along the line of the semicircular incision); (ii) easier skin thinning (Fig 6); (iii) shorter operative time.

Effective irrigation of contaminated open fractures

D Lin, V Reddy, A Vasireddy
Barts Health NHS Trust, UK

CORRESPONDENCE TO

David Lin, E: david.lin1@nhs.net
doi 10.1308/rcsann.2016.0140

BACKGROUND

The sound washout and debridement of an open fracture is fundamental to reducing the incidence of infection in open fracture management.¹

Figure 1 Irrigation set

High pressure pulse lavage has been contraindicated as there is the potential to push the contaminant further into the wound.² Using a giving set attached to an elevated fluid bag does not provide the required pressure and takes a long time to wash out the wound effectively. We describe a simple technique that allows for thorough, efficient wound washout.

TECHNIQUE

A warmed 3l saline bag is removed from its wrapping in a sterile fashion and cut at the drip insertion site. A sterile Yankauer sucker is attached to the end of the bag (Fig 1). The tip can now be used to direct the irrigation flow into the desired portions of the wound. By varying the pressure on the bag with one hand, the pressure of the stream can be varied. Use of a broad Mayo bag under the irrigated limb enables the washout fluid to be collected easily and stops contamination of the sterile operating field.

DISCUSSION

In our experience, this technique is very effective at washing out open fractures and the washout can be performed in under five minutes. This form of wound washout can be extended for use well beyond open fracture management and is very easy to set up as both components are universally available at all hospitals.

References

1. British Orthopaedic Association; British Association of Plastic, Reconstructive and Aesthetic Surgeons. *BOAST 4: The Management of Severe Open Lower Limb Fractures*. London: BOA; 2009.
2. Nanchahal J, Nayagam S, Khan U *et al*. *Standards for the Management of Open Fractures of the Lower Limb*. London: BAPRAS; 2009.