Published in final edited form as:

Arch Gerontol Geriatr. 2017 ; 69: 55–60. doi:10.1016/j.archger.2016.11.007.

Activity Engagement and Physical Function in Old Age Sample

Krupa N. Shah, MD MPH^{#1}, Feng V. Lin, PhD^{#2}, Fang Yu, PhD³, and James M. McMahon, PhD²

¹Divisions of Geriatrics and Aging, Department of Medicine, University of Rochester, Rochester, New York, USA

²School of Nursing, University of Rochester, Rochester, NY, USA

³School of Nursing University of Minnesota, Minneapolis, MN, USA

[#] These authors contributed equally to this work.

Abstract

Objectives—To describe the patterns of engagement in mental, physical, and social activity (MA, PA, and SA) and to examine the relationship between combined activity engagement and physical function among community-dwelling older adults.

Design—Cross-sectional correlational study.

Setting—Multiple communities.

Participants—A total of 466 individuals aged 55 years or older.

Measurements—Physical function was assessed using grip strength and gait speed. Engagement in PA, MA and SA was obtained from self-report questionnaires.

Results—We identified four classes (“Active PA and MA”, “Active MA”, “Active PA”, and “Inactive”) that significantly differed in the frequency of engagement in MA and PA using latent class analysis. SA didn't differ across classes. Controlling for age, the “Active PA and MA”, “Active MA”, “Active PA” groups displayed similar grip strength that was superior to the “Inactive” group. “Active PA and MA” group had best gait speed relative to other groups, especially “Active MA” and “Inactive” group, while the “Active PA”, “Active MA”, and “Inactive” group were similar in gait speed.

Conclusion—Combined physical and mental activity engagement was associated with better physical function, especially in gait speed. Future interventional research should investigate the

Corresponding author: Krupa N. Shah, MD MPH, Highland Hospital, University of Rochester School of Medicine and Dentistry, 1000 South Ave, Rochester, NY 14692, Phone: 584-341-0762, Fax: 584-341-8035, krupa_shah@urmc.rochester.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Author Contributions

The authors' role are as follows: study concept and design (FVL, KNS), data analysis and interpretation of data (FVL, KNS, FY, JMM), and preparation of manuscript (FVL, KNS, FY, JMM).

Conflicts of Interest

Financial Conflict: None

combination of both physical and cognitive training to prevent decline of physical function in older adults.

Keywords

physical function; physical; mental and social activity

INTRODUCTION

Physical function is defined as the ability to perform the basic actions (i.e., mobility, strength, and endurance) that are essential for maintaining independence and carrying out more complex activities (Painter et al., 1999). Decline in physical function is common among older adults and has been shown to increase the risk of falls, hospitalization, nursing home admissions, dependence, and poor quality of life (Brown and Flood, 2013). Hence, promoting physical function is critical for preventing its associated adverse health outcomes and constitutes one of the hallmark signs of successful aging.

One way to prevent functional decline is to engage in physical, mental, and/or social activities, although the amount and level of evidence for each activity varies. Physical activity (PA) is the most studied with accumulating evidence indicating that engaging in PA improved strength, endurance, balance and overall physical functions in older adults (Gomes Neto et al., 2013; Gomes-Neto et al., 2013; Taylor, 2014). The relationship between mental activities (MA) and physical function has been less studied. However, emerging interventional research suggests that engaging in MA (e.g., cognitive stimulation) can enhance physical function by improving gait and balance (Smith-Ray et al., 2015; Smith-Ray et al., 2014). Further, older adults who are socially active experienced less decline in physical function (e.g., as measured by their ability to perform daily tasks) compared to socially inactive counterparts (Mendes de Leon et al., 2003; Rosso et al., 2013). Overall, more studies are needed to examine the relationship between engagement in these activities and physical function.

Moreover, it remains unclear whether combined activity engagement would be associated with better physical function. Learning from the cognition literature and the Enriched Environment Theory suggest a simultaneously effect of PA, MA, and SA on cognitive function in old age (Hertzog et al., 2008). Combined activity engagement such as PA and MA (Theill et al., 2013) or MA and SA (Wang et al., 2002) indeed led to significantly greater cognitive improvement compared to engaging in a single activity. However, it is unknown whether there is any similar synergistic impact of combined activity engagement on physical function among older adults.

This study addresses the aforementioned gaps. We evaluated the relationship of engagement in individual and combined PA, MA, and SA and physical function. First, we described the patterns of engagement in PA, MA, and SA among older adults. Next, we identified the latent classes of activity engagement (clusters of participants with similar activity engagement). Last, we examined the association between activity engagement and physical function. We hypothesized that combined activity engagement would link to better physical functions than any single activity.

METHODS

Design and Procedure

The present study used a cross-sectional design to analyze data from the Survey of Midlife Development in the United States (MIDUS) II. MIDUS is a longitudinal study of community-dwelling adults' well-being, including two waves that were 10 years apart (MIDUS I, 1995-1996; MIDUS II, 2004-2009). Of note, a twenty-year follow-up was just completed (MIDUS III), but is not yet available publicly. MIDUS I only collected psychosocial behavioral data, which had no physical function assessments needed in the study. MIDUS II consisted of five projects covering different domains: 1) psychosocial and health, 2) daily diary, 3) cognitive assessment, 4) clinical and biomarker assessment, and 5) neuroscience. Data used in the present study were drawn from three of the projects: Project 1: participants self-administered questionnaires on socio-demographic and health information; Project 3: a series of cognitive tests were administered to participants over the telephone; and Project 4: physical function assessment obtained from a two-day visit to one of the participating General Clinical Research Centers (GCRCs). Data from individuals 55 years or older who have data from all three projects were used for the current study ($n = 466$). Of note, the sample from MIDUS I was considered a nationally representative sample, while participants who remained in MIDUS II (75% from MIDUS I) tended to have socioeconomic bias (e.g., being white, female, married, more educated), but not necessarily healthy status bias (Radler and Ryff, 2010). Institutional Review Boards from each study site approved the study. Institutional Review Boards from each study site approved the study.

Measures

Physical function was operationalized using two of the clinical assessment variables from MIDUS II Project 4: grip strength and gait speed. Grip strength was determined using a hand grip dynamometer, while subjects held the dynamometer in the hand to be tested with the elbow positioned at 90 degrees on the side of the body. An average of three readings (in Kg/force) of the dominant hand was taken. Hand grip dynamometer is a reliable instrument to measure grip strength with interclass correlation coefficient greater than 0.80 (Guerra and Amaral, 2009; Hamilton et al., 1992).

Gait speed was assessed using the time in seconds required for an individual to walk 50 feet in their usual pace without an assistive device. An average of two readings (in feet/second) was taken. The interclass correlation coefficient was 0.97 in previous studies (Dobson et al., 2013; Unver et al., 2015).

Activity engagement was measured using three distinct questionnaires: physical, mental and social. Participants were asked about frequency of engagement of these activities as part of the MIDUS II Project 1. The questionnaire of moderate PA asked about the frequency of engaging in 6 types of leisurely sports (i.e., light tennis, slow or light swimming, low impact aerobics, golfing without a power cart, brisk walking, mowing the lawn with a walking lawnmower). Participants were asked to respond to each item on a 6-point ordinal scale ranging from 1 (several times a week) to 6 (never). In the MIDUS II survey, three types of PA were available – light, moderate, and vigorous, which are highly correlated ($r = .34 - .69$)

(Lee et al., 2015). We decided to use the moderate type PA to balance the amount of exercise and feasibility of doing the exercise in old age. MA engagement included 6 types of activities (i.e., read, do word games, play card, attend lectures, do writing, use a computer). Each participant indicated the frequency of engaging in these activities using a 6-point ordinal scale ranging from 1 (daily) to 6 (never). For PA and MA, the ratings were reversely coded and averaged with higher scores indicating more frequent engagement. SA engagement was measured with three items quantifying the frequency of attending meetings and group gatherings (i.e., union, sports, or social groups) outside the workplace in a typical month. The total number of times involved in these SA was calculated. MA engagement scale was significantly correlated to SA engagement scale ($r = .29, p < .001$). PA engagement scale did not correlate to MA or SA engagement scales. The three activity engagement scales were created in the MIDUS II, although formal validation evaluation has not been conducted, evidence has been accumulated to support the validity in outcomes related to the present study. For example, the three scales have been correlated to memory complaints, cognitive function and education (Lachman et al., 2011; Lin et al., 2013), or perceived physical health (Fujiwara and Kawachi, 2008; Lee et al., 2015) in previous studies of older adults.

Demographic and health characteristics included age, sex, education, medications, and smoking behaviors. Education was grouped into three categories: “high school graduate or less”, “some college” and “college graduate or more”. The use of medications such as anti-hypertension, anti-depressants and/or corticosteroids were recorded based on the original medication bottles the participants brought with them to the GCRC. Data on smoking was collected using a single question on whether the participant had ever smoked regularly. Abdominal obese was defined as waist circumference >102 cm in men and >88 cm in women.

Data analysis

Across the entire sample, descriptive data on all main and background variables were analyzed, and the correlations between activity engagement and physical function was analyzed with Pearson's correlation in IBM SPSS 22.0.

Latent class analysis (LCA), a form of mixture modeling, was performed using Mplus version 7.0 to find the smallest number of classes (participants with similar activity engagement). LCA is a method of identifying unique classes within a set of heterogeneous individuals by examining the mean of individual cases. In the present study, we determined the number of classes controlling for the influence of age on each activity engagements. A series of models were evaluated beginning with a 1-class solution and ending with a 5-class solution. The optimal number of classes was decided based on Bayesian, Akaike, and Adjusted Bayesian Information Criteria in which lower values indicate a more parsimonious model; entropy in which higher values indicate a better model, and the Lo-Mendell-Rubin (LMR) adjusted likelihood ratio test which compares the current class solution with the class - 1 solution to determine whether the two solutions are similar ($p > .05$) or different ($p < .05$) (Ram and Grimm, 2009). Each class should have more than 1% of the total sample

(Jung and Wickrama, 2008). Each class was described by the mean and standard deviation (SD) of relevant variables and the number of participants belonging to the class.

After deciding the number of latent classes ($n = 4$ in the present study), remaining analyses were performed in IBM SPSS 22.0. Analysis of variance controlling for covariates (ANCOVA) was used to assess differences across latent classes on continuous variables, and Chi-square tests were applied to compare categorical variables. Post-hoc analysis was based on the Fisher's least significant difference adjustment for multiple comparisons with equal variance. Of note, for the ANCOVA, estimating a small to moderate effect size ($f = 0.25$), alpha at 0.01 (considering multiple outcomes), power at 0.90, 4 groups/latent class, and 1 covariate (age), the total sample size should be 444, which is similar to our sample size at 466.

RESULTS

Activity engagement across the entire sample

Across the entire sample, participants engaged in MA ($M = 2.27$, $SD = 0.85$) and PA ($M = 2.42$, $SD = 1.37$) once to several times a month; and attended SA an average of 1.15 times a month ($SD = 1.02$). PA was related to both grip strength ($r = .14$, $p = .003$) and gait speed ($r = .22$, $p < .001$), while MA was associated with grip strength ($r = .12$, $p = .009$). SA was not related to either physical function measure.

Latent class of activity engagement

Table 1 summarizes the series of model fit statistics for LCA. Synthesizing the model fit indices and the number of participants in each class, the four-class model was considered the best solution. Figure 1A and Table 2 present the data from the four classes. Class I ($n = 50$, 12.7%) was characterized by relatively frequent engagement in PA and MA, labeled as “Active PA and MA”. Class II ($n = 151$, 32.4%) was characterized by relatively frequent engagement in MA but not PA, labeled as “Active MA”. Class III ($n = 172$, 36.9%) was characterized by relatively frequent engagement in PA but not MA, labeled as “Active PA”. Class IV ($n = 93$, 20.0%) was inactive in both MA and PA, labeled as “Inactive”. Of note, variation in SA contributed little to class designation. That is, the four classes were significantly different in MA ($F = 2.99$, $p = .031$) and PA ($F = 2014.01$, $p < .001$) but not SA ($F = 1.99$, $p = .12$), controlling for age.

Physical functions by latent class of activity engagement

Table 2 displays descriptive data for all variables by latent class. We found class differences in age and level of education but not in health history. Grip strength ($F = 2.86$, $p = .036$, $\eta_p^2 = 0.018$) and gait speed ($F = 3.57$, $p = .014$, η_p^2 differed significantly by latent class controlling for age. For post-hoc analysis, in “Active PA and MA”, “Active PA”, and “Active MA” classes all had significantly better grip strength than “Inactive” class, while the three classes were similarly in grip strength. “Active PA and MA” class had significantly better gait speed than “Active MA” or “Inactive” class, while there was no evidence of a difference between “Active PA and MA” and “Active PA” classes, or between “Active PA”, “Active MA”, and “Inactive” class (also see Figure 1B).

DISCUSSION

In the present study we have identified four groups of individuals that were characterized by distinctly different patterns of activity engagement in PA and MA. SA didn't differ among groups. The first group included individuals who were active in both PA and MA ("Active PA and MA"). The second and third groups included individuals who were active in either activity ("Active PA" or "Active MA"). The fourth group included individuals who were active in neither activity ("Inactive"). Controlling for age, the first three groups displayed similar grip strength that was superior to the "Inactive" group. "Active PA and MA" group had the better gait speed than other groups, especially "Active MA" and "Inactive" group, while the "Active PA", "Active MA", and "Inactive" groups were similar in their gait speed. Noticeably, the four groups were similar in their health status. To our best knowledge this is one of the first studies to assess the relationship between combined activity engagement and physical function.

Previous studies that have evaluated combined activity engagement in the aging literature have focused on the cognitive outcomes. Our findings suggest that combined activity engagement is associated with better physical function, especially in the domain of gait speed. The difference in gait speed between "Active PA and MA" and other groups (e.g., "Active MA", "Inactive") is 0.30 feet/second or greater (equal to 0.09 m/s or greater), which is greater than the clinically meaningful change indicated in the literature (0.04 to 0.06 m/s) (Perera et al., 2006). On the other hand, the difference in grip strength between the activity engagement groups and the inactive group is ~5kg/force. This change is statistically significant but is slightly less clinically meaningful difference than what is indicated in literature (i.e., 6 Kg/force) (Kim et al., 2014; Nitschke et al., 1999).

We found that maintaining or improving gait speed in old age relies on the engagement of both mental and physical activities. There has been a consistent literature supporting the positive impact of physical intervention or PA on gait speed (Nadkarni et al., 2013; Pahor et al., 2014; VanSwearingen et al., 2011; Villareal et al., 2011). More emerging literature suggests that cognitive intervention such as MA leads to improvement in gait speed (Dodge et al., 2008; Dumas et al., 2009; Li et al., 2010; Smith-Ray et al., 2015; Verghese et al., 2010). Noticeably, gait is not a unitary concept, which is related to multiple cognitive capacities (e.g., executive function, attention, episodic memory and processing speed) (Bolanzadeh et al., 2014; Parihar et al., 2013; Yogev et al., 2008). Our results differ from those of Ng et al.'s intervention study reporting that in frail Asian older adults PA alone had better effects on gait speed than combined PA and MA engagement (Ng et al., 2015). In contrast, we found similar relationship between combined PA and MA, as well as PA alone, and gait speed. Different from gait speed, grip strength seems to have a nonselective acceptability to engagement in PA or/and MA. Grip strength is known to be modifiable by increasing the level of PA (Cadore et al., 2014; Justine et al., 2012). Our previous study on a different older American cohort suggests that grip strength may be particularly sensitive to cognitive deficits (e.g., speed of processing, attention), compared to some other measures of physical function (e.g., overall physical function, activity of daily living) (Lin et al., 2012). Also, grip strength may have a stronger relationship with cognitive function and MA than gait speed (Atkinson et al., 2010). Synthesizing the different findings and evidence on grip

strength and gait speed and their relationship to activity engagement, we urge the further validation and comprehensive comparison of different activity engagement on different aspects of physical function.

In this study, we observed that the strength of correlations between physical function (muscle strength and gait) and engagement in PA and MA are statistically significant, but relatively low. Previous observational studies do not always find positive associations between PA and muscle strength (Cooper et al., 2011; Scott et al., 2011). Several factors besides PA engagement influence grip strength such as fatigue, time of day, nutritional status, restricted motion, and pain. Similarly in meta-analysis of PA interventions, engaging in PA has small effect on gait speed, suggesting that other factors may be involved in supporting gait speed (Chou et al., 2012; de Vries et al., 2012). Of note, we did not observe an association between the engagement in SA and physical function. The difference in the measures of functional outcome may explain the different findings. Compared to MA and PA that may directly modify basic upper or lower extremity function, SA may contribute to the self-perception of functions in more complex activities under social context (e.g., instrumental activities of daily living, disability) (Mendes de Leon et al., 2003; Rosso et al., 2013; Thomas, 2011).

We acknowledge some limitations of our study. First, the cross-sectional study design can only identify associations instead of causality. Second, the three activities (PA, MA and SA) were assessed using frequency of activity engagement based on self-report questionnaires without attention to the duration and intensity of activity engagement and did not capture other activities that were not listed in the study activity engagement questionnaires. Third, the four groups were similar in SA, which may be due to the different format the questions were asked compared to the questions for PA and MA. Fourth we did not exclude acceleration and deceleration phases in our gait speed measurement to obtain stable and comparable gait speed (Wang et al., 2012). Finally, our findings may not be applicable to other groups of older adults such as older men, or older adults with lower education, racial or ethnic minorities. Regardless, the strengths of our study also should be noted. First, our study is among the first to examine relationships between different type of activity engagement, especially the combined activity, and physical function. Second, our outcome data uses objective measures of physical function rather than subjective measures; performance based measures are better suited to capture early stages of functional decline and have greater validity and reliability compared to using self-reports (Brown et al., 2000; Guralnik et al., 1994).

In conclusion, our study provides evidence that engagement in a combined PA and MA compared to engagement in only one type of activity is associated with better physical function (particularly walking ability). Our study not only reinforces the importance of PA in improving physical function, but also underscores the importance of MA in improving walking ability in older adults. Lastly, the significance of increasing MA (e.g. cognitive stimulation) to prevent functional decline is potentially even more important in older adults, particularly those with underlying cognitive impairments. Future interventional research should investigate the combination of both physical and cognitive training to prevent functional decline and frailty in older adults.

ACKNOWLEDGMENTS

Funding Source: NIH K23 AG043319 to K. Shah, Alzheimer's Association New Investigator Grant (NIRG-14-317353) and NIH R01 grant (NR015452) to F. Lin

Sponsor's Role: None

REFERENCES

- Atkinson HH, Rapp SR, Williamson JD, Lovato J, Absher JR, Gass M, Henderson VW, Johnson KC, Kostis JB, Sink KM, Mouton CP, Ockene JK, Stefanick ML, Lane DS, Espeland MA. The relationship between cognitive function and physical performance in older women: results from the women's health initiative memory study. *J. Gerontol. A Biol. Sci. Med. Sci.* 2010; 65:300–306. [PubMed: 19789197]
- Bolanzadeh N, Liu-Ambrose T, Aizenstein H, Harris T, Launer L, Yaffe K, Kritchevsky SB, Newman A, Rosano C. Pathways linking regional hyperintensities in the brain and slower gait. *Neuroimage.* 2014; 99:7–13. [PubMed: 24841418]
- Brown CJ, Flood KL. Mobility limitation in the older patient: a clinical review. *JAMA.* 2013; 310:1168–1177. [PubMed: 24045741]
- Brown M, Sinacore DR, Binder EF, Kohrt WM. Physical and performance measures for the identification of mild to moderate frailty. *J. Gerontol. A Biol. Sci. Med. Sci.* 2000; 55:M350–M355. [PubMed: 10843356]
- Cadore EL, Casas-Herrero A, Zambom-Ferraresi F, Idoate F, Millor N, Gómez M, Rodríguez-Mañas L, Izquierdo M. Multicomponent exercises including muscle power training enhance muscle mass, power output, and functional outcomes in institutionalized frail nonagenarians. *Age.* 2014; 36:773–785. [PubMed: 24030238]
- Chou CH, Hwang CL, Wu YT. Effect of exercise on physical function, daily living activities, and quality of life in the frail older adults: a meta-analysis. *Arch. Phys. Med. Rehabil.* 2012; 93:237–244. [PubMed: 22289232]
- Cooper R, Mishra GD, Kuh D. Physical activity across adulthood and physical performance in midlife: findings from a British birth cohort. *Am. J. Prev. Med.* 2011; 41:376–384. [PubMed: 21961464]
- de Vries NM, van Ravensberg CD, Hobbelen JS, Olde Rikkert MG, Staal JB, Nijhuis-van der Sanden MW. Effects of physical exercise therapy on mobility, physical functioning, physical activity and quality of life in community-dwelling older adults with impaired mobility, physical disability and/or multi-morbidity: a meta-analysis. *Ageing research reviews.* 2012; 11:136–149. [PubMed: 22101330]
- Dobson F, Hinman RS, Roos EM, Abbott JH, Stratford P, Davis AM, Buchbinder R, Snyder-Mackler L, Henrotin Y, Thumboo J, Hansen P, Bennell KL. OARSI recommended performance-based tests to assess physical function in people diagnosed with hip or knee osteoarthritis. *Osteoarthritis Cartilage.* 2013; 21:1042–1052. [PubMed: 23680877]
- Dodge HH, Kita Y, Takechi H, Hayakawa T, Ganguli M, Ueshima H. Healthy Cognitive Aging and Leisure Activities Among the Oldest Old in Japan: Takashima Study. *J. Gerontol. A Biol. Sci. Med. Sci.* 2008; 63:1193–1200. [PubMed: 19038834]
- Doumas M, Rapp MA, Krampe RT. Working memory and postural control: adult age differences in potential for improvement, task priority, and dual tasking. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 2009; 64:193–201. [PubMed: 19255088]
- Fujiwara T, Kawachi I. Social capital and health. A study of adult twins in the U.S. *Am J Prev Med.* 2008; 35:139–144. [PubMed: 18617082]
- Gomes Neto M, Ogalha C, Andrade AM, Brites C. A systematic review of effects of concurrent strength and endurance training on the health-related quality of life and cardiopulmonary status in patients with HIV/AIDS. *BioMed research international.* 2013; 2013:319524. [PubMed: 23691497]
- Gomes-Neto M, Conceicao CS, Oliveira Carvalho V, Brites C. A systematic review of the effects of different types of therapeutic exercise on physiologic and functional measurements in patients with HIV/AIDS. *Clinics (Sao Paulo, Brazil).* 2013; 68:1157–1167.

- Guerra RS, Amaral TF. Comparison of hand dynamometers in elderly people. *J. Nutr. Health Aging.* 2009; 13:907–912. [PubMed: 19924352]
- Guralnik JM, Simonsick EM, Ferrucci L, Glynn RJ, Berkman LF, Blazer DG, Scherr PA, Wallace RB. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J. Gerontol.* 1994; 49:M85–M94. [PubMed: 8126356]
- Hamilton GF, McDonald C, Chenier TC. Measurement of grip strength: validity and reliability of the sphygmomanometer and jamar grip dynamometer. *J. Orthop. Sports Phys. Ther.* 1992; 16:215–219. [PubMed: 18796752]
- Hertzog C, Kramer AF, Wilson RS, Lindenberger U. Enrichment Effects on Adult Cognitive Development: Can the Functional Capacity of Older Adults Be Preserved and Enhanced? *Psychol. Sci. Public Interest.* 2008; 9:1–65. [PubMed: 26162004]
- Jung T, Wickrama KAS. An introduction to latent class growth analysis and growth mixture modeling. *Social and Personality Psychology Compass.* 2008; 2:302–317.
- Justine M, Hamid TA, Mohan V, Jagannathan M. Effects of Multicomponent Exercise Training on Physical Functioning among Institutionalized Elderly. *ISRN Rehabilitation.* 2012; 2012:7.
- Kim JK, Park MG, Shin SJ. What is the minimum clinically important difference in grip strength? *Clin. Orthop. Relat. Res.* 2014; 472:2536–2541. [PubMed: 24817380]
- Lachman ME, Agrigoroaei S, Murphy C, Tun PA. Frequent cognitive activity compensates for education differences in episodic memory. *Am J Geriatr Psychiatry.* 2011; 18:4–10.
- Lee B, Lawson KM, Chang PJ, Neuendorf C, Dmitrieva NO, Almeida DM. Leisure-time physical activity moderates the longitudinal associations between work-family spillover and physical health. *Journal of leisure research.* 2015; 47
- Li KZ, Roudaia E, Lussier M, Bherer L, Leroux A, McKinley PA. Benefits of cognitive dual-task training on balance performance in healthy older adults. *J. Gerontol. A Biol. Sci. Med. Sci.* 2010; 65:1344–1352. [PubMed: 20837662]
- Lin F, Chen D, Vance DE, Mapstone M. Trajectories of Combined Laboratory- and Real World-based Speed of Processing in Community-Dwelling Older adults. Predictors and Functional Outcomes *Journal of Gerontology: Psychological Sciences and Social Sciences.* 2012 Epub ahead of print.
- Lin F, Heffner K, Mapstone M, Chen DG, Porsteisson A. Frequency of Mentally Stimulating Activities Modifies the Relationship Between Cardiovascular Reactivity and Executive Function in Old Age. *Am J Geriatr Psychiatry.* 2013
- Mendes de Leon CF, Glass TA, Berkman LF. Social engagement and disability in a community population of older adults: the New Haven EPESE. *Am. J. Epidemiol.* 2003; 157:633–642. [PubMed: 12672683]
- Nadkarni NK, Studenski SA, Perera S, Rosano C, Aizenstein HJ, Brach JS, Van Swearingen JM. White matter hyperintensities, exercise, and improvement in gait speed: does type of gait rehabilitation matter? *J. Am. Geriatr. Soc.* 2013; 61:686–693. [PubMed: 23590257]
- Ng TP, Feng L, Nyunt MS, Feng L, Niti M, Tan BY, Chan G, Khoo SA, Chan SM, Yap P, Yap KB. Nutritional, Physical, Cognitive, and Combination Interventions and Frailty Reversal Among Older Adults: A Randomized Controlled Trial. *Am. J. Med.* 2015; 128:1225–1236.e1221. [PubMed: 26159634]
- Nitschke JE, McMeeken JM, Burry HC, Matyas TA. When is a change a genuine change? A clinically meaningful interpretation of grip strength measurements in healthy and disabled women. *J. Hand Ther.* 1999; 12:25–30. [PubMed: 10192632]
- Pahor M, Guralnik JM, Ambrosius WT, et al. Effect of structured physical activity on prevention of major mobility disability in older adults: The life study randomized clinical trial. *JAMA.* 2014; 311:2387–2396. [PubMed: 24866862]
- Painter P, Stewart AL, Carey S. Physical functioning: definitions, measurement, and expectations. *Advances in renal replacement therapy.* 1999; 6:110–123. [PubMed: 10230878]
- Parihar R, Mahoney JR, Verghese J. RELATIONSHIP OF GAIT AND COGNITION IN THE ELDERLY. *Current translational geriatrics and experimental gerontology reports.* 2013; 2

- Perera S, Mody SH, Woodman RC, Studenski SA. Meaningful change and responsiveness in common physical performance measures in older adults. *J. Am. Geriatr. Soc.* 2006; 54:743–749. [PubMed: 16696738]
- Radler BT, Ryff CD. Who participates? Accounting for longitudinal retention in the MIDUS national study of health and well-being. *J Aging Health.* 2010; 22:307–331. [PubMed: 20103686]
- Ram N, Grimm KJ. Growth Mixture Modeling: A Method for Identifying Differences in Longitudinal Change Among Unobserved Groups. *International journal of behavioral development.* 2009; 33:565–576. [PubMed: 23885133]
- Rosso AL, Taylor JA, Tabb LP, Michael YL. Mobility, disability, and social engagement in older adults. *J. Aging Health.* 2013; 25:617–637. [PubMed: 23548944]
- Scott D, Blizzard L, Fell J, Jones G. Prospective associations between ambulatory activity, body composition and muscle function in older adults. *Scand. J. Med. Sci. Sports.* 2011; 21:e168–175. [PubMed: 20874860]
- Smith-Ray RL, Hughes SL, Prohaska TR, Little DM, Jurivich DA, Hedeker D. Impact of Cognitive Training on Balance and Gait in Older Adults. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 2015; 70:357–366. [PubMed: 24192586]
- Smith-Ray RL, Makowski-Woidan B, Hughes SL. A randomized trial to measure the impact of a community-based cognitive training intervention on balance and gait in cognitively intact Black older adults. *Health Educ. Behav.* 2014; 41:62s–69s. [PubMed: 25274713]
- Taylor D. Physical activity is medicine for older adults. *Postgrad. Med. J.* 2014; 90:26–32. [PubMed: 24255119]
- Theill N, Schumacher V, Adelsberger R, Martin M, Jancke L. Effects of simultaneously performed cognitive and physical training in older adults. *BMC Neurosci.* 2013; 14:103. [PubMed: 24053148]
- Thomas PA. Trajectories of social engagement and limitations in late life. *J. Health Soc. Behav.* 2011; 52:430–443. [PubMed: 22144732]
- Unver B, Kalkan S, Yuksel E, Kahraman T, Karatosun V. Reliability of the 50-foot walk test and 30-sec chair stand test in total knee arthroplasty. *Acta Ortopedica Brasileira.* 2015; 23:184–187. [PubMed: 26327798]
- VanSwearingen JM, Perera S, Brach JS, Wert D, Studenski SA. Impact of exercise to improve gait efficiency on activity and participation in older adults with mobility limitations: a randomized controlled trial. *Phys. Ther.* 2011; 91:1740–1751. [PubMed: 22003158]
- Verghese J, Mahoney J, Ambrose AF, Wang C, Holtzer R. Effect of cognitive remediation on gait in sedentary seniors. *J. Gerontol. A Biol. Sci. Med. Sci.* 2010; 65:1338–1343. [PubMed: 20643703]
- Villareal DT, Chode S, Parimi N, Sinacore DR, Hilton T, Armamento-Villareal R, Napoli N, Qualls C, Shah K. Weight loss, exercise, or both and physical function in obese older adults. *N Engl J Med.* 2011; 364:1218–1229. [PubMed: 21449785]
- Wang CY, Chen TR, Lin YH, Liu MH, Chen YC. Gait speed measure: the effect of different measuring distances and the inclusion and exclusion of acceleration and deceleration. *Percept. Mot. Skills.* 2012; 114:469–478. [PubMed: 22755452]
- Wang HX, Karp A, Winblad B, Fratiglioni L. Late-life engagement in social and leisure activities is associated with a decreased risk of dementia: a longitudinal study from the Kungsholmen project. *Am. J. Epidemiol.* 2002; 155:1081–1087. [PubMed: 12048221]
- Yogev G, Hausdorff JM, Giladi N. The Role of Executive Function and Attention in Gait. *Movement disorders : official journal of the Movement Disorder Society.* 2008; 23:329–472. [PubMed: 18058946]

Highlights

- We evaluated the relationship of engagement in individual and combined physical activity, mental activity, and social activity and physical function
- Combined physical and mental activity engagement was associated with better physical function, especially in gait speed.
- Future interventional research should investigate the combination of both physical and cognitive training to prevent decline of physical function in older adults.

Figure 1.

Class Difference in Activity Engagement (A) and Physical Function (B). Note. All data were Z-transformed. Higher value in activity engagement indicates more engagement; higher value in mobility outcome indicates better function.

Table 1

Latent Class Analysis Fit Statistics for Different Class Solutions of Activity Engagement.

Model	Latent Class	N (%)	AIC	BIC	Adjusted BIC	Entropy	LMR adjusted Likelihood ratio test, χ^2 test (p)
One-class	1	466 (100)					
Two-class	1	466 (100)	4079.39	4149.84	4095.89	1.00	0 (.50)
	2	0 (0)					
Three-class	1	59 (12.7)	4022.46	4113.63	4043.81	0.846	55.50 (.001)
	2	302 (64.8)					
	3	105 (22.5)					
Four-class	1	50 (10.7)	3963.30	4075.20	3989.51	0.880	66.98 (< .001)
	2	151 (32.4)					
	3	172 (36.9)					
	4	93 (20.0)					
Five-class	1	8 (1.7)	3887.56	4020.18	3918.62	0.768	2.81 (.57)
	2	148 (31.8)					
	3	143 (30.7)					
	4	45 (9.7)					
	5	122 (2.6)					

Table 2

Background Characteristics, Activity Engagement, Mobility Outcomes, and Cognitive Function as Total Sample and by Activity Engagement Class.

Variable	Total N = 466	Class				F or χ^2 value (p)
		“Active PA and MA” n = 50	“Active MA” n = 151	“Active PA” n = 172	“Inactive” n = 93	
Age, M(SD)	67.6 (7.51)	65.80 (6.91) ^a	67.75 (7.39) ^a	66.13 (6.84) ^a	71.04 (8.12) ^b	10.27 (< .001)
Male, n (%)	219 (47.0)	28 (56.0)	71 (47.0)	87 (50.6)	33 (35.5)	7.46 (.059)
Education, n (%)						17.80 (.007)
- high school or less	119 (25.6)	12 (24.0) ^{a, b}	28 (18.7) ^b	44 (25.7) ^{a, b}	35 (37.6) ^a	
- some college	137 (29.5)	18 (36.0) ^a	44 (29.3) ^a	44 (25.7) ^a	31 (33.3) ^a	
- college graduate or more	208 (44.8)	20 (40.0) ^{a, b}	78 (52.0) ^b	83 (48.5) ^b	27 (29.0) ^a	
Smoker, n (%)	40 (8.6)	6 (12.0)	11 (7.3)	13 (7.6)	10 (10.8)	1.86 (.60)
Taking antidepressant, n (%)	60 (12.9)	4 (8.0)	17 (11.3)	24 (14.0)	15 (16.1)	2.47 (.48)
Taking corticosteroid, n (%)	73 (15.7)	6 (12.0)	23 (15.2)	28 (6.0)	16 (3.4)	0.75 (.86)
Taking hypertension medication, n (%)	222 (47.6)	17 (34.0)	70 (46.4)	83 (48.3)	52 (55.9)	6.41 (.10)
Abdominal obese, n (%)	255 (54.7)	24 (48.0)	79 (52.3)	94 (54.7)	58 (62.4)	3.46 (.33)
MA [#] , M(SD)	2.27 (0.85)	2.36 (1.00) ^{a, b}	2.42 (0.80) ^b	2.18 (0.79) ^a	2.15 (0.93) ^a	2.99 (.031)
PA [#] , M(SD)	2.42 (1.37)	4.86 (0.25) ^a	1.99 (0.35) ^b	3.19 (0.37) ^c	0.51 (0.40) ^d	2014.01 (< .001)
SA [#] , M(SD)	1.15 (1.02)	1.10 (1.07)	1.29 (1.00)	1.13 (1.03)	1.01 (0.99)	1.99 (.12)
Grip strength (Kg/force) [#] , M(SD)	34.04 (11.24)	35.24 (12.26) ^a	34.29 (11.24) ^a	35.62 (11.51) ^a	30.05 (9.18) ^b	2.86 (.036)
Gait speed (feet/second) [#] , M(SD)	3.34 (0.72)	3.61 (0.69) ^{a, b}	3.32 (0.72) ^c	3.43 (0.72) ^{b, c}	3.09 (0.67) ^c	3.57 (.014)

Note.

[#] controlled for age; Letters indicate significant difference at p = 0.05 level in post-hoc analysis.