Published in final edited form as:

Angew Chem Int Ed Engl. 2016 August 16; 55(34): 10093–10097. doi:10.1002/anie.201605199.

Mixed Aggregates of the Dilithiated Koga Tetraamine: NMR Spectroscopic and Computational Studies

Dr. Yun Ma[‡], Kyle A. Mack[‡], Dr. Jun Liang[‡], Dr. Ivan Keresztes[‡], Prof. David B. Collum[‡], and Prof. Armen Zakarian[‡]

[‡]Chemistry and Chemical Biology, Cornell University, Ithaca, NY 14853-1301 (USA)

[‡]Chemistry and Chemical Biology, University of California, Santa Barbara, Santa Barbara, CA 93106 (USA)

Abstract

A combination of ^1H , ^6Li , ^{13}C , and ^{15}N NMR spectroscopies and density functional theory computations explores the formation of mixed aggregates by a dilithium salt of a C_2 -symmetric chiral tetraamine (Koga's base). Lithium halides, acetylides, alkoxides, and monoalkylamides form isostructural trilithiated mixed aggregates with few exceptions. ^6Li – ^{13}C and ^6Li – ^{15}N couplings reveal heretofore undetected transannular contacts (laddering) with lithium acetylides and lithium monoalkylamides. Marked temperature-dependent ^{15}N chemical shifts seem to be affiliated with this laddering. Computational studies shed light on the general structures, penchant for laddering, and stereochemical consequences of aggregation.

Graphical Abstract

Koga's chiral dilithium amide used in asymmetric synthesis is shown to promiscuously form mixed aggregates with a wide variety of lithium salts including those containing halides, alkoxides, acetylides, and amides. Although largely isostructural, the presence or absence of Li–X transannular contacts distinguishes ladder from cyclic motifs.

Keywords

Lithium; NMR Spectroscopy; Aggregation; Synthetic Methods; Enantioselectivity

C_2 -symmetric dilithiated tetraamine **1**, developed by Koga and coworkers^[1] as a chiral Brønsted base, has proven to be a versatile, non-covalent auxiliary for enantioselective synthesis.^[2,3] Amgen^[2] effected asymmetric alkylations of simple enolates en route to

Correspondence to: David B. Collum; Armen Zakarian.

Supporting information for this article is given via a link at the end of the document.

steroid dehydrogenase inhibitors [Eq. (1)], and one of us (AZ)^[3] developed enantioselective asymmetric alkylations and 1,4-additions of enediolates (Scheme 1).^[4] The exceptional selectivities prompted a collaboration to study the structures of putative mixed aggregates derived from dilithiated enediolates and dianion **1**.^[5] Despite the potential complexities of characterizing mixtures of *two* dianions, mixed aggregates **2** (containing *n*-BuLi) and **3** (containing dilithium enediolate) proved structurally tractable and yielded a computationally viable and predictive stereochemical model.^[5]

(1)

Instances in which highly stereoselective organolithium reactions have been traced to specific solvation and aggregation effects,^[6] control of aggregate structure appears to accompany—and possibly be a prerequisite for—high stereocontrol. With that hypothesis in mind, we surveyed the structures of mixed aggregates of **1** with a variety of lithium salts (Chart 1). Dianion **1** is a promiscuous pairing partner, affording isostructural mixed aggregates **4** or the corresponding ladder structures **5** (distinguished by a transannular Li–X contact).^[7] Spectroscopic and computational studies revealed structural nuances, including the stereochemical consequences of aggregation.^[8–10]

Mixed aggregation of **1** was examined with ⁶Li NMR spectroscopy. We routinely used minor excesses of *n*-BuLi-derived mixed aggregate **2** to generate the new mixed aggregates from LiX precursors, which caused low concentrations of **2** to persist in most recorded spectra. Otherwise, the spectra were routinely quite clean, as documented in the supporting information. The LiX salts **A–HH** (Chart 1) show a three-resonance fingerprint (1:1:1) characteristic of **4** or **5**. Throughout the paper, we refer to the mixed aggregates as **4**, **5**, or **4/5**, depending on context and the confidence levels of our assignments. The default attribution is **4**. We occasionally detected minor species—possibly other mixed aggregates—that were not pursued but are documented in the supporting information.

We balanced the depth of study against the scope of the survey. Five mixed aggregates prepared from salts **A–E** were deemed to be of special importance because of their role in current synthesis applications or because isotopically labeled LiX offered useful

spectroscopic probes. In these cases, we used the relatively precious labeled dilithium amide $[^{15}\text{N}_4]\mathbf{1}$ ^[5] to confirm the connectivities within the full aggregate structures^[11] and exploited a combination of COSY, TOCSY, HSQC, HMBC, and ROESY spectroscopies to obtain three-dimensional structures. Unexpected insights into the complex interplay between solvation and laddering came from temperature- and tetrahydrofuran (THF)-concentration-dependent ^6Li and ^{15}N chemical shifts. Assignments for salts **F–HH** were based on the presence of the characteristic three-peak pattern and analogy to the detailed studies. Dialkylamides **II–SS** failed to cleanly form mixed aggregates, displaying a variety of pathologies.

Density functional theory (DFT) calculations at the B3LYP/6–31G(d) level of theory^[12] with single-point calculations at the MP2 level of theory examined conformational properties and solvation. The results concurred with those described in detail elsewhere^[5] and are largely relegated to supporting information. A few especially pertinent computational results are salted throughout the text.

We begin the survey with a description of LiCl-derived mixed aggregate type **4A** (or **5A**; see Chart 1) owing to the key role of lithium halides as alkylation byproducts. ^6Li and ^{15}N NMR spectra of samples of **4A** showed ^6Li – ^{15}N coupling patterns akin to those noted for mixed aggregate **2** in a previous study (Figure 1A).^[5] Owing to slow aggregate exchanges, samples required aging; times varied depending on the conditions. The corresponding ^6Li spectrum with broadband ^{15}N decoupling revealed three resonances (1:1:1) along with the three resonances of residual **2** (Figure 1B). Single-frequency ^{15}N decoupling readily deconvoluted the splitting (supporting information) and provided the gross structure as **4A** (or **5A**). A combination of COSY, TOCSY, HSQC, HMBC, and ROESY spectroscopies of **4A** in THF- d_8 showed a three-dimensional structure consistent with the lowest-energy computed structure of **4A** (Figure 2.) Of the two possible diastereomers resulting from the chelate orientations [Eq. (2)], the observed form was the same as that noted previously.^[5] The preference illustrated in Equation (2) has proven immutable in all mixed aggregates prepared from dianion **1**.

(2)

Variable-temperature NMR spectroscopic studies revealed moderately temperature-dependent ^6Li chemical shifts and strikingly temperature-dependent ^{15}N chemical shifts (Figure 3). The two anionic amido ^{15}N resonances changed markedly with temperature, whereas the chelating piperidino resonances were unaffected. Given the results of studies of the lithium acetylides described in the next section, we suspect that a temperature-dependent **4A–5A** equilibrium with affiliated solvation number changes is operative.

Prompted by the temperature-dependent shifts and results obtained with lithium acetylides (below), we focused DFT computations on the relationship between laddering and solvation. We found a single minimum rather than discrete structures corresponding to **4A** and **5A**. The di- and trisolvated forms were, within the error of the computations, equal energy [Eq. (3)]. We noticed, however, that compared with the disolvated form (2.83 Å), the trisolvate showed a shortening of the transannular Li–Cl contact (2.63 Å). Mixed aggregates bearing a single THF are much less stable, with no evidence of Li–Cl contact. These issues become more prominent and spectroscopically tractable in the context of lithium acetylides discussed below. It may seem counterintuitive that laddering is affiliated with solvation, but laddering opens up the coordination sphere on the right-hand lithium.

(3)

Lithium acetylides brought laddering into full view. The mixed aggregate derived from lithium phenylacetylide (**B**) afforded the characteristic three resonances. Analogous spectra recorded with ^{15}N -labeled substrate showed the expected coupling for **4B** or **5B**. Spectra obtained with $[^6\text{Li}, ^{13}\text{C}]\text{PhCCLi}$ showed strong ^6Li – ^{13}C coupling, affording three ^6Li doublets and a ^{13}C heptet consistent with ladder structure **5B**. The transannular ^6Li – ^{13}C coupling ($J_{\text{Li-C}} = 5.2 \text{ Hz}$) is comparable to those of the other ($J_{\text{Li-C}} = 5.0$ and 5.3 Hz). This arrangement appears to stem from a static structure. Warming the sample, however, caused the transannular coupling to drop monotonically ($J_{\text{Li-C}} = 3.8 \text{ Hz}$ at -20°C while the other two doublets coalesced.^[13]

Variable-temperature NMR spectroscopic studies revealed temperature-dependent ^6Li and ^{15}N chemical shifts analogous to those noted for the LiCl-containing mixed aggregate (Figure 3). The two anionic amido ^{15}N resonances migrated downfield with warming, whereas the chelating piperidino resonances (N_1 and N_4) were unaffected. The two ^6Li resonances (Li_1 and Li_2) suggested to be THF solvated also shifted downfield with warming (although to quite different degrees), whereas the THF-free ^6Li resonance (Li_3) remained invariant.

To probe the role of solvation–desolvation, we carried out two types of experiments. Monitoring chemical shift versus THF concentration by using hexane as a cosolvent showed little change in chemical shift. Although pyridine is often a particularly useful chemical shift reagent for ^6Li NMR spectroscopy,^[14] preliminary studies of samples containing added pyridine were complex and unhelpful.

Computational studies uncovered a single discrete structure corresponding to ladder **5B**. The acetylide showed a distinct cant toward the three-coordinate lithium bearing one THF (no chelated piperidine). Whether this result derives from a stabilizing acetylide–lithium interaction or simply avoidance of steric problems is unknown, but arguments for steric bias stem from similar canting by lithium monoalkylamides and alkoxides (*vide infra*).

A low-resolution (1.04 Å) crystal structure was obtained by chance *after all spectroscopic and computational studies were complete*. Comparison of the solid-state structure with the computed lowest-energy form shows remarkable similarities, including a highly puckered ladder, a canted acetylide, and two THF ligands (Figure 4). A cursory survey of lithium acetylides **P–U** afforded spectra suggesting analogy to lithium phenylacetylide.

We focused studies of lithium alkoxides and related O-lithiated species on *n*-BuOLi as a potential (albeit unobserved) contaminant in *n*-BuLi. ^6Li NMR spectroscopy showed a three-resonance pattern characteristic of mixed aggregate **4C** or ladder **5C** along with low concentrations of *n*-BuLi-derived mixed aggregate **2** resulting from the method of preparation. [^6Li , ^{15}N]**1** showed multiplets and coupling patterns consistent with **4C/5C** (supporting information). Two-dimensional ^1H and ^{13}C NMR spectroscopies showed spatial relationships more consistent with **5C**, although they did not exclude **4C**. Temperature-dependent ^{15}N resonances akin to those observed for both lithium phenylacetylide and LiCl were observed. Computations converged on disolvated ladder-type aggregate **5C** with a 2.10 Å transannular Li–O contact irrespective of the starting geometry.

Cursory examination of assorted lithium alkoxides (**F–J**) and lithium phenolates (**K–N**) showed the characteristic three-resonance pattern. The only exception was highly hindered phenolate **4O**, which incorporates into a mixed aggregate to only a limited extent.

Studies of monoalkylamides first focused on lithium cyclohexylamide (**D**). ^6Li and ^{15}N NMR spectroscopy using mixed aggregates prepared from [$^{15}\text{N}_4$]**1** showed the usual three resonances and affiliated coupling patterns. Samples prepared from [^{15}N]**D**^[5] showed the expected strong coupling within the Li_3N_3 ring ($J_{\text{Li–N}} = 4.6$ and 3.7 Hz) as well as weak transannular coupling ($J_{\text{Li–N}} = 1.1$ Hz) indicative of laddering. Whether the weak coupling attested to a long contact or a dynamic exchange between the ladderized and unladderized forms could not be determined experimentally. Computations showed a single species with a long (2.87 Å) N–Li transannular contact for the lowest-energy disolvated form. (Three THF ligands would not bind.) Of the two possible orientations of the H and Cy substituents on nitrogen, the computationally most viable structure placed the N–H on the β -face and formed a ladder [Eq. (4)]. That orientation was confirmed with ^1H and ^{13}C NMR spectroscopies.

(4)

A complete spectroscopic analysis of mixed aggregates with both [$^{15}\text{N}_4$]**1** and [^{15}N]**E** showed the same patterns observed for lithium cyclohexylamide. In addition, spectra recorded at -100°C showed decoalesced resonances consistent with a 2:1 mixture of two isomers presumed to be isomeric chelates. Computations showed the relative energies of two logical isomers in [Eq. (5)]. Detailed ^1H and ^{13}C NMR spectroscopies failed to distinguish the two.

(5)

Monoalkylamides **V-HH** all formed discrete structures **4** or **5**. We briefly probed the stereochemical consequences of the stereogenic centers in phenethylamides **DD** and **EE**. Samples containing 1.0 equiv of each displayed a 10:1 preference for forming **5EE** [Eq. (6)]. This selectivity was good considering that the distinction of the two is based on subtle steric effects. One could envision schemes in which such distinctions might be used for kinetic resolutions.

Presumably owing to steric hindrance, the dialkylamides did not cleanly form mixed aggregates of type **4** or **5**. In some cases they may have been detected, but they were part of complex mixtures that were insufficiently important to pursue further.

(6)

Mixed aggregates of dilithiated tetraamine **1** have provided interesting structural comparisons through NMR spectroscopy, DFT computational chemistry, and X-ray

crystallography. The computationally and spectroscopically determined structures matched remarkably well. A crystal structure provided a pleasing correlation with the solid-state, solution, and computed structures. The results also offer a guide for exploiting mixed aggregates of Koga's base in enantioselective synthesis. A shocking reversal in selectivity with subtle changes in the ligand structure^[3a]—replacing the piperidines of **1** with pyrrolidines—underscores how little we understand the influence of aggregate structure and reaction mechanism on enantioselective transformations.^[14]

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

AZ thanks the National Institutes of Health (NIGMS GM077379), Eli Lilly, and Amgen for direct support of this work. DBC thanks the National Institutes of Health (NIGMS GM39764 and NIGMS GM077167). Both thank Amgen for generous supplies of Koga tetraamine.

References

1. a) Yamashita Y, Odashima K, Koga K. *Tetrahedron Lett.* 1999; 40:2803–2806. b) Yamashita Y, Emura Y, Odashima K, Koga K. *Tetrahedron Lett.* 2000; 41:209–213.
2. Frizzle MJ, Nania RR, Martinelli MJ, Moniz GA. *Tetrahedron Lett.* 2011; 52:5613–5616.
3. Stivala CE, Zakarian A. *J Am Chem Soc.* 2011; 133:11936–11939. [PubMed: 21744818] b) Lu P, Jackson JJ, Eickhoff JA, Zakarian A. *J Am Chem Soc.* 2015; 137:656–659. [PubMed: 25562717] c) Jackson JJ, Kobayashi H, Steffens SD, Zakarian A. *Angew Chem Int Ed.* 2015; 54:9971–9975. *Angew Chem.* 2015; 127:10109–10113.
4. For other applications of the Koga diamine in asymmetric synthesis, see: Swingle NM, Reddy KV, Rossiter BE. *Tetrahedron.* 1994; 50:4455–4466. Fehr C, Galindo J, Farris I, Cuenca A. *Helv Chim Acta.* 2004; 87:1737–1747.
5. Ma Y, Stivala CE, Wright AM, Hayton T, Liang J, Keresztes I, Lobkovsky E, Collum DB, Zakarian A. *J Am Chem Soc.* 2013; 135:16853–16864. [PubMed: 23654300]
6. a) Harrison-Marchand A, Mongin F. *Chem Rev.* 2013; 113:7470–7562. [PubMed: 23952819] b) Reich HJ. *Chem Rev.* 2013; 113:7130–7178. [PubMed: 23941648]
7. For a general discussion of the ladder principle in lithium amides, see: Mulvey RE. *Chem Soc Rev.* 1991; 20:167–209.
8. For leading references and discussions of mixed aggregation effects and their applications in synthesis, see: Seebach D. *Angew Chem Int Ed Engl.* 1988; 27:1624–1654. *Angew Chem.* 1988; 100:1685–1715. Reich HJ. *J Org Chem.* 2012; 77:5471–5491. [PubMed: 22594379]
9. Dilithium amides showing structural analogies with **4** and their mixed aggregates have been studied computationally: Pratt LM, Mu R, Carter C, Woodford B. *Tetrahedron.* 2007; 63:1331–1338.
10. For references to crystallographic studies of dilithium amides, see: reference 5.
11. Collum DB. *Acc Chem Res.* 1993; 26:227–234.
12. Frisch, MJ., et al. *GaussianVersion 3.09*; revision A.1. Gaussian, Inc; Wallingford, CT: 2009.
13. For theoretical investigations of ⁶Li–¹⁵N coupling constants, see: Parisel O, Fressigné C, Maddaluno J, Giessner-Prettre C. *J Org Chem.* 2003; 68:1290–1294. [PubMed: 12585867] Koizumi T, Morihashi K, Kikuchi O. *Bull Chem Soc Jpn.* 1996; 69:305–309.
14. Tallmadge EH, Collum DB. *J Am Chem Soc.* 2015; 137:13087–13095. and references therein. [PubMed: 26437278]

Figure 1. ^6Li NMR spectra of 0.10 M $[^6\text{Li}, ^{15}\text{N}]\mathbf{4/5A}$ prepared from 0.10 M *n*-BuLi-derived mixed aggregate $[^6\text{Li}, ^{15}\text{N}]\mathbf{2}$ and 0.10 M $[^6\text{Li}]\text{LiCl}/i\text{-Pr}_2\text{NH}$ in neat THF- d_8 . The spectra were recorded at -70°C after aging at -78°C for 12 h: (A) ^6Li spectrum; (B) $^6\text{Li}\{^{15}\text{N}\}$ spectrum. $\text{LiCl}/i\text{-Pr}_2\text{NH}$ was generated in situ from $\text{Et}_3\text{N}\cdot\text{HCl}$ and $[^6\text{Li}]\text{LDA}$ (LDA = lithium diisopropylamide), with regeneration of the visible LDA by *n*-BuLi as denoted by *.

Figure 2.
DFT calculated structure of **4/5A**.

Figure 3.

Variable temperature ^{15}N NMR spectra of 0.10 M $[\text{}^6\text{Li},^{15}\text{N}]\mathbf{4/5A}$ prepared from 0.10 M *n*-BuLi-derived mixed aggregate $[\text{}^6\text{Li},^{15}\text{N}]\mathbf{2}$ and 0.10 M $[\text{}^6\text{Li}]\text{LiCl}/i\text{-Pr}_2\text{NH}$ in neat $\text{THF-}d_8$ after aging at $-78\text{ }^\circ\text{C}$ for 12 h. $\text{LiCl}/i\text{-Pr}_2\text{NH}$ was generated in situ from $\text{Et}_3\text{N}\cdot\text{HCl}$ and $[\text{}^6\text{Li}]\text{LDA}$.

Figure 4.
Overlap of DFT-computed and crystallographically characterized structures of **5B**.

Scheme 1.