Commentary: patient's view

Maria Tomasa with the support of Giorgina Barbara Piccoli

My nickname is Tommy. At the time of the story I was 35 years old.¹ I have been diabetic almost all my life, and this has had a big effect on me. I have wanted a baby since the day of my marriage, or even before that. Someone asked me if I wanted a baby to feel more "normal." That is not the reason. I wanted a baby because it is a fundamental part of life and because I needed a child. I wanted it. When you deeply want something you have no particular reasons for doing it; you just want it day and night.

Over the past few years, I have spent money and time trying to have a baby. Most of the doctors I met said that it was too risky and that I shouldn't even think about it. I therefore changed doctors several times, trying to find one who would say "yes," until I found Giorgina and her father. They finally understood my opinions. They are special people.

I don't know, but maybe it's also because Giorgina has two children (and when I first met her, her first child was a baby). None of the previous doctors, for example, was able to answer this question: "What does high risk, or too risky mean?"

I knew that I would be risking my life and the life of the baby, but I also knew that I could make it. I was sure of that, and you have to follow your feelings. Even if the risk was as high as 70%, I wanted to try.

I understood why the doctors explained all the risks, but my wish was much stronger than they could understand. I think that the doctors should have

supported my decision instead of trying to make me change my mind, sometimes too forcefully. Now my child is 2 years old. I call him many things: *topolino*, *ciplollino* (little mouse, little onion); he is very sweet. The first time I saw him I thought: "It's impossible it's my child. He's too beautiful." And the nurses laughed and said: "No, of course he's yours." He was small and was in the emergency room.

The first time I touched him, he had a tachycardic crisis. The doctor said that he probably needed a surgical intervention for his heart; the surgeon came over, and he talked to me. After 24 hours' observation, they told me that it was only the contact with my hands that had excited him. That day I thought I was dying, and I thought, "God, take me, not him." He was well by the late evening.

Last week I went on the waiting list for a pancreas-kidney transplant. I'm scared, even though I know it's the best thing to do. I'm scared for my child, and I hope that this time too everything will be all right. I've been happy to let Giorgina tell my story, because I hope it will be useful for somebody else, perhaps someone who has diabetes and wants a baby.

Competing interests: Giorgina Piccoli is Tommy's doctor and is godmother to Tommy's son. Both of them are very happy.

1 Piccoli GB, Mezza E, Grassi G, Burdese M, Todros T. A 35 year old woman with diabetic nephropathy who wants a baby: case outcome. *BMJ* 2004;329:900.

Chair of
Nephrology,
Department of
Internal Medicine,
University of Turin,
Corso Dogliotti,
14-10126 Turin,
Italy

Giorgina Barbara
Piccoli
nephrologist

Correspondence to:
G B Piccoli
gbpiccoli@
hotmail.com

Commentary: when do we say "no"?

Thomas R Aust

Tommy's doctors made the right decision to support her in trying for a natural pregnancy by optimising her diabetic treatment and advising her of the risks of becoming pregnant.¹ This is because many patients will carry on and conceive with or without their doctors' blessing. After all, we would prefer it if drug misusers didn't inject heroin, but if they must we would rather they use clean needles.

Tommy's request for assisted conception was a different matter because all complications of assisted conception are, by definition, iatrogenic. Her doctors had to consider whether they would be prepared to help her produce a potentially damaged child or even kill herself in the process. As reproductive physicians we are never absolved from this moral responsibility, even when patients have given their consent and fully understand the risks.

One respondent felt that Tommy's (fetotoxic) angiotensin converting enzyme inhibitor should not have been stopped while she was trying for a natural conception but continued until she became pregnant.² I feel that if her remaining fallopian tube had been checked for patency and found to be blocked, her

angiotensin converting enzyme inhibitor could have been continued to optimise her blood pressure control until she started treatment for in vitro fertilisation.

One rapid respondent suggested that only two rather than three embryos should be transferred to reduce the chance of multiple pregnancy and thus the risks of prematurity, pre-eclampsia, and miscarriage (among others).³ One could even have considered replacing a single embryo (and freezing any remaining embryos) to reduce this risk even further. In Sweden, where mandatory single embryo transfer has been introduced, excellent pregnancy rates have been reported but with a much reduced twinning rate.³

The ovarian response at the time of egg collection often gives an indication of the risk of developing ovarian hyperstimulation syndrome. The decision to freeze or replace embryos (and risk worsening ovarian hyperstimulation syndrome if a pregnancy occurs) around this time is difficult as the success rates after replacement of frozen embryos tend to be poorer.⁴ Unfortunately, recent legislation changes in Italy have removed this option for safety⁵ and would drastically alter Tommy's management if she were treated today.

Hewitt Centre for
Reproductive
Medicine, Liverpool
Women's Hospital,
Liverpool L8 7SS
Thomas R Aust
research fellow

tomaust@
doctors.org.uk