Firstly, the imperative to “do no harm” is not straightforward in this case. As Trisha Greenhalgh points out, there are risks involved whatever choice is made, and the weight that should be given to these risks is a personal assessment.² So the imperative not to harm is not clearly in contradiction with patient self determination.³ Tommy knows best what would be most harmful to her: the possible adverse outcomes of pregnancy or not having this child.

Commentary: beyond the evidence

Gillian Hawthorne, Maggie Blott

Tommy's story shows the difficulty that is often encountered when looking after women with complex medical problems in pregnancy.¹ There is simply insufficient evidence to customise the personal risk for a woman with diabetic complications who wishes to become pregnant. Women like Tommy need consensual advice and support from a highly skilled, well informed, multidisciplinary team with a good understanding of the complexity of the issues and the necessary skills to provide the required care. Good preconception care should, and in this case did, underpin the management and must include counselling about risk. This is not a generalist case, and we are not surprised that there were few rapid responses to this case.

Tommy changed her diabetes team in search of support for her desire for a baby. Once she made her decision, the ongoing support of the care team was paramount, whether or not they fully agreed with her. The failure to refer Tommy earlier for assisted conception when it was clear that there was a mechanical barrier to conception may have been a symptom of the team's ambivalence but not of their lack of commitment.

Assessment of diabetic control, diabetic retinopathy, hypertension, cardiac status, and renal function are vital before pregnancy. Glycaemic control can be optimised, blood pressure stabilised, and folic acid started. Ischaemic heart disease is an absolute contraindication to pregnancy, as suggested by the rapid responses. Angiotensin converting enzyme inhibitors can be continued until the pregnancy is diagnosed; they are not known to be teratogenic.²

Reports suggest that outcome of pregnancy is generally good in women with mild to moderate

This also reminds us that evidence based medicine is a source of factual knowledge. No less, but no more. It will not, by itself, solve ethical difficulties. Knowing what should be done certainly includes knowing the facts—the team was correct to go to the literature with such care—but the difficulties do not end there.⁴

This difficulty with medical decision making is not, of course, Tommy's problem. Her decision, whether to endanger her body in order to give birth, is one that women were already facing long before people started writing about it. Only recently did inventions of the human mind make this risk more distant and less generally accepted.

Contributors: SAH wrote the first draft of this commentary. Both authors contributed to the intellectual content and approved the final version. SAH is the guarantor.

Competing interests: None declared.

- 1 Piccoli GB, Mezza E, Grassi G, Burdese M, Todros T. A 35 year old woman with diabetic nephropathy who wants a baby: case outcome. *BMJ* 2004;329:900.
- 2 Greenhalgh T. How strong is the evidence? [electronic response to Piccoli GB, Mezza E, Grassi G, Burdese M, Todros T. A 35 year old woman with diabetic nephropathy who wants a baby: case progression]. *BMJ* 2004 <http://bmj.com/cgi/eletters/329/7468/7293>.
- 3 Buchanan AE, Brock DW. *Deciding for others: the ethics of surrogate decision making*. Cambridge: Cambridge University Press, 1990.
- 4 Eddy DM. Clinical decision making: from theory to practice. *Anatomy of a decision*. *JAMA* 1990;263:441-3.

diabetic renal insufficiency, with a 90% take home baby rate.³ The woman would be at risk from pre-eclampsia, with attendant risks of intrauterine growth restriction and iatrogenic prematurity for the fetus. Aggressive control of blood pressure is therefore essential. No consensus exists on the effect of pregnancy on progression of diabetic nephropathy to end stage renal failure, although women with more severe disease are less likely to recover renal function after delivery of the baby.^{4 5}

The risks of in vitro fertilisation in women with diabetic nephropathy have not been evaluated. As women with diabetic nephropathy have a shortened life expectancy, however, issues surrounding the welfare of the child must be discussed before treatment.

Pregnancy is not without risk. Care must be tailored to the individual and delivered by a team with sufficient expertise to provide the skilled support that women need.

Contributors: GH and MB discussed the case and jointly wrote the commentary.

Competing interests: GH and MB hold a preconception clinic at Newcastle Diabetes Centre

- 1 Piccoli GB, Mezza E, Grassi G, Burdese M, Todros T. A 35 year old woman with diabetic nephropathy who wants a baby: case outcome. *BMJ* 2004;329:900.
- 2 Lip GYH, Churchill D, Beevers M, Auckett A, Beevers DG. Angiotensin converting enzyme inhibitors in early pregnancy. *Lancet* 1997;350:1446-7.
- 3 Dunne FP, Chowdhury TA, Hartland A, Smith T, Brydon PA, McConkey C, Nicholson HO. Pregnancy outcome in women with insulin-dependent diabetes mellitus complicated by nephropathy. *Q J Med* 1999;92:451-4.
- 4 Biesenbach G, Grafinger P, Stoger H, Zazgornik J. How pregnancy influences renal function in nephropathic type 1 diabetic women depends on their pre-conceptual creatinine clearance. *J Nephrol* 1999;12:41-6.
- 5 Rossing K, Jacobsen P, Hommel E, Mathiesen E, Svenningsen A, Rossing P, et al. Pregnancy and progression of diabetic nephropathy. *Diabetologia* 2002 45:36-41.

Newcastle Diabetes Centre, Newcastle General Hospital, Newcastle upon Tyne NE4 6BE

Gillian Hawthorne
consultant
diabetologist

Department of Obstetrics, Royal Victoria Infirmary, Newcastle upon Tyne NE1 4LP

Maggie Blott
consultant
obstetrician

Correspondence to:
G Hawthorne
gillian.hawthorne@
newcastle-pct.nhs.uk