Published in final edited form as:

Med Care. 2017 February ; 55(2): 173–181. doi:10.1097/MLR.0000000000000638.

Perceived Discrimination in Healthcare and Mental Health/ Substance Abuse Treatment Among Blacks, Latinos, and Whites

Vickie M. Mays, Ph.D., MSPH [Professor],

Psychology and Health Policy and Management, UCLA Fielding School of Public Health and UCLA Center for Bridging Research, Innovation, Training and Education for Minority Health Disparities Solutions, UCLA; 405 Hilgard Avenue, Box 951563 Los Angeles, CA 90095-1563; mays@ucla.edu; Telephone: 310-206-5159; Fax: 310-206-5895

Audrey Jones, Ph.D.,

Center for Health Equity Research and Promotion, VA Pittsburgh Healthcare System— and UCLA Center for Bridging Research Innovation, Training and Education for Minority Health Disparities Solutions, Address: University Drive 151C, Building 30, Pittsburgh, PA 15240-1001

Ayesha Delany-Brumsey, Ph.D.,

Vera Institute of Justice and UCLA Center for Bridging Research Innovation, Training and Education for Minority Health Disparities Solutions, Address: 233 Broadway 12th Floor, New York, NY 10279

Courtney Coles, MPH, and

Department of Health Policy and Management, UCLA Fielding School of Public Health and UCLA Center for Bridging Research Innovation, Training and Education for Minority Health Disparities Solutions, Address: 31-269 Center for the Health Sciences, Box 951772, Los Angeles, CA 90095-1772

Susan D. Cochran, Ph.D., MS [Professor]

Epidemiology and Statistics, UCLA Fielding School of Public Health and UCLA Center for Bridging Research Innovation, Training and Education for Minority Health Disparities Solutions, Address: UCLA Department of Epidemiology, 650 Charles E. Young Dr., UCLA School of Public Health, Los Angeles, CA 90095

Abstract

Background—Experiences of discrimination in healthcare settings may contribute to disparities in mental health outcomes for Blacks and Latinos. We investigate whether perceived discrimination in mental health/substance abuse visits contributes to participants' ratings of treatment helpfulness and stopped treatment.

Research Methods—We used data from three waves of the California Quality of Life Survey, a statewide population-based telephone survey assessing mental health/substance disorders and their treatment. In a sample of 1,099 adults (age 18–72) who indicated prior year mental health/

Corresponding Author: Vickie M. Mays Ph.D., MSPH; Address: UCLA, Department of Psychology, 405 Hilgard Avenue, Box 951563 Los Angeles, CA 90095-1563, mays@ucla.edu, Telephone: 310-206-5159.

No potential conflicts of interest exist.

substance abuse visits, we examined: experiences of discrimination that occurred during healthcare and mental health/substance abuse visits, ratings of treatment helpfulness, and reports of stopping treatment early.

Results—Fifteen percent of California adults reported discrimination during a healthcare visit and 4% specifically during mental health/substance abuse visits. Latinos, the uninsured, and those with past-year mental disorders were twice as likely as others to report healthcare discrimination (Adjusted Odds Ratio [AORs]=2.08, 2.77 and 2.51). Uninsured patients were seven times more likely to report discrimination in mental health/substance abuse visits (AOR=7.27, $p<.01$). The most commonly reported reasons for healthcare discrimination were race/ethnicity for Blacks (52%) and Latinos (31%), and insurance status for Whites (40%). Experiences of discrimination in mental health/substance abuse visits were associated with less helpful treatment ratings for Latinos (AOR=0.09, $p<.05$) and Whites (AOR=0.25, $p<.01$), and early treatment termination for Blacks (AOR=13.38, $p<.05$).

Conclusions—Experiences of discrimination are associated with negative mental health/substance abuse treatment experiences and stopped treatment, and could be a factor in mental health outcomes.

Keywords

race/ethnicity; provider discrimination; mental health; substance abuse

Introduction

Significant attention is focused on racial/ethnic disparities in mental health status and mental health services use.^{1,2} Blacks and Latinos are less likely than Whites to receive mental health or substance abuse services and are more likely than Whites to delay seeking care.^{3–5} Even when Blacks and Latinos do receive mental health/substance abuse services, they are more likely than Whites to obtain inappropriate diagnoses,^{6,7} drop out of treatment early,⁸ report less satisfaction with treatment,⁹ and receive inadequate or substandard care.¹⁰ Untreated, mental health/substance use disorders lead to economic instability,¹¹ disruptions in education and employment,^{12,13} and premature mortality due to suicide, chronic disease and poor physical health outcomes.^{14,15} For Blacks and Latinos specifically, disparities in treatment may be a factor in why these groups experience elevated levels of psychological distress,¹⁶ more chronic psychiatric disorders,¹⁷ and more frequent use of emergency psychiatric services¹⁸ compared to Whites. Given the significant personal and societal costs of untreated mental health/substance use disorders, there is a need to determine modifiable factors within the healthcare system that may be leveraged to better meet the care needs of racial/ethnic minorities.

To date, researchers have identified a number of barriers that contribute to higher levels of unmet behavioral health needs among racial/ethnic minorities.¹⁹ These include access due to a lack of insurance,²⁰ unavailability of services in their native language,²¹ greater burden of stigma^{22,23} and historical mistreatment of in health care settings which may engender distrust of mental health professionals.²⁴ Another possible barrier to care that has received relatively less attention is whether experiences of discrimination in mental health/substance

abuse services contribute to the greater level of unmet mental health needs of racial/ethnic minorities.

Previous research on discrimination in physical health care settings has shown that racial/ethnic minorities report experiencing discrimination more frequently than Whites,^{25–27} with common reasons being race/ethnicity, insurance type, ability to pay, and language.^{28,29} These experiences of discrimination in the medical setting are associated with negative ratings of provider communication,^{26,30} treatment dissatisfaction³⁰, avoiding or delaying treatment,^{25,31} and poor physical health outcomes.^{25,32} Two studies in the United Kingdom found that experiences of discrimination occur frequently in mental health service settings, contributing to patients' mistrust with mental health services and disruptions to the therapeutic relationship.^{33,34} Less known is the extent to which negative healthcare outcomes occur as a function of experiences of discrimination in mental health/substance abuse treatment for Blacks and Latinos. Therefore, one area ripe for exploration is to determine whether, for Blacks and Latinos, there is a relationship between perceived experiences of discrimination, treatment noncompliance, dissatisfaction, dropout, and seeking of future treatment.

The purpose of this study is to investigate the prevalence of discrimination occurring in mental health/substance abuse service visits and to examine associations of perceived discrimination in mental health/substance abuse visits with treatment experiences in an ethnically diverse population-based sample of adults. We hypothesize that racial/ethnic minorities will report more experiences of discrimination in the general healthcare and mental health/substance abuse visits than Whites. Moreover, we hypothesize that experiences of discrimination in a mental health/substance abuse service visit will be associated with racial/ethnic minorities reporting less helpful treatment ratings and stopping treatment early.

Methods

Data

We use data from the California Quality of Life Surveys (Cal-QOL) I–III, three cross-sectional population-based mental health surveys of English and Spanish speaking adults in California.³⁵ The Cal-QOLs are follow-backs to the 2003, 2007, and 2011 waves of the California Health Interview Survey (CHIS), a biennial population-based random-digit dial telephone survey of more than 40,000 adults in California, age 18 or older.³⁶ The CHIS collects data on a range of health-related topics including health status, chronic conditions, health-related behaviors, healthcare access and use of health services. The Cal-QOLs interviewed a subset of CHIS participants to obtain detailed information on mental health and substance abuse disorders, treatment, and experiences of discrimination. The Cal-QOL sampling frames selected by probability methods are a subset of adults who are age 18 to 72 at the time of their CHIS participation. Stratification methods were used to oversample African Americans in the last two waves. All eligible CHIS respondents were originally interviewed in English or Spanish and agreed to re-contact. The Cal-QOL collected data between October 2004 and February 2005 (Wave I, n=2,322), October 2007 until February

2008 (Wave II, n=2,815), and January 2012 until February 2013 (Wave III, n=2,449). The response rate for each survey was 56%.

Sample

For this study, participants included adults, age 18 to 72, who indicated they used mental health or substance abuse services in the past year and answered a series of questions on discrimination in healthcare (see Figure 1). Overall, 7,586 adults participated in the Cal-QOL I–III. We excluded 356 participants who received a shortened interview lacking critical study variables. Of the remaining 7,228 participants, 15% reported past year use of mental health or substance abuse services, leaving a final sample of 1,099 individuals: 817 Non-Latino White, 136 Latino, 89 Non-Latino Black, and 57 participants classified as ‘Other’ race/ethnicity (40 Asian American and 17 Native American).

Measures

Discrimination in Healthcare Visits—Three items assessed discrimination occurring in a healthcare setting visit in the 12 months prior to the interview. Participants were asked whether: 1) a healthcare worker or provider acted as if they were uncomfortable with the respondent, 2) a healthcare worker or provider made negative, hostile, or disrespectful comments, or 3) the respondent felt discriminated against, treated disrespectfully, or given less good care than other people. These items originated from the Healthcare for Communities Survey, a Robert Wood Johnson Foundation-sponsored national study of mental health and substance abuse services.³⁷

Individuals who endorsed any of the discrimination items were asked what they thought was the reason for the disparate treatment, and whether any of their experiences with discrimination occurred at a visit for emotional, mental health, or alcohol/drug problems. We coded respondents who did not report any discrimination in the initial three questions as also not experiencing discrimination in services directed toward health, mental health or substance use concerns.

Treatment Experiences—Individuals receiving mental health or substance abuse services in the year prior to the interview were also asked to rate the helpfulness of their care on a Likert scale: “very helpful”, “somewhat helpful”, “neither helpful nor not helpful”, or “not helpful at all”.³⁷ Early treatment termination was assessed through self-report. Patients who reported that they stopped or chose to get less treatment than recommended by their provider for their mental health or substance use concerns were considered to have stopped treatment early. These items were administered in Cal-QOL II and III only (n=745).

Participant Characteristics—Several demographic characteristics were measured including age, gender, education, income, and insurance status. We divided age into four cohorts (18–29, 30–44, 45–59, 60–72), and education into four levels (less than high school education, high school degree or some college, four or more years college), and households with low income were determined by those earning less than 200 percent of the Federal Poverty Level adjusted for survey year. We coded insurance coverage into three categories: public, private, or uninsured. Public insurance included Medicare, MediCal, and Healthy

Families, as well as the military care programs (VA, Tricare). Private insurance included employer-based or privately purchased coverage. As described elsewhere, the Cal-QOL used the Composite International Diagnostic Interview Short Form (CIDI-SF) and other screening measures to determine the prevalence of eight past year mental health and substance use disorders: Major Depressive Disorder, Generalized Anxiety Disorder, Panic Attack, Post-Traumatic Stress Disorder, Alcohol Abuse or Dependence, or Drug Abuse or Dependence.³⁸

Statistical Analyses

We used Stata version 13.1 to conduct all statistical analyses. Sampling weights were applied to adjust for the probability of selection, nonresponse, and for post-stratification to generate estimates representative of the California population.

We first used cross-tabulations to examine the overall prevalence of any healthcare discrimination and then experiences of discrimination in mental health/substance abuse visits. Next, to test our hypothesis that racial/ethnic minorities will be more likely to report experiencing discrimination compared to Whites, we used logistic regression models to examine the association between race/ethnicity and reports of discrimination in any healthcare and in a mental health/substance abuse visit. Using multivariate methods, we controlled for age, gender, sexual orientation, education, income, insurance status, and positive screenings for a mental health/substance abuse disorder to obtain adjusted estimates.

We employed multivariate methods to test our hypothesis that perceived discrimination during a mental health/substance abuse visit would be associated with ratings of treatment as less helpful and early termination for Blacks and Latinos. To test the association between experiences of discrimination and ratings of treatment helpfulness we used multivariate ordinal regression, and multivariate logistic regression were used to test the association between experiences of discrimination and stopping treatment early. We ran these models separately for Blacks, Latinos, and Whites. Also, because the treatment questions were only administered in Cal-QOL II–III (n=745), these final analyses were restricted to this group only.

Since only five percent of participants (n=59) had missing data on one of the variables of interest, we used multiple imputation procedures to estimate the missing data with the assumption that missing values were missing at random. Analyses were conducted using the *mim* package in Stata which averages the model estimates across imputed datasets and produces pooled standard errors according to Rubin's rules.³⁹

Results

Prevalence of Perceived Discrimination in Healthcare or Mental Health/Substance Abuse Visits

We began by investigating the prevalence of perceived discrimination during any healthcare or mental health/substance abuse visits (see Table 1). The results showed that, overall, approximately one in seven participants (14.9%) reported that they had experienced discrimination in a healthcare visit in the past year. Approximately one in four Black and Latino patients reported experiences of discrimination in the healthcare setting (24% and

25% respectively) compared to 1 in 9 Whites (12%) and participants of other race/ethnicity (11%), $p = .02$. Discrimination in a healthcare visit was also frequently reported among those with less than high school education (24%), some college education (21%), and those with no insurance (27%) or public insurance (19%). Moreover, participants who met the criteria for a psychiatric disorder were more likely to report healthcare discrimination than those without a psychiatric disorder (19% vs 7%, $p < .001$). In particular, we found elevated reports of healthcare discrimination among adults who met criteria in the past year for Generalized Anxiety Disorder (27%), Panic Attack (24%), Post-traumatic Stress Disorder (24%), or Major Depressive Disorder (21%).

Among participants who reported any healthcare discrimination, 27% also reported that the discrimination occurred in a mental health/substance abuse visit. Thus, four percent of adults who used mental health or substance abuse services in the past year reported an experience of discrimination in a visit for mental health or substance abuse concerns. As shown in Table 1, the prevalence of discrimination in a mental health/substance abuse visit was three times higher for Blacks (9%) and Latinos (8%) than Whites (3%), $p < .01$. Discrimination in a mental health/substance abuse visit was more frequently reported by uninsured participants (16%) than by those with public (4%) or private insurance (2%), $p < .01$. In addition, participants who met screening criteria for a psychiatric disorder reported discrimination in a mental health/substance abuse visit more frequently than participants who did not meet screening criteria (6% vs 1%, $p < .01$). Higher reporting of discrimination in a mental health/substance abuse visit was found in those with Generalized Anxiety Disorders (8%) and Post Traumatic Stress Disorder (9%).

Predictors of Perceived Discrimination in Healthcare and Discrimination in a Mental Health/Substance Abuse Visit

Multivariate models were used to test for racial/ethnic disparities in healthcare discrimination after controlling for other sociodemographic characteristics (see Table 2). In the adjusted analyses, the only significant race/ethnicity findings was for Latinos who had two times greater odds of experiencing healthcare discrimination than Whites, $AOR = 2.08$ (95% CI = 1.02–4.25). Significant differences in experiences of healthcare discrimination were also observed for the uninsured, $AOR = 2.51$ (95% CI = 1.19–5.29), and those with a mental health disorder in the past year, $AOR = 2.77$ (95% CI = 1.50–5.11).

We did not find support for the hypothesis of Black-White or Latino-White differences in discrimination in a mental health/substance abuse visit after controlling for sociodemographic characteristics. However, we did find that patients of other races compared to Whites had a significantly lower odds of reporting discrimination experiences in a mental health/substance abuse visit ($AOR = 0.01$, 95% CI = 0.00–0.12). The uninsured had seven times greater odds of experiencing discrimination in a mental health/substance abuse visit than those with private insurance ($AOR = 7.27$, 95% CI = 2.27–23.30).

Perceived Basis of Discrimination in Healthcare Visits

Racial/ethnic differences were found in perceived attributes for the discrimination (see Table 3). While specific reasons for discrimination during a mental health/substance abuse were

not asked in this study, we did, however, find that among Blacks the most commonly reported reason for discrimination during a healthcare visit was race/ethnicity (52%), followed by insurance type (30%) and income (29%). Latinos reported race/ethnicity (31%) most often, followed by income (22%), and insurance (15%). Among Whites, a different pattern emerged with insurance (40%) as the most reported reason for discrimination, followed by not-specified reasons (22%) and age (11%).

Associations of Perceived Discrimination in Mental Health/Substance Abuse Visits with Treatment Helpfulness and Stopped Treatment

In determining the role of discrimination in participants treatment behaviors using data from only Cal-QOL II and III, nine percent of the participants rated their mental health/substance abuse treatment as “not helpful at all” or “neither helpful nor not helpful”: 25% Blacks, 8% Latinos, 7% of Whites. Those Latinos and Whites who reported experiencing discrimination in their mental health/substance visit rated their treatment as less helpful than those who did not report discriminatory experiences (AOR = 0.09, 95% CI = 0.01–0.84, AOR = 0.25, 95% CI = 0.09–0.67) (see Table 4). For Blacks, no significant relationship was found relative to their experiences of discrimination in mental health/substance abuse visits (AOR=0.19, 95% CI = 0.03–1.48).

Last, we examined participants’ reports of stopping treatment early. One in five participants (19.6%) in Cal-QOL II–III reported that they stopped their mental health/substance abuse treatment in the past year earlier than recommended: 21% for Blacks, 21% for Latinos, and 18% for Whites. Blacks who reported discrimination during a mental health/substance abuse visit times had 13 times greater odds of stopping treatment early compared to Blacks who did not report such experiences of discrimination (AOR = 13.38, 95% CI = 1.36–131.99). These same types of experiences of discrimination were not found to be statistically associated with stopping treatment early for Latinos (AOR = 2.06, 95% CI = 0.08–51.97) or Whites (AOR = 0.66, 95% CI = 0.09–5.09).

Discussion

The results of our study, which investigated patients reporting of discrimination in general healthcare and mental health/ substance abuse visits in a population-based sample of California adults, found support for our hypothesis that patient reporting of discrimination is associated with stopping treatment early and perceptions that treatment is not helpful. The most frequently given attributions for Blacks and Latinos for the basis of their discrimination was race/ethnicity, while for Whites it was their insurance status. Within the context of mental health/substance abuse services, our results demonstrate that experiences of discrimination are associated with negative perceptions of treatment helpfulness for Latinos and Whites and early treatment termination for Blacks. Importantly, our findings suggest that health care discrimination may contribute to disparities in mental health/substance abuse treatment outcomes for marginalized populations.

While the overall rates of discrimination in a mental health or substance abuse visit appear relatively low, these experiences occur two to three times more frequently in populations who already experience inequities in the access to and quality of mental health/substance

abuse services.^{19,20} Indeed, Latinos, individuals who were uninsured, and those who suffered from a mental health disorder, such as Generalized Anxiety Disorder, experienced the highest prevalence of healthcare discrimination. In addition, the results suggest that the uninsured, many of whom are racial and ethnic minorities, are at greater risk for experiencing discrimination specifically during a mental health/substance abuse service visit. As increasing numbers of Blacks and Latinos enter integrated primary care services through the Affordable Care Act (ACA)'s Medicaid expansion,⁴⁰ our results caution that the hoped for reduction in mental health disparities could be compromised in the face of perceived discrimination in the healthcare visit.

Prior studies of experiences of discrimination during physical health services indicate that discriminatory experiences damage the patient/provider relationship,^{26,30} contribute to patients' treatment dissatisfaction,³⁰ medication noncompliance,²⁹ and contribute to poor health outcomes.^{25,32} In mental health, our findings indicate that experiences of discrimination in a mental health/substance abuse visit also can contribute to negative perceptions of treatment for Latinos and Whites, and even worse to early treatment discontinuation for Blacks. This has important implications for possible health outcomes as prior research indicates that mental health treatment satisfaction and treatment continuation are associated with better physical and mental health outcomes.^{41,42} Therefore, improving the treatment experiences of vulnerable populations in mental health and substance abuse services may be a critical step in reducing mental health disparities, particularly for uninsured and racial/ethnic minority patients.

Our findings are also important in the context of the ACA where patient satisfaction is an important measure of the quality of care, particularly in the hospital setting where 30 percent of hospitals' Medicare reimbursements are tied to patient satisfaction survey scores.⁴³ Future research is needed to better understand patient preferences and systems of care dynamics that contribute to enhanced treatment experiences for racial/ethnic minorities, those with public insurance, and those with psychiatric disorders.⁴⁴ Incorporating patient-centered approaches that are culturally sensitive to the treatment of mental disorders in primary and specialty mental health care settings may be one strategy to improve treatment experiences and mental health outcomes for these vulnerable populations.⁴⁵ In addition, implementing training programs to increase the cultural competence of providers has been shown to improve patient satisfaction with care.⁴⁶ Educating providers about the negative effects of discrimination that occurs during mental health/substance abuse services delivery, as well as the subtle and often unconscious nature of bias, should be incorporated into such training programs.⁴⁷ In addition to cultural competency and bias reduction training, increasing the number of providers from diverse backgrounds, including racial/ethnic minorities, may increase utilization of treatment services and decrease rates of perceived discrimination and premature treatment drop out.⁴⁸ Collectively, these efforts may hold promise for reducing mental health/ substance abuse services disparities not only for racial/ethnic minorities for but also for others who encounter discrimination in mental health/substance abuse services.

Study limitations deserve consideration in contextualizing our results. First, we note factors that limit generalizability. CHIS (and, therefore, the Cal-QOL) excluded adults unreachable by telephone (e.g., homeless, incarcerated, or institutionalized) at the time of the CHIS

survey. Consequently, our estimates do not generalize to these marginalized populations who are known to experience discrimination in healthcare services.⁴⁹ Further, individuals who changed phone numbers in the intervening time between CHIS and Cal-QOL interview were generally lost to follow-up. While we used a weighting approach to adjust for this occurrence, it may not have fully remedied loss to follow-up effects. Second, despite the population base of our sample, we had relatively small sample sizes for all of the racial/ethnic minority groups who reported use of mental health or substance abuse services in the past year. While the small sample sizes are not surprising given the well-known racial/ethnic disparities in access to and use of mental health services,²⁰ this may have limited our ability to observe effects that would have been more apparent with a larger sample size. Moreover, the relatively small number of minorities reduces our ability to make statements specific to some racial/ethnic subgroups. For example, we were unable to examine separately the service experiences of Asians, American Indians, and Latino sub-populations in California. Third, some experiences of discrimination were not measured in the Cal-QOL that would inform our understanding of the mental health and substance abuse services of racial/ethnic minorities. Thus additional research is needed that focuses on both the experiences and consequences of discrimination associated with criminal justice involvement, homelessness, or serious mental illness.⁵⁰ And finally, due to the cross-sectional nature of the data, we are unable to establish a causal link between discrimination in mental health/substance abuse services and reports of treatment dissatisfaction and stopped treatment. It may be the case that clients who are unhappy with their care are more likely to perceive that the provider is discriminating against them and to stop treatment early.

Despite these limitations this study is among the few that provides empirically-based insights into how perceptions of discrimination may be related to mental health/substance abuse treatment outcomes in a population-based sample. Our findings represent an important consideration for the implementation of ACA with its requirements to monitor patient satisfaction and well-being and to employ comparative effectiveness research to aid in eliminating health disparities in those who use mental health or substance abuse services.

Acknowledgments

This research was supported by funding from the National Institute of Drug Abuse (DA 15539; DA 20826), and National Institute of Minority Health Disparities (MD 00508, MD006923)

References

1. Substance Abuse and Mental Health Services Administration. Racial/Ethnic Differences in Mental Health Service Use Among Adults. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2015.
2. Barnes DM, Keyes KM, Bates LM. Racial differences in depression in the United States: how do subgroup analyses inform a paradox? *Soc Psychiatry Psychiatr Epidemiol*. 2013; 48(12):1941–1949. [PubMed: 23732705]
3. Cook BL, Zuvekas SH, Carson N, et al. Assessing racial/ethnic disparities in treatment across episodes of mental health care. *Health Serv Res*. 2013; 49(1):206–229. [PubMed: 23855750]
4. Wang PS, Berglund P, Olfson M, et al. Failure and delay in initial treatment contact after first onset of mental disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005; 62(6):603–613. [PubMed: 15939838]

5. Nowotny KM. Race/ethnic disparities in the utilization of treatment for drug dependent inmates in U.S. state correctional facilities. *Addict Behav.* 2015; 40:148–153. [PubMed: 25270722]
6. Strakowski SM, Keck PE Jr, Arnold LM, et al. Ethnicity and diagnosis in patients with affective disorders. *J Clin Psychiatry.* 2003; 64(7):747–754. [PubMed: 12934973]
7. Minsky S, Vega W, Miskimen T, et al. Diagnostic patterns in Latino, African American, and European American psychiatric patients. *Arch Gen Psychiatry.* 2003; 60(6):637–644. [PubMed: 12796227]
8. Saloner B, Le Cook B. Blacks and Hispanics are less likely than whites to complete addiction treatment, largely due to socioeconomic factors. *Health Aff.* 2013; 32(1):135–145.
9. Wells K, Klap R, Koike A, et al. Ethnic disparities in unmet need for alcoholism, drug abuse, and mental health care. *Am J Psychiatry.* 2001; 158(12):2027–2032. [PubMed: 11729020]
10. Ault-Brutus AA. Changes in racial-ethnic disparities in use and adequacy of mental health care in the United States, 1990–2003. *Psychiatr Serv.* 2012; 63(6):531–540. [PubMed: 22422014]
11. Levinson D, Lakoma MD, Petukhova M, et al. Associations of serious mental illness with earnings: results from the WHO World Mental Health surveys. *Br J Psychiatry.* 2010; 197(2):114–121. [PubMed: 20679263]
12. Fletcher JM. Adolescent depression and educational attainment: results using sibling fixed effects. *Health Econ.* 2010; 19(7):855–871. [PubMed: 19582699]
13. Chatterji P, Alegria M, Takeuchi D. Racial/ethnic differences in the effects of psychiatric disorders on employment. *Atl Econ J.* 2009; 37(3):243–257. [PubMed: 19898677]
14. Zivin K, Ilgen MA, Pfeiffer PN, et al. Early mortality and years of potential life lost among veterans affairs patients with depression. *Psychiatr Serv.* 2012; 63(8):823–826. [PubMed: 22854727]
15. Druss BG, Zhao L, Von Esenwein S, et al. Understanding excess mortality in persons with mental illness: 17-year follow up of a nationally representative US survey. *Med Care.* 2011; 49(6):599–604. [PubMed: 21577183]
16. Krieger N, Kosheleva A, Waterman PD, et al. Racial discrimination, psychological distress, and self-rated health among US-born and foreign-born Black Americans. *American journal of public health.* 2011; 101(9):1704–1713. [PubMed: 21778504]
17. Breslau J, Aguilar-Gaxiola S, Kendler KS, et al. Specifying race-ethnic differences in risk for psychiatric disorder in a USA national sample. *Psychological medicine.* 2006; 36(1):57–68. [PubMed: 16202191]
18. Snowden LR, Catalano R, Shumway M. Disproportionate use of psychiatric emergency services by african americans. *Psychiatric services (Washington, D.C.)* 2009; 60(12):1664–1671.
19. McGuire TG, Alegria M, Cook BL, et al. Implementing the Institute of Medicine definition of disparities: an application to mental health care. *Health Serv Res.* 2006; 41(5):1979–2005. [PubMed: 16987312]
20. Alegria M, Lin J, Chen CN, et al. The impact of insurance coverage in diminishing racial and ethnic disparities in behavioral health services. *Health Serv Res.* 2012; 47(3 Pt 2):1322–1344. [PubMed: 22568675]
21. Bauer AM, Chen CN, Alegria M. English language proficiency and mental health service use among Latino and Asian Americans with mental disorders. *Med Care.* 2010; 48(12):1097–1104. [PubMed: 21063226]
22. Jimenez DE, Bartels SJ, Cardenas V, et al. Stigmatizing attitudes toward mental illness among racial/ethnic older adults in primary care. *Int J Geriatr Psychiatry.* 2013
23. Nadeem E, Lange JM, Edge D, et al. Does stigma keep poor young immigrant and U.S.-born Black and Latina women from seeking mental health care? *Psychiatr Serv.* 2007; 58(12):1547–1554. [PubMed: 18048555]
24. Suite DH, La Bril R, Primm A, et al. Beyond misdiagnosis, misunderstanding and mistrust: relevance of the historical perspective in the medical and mental health treatment of people of color. *J Natl Med Assoc.* 2007; 99(8):879–885. [PubMed: 17722664]
25. Lee C, Ayers SL, Kronenfeld JJ. The association between perceived provider discrimination, healthcare utilization and health status in racial and ethnic minorities. *Ethn Dis.* 2009; 19(3):330–337. [PubMed: 19769017]

26. Hausmann LR, Hannon MJ, Kresevic DM, et al. Impact of perceived discrimination in healthcare on patient-provider communication. *Med Care*. 2011; 49(7):626–633. [PubMed: 21478769]
27. Sorkin DH, Ngo-Metzger Q, De Alba I. Racial/ethnic discrimination in health care: impact on perceived quality of care. *J Gen Intern Med*. 2010; 25(5):390–396. [PubMed: 20146022]
28. Trivedi AN, Ayanian JZ. Perceived discrimination and use of preventive health services. *J Gen Intern Med*. 2006; 21(6):553–558. [PubMed: 16808735]
29. Bird ST, Bogart LM, Delahanty DL. Health-related correlates of perceived discrimination in HIV care. *AIDS Patient Care STDS*. 2004; 18(1):19–26. [PubMed: 15006191]
30. Perez D, Sribney WM, Rodriguez MA. Perceived discrimination and self-reported quality of care among Latinos in the United States. *J Gen Intern Med*. 2009; 24(Suppl 3):548–554. [PubMed: 19842005]
31. Greer TM. Perceived racial discrimination in clinical encounters among African American hypertensive patients. *J Health Care Poor Underserved*. 2010; 21(1):251–263. [PubMed: 20173267]
32. Peek ME, Wagner J, Tang H, et al. Self-reported racial discrimination in health care and diabetes outcomes. *Med Care*. 2011; 49(7):618–625. [PubMed: 21478770]
33. Clement S, Williams P, Farrelly S, et al. Mental health-related discrimination as a predictor of low engagement with mental health services. *Psychiatr Serv*. 2015; 66(2):171–176. [PubMed: 25642612]
34. Gabbidon J, Farrelly S, Hatch SL, et al. Discrimination attributed to mental illness or race-ethnicity by users of community psychiatric services. *Psychiatr Serv*. 2014; 65(11):1360–1366. [PubMed: 25081996]
35. Cochran SD, Mays VM. Physical health complaints among lesbians, gay men, and bisexual and homosexually experienced heterosexual individuals: results from the California Quality of Life Survey. *Am J Public Health*. 2007; 97(11):2048–2055. [PubMed: 17463371]
36. California Health Interview Survey. CHIS 2007 Methodology Report Series: Report 1 - Sample Design. Los Angeles, CA: UCLA Center for Health Policy Research; 2009.
37. Sturm R, Gresenz C, Sherbourne C, et al. The design of Healthcare for Communities: a study of health care delivery for alcohol, drug abuse, and mental health conditions. *Inquiry*. 1999; 36(2): 221–233. [PubMed: 10459376]
38. Grella CE, Greenwell L, Mays VM, et al. Influence of gender, sexual orientation, and need on treatment utilization for substance use and mental disorders: findings from the California Quality of Life Survey. *BMC Psychiatry*. 2009; 9:52. [PubMed: 19682355]
39. White IR, Royston P, Wood AM. Multiple imputation using chained equations: Issues and guidance for practice. *Stat Med*. 2011; 30(4):377–399. [PubMed: 21225900]
40. McMorrow S, Long SK, Kenney GM, et al. Uninsurance Disparities Have Narrowed For Black And Hispanic Adults Under The Affordable Care Act. *Health Aff*. 2015; 34(10):1774–1778.
41. Katon WJ, Lin EH, Von Korff M, et al. Collaborative care for patients with depression and chronic illnesses. *N Engl J Med*. 2010; 363(27):2611–2620. [PubMed: 21190455]
42. Deen TL, Fortney JC, Pyne JM. Relationship between satisfaction, patient-centered care, adherence and outcomes among patients in a collaborative care trial for depression. *Adm Policy Ment Health*. 2011; 38(5):345–355. [PubMed: 20978932]
43. Anhang Price R, Elliott MN, Zaslavsky AM, et al. Examining the role of patient experience surveys in measuring health care quality. *Med Care Res Rev*. 2014; 71(5):522–554. [PubMed: 25027409]
44. Fenton JJ, Jerant AF, Bertakis KD, et al. The cost of satisfaction: a national study of patient satisfaction, health care utilization, expenditures, and mortality. *Arch Intern Med*. 2012; 172(5): 405–411. [PubMed: 22331982]
45. van Schaik DJ, Klijn AF, van Hout HP, et al. Patients' preferences in the treatment of depressive disorder in primary care. *Gen Hosp Psychiatry*. 2004; 26(3):184–189. [PubMed: 15121346]
46. Weech-Maldonado R, Elliott M, Pradhan R, et al. Can hospital cultural competency reduce disparities in patient experiences with care? *Med Care*. 2012; 50(Suppl):S48–S55. [PubMed: 23064277]

47. Hall WJ, Chapman MV, Lee KM, et al. Implicit Racial/Ethnic Bias Among Health Care Professionals and Its Influence on Health Care Outcomes: A Systematic Review. *Am J Public Health*. 2015; 105(12):e60–e76.
48. Cabral RR, Smith TB. Racial/ethnic matching of clients and therapists in mental health services: a meta-analytic review of preferences, perceptions, and outcomes. *J Couns Psychol*. 2011; 58(4): 537–554. [PubMed: 21875181]
49. Zerger S, Bacon S, Corneau S, et al. Differential experiences of discrimination among ethnoracially diverse persons experiencing mental illness and homelessness. *BMC Psychiatry*. 2014; 14:353. [PubMed: 25496296]
50. Skosireva A, O'Campo P, Zerger S, et al. Different faces of discrimination: perceived discrimination among homeless adults with mental illness in healthcare settings. *BMC Health Serv Res*. 2014; 14:376. [PubMed: 25196184]

Figure 1.
Discrimination in California Quality of Life Survey Waves I–III Healthcare and Mental Health/Substance Abuse Service Visits Study Eligibility Criteria

Table 1

Participant Characteristics and Prevalence Of Perceived Discrimination In Healthcare and Mental Health/Substance Abuse Service Visits In The California Quality Of Life Survey, I – III (n=1,099).

	Wave I–III N=1,099	Wave II–III N=745	Percent Who Reported Any Healthcare Discrimination, Waves I–III		Percent Who Reported Discrimination In MH/SA Visit, Waves I–III	
	Weighted N (%)		Weighted % (SE)	F	Weighted % (SE)	F
Race / Ethnicity						
Latino	225 (20.5%)	138 (18.5%)	24.1 (5.4)		7.9 (3.0)	
Non-Latino White	723 (65.8%)	487 (65.4%)	12.0 (1.6)		3.0 (0.8)	
Non-Latino Black	52 (4.8%)	48 (6.4%)	24.6 (6.9)		8.8 (4.3)	
Other	99 (9%)	72 (9.7%)	10.7 (5.1)	3.33*	0.1 (0.1)	8.72***
Age						
18–29	211 (19.2%)	157 (21%)	13.5 (3.8)		5.7 (2.4)	
30–44	363 (33%)	246 (33%)	16.1 (3.0)		3.7 (1.5)	
45–59	399 (36.4%)	259 (34.7%)	16.2 (2.8)		4.2 (1.4)	
60–72	126 (11.4%)	84 (11.2%)	10.2 (3.8)	0.54	1.4 (1.2)	0.78
Gender						
Male	414 (37.6%)	185 (38.2%)	13.4 (2.8)		3.0 (1.1)	
Female	685 (62.4%)	460 (61.8%)	15.6 (2.1)	0.26	4.7 (1.2)	0.99
Sexual Orientation						
Heterosexual	981 (89.2%)	665 (89.2%)	14.4 (1.8)		4.1 (1.0)	
Sexual Orientation Minority	118 (10.8%)	80 (10.8%)	19.5 (3.6)	1.76	3.7 (0.9)	0.06
Education						
Less Than High School	70 (9.2)	38 (6.8%)	23.8 (7.2)		7.7 (3.9)	
High School	166 (19.2%)	119 (19.5%)	14.4 (3.8)		1.6 (1.4)	
Some College	303 (31.2%)	202 (30.8%)	21.3 (3.4)		8.0 (2.2)	
Four Or More Years College	560 (40.4%)	386 (42.9%)	8.2 (1.8)	4.85***	1.3 (0.6)	4.93***
Household Income						
Above 200% FPL ²	779 (70.9%)	507 (68.1%)	13.0 (1.8)		3.7 (1.0)	
Below 200% FPL	320 (29.1%)	238 (31.9%)	19.6 (3.6)	3.10	4.7 (1.8)	0.31
Insurance Status						

	Wave I-III N=1,099		Wave II-III N=745		Percent Who Reported Any Healthcare Discrimination, Waves I-III		Percent Who Reported Discrimination In MH/SA Visit, Waves I-III	
	Weighted N (%)	Weighted % (SE)	F	Weighted % (SE)	F	Weighted % (SE)	F	
Uninsured	128 (11.7%)	78 (10.5%)		27.4 (6.4)		15.5 (5.2)		
Private Insurance Only	648 (58.9%)	428 (57.4%)		10.5 (1.7)		1.8 (0.7)		
Any Public Insurance	323 (29.4%)	239 (32.1%)		19.0 (3.5)	6.04**	3.9 (1.4)	8.77***	
Screening For Psychiatric Disorder								
Screened Negative For All Disorders	397 (36.1%)	276 (37%)		7.1 (1.9)		1.2 (0.6)		
Screened Positive For Any Disorder	702 (63.9%)	469 (63%)		19.4 (2.3)	13.05***	5.6 (1.3)	7.00**	
Psychiatric Disorder ³								
Major Depressive Disorder	436 (39.6%)	286 (38.4%)		21.4 (3.0)	9.59**	6.0 (1.7)	3.52	
Generalized Anxiety Disorder	302 (27.5%)	191 (25.7%)		26.9 (3.9)	18.92***	7.7 (2.3)	6.25*	
Panic Attack	209 (19%)	136 (18.2%)		23.8 (4.7)	6.38*	6.3 (2.4)	1.55	
Post-Traumatic Stress Disorder	189 (17.2%)	123 (16.5%)		24.2 (4.8)	6.44*	9.4 (3.2)	7.07**	
Alcohol or Drug Use Disorder	190 (17.3%)	124 (16.6%)		16.4 (4.0)	0.17	5.4 (2.2)	0.59	

¹ MH/SA = Mental Health or Substance Abuse;

² FPL = Federal Poverty Level;

³ Some participants met criteria for more than one psychiatric disorder. Participants who met criteria for each disorder are included and therefore do not add to 100 percent.

Table 2

Predictors Of Perceived Healthcare Discrimination Reported By Users Of Any HealthCare and Mental Health/ Substance Abuse Services In The Past Year, California Quality of Life Survey Waves I – III (n=1,099).

	Any Past Year Healthcare Discrimination		Discrimination Occurring During A Mental Health or Substance Abuse Visit	
	AOR	(95% CI)	AOR	(95% CI)
Race / Ethnicity				
Non-Latino White (ref)	1.0		1.0	
Latino	2.08 *	(1.02, 4.25)	2.38	(0.77, 7.34)
Non-Latino Black	2.02	(0.80, 5.10)	2.97	(0.86, 10.26)
Other	0.81	(0.26, 2.53)	0.01 ***	(0.00, 0.12)
Age				
18–29 (ref)	1.0		1.0	
30–44	1.75	(0.80, 3.80)	1.05	(0.31, 3.64)
45–59	1.72	(0.79, 3.76)	1.34	(0.48, 3.75)
60–72	1.09	(0.35, 3.38)	0.58	(0.08, 3.94)
Gender				
Male	1.0		1.0	
Female	1.02	(0.5, 1.91)	1.31	(0.51, 3.34)
Sexual Orientation				
Heterosexual	1.0		1.0	
Non-Heterosexual	1.44	(0.78, 2.66)	0.81	(0.33, 2.01)
Education				
High School or Less (ref)	1.0		1.0	
Some College	1.47	(0.75, 2.90)	2.52	(0.67, 9.49)
Four Year College Degree	0.65	(0.30, 1.44)	0.67	(0.15, 2.94)
Household Income				
Above 200% FPL	1.0		1.0	
Below 200% FPL	0.93	(0.52, 1.68)	0.80	(0.31, 2.08)
Insurance Status				
Private Insurance (ref)	1.0		1.0	
Public Insurance	1.45	(0.73, 2.90)	1.74	(0.52, 5.79)
Uninsured	2.51 *	(1.19, 5.29)	7.27 **	(2.27, 23.30)
Past Year Mental Health Disorder				
No	1.0		1.0	
Yes	2.77 **	(1.50, 5.11)	3.37	(0.98, 11.55)
Past Year Substance Use Disorder				
No	1.0		1.0	
Yes	1.07	(0.51, 2.22)	1.37	(0.46, 4.03)

AOR = Adjusted Odds Ratio; CI = Confidence Interval; FPL= Federal Poverty Level

* p < .05,

**
p<.01,

p < .001

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 3

Reasons for Perceived Healthcare Discrimination Reported By Black, Latino and White Users of Mental Health/Substance Abuse Services In The Past Year, California Quality of Life Surveys Waves I – III (n=193)

	Non-Latino Black	Latino	Non-Latino White
Reason For Discrimination	Weighted % (SE)		
Race or Ethnic Origin	52.2 (16.7)	30.5 (10.9)	5.7 (3.1)
Insurance Type	30.2 (16.0)	15.3 (10.1)	40.0 (7.2)
Income Level	29.1 (16.6)	22.4 (11.5)	8.0 (3.6)
Health Or Disability	0.2 (0.2)	0.3 (0.3)	6.6 (3.1)
Age	19.0 (16.0)	0.2 (0.2)	11.3 (5.0)
Gender	13.8 (11.5)	1.9 (2.2)	3.1 (1.9)
Language / Accent	0	1.6 (4.2)	3.0 (2.8)
Religion	0	6.7 (7.1)	2.0 (1.8)
Sexual Orientation	0	2.1 (1.4)	2.3 (0.8)
Body Weight	0.6 (0.6)	1.3 (1.0)	9.5 (4.5)
Physical Appearance	0	0.3 (0.3)	5.1 (4.7)
Other	11.3 (10.7)	11.1 (7.6)	22.3 (6.1)

¹ Some participants reported more than one reason for discrimination. All participant responses are included and therefore do not add to 100 percent.

Associations of Perceived Discrimination in Past Year Mental Health/Substance Abuse Service Visits with Treatment Helpfulness and Stopped Treatment, California Quality Of Life Survey II – III (n=745).

Table 4

	Ratings Of Treatment Helpfulness ¹					
	Non-Latino Black			Latino		
	AOR	(95% CI)	AOR	(95% CI)	AOR	(95% CI)
Discrimination In MH/SA Visit	0.19	(0.03, 1.48)	0.09 [*]	(0.01, 0.84)	0.25 ^{**}	(0.09, 0.67)
Age						
18–44 (ref)	1.0		1.0		1.0	
45–72	1.30	(0.39, 4.36)	4.31	(1.00, 18.60)	0.63	(0.38, 1.03)
Female Gender	1.12	(0.22, 5.62)	2.88	(0.81, 10.33)	1.75 [*]	(1.05, 2.92)
High School or Less Education	1.63	(0.42, 6.34)	0.78	(0.15, 4.01)	0.61	(0.32, 1.15)
Private Insurance	2.18	(0.53, 8.96)	0.42	(0.08, 2.09)	1.43	(0.85, 2.39)
Past Year Psychiatric Disorder	0.95	(0.24, 3.81)	1.83	(0.46, 7.25)	0.57 [*]	(0.32, 0.99)
Stopped Treatment Early ²						
	Non-Latino Black			Latino		
	AOR	(95% CI)	AOR	(95% CI)	AOR	(95% CI)
Discrimination In MH/SA Visit	13.38 [*]	(1.36, >100)	2.06	(0.08, 51.97)	0.66	(0.09, 5.09)
Age						
18–44 (ref)	1.0		1.0		1.0	
45–72	0.75	(0.20, 2.84)	1.18	(0.23, 6.11)	0.58	(0.28, 1.20)
Female Gender	2.58	(0.67, 9.99)	45.47	(0.20, >100)	1.25	(0.60, 2.63)
High School or Less Education	0.46	(0.07, 2.37)	0.93	(0.15, 5.82)	0.64	(0.24, 1.73)
Private Insurance	5.46 [*]	(1.34, 22.19)	1.74	(0.26, 11.79)	0.59	(0.28, 1.25)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Past Year Psychiatric Disorder	2.96	(0.67, 13.03)	3.22	(0.49, 21.06)	1.86	(0.85, 4.08)
Treatment Rated Non-Helpful	2.80	(0.68, 11.42)	8.83	(0.14, > 100)	3.14 *	(1.02, 9.62)

AOR = Adjusted Odds Ratio; SE = Standard Error; MH/SA = Mental Health or Substance Abuse

1 Ordered logistic regression of treatment helpfulness on discrimination during a mental health or substance abuse visit. A coefficient greater than 1.0 indicates increased odds of rating treatment as more helpful

2 Logistic regression of stopped treatment on discrimination during a mental health or substance abuse visit. A coefficient greater less than 1.0 indicates increased odds of stopping treatment early.

* p < .05,
** p<.01