Published in final edited form as:

Genes Brain Behav. 2017 January ; 16(1): 101–117. doi:10.1111/gbb.12324.

Structural and functional plasticity of dendritic spines – root or result of behavior?

Cassandra D. Gipson and M. Foster Olive

Department of Psychology, Arizona State University. PO Box 871104, Tempe AZ 85287

Abstract

Dendritic spines are multifunctional integrative units of the nervous system and are highly diverse and dynamic in nature. Both internal and external stimuli influence dendritic spine density and morphology on the order of minutes. It is clear that the structural plasticity of dendritic spines is related to changes in synaptic efficacy, learning and memory, and other cognitive processes. However, it is currently unclear whether structural changes in dendritic spines are primary instigators of changes in specific behaviors, a consequence of behavioral changes, or both. In this review, we first review the basic structure and function of dendritic spines in the brain, as well as laboratory methods to characterize and quantify morphological changes in dendritic spines. We then discuss the existing literature on the temporal and functional relationship between changes in dendritic spines in specific brain regions and changes in specific behaviors mediated by those regions. Although technological advancements have allowed us to better understand the functional relevance of structural changes in dendritic spines that are influenced by environmental stimuli, the role of spine dynamics as an underlying driver or consequence of behavior still remains elusive. We conclude that while it is likely that structural changes in dendritic spines are both instigators and results of behavioral changes, improved research tools and methods are needed to experimentally and directly manipulate spine dynamics in order to more empirically delineate the relationship between spine structure and behavior.

Keywords

Dendritic spine; plasticity; morphology; glutamate; actin; cytoskeleton; behavior; Golgi; histology; dynamics

Introduction

Dendritic spines are key specialized structures of neuronal connectivity and signaling in the nervous system. Dendritic spines are also highly dynamic in nature, having the capacity to change their morphology, number, density, and motility within relatively short timeframes. For the purposes of this review, we utilize the term dendritic spine “dynamics” to refer to changes in one or more of the aforementioned structural properties, as opposed to

Author contact information: Cassandra D. Gipson, Ph.D., Phone: 480-727-5052, cgipsonr@asu.edu, M. Foster Olive, Ph.D., Phone: 480-727-9557, foster.olive@asu.edu.

The authors have no conflicts of interest to declare.

subcellular phenomena such as biochemical signaling or dendritic mRNA translation. The dynamic nature of dendritic spines has recently become of great interest in the study of the neurobiological underpinnings of behavior. For example, synaptic plasticity, which is thought to underlie learning processes and the formation and consolidation of memory (Durand *et al.*, 1996; Fortin *et al.*, 2012; Oe *et al.*, 2013), involves de novo dendritic spine formation (spinogenesis) and remodeling of existing spines to sustain neuronal connections that result from input from the environment or other brain regions. Synaptic strengthening and long-term potentiation (LTP) are associated with increases in spine density and/or spine head diameter, while synaptic weakening induced by long-term depression (LTD) causes spine retraction (Alvarez & Sabatini, 2007; De Roo *et al.*, 2008a; Sala & Segal, 2014; Segal, 2005; van der Zee, 2015). However, it should be noted that synaptic overstimulation can also lead to spine shrinkage to prevent damage from excess calcium influx (Paulin *et al.*, 2016).

The involvement of spine dynamics in behavior and the impact of environmental stimuli on spines have become topics of interest across various fields of research. However, it remains unclear whether spine changes *precede* environmental input and/or behavioral output, or if environmental input and/or behavioral output in and of themselves *lead to* spine changes. In this review, we will examine the function of dendritic spines as well as technological advancements in the neuroanatomical methods used to analyze them. In doing so, we will discuss whether changes in spine morphology are an accurate correlate for changes in synaptic strength. We will then discuss structural changes in dendritic spines in relation to different types of environmental input or behavioral output including learning and memory processes, addiction-related behaviors, stress, depression, and aging. Finally, we will attempt to address the question: do changes in dendritic spine dynamics lead to behavioral changes, specifically when behavior becomes aberrant, or is behavior itself the root cause of changes in dendritic spines? In other words, do spine dynamics underlie pathological behaviors and therefore constitute a neurobiological substrate of certain brain disorders?

Structure and function of dendritic spines

Structure

Dendritic spines were discovered more than century ago by renowned Spanish anatomist and Nobel laureate Santiago Ramón y Cajal. In his investigations of the microarchitecture of the avian cerebellar cortex, Cajal noted that the “surface of the Purkinje cells' dendrites appear ruffled with thorns or short spines, which on the terminal dendrites look like light protrusions” (Ramón y Cajal, 1888). Cajal subsequently revealed dendritic spines to be present in numerous neuronal subtypes and brain regions, and also relatively consistent across various animal species (Ramón y Cajal, 1899). Incorporating these observations into his neuron doctrine, Cajal hypothesized that dendritic spines served to increase the surface area of dendrites to accommodate the vast complexity and number of neural connections in the brain (Ramón y Cajal, 1899; Yuste, 2015).

The subsequent development of high resolution microscopy techniques such as confocal microscopy, two-dimensional transmission electron microscopy, and three-dimensional reconstruction of serial confocal and electron microscopy images, has revealed substantial heterogeneity in dendritic spine morphology and associated metrics. In general, dendritic

spines are 1–3 μm in length, 0.01–0.8 μm^3 in volume, and typically consist of a rounded “head” (which receives primarily excitatory synaptic input) atop a thinner “neck” apparatus emanating from the dendritic shaft (Harris & Spacek, 2016; Yuste, 2010) (Fig. 1A). However, neurons exhibit many variations of dendritic spine subtypes based on their morphological and physiological properties, each with unique putative functions. Spine-containing (“spiny”) neurons are found in numerous brain regions, the most widely studied being pyramidal cells of the cerebral cortex, medium spiny neurons of the dorsal and ventral striatum, and Purkinje cells of the cerebellum. Recently, advancements in cellular imaging techniques have allowed three-dimensional analyses of dendritic spines across different cell types, brain regions, and time. Importantly, it should be noted that although these advancements have yielded a wealth of information regarding the dynamic structure of dendritic spines, they have not yet fully defined the exact functions of changes in dendritic spine morphology, particularly with respect to behavior.

More than 100,000 dendritic spines can populate the dendritic arbor of a single neuron (Yuste, 2010). During the development of the nervous system, dendritic spines tend to have a long ($>2 \mu\text{m}$), “filopodia-like” morphology containing relatively few organelles and no discernible head apparatus. This type of dendritic spine is highly dynamic in its ability to retract or extend within minutes of chemical stimulation (Fischer *et al.*, 1998; Harris, 1999). However, its functionality in synaptic transmission, particularly in adulthood, appears limited, since it is largely devoid of synaptic inputs and thus is often considered an “immature” or “transient” phenotype. Yet following increased synaptic input, immature dendritic spines can transform into more “mature” and “stable” phenotypes (Bourne & Harris, 2007).

Dendritic spines in adult neurons can have simple or branched morphologies (Fig. 1B), and over the years, investigators have developed various criteria for classifying spines into specific categories based on their overall morphology. However, as we caution below, such categorical approaches to examining spine dynamics may be inherently self-limiting and potentially flawed. Some spine types are sessile in nature, being devoid of any discernible neck apparatus and relatively short in length ($<0.5 \mu\text{m}$), which are often referred to as “stubby” spines (Harris & Spacek, 2016). Other mature types of spines are pedunculated, containing a head apparatus that is wider in diameter than that of the neck, often referred to as “mushroom” spines. Spines that retain this distinct head and neck apparatus but with much narrower diameters are often referred to as “thin” spines. Less frequently, mature spines can have two (bifurcated) or more (multi-branched) processes with fully functional heads, thus allowing multiple excitatory synaptic inputs onto a single spine. It should be noted that aside from these classical dendritic spine morphologies, other dendritic specializations for receiving inputs include more complicated architectures such as synaptic crests, claw-like glomerular endings, brush endings, and thorny and coralline excrescences. Yet the occurrence of these structures tends to be far less frequent than that of simple dendritic spines (Harris & Spacek, 2016).

As alluded to above, many studies investigating the structural plasticity of dendritic spines in normal and diseased states have utilized categorical approaches to quantifying spine plasticity. For example, observed increases in the numbers of “mushroom-shaped” spines following a specific experimental manipulation, concomitant with decreases in the numbers

of “filopodia-like” spines, are often interpreted as being suggestive of increased spine maturation. We argue, however, that many metrics of spine morphology, including head and neck diameter and volume, neck length, ratio of head to neck diameter, etc., occur along various continuums. As a result, categorical approaches to assessing structural plasticity of dendritic spines may overlook subtle differences in spine morphology that do not meet pre-defined classifications (Arellano *et al.*, 2007). For example, putative thin and filopodia-like spines often have similar diameters, and the length of the spine that differentiates between these two subtypes is not universally agreed upon. Similarly, spines that have discernible yet relatively small head apparatuses located on spine necks with large diameters may be mistakenly classified as thin spines. The conundrum of discrete morphological spine classification is further exemplified by recent findings that indicate spine neck width and length are more indicative of synaptic efficacy and compartmentalization than spine head measurements (Tonnesen *et al.*, 2014). As suggested by others (Arellano *et al.*, 2007; Yuste & Majewska, 2001), we assert that dendritic spine dynamics are more accurately assessed by continuous numerical measures, such as spine head diameter and volume, neck width and length, and so forth, rather than discrete morphology-based classifications.

Hypothesized functions of dendritic spines

As noted earlier, Cajal hypothesized that dendritic spines served to increase the surface area of dendrites, thereby increasing their capacity to receive synaptic inputs (Ramón y Cajal, 1899). While this hypothesis continues to be supported today, recent advances in electrophysiology, microscopy, and biochemical techniques have allowed additional hypotheses to be formed regarding the function of dendritic spines. These functions include, but are not limited to, (1) a means for regulating the efficacy of single synapses at the pre- and post-synaptic levels; (2) expanding the computational capacity of neurons; and (3) compartmentalization of postsynaptic biochemical signaling and gene expression (Crick, 1982; Koch & Zador, 1993; Lee *et al.*, 2012; Rall, 1978; Shepherd, 1996; Yuste & Majewska, 2001). Indeed, all of these hypothesized functions are substantiated by a considerable body of empirical studies.

One of the most widely studied aspects of structural plasticity of dendritic spines is the influence of learning and memory related processes. Early studies showed that tetanic stimulation of hippocampal afferents, with parameters similar to those used for inducing long-term potentiation (LTP), resulted in increased spine density and enlarged spine heads and necks in granule cells of the dentate gyrus (Chang & Greenough, 1984; Desmond & Levy, 1986; Fifkova & Anderson, 1981; Fifkova & Van Harreveld, 1977; Van Harreveld & Fifkova, 1975). Since dendritic spine head diameter and volume have been positively correlated with synaptic strength (De Roo *et al.*, 2008a; Sala & Segal, 2014), it would seem reasonable to conclude that these morphological changes in dendritic spines are anatomical substrates of increased synaptic efficacy. Similarly, increases in dendritic spine density in the dentate gyrus have been observed following optogenetic activation of entorhinal inputs to these cells, paralleled by increases in membrane capacitance (Ryan *et al.*, 2015). On the other hand, chemically induced synaptic de-potentialization (e.g., LTD) can cause spine retraction and shrinkage (Nagerl *et al.*, 2004; Oh *et al.*, 2013; Wang *et al.*, 2007; Zhou *et al.*, 2004). It should be noted, however, that LTP- or LTD-induced changes in spine dynamics

are not necessarily uniform within a specific brain region, population of neurons, or even along the length of a single dendritic shaft, but rather appeared clustered to specific dendritic segments (Bosch & Hayashi, 2012; De Roo *et al.*, 2008a; Fortin *et al.*, 2012; Yuste & Bonhoeffer, 2001).

The ability of LTP to induce increases in spine density raises some interesting questions. For example, does LTP actually induce the de novo growth of dendritic spines (spinogenesis), or merely enlarge existing spines? Are these changes merely transient, or do they become stabilized and mature to establish new synaptic connections? The answers to these questions appear to be in the affirmative. It is now clear that de novo spinogenesis occurs under conditions of synaptic potentiation both in vitro (Engert & Bonhoeffer, 1999; Maletic-Savatic *et al.*, 1999; Toni *et al.*, 1999) and in vivo (Hayashi-Takagi *et al.*, 2015; Holtmaat *et al.*, 2005; Kopec *et al.*, 2006; Matsuzaki *et al.*, 2004; Ryan *et al.*, 2015; Trachtenberg *et al.*, 2002; Yang *et al.*, 2009). Glutamatergic signaling appears to be critical for this phenomenon (Fischer *et al.*, 2000; Kwon & Sabatini, 2011; Mattison *et al.*, 2014; Richards *et al.*, 2005; Ultanir *et al.*, 2007), although other neurotransmitter such as dopamine also likely play a role (Yagishita *et al.*, 2014). Ultrastructural studies have shown that at least some of these newly formed spines are capable of becoming mature and stabilized to form intact synapses, albeit hours to days following de novo spine growth (Knott *et al.*, 2006; Nagerl *et al.*, 2007; Toni *et al.*, 1999).

Another interesting question raised by these observations is whether dendritic spine growth, plasticity, or even elimination occurs on a regular basis in the absence of significant synaptic input. This appears to be the case especially during brain development, where spine turnover (protrusion, maturation, and elimination) occurs in as many as 10–15% of dendritic spines in a 24-hour period, whereas in adulthood this number declines to ~1–2% (Attardo *et al.*, 2015; Grutzendler *et al.*, 2002; Trachtenberg *et al.*, 2002; Zuo *et al.*, 2005). While many “stabilized” spines appear to remain plastic throughout the lifespan, only a small subset of dendritic spines actually undergo complete turnover. Interestingly, under pathological developmental conditions such as in Fragile X syndrome, the abolishment of the fragile X mental retardation protein (FMRP) impacts both the stability of spines (Cruz-Martin *et al.*, 2010; Pan *et al.*, 2010) and proper spine pruning during development, resulting in an excess of small diameter and longer length spines on pyramidal neuron apical dendrites within the cerebral cortex. For a more detailed summary of dendritic spine abnormalities associated with neurodevelopmental or other neuropsychiatric disorders, the reader is directed to several recent reviews (Glausier & Lewis, 2013; He & Portera-Cailliau, 2013; Leuner & Shors, 2013; van Spronsen & Hoogenraad, 2010). Although these observations in Fragile X syndrome highlight the potential importance of proper synaptic pruning and spine stabilization in normal brain development, care should be taken when interpreting dendritic spine function based solely on measurements such as head diameter and neck length, as these may lead to incorrect assumptions regarding spine function. Additional measurements such as receptor insertion, postsynaptic density quantification, or synaptic efficacy should be utilized, when possible, to confirm functionality of changes in spine dynamics. Finally, traditional classification of spines into different categories has recently been suggested to be obsolete, as a time lapse imaging study using high resolution stimulated emission depletion microscopy found that spines previously categorized as “stubby” are actually “mushroom”

shaped (Tonnesen *et al.*, 2014). This discrepancy is attributable to the limits of less evolved imaging technologies that measure spine neck diameter inaccurately. These findings also suggest that the proportion of putative “stubby” spines may be overestimated in earlier studies (Segal, 2016), which leads to a concern regarding the interpretation of spine function based on categorizations across different fields of research.

Molecular mechanisms of dendritic spine dynamics

The dynamic nature of dendritic spines suggests a substantial amount and sophistication of cellular machinery present locally in the dendrite that subserves the demands of synaptic homeostasis. In addition to the presence of polyribosomes for local mRNA translation in dendrites (Martin & Zukin, 2006), hundreds of different molecular entities have been identified as mediators of dendritic spine formation, structure, stability, motility, and elimination. These can be classified into various functional categories including: actin binding and cytoskeletal proteins (e.g., actin, myosin, profilin, cofilin, etc), small guanosine exchange factors and triphosphatases (GEFs and GTPases, respectively) and associated proteins (including RhoA, Kalrin-7, and Cdc42), cell surface receptors and adhesion molecules (e.g., glutamate receptors, integrins and cadherins), receptor tyrosine and other kinases (e.g., TrkB receptors, calcium/calmodulin-dependent protein kinase II, etc.), postsynaptic scaffold and adaptor proteins (e.g., Homer, Shank, PSD, etc.), microRNAs, microRNA binding proteins and transcription factors, and steroid hormones (e.g., estrogen and glucocorticoids). For an in depth discussion of the function of all of these classes of spine-associated proteins, the reader is directed to several other excellent recent reviews (Bellot *et al.*, 2014; Frankfurt & Luine, 2015; Murakoshi & Yasuda, 2012; Sala & Segal, 2014; Tada & Sheng, 2006; Uchoa *et al.*, 2014; Wang & Zhou, 2010).

The most rapid events in initiating structural plasticity in dendritic spines are changes in intracellular calcium levels (which occurs on a millisecond time scale), followed by activation of various kinases and GTPases (on the order of seconds), activation of GEFs, actin and other cytoskeletal reorganization, and insertion of various neurotransmitter receptors (on the order of minutes). Lastly, phosphorylation and nuclear translocation of various transcription factors to initiate gene expression (on the order of minutes to hours), culminates in spinogenesis or structural changes and accompanying synaptic potentiation or de-potentiation (Yasuda, 2016).

Since this review focuses on the structural alterations of dendritic spines, it is worth briefly mentioning the basic aspects of cytoskeletal dynamics. The cytoskeleton of dendritic spines is composed primarily of smaller globular (G) actin building blocks that assemble into paired and twisted filaments that constitute filamentous (F) actin. F-actin is the primary cytoskeletal component of the spine neck and latticework of the spine head (see Fig. 1A). Changes in spine morphology during cytoskeletal restructuring are thought to be accompanied by changes in AMPA receptor expression (Cingolani & Goda, 2008). During dendritic spine restructuring, F-actin can become depolymerized into G-actin by direct actin-binding molecules such as cofilin-1 and destrin. Alternatively, F-actin can be bound and either contracted or stabilized by members of the myosin family of molecular motors, such as non-muscle myosin IIB. Other actin binding protein that promote stabilization are drebrin

A, Ras, profilin, and gelsolin. Actin polymerization and de-polymerization are also influenced by proteins such as Shank, PSD, cortactin, Rac1, and various capping proteins. As a side note and as discussed below, there are recent reports of the development of a photoactivatable form of the GTPase Rac1 that allows for optogenetic induction of spine shrinkage (Hayashi-Takagi *et al.*, 2015). There is also recent evidence that microtubule-associated proteins also participate in dendritic spine dynamics, as do epigenetic phenomena (see below). For a more detailed description of actin and other proteins involved in dendritic spine morphogenesis and plasticity, see (Bellot *et al.*, 2014; Bhatt *et al.*, 2009; Bramham, 2008; Fortin *et al.*, 2012; Halpain, 2000; Hotulainen & Hoogenraad, 2010; Koleske, 2013; Lamprecht, 2014; Murakoshi & Yasuda, 2012; Sala & Segal, 2014; Shirao & Gonzalez-Billault, 2013; Tada & Sheng, 2006).

Methods for assessing dendritic spine morphology

Assessment of dendritic spine dynamics has traditionally been performed in postmortem brain tissue using various methods, most frequently using Golgi-Cox or fluorescent dye staining. However, these methods are inherently limited in their ability to reveal information about changes in dendritic spines over time and/or in conjunction with specific behaviors, in that they allow only a “snapshot” in time rather than real-time imaging of spine dynamics. In addition, they typically involve using fixed tissue to allow for between-subjects comparisons. Fortunately, technological advances in microscopy and genetic engineering have now made imaging dendritic spine dynamics in the living brain a distinct reality. However, as mentioned below, imaging spine dynamics in deep brain structures with sufficient resolution remains a challenge.

Golgi-Cox staining

The first method utilized for assessing the fine structure of cells of the nervous system was the Golgi stain. This method was developed by renowned anatomist Camillo Golgi (Golgi, 1873), utilized by Cajal in his seminal studies described earlier (Ramón y Cajal, 1888; Ramón y Cajal, 1899), and refined by Cox and other groups of investigators to produce what is now widely referred to as the modified Golgi-Cox stain (Cox, 1891; Gibb & Kolb, 1998; Glaser & Van der Loos, 1981). In this procedure, post-mortem brain tissue is cut into relatively thick (~100–300 µm) sections, and immersed in a solution containing potassium chromate and dichromate and chloride salts of heavy metals (silver or mercury) for several weeks. Following tissue dehydration, clearing, and mounting onto microscope slides, Golgi-impregnated tissue can be viewed under brightfield microscopy to reveal detailed dendritic arborizations as well as high resolution of individual dendritic spines (Fig. 2A), which can then be quantified manually or semi-automatically using various computer-based algorithms (Orlowski & Bjarkam, 2012). Advantages of the Golgi-Cox method for assessing dendritic spine dynamics are its relative low cost and resistance to fading or photobleaching over time. Often cited disadvantages of Golgi-Cox staining are the limited ability to determine the neurochemical phenotypes of impregnated neurons, overlapping and out-of-focus dendritic segments, and underestimation of spine numbers due to the two-dimensional nature of the obtained images. However, methods to minimize these disadvantages have been described, including the use of fluorescent Golgi stains (Koyama & Tohyama, 2013), confocal laser

scanning microscopy to view Golgi-impregnated neurons in reflected mode to allow for subsequent three-dimensional reconstruction of serial images (Tredici *et al.*, 1993), and introducing immunofluorescence procedures to allow for identification of labeled neurons (Pinto *et al.*, 2012; Spiga *et al.*, 2011). Despite these advances, Golgi-Cox staining yields comparatively less morphological detail than newer fluorescence-based methods.

Fluorescent dyes

Another approach to examining neuronal microarchitecture is the use of fluorescent dyes that fill the cell body and its processes to reveal the fine structure of dendritic spines. Following labeling with fluorescent dyes, serial sections of images are obtained on a confocal laser scanning microscope and processed for three-dimensional reconstruction and analysis of dendritic spine density and morphology. One such fluorescent dye is Lucifer Yellow (Belichenko & Dahlstrom, 1995) which produces robust spine labeling (Fig. 2B). A disadvantage of this dye, however, is that cells must be filled manually via intracellular injections. Yet this method allows for specific cell impregnation, and is useful in the analysis of spine morphology following electrophysiological recordings, thus permitting structure and function to be measured from the same cell rather than from cells from different experimental cohorts. Another advantage of Lucifer Yellow staining is that fewer cells within a tissue section are labeled, allowing for less interference from overlapping cells and improved imaging and analysis.

An alternative method to Lucifer Yellow is the use of carbocyanine dyes such as 1,1'-dioctadecyl-3,3',3'-tetramethylindocarbocyanine (DiI; see Fig 2C), 3,3'-dioctadecyloxacarbocyanine (DiO), or 1,1'-dioctadecyl-3,3',3'-tetramethylindodicarbocyanine (DiD). The lipophilic nature of these dyes makes them particularly suitable for the assessment of dendritic spine morphology, as they are easily incorporated into the lipid bilayer and diffuse laterally to fill the neuronal microarchitecture. Methods have been developed to inject these dyes into tissue sections via helium-pressured ballistic “gene guns”, which results in cell labeling efficiency roughly equivalent to that achieved by Golgi-Cox staining. In addition, various investigators have introduced immunohistochemical methods to identify fluorescently labeled neurons (Dunaevsky, 2013; Gan *et al.*, 2009; Gan *et al.*, 2000; O'Brien & Lummis, 2006; Seabold *et al.*, 2010; Staffend & Meisel, 2011a; Staffend & Meisel, 2011b). As with Golgi-Cox staining, computer-based algorithms can be used for semi-automatic detection and classification of dendritic spines (Mancuso *et al.*, 2013; Parekh & Ascoli, 2013; Shen *et al.*, 2008). However, like most methods employing fluorescent labels, Lucifer Yellow and carbocyanine dye are susceptible to fading and photobleaching with prolonged imaging, as well as loss of anchoring to the labeled cell, especially when stored for extended periods of time. Generally, in our experience, DiI-labeled cells can only be imaged for up to ~6 months following tissue preparation prior to the leaching of the fluorescent dye from the membranes.

Other methods for assessing dendritic spine morphology include immunohistochemical staining for markers of dendritic spines such as PSD. However, this approach is inherently limited by poor and non-uniform antibody penetration into thicker tissue sections (Mancuso *et al.*, 2013). Similar methods utilized in ultra-thin sections, such as array tomography

followed by three-dimensional reconstruction of serially collected images, have also been utilized and appear to yield improved results (Micheva *et al.*, 2010). Unfortunately, this method requires laborious sample preparation and image reconstruction methods, limiting its utility in larger data sets.

In vivo imaging of spine plasticity

All of the aforementioned methods are performed in post-mortem tissue, and thus lack the ability to assess one of the most critical aspects of dendritic spines with respect to their behavioral relevance: dynamic changes in spine number and morphology over time, particularly during specific behaviors. This limitation has only recently been overcome with advanced imaging methods involving two-photon microscopy in awake and behaving animals. In this approach, either viral vectors or transgenic animals are employed to express fluorescent proteins such as enhanced green fluorescent protein (EGFP) or yellow fluorescent protein (EYFP) under the control of a ubiquitous or cell-type specific promoter (Grutzendler & Gan, 2006; Grutzendler *et al.*, 2011; Hayashi-Takagi *et al.*, 2015; Majewska *et al.*, 2006; Matsuzaki *et al.*, 2004; Ryan *et al.*, 2015; Trachtenberg *et al.*, 2002; Yang *et al.*, 2009; Yasumatsu *et al.*, 2008). Fluorescence is then imaged *in vivo* using a thinned skull, cranial window, or microendoscopic camera connected to a two-photon microscope system (Isshiki & Okabe, 2014). A particular advantage of this approach is that not only can cellular morphology be determined in an intact living brain, but other experimental methods such as fluorescent calcium or other ionic dye indicators as well as pharmacological uncaging (e.g., glutamate) can be incorporated, as can near infrared excitation wavelengths that penetrate deeper into tissue with reduced light scattering and absorption (Holtmaat *et al.*, 2012; Murayama & Larkum, 2012; Swanger *et al.*, 2011; Yuste, 2010). A disadvantage of this approach, however, is that the quality of the image through a thinned skull, without the incorporation of other methods such as microendoscopy, degrades at distances deeper than 50 μm from the brain surface (Isshiki & Okabe, 2014).

Do changes in spine morphology have behavioral relevance?

Learning and Memory Processes

Dendritic spines receive synaptic contacts that can be altered following new experiences, and are necessary in the processes of learning and memory. The overarching hypotheses regarding the role of dendritic spines in learning and memory have changed over time, beginning with the notion that spine density correlates with learning, eventually changing to the idea that morphological changes within the spine itself are more important for new learning and memory formation than total spine number or density (van der Zee, 2015). Some have postulated that spines with small diameters and/or lengths are unstable, easily eliminated, and responsible for acquisition of memory, whereas spines with larger head diameters are more stable and necessary for long-term memory formation (Kasai *et al.*, 2010; Kasai *et al.*, 2003). In line with this notion, it has been postulated that these spines with large head diameters arise from the maturation of thinner spines with small or absent head apparatuses (Bourne & Harris, 2007).

However, there is not universal agreement on the function of dendritic spines and their morphology with respect to learning and memory, particularly when discordant results are reported in studies that have utilized similar model systems of learning and memory. For example, increases in dendritic spine density in apical and basilar dendrites of hippocampal neurons have been observed following trace eyeblink conditioning (Leuner *et al.*, 2003), while others have failed to observe such changes on apical dendrites of hippocampal neurons (Geinisman *et al.*, 2000), though this latter study did report increases in total PSD area. A subsequent study by Geinisman and colleagues found that trace conditioning induced increases in the formation of multi-synapse boutons (MSBs), in which two or more postsynaptic elements are associated with a single presynaptic element (Geinisman *et al.*, 2001). These examples illustrate that discordant reports on effects of learning on spine dynamics, even those utilizing similar learning paradigms, may be a reflection of different experimental approaches (Golgi staining vs. spine ultrastructure via electron microscopy), different dendritic domains analyzed (apical vs. apical and basilar dendrites), or even different dependent measures (spine density vs. PSD area and number of MSBs). In light of this example, it is clear why it is difficult to come to a consensus regarding cause-and-effect relationships between spine dynamics and learning and memory.

Fortunately, more recent studies utilizing *in vivo* spine imaging have increased our understanding of this complex relationship. In one such study, Yang and colleagues (Yang *et al.*, 2009) utilized EYFP-expressing mice to examine spine dynamics in the primary motor cortex following a motor (rotarod) learning task, or in the barrel cortex following sensory experience (environmental enrichment). Experience-induced de novo spine formation was observed within two 2 days of task performance, and a small percentage of these newly formed spines persisted for months, while other spines were eliminated by subsequent exposure to novel experiences. Interestingly, in this study, control experiments demonstrated that spinogenesis was not a result of non-specific motor activity or sensory experience. These findings strongly suggest that sensory experiences result in long-lasting synaptic alterations that underlie persistent sensory or motor memories. However, since spine dynamics were not directly manipulated in this study, it is difficult to determine precisely whether spinogenesis and its persistence were induced by environmental input, or were a result of specific behavior output.

A more recent study (Hayashi-Takagi *et al.*, 2015) used direct experimental manipulations of spine dynamics to examine their role in learning and memory processes. In this study, the authors developed a photoactivatable form of the GTPase Rac1, which under conditions of prolonged activation results in spine shrinkage. The expression of this photoactivatable form of Rac1 (PaRac1) was targeted to recently activated synapses (AS-PaRac1) via use of the promoter for the activity-dependent immediate early gene Arc/Arg3.1. Using this technology, the authors demonstrated that motor learning in a rotarod task resulted in de novo spinogenesis and enlargement of existing spines in the primary motor cortex. Upon photoactivation of AS-PaRac1, which resulted in shrinkage of previously potentiated spines (estimated to be ~410,000 spines localized to ~4,700 neurons), the authors were able to disrupt (“erase”) learning in the rotarod task during a critical time period in the experiment. These highly novel findings suggest that not only are dendritic spines potentiated by learning experiences, but their disruption can adversely impact motor learning ability. Thus, changes

in dendritic spines can not only result from learning experiences, but also regulate behavioral output that is indicative of learning.

During early life development, when neuronal plasticity is arguably at its peak, it is thought that dendritic spines and synaptic contacts are formed or pruned via mechanisms that rely on an interplay between learning-based environmental input and behavioral output. Learning-related patterns of neuronal activity induce LTP and lead to stabilization and localization of spines (De Roo *et al.*, 2008a; De Roo *et al.*, 2008b). In very young animals (postnatal day 15 rats), a time of immense synaptogenesis, it has been demonstrated that theta burst stimulation (TBS)-induced LTP activates existing synapses and leads to the formation of new spines, illustrating robust plasticity of neuronal networks during development (Watson *et al.*, 2016). Interestingly, hippocampal synaptic growth and pruning during the early life postnatal period induces remodeling of neuronal networks which is regulated by activity. In one recent study, the impact of synaptic adhesion molecules on synaptic maturation and survival was assessed in 7 to 9 week old mice (Krzisch *et al.*, 2016). In this study, it was demonstrated that overexpression of the synaptic adhesion molecule SynCAM1 increased maturation of dendritic spines in the hippocampus, whereas overexpression of another adhesion molecule Neuroligin-1B increased spine density in the hippocampus. In contrast, overexpression of Neuroligin-2A increased spine density and GABAergic innervation, which resulted in significantly higher levels of neuronal survival. In addition, mice overexpressing Neuroligin-2A specifically in new neurons, showed impaired spatial learning performance in the Morris water maze. Although higher levels of spine survival appear to be linked to impaired performance in this particular task, suggesting that overall spine density may not be the most important neurobiological substrate for enhancing cognitive ability, it should be noted that the role of each adhesion molecule was assessed by cell-autonomous overexpression which may have led to other compensatory mechanisms and/or neuronal organization that resulted in altered cognitive performance. Regardless, these results show a link between synaptic maturation, increased neuronal survival during a period of high levels of synaptic reorganization, and learning processes.

Fear conditioning is a form of associative learning in which subjects exposed to an aversive stimulus associate a neutral stimulus with defense responses, after which this neutral stimulus thus becomes a conditioned stimulus which elicits a conditioned response (Blair *et al.*, 2001). In the preclinical literature, the neurobiological mechanisms of fear conditioning have been studied in detail. Alterations in spine morphology in the amygdala have been shown to occur following fear conditioning, such that profilin, an actin polymerization regulatory protein, impregnates dendritic spines which then undergo post-synaptic density enlargement (Lamprecht *et al.*, 2006). Interestingly, one study found that recruitment of calcium-permeable (GluA2-lacking) AMPA receptors was specific to mushroom-type spines in hippocampal CA1 neurons after fear conditioning (Matsuo *et al.*, 2008). Specifically, transgenic mice showed enhanced GFP-GluA1 recruitment into spines with large head diameters 24 hours after fear conditioning, which then reversed by 72 hours post-conditioning. This reversal may indicate replacement of AMPA receptor subunits to others such as GluA2, which are more stable and calcium impermeable. Here, dynamic changes in spine morphology seem to have functional relevance and occur as a result of a learning mechanism. Alterations in spine morphology have also been found in the lateral amygdala

(LA) in a fear conditioning paradigm (Radley *et al.*, 2006a), as conditioned and unconditioned stimuli activate neurons within this region, and damage to the LA inhibits acquisition of fear conditioning (Quirk *et al.*, 1997; Romanski *et al.*, 1993; Schafe *et al.*, 2001). Specifically, fear conditioning increased spine density measured via spinophilin-immunoreactivity in spine heads (Radley *et al.*, 2006a). These studies suggest that processes of learning and memory induce alterations in spine morphology, specifically with respect to changes underlying acquisition of Pavlovian associations.

In another study, Ryan and colleagues found that both optogenetic activation of entorhinal cortex as well as contextual fear conditioning increased dendritic spine densities in putative “engram” cells in the dentate gyrus, which encode context-dependent memories (Ryan *et al.*, 2015). The increases in spine density were paralleled by increases in membrane capacitance, and were abolished by administration of anisomycin, indicating the necessity of de novo protein synthesis for this phenomenon to occur. These findings were among the first to identify structural spine plasticity in optogenetically and behaviorally defined memory engram cells, and that structural plasticity in these specific cell populations was required for memory to occur. These findings support the notion that contextual conditioning drives spine changes in specific cell populations, rather than being induced by the behavior itself.

Finally, there is a wealth of evidence that maladaptive spine dynamics may contribute to, and/or be a result of, brain disease processes. In the case of Alzheimer's disease (AD), evidence suggests that the loss of dendritic spines is one of the first structural changes that occurs in neurons of AD patients. This dendritic spine loss is directly correlated with the loss of synaptic function, and is likely a result of β -amyloid accumulation, tau hyperphosphorylation, excitotoxicity, neuroinflammation, dysfunction of intracellular cytoskeletal regulators, or a combination of these factors (Dorostkar *et al.*, 2015; Fiala *et al.*, 2002; Knobloch & Mansuy, 2008). Most research efforts on slowing the progress of AD have focused on amyloid clearance, neuroprotection, and delay of cognitive decline. However, new evidence suggests that promoting synaptogenesis and/or spinogenesis, perhaps via epigenetic mechanisms, may be of potential benefit. This was demonstrated recently in mouse models of AD, where histone deacetylase inhibitors were found to promote synaptogenesis, dendritic sprouting, and regaining of long-term memories especially when combined with environmental enrichment (Fischer *et al.*, 2007). Thus, both intrinsic and extrinsic factors affecting spine dynamics may provide more effective avenues for treating age-related neurodegenerative diseases.

Addiction

Exposure to drugs of abuse such as nicotine (Brown & Kolb, 2001), cocaine and amphetamines (Robinson & Kolb, 1999a), alcohol (Carpenter-Hyland & Chandler, 2006; Spiga *et al.*, 2014a; Spiga *et al.*, 2014b), cannabis (Spiga *et al.*, 2010) and opiates (Robinson & Kolb, 1999b) can alter dendritic spine morphology (such as spine density or size) in various brain regions including the prefrontal cortex (PFC) and NA (Russo *et al.*, 2010). As a specific example, exposure to cocaine leads to differential changes in NA spine dynamics when measured proximally or distally from the cell soma, illustrating that spine morphology varies according to location on the dendrite (Dumitriu *et al.*, 2012). In many of these studies,

animals received experimenter-delivered injections of various drugs of abuse, and dendritic spine structure was characterized after a period of withdrawal, indicating drug-induced alterations in constitutive spine morphology. In addition to experimenter-delivered exposure, however, dendritic spines have been shown to be important in behaviors motivated by drugs of abuse. The development of addiction and relapse vulnerability involves complex learning and memory processes, as well as associations with constellations of environmental stimuli that become powerful cues in initiating craving and relapse behaviors (Beckmann *et al.*, 2011; Gipson *et al.*, 2014; Marchant *et al.*, 2015). Addiction is a complex, progressing disorder involving learning and memory processes (Hyman *et al.*, 2006; Nestler, 2001). Animal studies have long established that previously neutral environmental stimuli consistently and contiguously paired with drugs of abuse begin to take on an important associative meaning. These cues enhance drug self-administration thus illustrating the cue-dependency of drug use (Caggiula *et al.*, 2001; Schenk & Partridge, 2001). Once these stimuli become conditioned or discriminative stimuli predicting drug presentation, they consistently produce drug-seeking in animal models of cue-induced relapse (Bossert *et al.*, 2013; Crombag *et al.*, 2008; See, 2002; Shaham *et al.*, 2003) and can also elicit structural changes in dendritic spines within the nucleus accumbens (NA). Specifically, exposure to an amphetamine-paired environmental context elicited an increase in spine density and frequency in activated (c-fos positive) cells in the NA compared to non-paired controls (Singer *et al.*, 2016).

Interestingly, exposure to different drug classes can induce differential changes in spine morphology (Dumitriu *et al.*, 2012; Gipson *et al.*, 2014; Gipson *et al.*, 2013a; Shen *et al.*, 2011), however there are also overlapping patterns of change that occur as well (Mulholland *et al.*, 2016; Scofield *et al.*, 2016). Drug-induced spine changes are also highly dynamic and governed by glutamatergic and dopaminergic input from mesocorticolimbic circuit afferents (Shen *et al.*, 2014; Yagishita *et al.*, 2014). For example, Shen and colleagues (Shen *et al.*, 2009) demonstrated that acute passive cocaine exposure increases NA spine density accompanied by increases in excitatory transmission as well as levels of proteins involved in actin regulation, protein catabolism, and glutamatergic transmission on the order of minutes to hours. These changes appear to return to pre-drug levels by approximately by approximately 24 hours following cocaine exposure. However, in animals with a history or repeated cocaine exposure, such changes in protein levels follow a different time course and yield bi-direction changes in spine head diameters, with enlarged spine heads followed by reductions in spine head diameters (Shen *et al.*, 2009). These findings suggest a considerable degree of metaplasticity of dendritic spine dynamics in the NA that depend on the prior history of drug exposure. While such drug-induced changes in NA dynamics are largely attributable to alterations in excitatory glutamatergic transmission, recent optogenetic studies have revealed that synaptically released dopamine also contributes to NA spine dynamics (specifically, spine enlargement), although only within a narrow time window that follows glutamate-induced changes (Yagishita *et al.*, 2014).

These results highlight important mechanisms underlying spine dynamics, specifically the role of dopaminergic innervation of NA MSNs. Older studies revealed that dopamine depletion via 6-hydroxydopamine lesions of the NAcore or shell induced region-specific structural changes such as reduced dendritic length and spine density, and increased

dendritic tortuosities (Meredith *et al.*, 1995). More recent findings indicate that subpopulations of NA MSNs, specifically those expressing dopamine D1 or D2 receptors, are differentially altered during cocaine-seeking (Heinsbroek *et al.*, 2014). Thus, the consequential changes in NA spines following exposure to drugs of abuse likely involve a complex and time-dependent interplay between glutamatergic and dopaminergic transmission. On an anatomical level, individual dendritic spines of NA MSNs often receive converging inputs, with glutamatergic afferents primarily forming synaptic contacts on spine heads and dopaminergic inputs forming synapses on spine necks (Sesack *et al.*, 2003). It is currently unclear what alterations in this convergent synaptic architecture occur across different classes of abused drugs, drug history, and phases of the addiction cycle, and subsequently influence addiction-related behaviors.

Dendritic spine dynamics may also be influenced by changes in dopamine and glutamatergic transmission that persist into withdrawal, such as hyperglutamatergic transmission during alcohol withdrawal (Holmes *et al.*, 2013), and hypoactivity of both glutamatergic and dopaminergic transmission following chronic exposure to cocaine (Baker *et al.*, 2003; Diana, 2011; Weiss *et al.*, 1992). In turn, glutamatergic and dopaminergic transmission in the NA become dysregulated during the reinstatement of cocaine-seeking behavior (McFarland *et al.*, 2003; Neisewander *et al.*, 1996). These drug-dependent fluctuations in both dopamine and glutamate signaling likely induce numerous transient or lasting changes in NA spine dynamics.

With regard to other drugs of abuse such as alcohol, proteomic analyses have revealed that changes in PSD-associated proteins are associated with alcohol-induced changes in synaptic morphology within the NA (Uys *et al.*, 2016). This study found that exposure to and withdrawal from chronic intermittent ethanol (CIE) induced an increase in NA spine density, dendritic shaft diameter, as well as spine head diameter. Additionally, this study found significant alterations in the expression of a multitude of scaffolding proteins, glycoproteins, kinases, neuropeptides, among many others within the NA after CIE. While alterations in spine dynamics occurred following exposure to ethanol, it is unclear if these changes are important for or relevant to ethanol-related motivated behavior.

Various investigators have postulated that drug-induced spine changes create a neurobiological environment that is more plastic compared to drug-naïve conditions, and may lead to aberrant motivation to seek the drug (Mulholland & Chandler, 2007; Nyberg, 2014; Spiga *et al.*, 2014a; Young *et al.*, 2014; Young *et al.*, 2016). While most studies supporting this theory have examined changes in spine dynamics in mesocorticolimbic circuits, an eloquent series of studies by Young and colleagues have recently expanded this to include the amygdala, a key regulator of the emotional and associative components of addiction. Utilizing place conditioning procedures, it was demonstrated that methamphetamine increased the plasticity of spines in the amygdala, and that inhibition of actin polymerization with latrunculin A in this region (which inhibits actin depolymerization and thus promotes spine stability) disrupted the retrieval of methamphetamine-associated contextual memories (Young *et al.*, 2014). These effects appeared to be specific to methamphetamine, as they were not observed following contextual conditioning with saline, food, or footshock. The ability of spine stabilization to inhibit methamphetamine-associated

contextual memory were also independent of the strength of contextual training, and were observed following context-induced reinstatement in an operant paradigm. Unfortunately, due to its ubiquitous expression in numerous cell types throughout the brain and body, inhibition of actin polymerization with latrunculin A may not selectively modulate dendritic spines *per se*. To address this issue, Young and colleagues also demonstrated that intra-amygdala infusions of Blebbistatin, an inhibitor of the synaptically enriched actin polymerization promoter non-muscle myosin IIB, produces similar inhibitory effects on methamphetamine-induced contextual memories (Young *et al.*, 2014; Young *et al.*, 2016). Finally, Young *et al.* showed that systemic administration of Blebbistatin selectively disrupted the storage methamphetamine- but not fear-associated contextual memories, and also reversed the ability of methamphetamine-associated memories to increase spine density in the amygdala (Young *et al.*, 2016). Thus, specific targeting of synaptic actin dynamics may represent a novel approach to treating pathological memories associated with abused drugs.

Within dendritic spines, cell surface proteins such as AMPA and NMDA receptors are inserted or internalized regularly (Kopec & Malinow, 2006). Following extended periods of withdrawal from cocaine self-administration, craving for cocaine can “incubate”, and this has been associated with an increase in calcium-permeable (GluA2-lacking) AMPA receptors (Conrad *et al.*, 2008). Following extinction of cocaine self-administration, baseline increases in spine head diameter in the NA have been found (Gipson *et al.*, 2013a; Shen *et al.*, 2009), supporting the notion that cocaine causes NA spines to exist in a relatively potentiated state to increase surface area for protein insertion. It should be noted, however, that withdrawal from heroin self-administration results in an LTD-like state of NA core neurons, in which spines show reductions in head diameter compared to those of drug-naïve animals (Shen *et al.*, 2011). In addition, following heroin self-administration, there is an upregulation of NMDA receptors that has functional relevance in heroin-seeking behavior (Shen *et al.*, 2011). Interestingly, a hypothesis has recently emerged that the brain returns to more developmental, immature-like conditions following chronic exposure to drugs of abuse (Dong & Nestler, 2014). In this “neural rejuvenation” hypothesis, exposure to drugs of abuse lead to alterations in brain physiology that are comparable to developmental conditions at the molecular, cellular, and circuitry levels. In doing so, drugs of abuse open a window of plasticity that is not found in the adult brain, which may underlie habitual and pathological drug seeking behaviors. This notion is supported by the associated increase of calcium-permeable AMPA receptors in the NA as well as expression of the GluN2B subunit of NMDA receptors and silent synapses after withdrawal from different drugs of abuse (Conrad *et al.*, 2008; Dong, 2015; Gipson *et al.*, 2013b; Lee & Dong, 2011; Shen *et al.*, 2011). Interestingly, silent synapses, which lack AMPA receptors and thus are unable to remove the magnesium block of NMDA receptors, exist during development as well as after cocaine exposure. Although the behavioral relevance of these silent synapses in addiction processes is still not fully characterized, they have been associated with incubation of cocaine craving (Lee *et al.*, 2013). As well, it was recently found that re-silencing these synapses via optogenetic removal of calcium-permeable AMPA receptors, allowing environmental enrichment (EE; previously found to have anti-relapse beneficial effects (Thiel *et al.*, 2009))

to lead to insertion of calcium-impermeable AMPA receptors which inhibit relapse-like behaviors (Ma *et al.*, 2016).

In addition to baseline changes in dendritic spines after drug exposure and periods of withdrawal, drug-seeking behavior has been associated with altered dendritic spine morphology (Gipson *et al.*, 2013a; Gipson *et al.*, 2013b; Shen *et al.*, 2014; Stefanik *et al.*, 2016). Specifically, conditioned stimuli associated with self-administration of drugs of abuse such as cocaine and nicotine can elicit motivated behavior to seek the drug. These stimuli are also associated with rapid, transient increases in spine head diameters of NA core medium spiny neurons (MSNs). In the reinstatement model of drug relapse, exposure to drug-associated cues was associated with the rapid and transient growth of dendritic spines (within 15 min), with no alterations in dendritic spine density during this timeframe. Importantly, these changes in spines (as well alterations in the ratio of AMPA to NMDA receptor-mediated excitatory post-synaptic currents, a functional correlate of synaptic plasticity) were positively correlated with magnitude of cue reinstated drug seeking (Gipson *et al.*, 2013a). These rapid, transient changes in spine dynamics were found to be specific to drug seeking behavior rather than behavior motivated by food. However, since some overlap has been found between the neurobiological substrates of obesity and drug abuse (Brown *et al.*, 2015; Kenny, 2011a; Kenny, 2011b), it is thus possible that in a model of diet-induced obesity, synaptic plasticity may occur during food seeking behavior similar to that induced by drugs of abuse. It is also possible that in drug naïve animals, food seeking behavior (particularly in food restricted animals) induces rapid, transient plasticity on a much shorter timescale, thus going undetected at the 15 min time point assessed for drug-induced changes. In the drug-exposed system, the rapid, transient increase in dendritic spine head diameter found after 15 min of cue-induced cocaine seeking occurred specifically in dopamine D1-expressing (but not D2-expressing) MSNs (Heinsbroek *et al.*, 2014). These results illustrate cell-type specificity of reinstatement-associated spine dynamics. It still remains unclear, however, if drugs of abuse lead to these changes which then impact behavior, or if behavior itself (drug-seeking) drives these changes.

Stress, Depression, and Aging

Acute or repeated exposure to stressors such as restraint or social defeat can remodel dendritic spine morphology in multiple brain regions. Although different types of stressors likely impact dendritic spine dynamics in different ways, preclinical modeling of stress has evolved in an attempt to capture criterion, construct, and translational validity of human stress exposure. Stress exposure can impact numerous neurobiological systems, all of which can vary based on a number of factors: the type and duration of stressor exposure (Capriles & Cancela, 1999), developmental stage of the organism (Biala *et al.*, 2011; Romeo *et al.*, 2006; Romeo & McEwen, 2006), genetic and epigenetic factors (McEwen *et al.*, 2012; McEwen *et al.*, 2016; Pillai *et al.*, 2012), sex and hormones (McLaughlin *et al.*, 2009; McLaughlin *et al.*, 2005; Weinstock, 2011), anxiety levels (Adamec *et al.*, 2012; McEwen *et al.*, 2012), among others (Leuner & Shors, 2013). Because stress is a complex phenomenon and involves both external stimuli and internal physiological responses to them, experimental models of stress are highly varied. Commonly used models include restraint (Esparza *et al.*, 2012; Platt & Stone, 1982), footshock (Long & Fanselow, 2012), or social

defeat stress (Covington *et al.*, 2005) among others such as chronic variable or unpredictable stress (Lopes *et al.*, 2016). Although the translational validity of these models remains an important topic, these are typically intended to model aspects of anxiety, post-traumatic stress disorder (PTSD), and/or depression. Interestingly, acute restraint stress has been postulated to induce some neurobiological and behavioral alterations akin to that induced by stress disorders such as PTSD in humans. Specifically, one study found increased NA core dendritic spine density 3 weeks after one 2-hr session of restraint stress in rats (Garcia-Keller *et al.*, 2015). This acute stress exposure also increased acquisition of cocaine self-administration, suggesting stress-induced vulnerability to addiction-related behaviors. Stress and cocaine have been shown to differentially alter cell-type specific spine morphology when measured with two-photon laser scanning microscopy (Khibnik *et al.*, 2015). Specifically, animals given repeated cocaine injections showed decreased synaptic strength in spines localized to D1-expressing MSNs and increased synaptic responses in D2-expressing MSNs. In contrast, however, this study revealed that animals exposed to chronic social defeat stress, showed increased synaptic strength in D1-expressing and reduced synaptic strength in D2-expressing MSNs. These opposing results illustrate the importance of not only duration of stress exposure but also cell-type specificity in the effects of stress exposure on spine dynamics.

While the mechanisms underlying the impact of acute or chronic stress on spine dynamics is currently a subject of intense study, some important advancements has been made. As noted above, it remains an issue in the addiction field to determine whether spine changes occur prior to drug-induced behaviors, leading to increased relapse vulnerability, or if these changes occur as a result of motivated behavior. A significant issue with solving this question is that teasing apart spine dynamics from behavior is difficult within the context of addiction as a complex biobehavioral disorder involving learning components as well as circuitry, pharmacological, and multi-level neurobiological mechanisms that lead to drug-seeking and intake. In the stress literature, however, alterations in spine dynamics has been studied independently of actual stress exposure by passive administration of corticosterone, the predominant stress steroid released in rodents following exposure to a stressor. Passive administration of corticosterone itself can induce dendritic spine remodeling in the hippocampus by impacting cytoskeletal protein expression (Cereseto *et al.*, 2006). Further, it has been demonstrated that repeated exposure to corticosterone over the course of 3 weeks induced remodeling of dendritic spines in the medial prefrontal cortex (mPFC) (Anderson *et al.*, 2016; Gourley *et al.*, 2013; Radley *et al.*, 2008). Specifically, animals chronically exposed to corticosterone showed a decrease in spine volume that persisted for 3 weeks following cessation of treatment (although these effects were dependent on brain region; see (Gourley *et al.*, 2013)). These results demonstrate that spine remodeling that results from a biological response associated with stress (elevated corticosterone levels) that can be separated from the stress exposure itself. As of yet, however, it has been difficult to examine addiction-related changes in spine dynamics in isolation from the behavioral, learning and memory, and/or environmental components of the addiction process. The results reviewed above suggest that some aspects of stress-induced dendritic remodeling can be simulated by chronic, steady exposure to corticosterone. This has implications for determining not only mechanisms underlying some aspects of stress-induced alterations in spine dynamics, but

also if spine remodeling is sufficient and/or necessary to occur before or after a behavioral stress response. Although this is an important advancement in our understanding of the influence of stress-related hormones on spine dynamics, it is unclear if such spine alterations actually mimic those that occur in response to an actual stressor, and if such approaches have translational validity for examining the neurobiological substrates of stress disorders such as PTSD. As noted above, stress is complex and leads to various biological responses. Thus, mimicking the biological response of elevated corticosterone in rodents may only reveal some of the processes that occur in response to a more complex stress experience.

Some preclinical stress models (e.g., chronic restraint, social defeat) have been used to model various aspects of depression (Qiao *et al.*, 2016; Shimamoto *et al.*, 2011). Dendritic atrophy has been found both in the PFC and hippocampus following chronic restraint stress (Conrad, 2006; Conrad *et al.*, 1999; Magarinos & McEwen, 1995; McEwen *et al.*, 1997; Radley *et al.*, 2006b; Stewart *et al.*, 2005), paralleled by reductions in dendritic spine density (reviewed in (Duman & Duman, 2015; Licznarski & Duman, 2013). These findings are similar to those from postmortem examinations of brain tissue from depressed human subjects, where reduced spine density, overall synapse number on dendritic spines, as well as reduced synaptic protein levels have been observed in the dorsolateral prefrontal cortex and hippocampus, and opposite changes observed in the amygdala (Kang *et al.*, 2012; Licznarski & Duman, 2013). Thus, spine and synapse remodeling has been proposed to be a novel approach to treating depressive disorders (Duman & Duman, 2015; Licznarski & Duman, 2013). Other brain regions where spine morphology has been shown to be affected by chronic stress is the NA, where increases in the density of shorter spines with smaller PSDs have been observed, paralleled by increases in functional glutamatergic synapses (Christoffel *et al.*, 2011). It is unclear, however, if similar alterations exist in the NA of depressed human subjects.

Spine loss is also associated with brain aging, and there is some evidence of overlap between the mechanisms of spine loss in neurodegenerative diseases such as AD and those induced by chronic stress, namely a critical role for tau proteins (Lopes *et al.*, 2016; Sotiropoulos & Sousa, 2016). Thus, models of chronic stress may have cross-validity for other models of neurodegenerative diseases with significant cognitive impairment. Changes in dendritic spine morphology parallels age-related cognitive decline (Morrison & Hof, 1997), as neural pathways vulnerable in aging are comprised of glutamatergic PFC and hippocampal spiny pyramidal neurons (Dumitriu *et al.*, 2010; Morrison & Baxter, 2012; von Bohlen und Halbach *et al.*, 2006). In line with these findings, aged female rats show a decline in object recognition memory as well as PFC spine density (Wallace *et al.*, 2007). Formation of new dendritic spines occurs in clusters around activated synapses (De Roo *et al.*, 2008b) both in cell culture as well as *in vivo* following motoric learning (Fu *et al.*, 2012). Since clusters of dendritic spines that are important in fostering synaptic strength appear to be impacted by age-related decline, it is important to note that a recent study found that administration of riluzole, an activator of glutamate uptake that promotes synaptic over extrasynaptic NMDA receptor, rescued these clusters (Pereira *et al.*, 2014). Riluzole-treated animals also showed a positive correlation between memory-related behavior and density of spines on apical dendrites in the CA1 region of the hippocampus, suggesting that rescuing dendritic spine density in this area is important in enhancing memory processes.

Discussion and Conclusions

Dendritic spine plasticity appears to be critical not only for normal brain development, function, and aging, but also underlies the pathophysiology of several psychiatric, neurodevelopmental, and neurodegenerative disorders. Although newer advanced techniques have been developed to measure spine changes in three dimensions at high resolution, significant issues remain with measuring spine dynamics from one or more brain structures during behavior. Measuring dendritic spines in real time requires highly advanced imaging techniques which are more difficult when examining deep structures *in vivo*. In addition, much research has relied on categorization of the structure of dendritic spines based on distinct morphologies, yet often these categorizations are not accompanied by evidence for functional divergence between different spine types. The lack of analysis of both structure *and* function within a study may lead to inaccurate conclusions regarding the importance of dendritic spine dynamics in behaviors and/or cognitive processes. As well, given recent evidence that accurate spine categorization may suffer from limitations of currently and frequently used imaging tools (Tonnesen *et al.*, 2014), we suggest that categorizations based on structure alone be avoided without functional correlates. Rather, the distributions of various spine characteristics (e.g., neck width and length, head diameter) be analyzed and compared between groups (or within-subject when possible).

Importantly, it should be mentioned that the function of dendritic spines has been examined in various studies using the correlate of AMPA to NMDA post-synaptic excitatory current ratios recorded using whole cell patch clamp electrophysiology from the soma (Gipson *et al.*, 2013a; Shen *et al.*, 2014; Stefanik *et al.*, 2015). Although this yields an important amount of information and is a window into the potential summation of spine function across an entire neuron, it is imperative to differentiate whole cell function from individual spine function, especially when examining nuanced changes in spine structural characteristics and categorization. Technological advances are needed to image individual spine changes *in vivo* in awake, unrestrained, and behaving animals in deep structures to better characterize the underlying functional impact of dendritic spine dynamics on behavior.

Finally, the original question posed at the beginning of this review remains: do changes in dendritic spines lead to changes in behavior, or does behavior itself lead to changes in spine morphology? In this review, we have attempted to highlight potential answers to this question within the existing literature, depending on the type of environmental input or behavioral output under examination. It appears that in the case of stress, some biological responses such as alterations in spine dynamics can be mimicked using pharmacological manipulation (e.g., corticosterone administration). This contribution is important, as it opens the possibility that spine dynamics occur in *response* to a biological process associated with exposure to an environmental stimulus. However, interpretations of such results may reflect only a small part of larger and more complex stress responses. With regard to learning and memory and drug addiction (specifically, relapse-related behaviors), the ability to separate spine dynamics from behavior remains difficult and elusive. Thus, without more advanced and accessible *in vivo* imaging techniques, this may not be possible.

Along these lines, we argue that there is a great need to develop an experimental toolbox to selectively modulate dendrite spine dynamics, in order to observe direct effects on behavior. For example, several groups of investigators have utilized localized infusions of latrunculin A, an actin polymerization inhibitor (Coue *et al.*, 1987), to promote spine stabilization and investigate the effects of spine stabilization on drug-related behavior (Esparza *et al.*, 2012; Toda *et al.*, 2010; Toda *et al.*, 2006; Young *et al.*, 2014). However, such approaches lack cell-type specificity, and would potentially affect all local cytoskeletal cycling. With this limitation, other investigators have explored targeting more upstream regulators of actin cycling, such as non-muscle myosin IIB (Young *et al.*, 2016). These avenues of research, as well as the development of other approaches (e.g., optogenetic activation of GTPases or other modulators of spine morphology, chemogenetic, chemical uncaging), are critical for advancing our knowledge of the functional contributions of dendritic spine dynamics on behavior, and vice versa.

The most likely answer to our overall question is that dendritic spine dynamics are both a cause *and* consequence of behavior. We argue that it is most likely that environmental stimuli lead to constitutive changes in dendritic spine morphology, and these alterations create a neural environment in which an organism is rendered more likely to engage in certain behaviors. However, non-specific pharmacological manipulation of spine dynamics will likely not be a viable therapeutic option for treatment of brain disorders with underlying dendritic spine pathologies. Such an approach would also affect normal synaptic plasticity and likely lead to various adverse effects on normal cognition and learning and memory. A more viable option would be modulating dendritic spine dynamics via environmental influences, such as environmental enrichment (Gipson *et al.*, 2011; Rojas *et al.*, 2013; Thiel *et al.*, 2012; Turner *et al.*, 2003; Yang *et al.*, 2009) or approaches directed at engaging specific neural circuits to strengthen synapses that promote cognitive health or weaken synapses that promote maladaptive behaviors.

Acknowledgements

The authors would like to thank Drs. Sam Golden, Scott Russo, and Daniel Christoffel for the Lucifer-Yellow images in Figure 1B. The authors would also like to thank Dr. Cheryl Conrad for comments on an earlier version of this review. The authors are funded by NIH grants R00 DA036569 (CDG) and R01 DA025606, R01 DA024355, and R21 DA037741 (MFO).

References

- Adamec R, Hebert M, Blundell J, Mervis RF. Dendritic morphology of amygdala and hippocampal neurons in more and less predator stress responsive rats and more and less spontaneously anxious handled controls. *Behav Brain Res.* 2012; 226:133–146. [PubMed: 21925210]
- Alvarez VA, Sabatini BL. Anatomical and physiological plasticity of dendritic spines. *Annu. Rev. Neurosci.* 2007; 30:79–97. [PubMed: 17280523]
- Anderson RM, Glanz RM, Johnson SB, Miller MM, Romig-Martin S, Radley JJ. Prolonged corticosterone exposure induces dendritic spine remodeling and attrition in the rat medial prefrontal cortex. *J Comp Neurol.* 2016
- Arellano JI, Benavides-Piccione R, Defelipe J, Yuste R. Ultrastructure of dendritic spines: correlation between synaptic and spine morphologies. *Front Neurosci.* 2007; 1:131–143. [PubMed: 18982124]
- Attardo A, Fitzgerald JE, Schnitzer MJ. Impermanence of dendritic spines in live adult CA1 hippocampus. *Nature.* 2015; 523:592–596. [PubMed: 26098371]

- Baker DA, McFarland K, Lake RW, Shen H, Tang XC, Toda S, Kalivas PW. Neuroadaptations in cystine-glutamate exchange underlie cocaine relapse. *Nat. Neurosci.* 2003; 6:743–749. [PubMed: 12778052]
- Beckmann JS, Marusich JA, Gipson CD, Bardo MT. Novelty seeking, incentive salience and acquisition of cocaine self-administration in the rat. *Behav Brain Res.* 2011; 216:159–165. [PubMed: 20655954]
- Belichenko PV, Dahlstrom A. Studies on the 3-dimensional architecture of dendritic spines and varicosities in human cortex by confocal laser scanning microscopy and Lucifer yellow microinjections. *J. Neurosci. Methods.* 1995; 57:55–61. [PubMed: 7791365]
- Bellot A, Guivernau B, Tajés M, Bosch-Morato M, Valls-Comamala V, Munoz FJ. The structure and function of actin cytoskeleton in mature glutamatergic dendritic spines. *Brain Res.* 2014; 1573C:1–16.
- Bhatt DH, Zhang S, Gan WB. Dendritic spine dynamics. *Annu. Rev. Physiol.* 2009; 71:261–282. [PubMed: 19575680]
- Biala YN, Bogoch Y, Bejar C, Linial M, Weinstock M. Prenatal stress diminishes gender differences in behavior and in expression of hippocampal synaptic genes and proteins in rats. *Hippocampus.* 2011; 21:1114–1125. [PubMed: 20623763]
- Blair HT, Schafe GE, Bauer EP, Rodrigues SM, LeDoux JE. Synaptic plasticity in the lateral amygdala: a cellular hypothesis of fear conditioning. *Learn Mem.* 2001; 8:229–242. [PubMed: 11584069]
- Bosch M, Hayashi Y. Structural plasticity of dendritic spines. *Curr. Opin. Neurobiol.* 2012; 22:383–388. [PubMed: 21963169]
- Bossert JM, Marchant NJ, Calu DJ, Shaham Y. The reinstatement model of drug relapse: recent neurobiological findings, emerging research topics, and translational research. *Psychopharmacology.* 2013; 229:453–476. [PubMed: 23685858]
- Bourne J, Harris KM. Do thin spines learn to be mushroom spines that remember? *Curr. Opin. Neurobiol.* 2007; 17:381–386. [PubMed: 17498943]
- Bramham CR. Local protein synthesis, actin dynamics, and LTP consolidation. *Curr. Opin. Neurobiol.* 2008; 18:524–531. [PubMed: 18834940]
- Brown RM, Kupchik YM, Spencer S, Garcia-Keller C, Spanswick DC, Lawrence AJ, Simonds SE, Schwartz DJ, Jordan KA, Zhou TC, Kalivas PW. Addiction-like Synaptic Impairments in Diet-Induced Obesity. *Biological psychiatry.* 2015
- Brown RW, Kolb B. Nicotine sensitization increases dendritic length and spine density in the nucleus accumbens and cingulate cortex. *Brain Res.* 2001; 899:94–100. [PubMed: 11311869]
- Caggiula AR, Donny EC, White AR, Chaudhri N, Booth S, Gharib MA, Hoffman A, Perkins KA, Sved AF. Cue dependency of nicotine self-administration and smoking. *Pharmacology, biochemistry, and behavior.* 2001; 70:515–530.
- Capriles N, Cancela LM. Effect of acute and chronic stress restraint on amphetamine-associated place preference: involvement of dopamine D(1) and D(2) receptors. *Eur. J. Pharmacol.* 1999; 386:127–134. [PubMed: 10618462]
- Carpenter-Hyland EP, Chandler LJ. Homeostatic plasticity during alcohol exposure promotes enlargement of dendritic spines. *Eur. J. Neurosci.* 2006; 24:3496–3506. [PubMed: 17229098]
- Cereseto M, Reines A, Ferrero A, Sifonios L, Rubio M, Wikinski S. Chronic treatment with high doses of corticosterone decreases cytoskeletal proteins in the rat hippocampus. *Eur J Neurosci.* 2006; 24:3354–3364. [PubMed: 17229084]
- Chang FL, Greenough WT. Transient and enduring morphological correlates of synaptic activity and efficacy change in the rat hippocampal slice. *Brain Res.* 1984; 309:35–46. [PubMed: 6488013]
- Christoffel DJ, Golden SA, Russo SJ. Structural and synaptic plasticity in stress-related disorders. *Rev Neurosci.* 2011; 22:535–549. [PubMed: 21967517]
- Cingolani LA, Goda Y. Actin in action: the interplay between the actin cytoskeleton and synaptic efficacy. *Nat Rev Neurosci.* 2008; 9:344–356. [PubMed: 18425089]
- Conrad CD. What is the functional significance of chronic stress-induced CA3 dendritic retraction within the hippocampus? *Behav Cogn Neurosci Rev.* 2006; 5:41–60. [PubMed: 16816092]

- Conrad CD, LeDoux JE, Magarinos AM, McEwen BS. Repeated restraint stress facilitates fear conditioning independently of causing hippocampal CA3 dendritic atrophy. *Behav. Neurosci.* 1999; 113:902–913. [PubMed: 10571474]
- Conrad KL, Tseng KY, Uejima JL, Reimers JM, et al. Heng LJ, Shaham Y, Marinelli M, Wolf ME. Formation of accumbens GluR2-lacking AMPA receptors mediates incubation of cocaine craving. *Nature.* 2008; 454:118–121. [PubMed: 18500330]
- Coue M, Brenner SL, Spector I, Korn ED. Inhibition of actin polymerization by latrunculin A. *FEBS Lett.* 1987; 213:316–318. [PubMed: 3556584]
- Covington HE 3rd, Kikusui T, Goodhue J, Nikulina EM, Hammer RP Jr. Miczek KA. Brief social defeat stress: long lasting effects on cocaine taking during a binge and zif268 mRNA expression in the amygdala and prefrontal cortex. *Neuropsychopharmacology.* 2005; 30:310–321. [PubMed: 15496936]
- Cox WH. Imprgnation des centralen Nervensystems mit Quecksilbersalzen. *Arch Mikrosk Anat.* 1891; 37:16–21.
- Crick F. Do dendritic spines twitch? *Trends Neurosci.* 1982; 5:44–46.
- Crombag HS, Bossert JM, Koya E, Shaham Y. Review. Context-induced relapse to drug seeking: a review. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences.* 2008; 363:3233–3243. [PubMed: 18640922]
- Cruz-Martin A, Crespo M, Portera-Cailliau C. Delayed stabilization of dendritic spines in fragile X mice. *J Neurosci.* 2010; 30:7793–7803. [PubMed: 20534828]
- De Roo M, Klauser P, Garcia PM, Pogli L, Muller D. Spine dynamics and synapse remodeling during LTP and memory processes. *Prog. Brain Res.* 2008a; 169:199–207. [PubMed: 18394475]
- De Roo M, Klauser P, Muller D. LTP promotes a selective long-term stabilization and clustering of dendritic spines. *PLoS biology.* 2008b; 6:e219. [PubMed: 18788894]
- Desmond NL, Levy WB. Changes in the numerical density of synaptic contacts with long-term potentiation in the hippocampal dentate gyrus. *J. Comp. Neurol.* 1986; 253:466–475. [PubMed: 3025272]
- Diana M. The dopamine hypothesis of drug addiction and its potential therapeutic value. *Front Psychiatry.* 2011; 2:64. [PubMed: 22144966]
- Dong Y. Silent Synapse-based Circuitry Remodeling in drug addiction. *Int J Neuropsychopharmacol.* 2015
- Dong Y, Nestler EJ. The neural rejuvenation hypothesis of cocaine addiction. *Trends Pharmacol. Sci.* 2014; 35:374–383. [PubMed: 24958329]
- Dorostkar MM, Zou C, Blazquez-Llorca L, Herms J. Analyzing dendritic spine pathology in Alzheimer's disease: problems and opportunities. *Acta Neuropathol.* 2015; 130:1–19. [PubMed: 26063233]
- Duman CH, Duman RS. Spine synapse remodeling in the pathophysiology and treatment of depression. *Neurosci. Lett.* 2015; 601:20–29. [PubMed: 25582786]
- Dumitriu D, Hao J, Hara Y, Kaufmann J, Janssen WG, Lou W, Rapp PR, Morrison JH. Selective changes in thin spine density and morphology in monkey prefrontal cortex correlate with aging-related cognitive impairment. *J Neurosci.* 2010; 30:7507–7515. [PubMed: 20519525]
- Dumitriu D, Laplant Q, Grossman YS, Dias C, Janssen WG, Russo SJ, Morrison JH, Nestler EJ. Subregional, dendritic compartment, and spine subtype specificity in cocaine regulation of dendritic spines in the nucleus accumbens. *J. Neurosci.* 2012; 32:6957–6966. [PubMed: 22593064]
- Dunaevsky A. The gene-gun approach for transfection and labeling of cells in brain slices. *Meth Mol Biol.* 2013; 1018:111–118.
- Durand GM, Kovalchuk Y, Konnerth A. Long-term potentiation and functional synapse induction in developing hippocampus. *Nature.* 1996; 381:71–75. [PubMed: 8609991]
- Engert F, Bonhoeffer T. Dendritic spine changes associated with hippocampal long-term synaptic plasticity. *Nature.* 1999; 399:66–70. [PubMed: 10331391]
- Esparza MA, Bollati F, Garcia-Keller C, Virgolini MB, Lopez LM, Brusco A, Shen HW, Kalivas PW, Cancela LM. Stress-induced sensitization to cocaine: actin cytoskeleton remodeling within mesocorticolimbic nuclei. *Eur. J. Neurosci.* 2012; 36:3103–3117. [PubMed: 22882295]

- Fiala JC, Spacek J, Harris KM. Dendritic spine pathology: cause or consequence of neurological disorders? *Brain Res. Brain Res. Rev.* 2002; 39:29–54. [PubMed: 12086707]
- Fifkova E, Anderson CL. Stimulation-induced changes in dimensions of stalks of dendritic spines in the dentate molecular layer. *Exp. Neurol.* 1981; 74:621–627. [PubMed: 7297640]
- Fifkova E, Van Harreveld A. Long-lasting morphological changes in dendritic spines of dentate granular cells following stimulation of the entorhinal area. *J. Neurocytol.* 1977; 6:211–230. [PubMed: 856951]
- Fischer A, Sananbenesi F, Wang X, Dobbin M, Tsai LH. Recovery of learning and memory is associated with chromatin remodelling. *Nature.* 2007; 447:178–182. [PubMed: 17468743]
- Fischer M, Kaech S, Knutti D, Matus A. Rapid actin-based plasticity in dendritic spines. *Neuron.* 1998; 20:847–854. [PubMed: 9620690]
- Fischer M, Kaech S, Wagner U, Brinkhaus H, Matus A. Glutamate receptors regulate actin-based plasticity in dendritic spines. *Nat. Neurosci.* 2000; 3:887–894. [PubMed: 10966619]
- Fortin DA, Srivastava T, Soderling TR. Structural modulation of dendritic spines during synaptic plasticity. *Neuroscientist.* 2012; 18:326–341. [PubMed: 21670426]
- Frankfurt M, Luine V. The evolving role of dendritic spines and memory: Interaction(s) with estradiol. *Horm. Behav.* 2015; 74:28–36. [PubMed: 25993604]
- Fu M, Yu X, Lu J, Zuo Y. Repetitive motor learning induces coordinated formation of clustered dendritic spines in vivo. *Nature.* 2012; 483:92–95. [PubMed: 22343892]
- Gan WB, Grutzendler J, Wong RO, Lichtman JW. Ballistic delivery of dyes for structural and functional studies of the nervous system. *Cold Spring Harb Protoc.* 2009; 2009:pdb prot5202. [PubMed: 20147144]
- Gan WB, Grutzendler J, Wong WT, Wong RO, Lichtman JW. Multicolor “DiOlistic” labeling of the nervous system using lipophilic dye combinations. *Neuron.* 2000; 27:219–225. [PubMed: 10985343]
- Garcia-Keller C, Kupchik YM, Gipson CD, Brown RM, Spencer S, Bollati F, Esparza MA, Roberts-Wolfe DJ, Heinsbroek JA, Bobadilla AC, Cancela LM, Kalivas PW. Glutamatergic mechanisms of comorbidity between acute stress and cocaine self-administration. *Mol Psychiatry.* 2015
- Geinisman Y, Berry RW, Disterhoft JF, Power JM, Van der Zee EA. Associative learning elicits the formation of multiple-synapse boutons. *J. Neurosci.* 2001; 21:5568–5573. [PubMed: 11466428]
- Geinisman Y, Disterhoft JF, Gundersen HJ, McEchron MD, Persina IS, Power JM, van der Zee EA, West MJ. Remodeling of hippocampal synapses after hippocampus-dependent associative learning. *J. Comp. Neurol.* 2000; 417:49–59. [PubMed: 10660887]
- Gibb R, Kolb B. A method for vibratome sectioning of Golgi-Cox stained whole rat brain. *J. Neurosci. Methods.* 1998; 79:1–4. [PubMed: 9531453]
- Gipson CD, Beckmann JS, El-Maraghi S, Marusich JA, Bardo MT. Effect of environmental enrichment on escalation of cocaine self-administration in rats. *Psychopharmacology (Berl).* 2011; 214:557–566. [PubMed: 21057774]
- Gipson CD, Kupchik YM, Kalivas PW. Rapid, transient synaptic plasticity in addiction. *Neuropharmacology.* 2014; 76:276–286. [PubMed: 23639436]
- Gipson CD, Kupchik YM, Shen H, Reissner KJ, Thomas CA, Kalivas PW. Relapse induced by cues predicting cocaine depends on rapid, transient synaptic potentiation. *Neuron.* 2013a; 77:867–872. [PubMed: 23473317]
- Gipson CD, Reissner KJ, Kupchik YM, Smith AC, Stankeviciute N, Hensley-Simon ME, Kalivas PW. Reinstatement of nicotine seeking is mediated by glutamatergic plasticity. *Proc. Natl. Acad. Sci. U. S. A.* 2013b; 110:9124–9129. [PubMed: 23671067]
- Glaser EM, Van der Loos H. Analysis of thick brain sections by obverse-reverse computer microscopy: application of a new, high clarity Golgi-Nissl stain. *J. Neurosci. Methods.* 1981; 4:117–125. [PubMed: 6168870]
- Glausier JR, Lewis DA. Dendritic spine pathology in schizophrenia. *Neuroscience.* 2013; 251:90–107. [PubMed: 22546337]
- Golgi C. Sulla struttura della sostanza grigia della cervello. *Gazz Med Ital Lombardia.* 1873; 6:244–246.

- Gourley SL, Swanson AM, Koleske AJ. Corticosteroid-induced neural remodeling predicts behavioral vulnerability and resilience. *J Neurosci*. 2013; 33:3107–3112. [PubMed: 23407965]
- Grutzendler J, Gan WB. Two-photon imaging of synaptic plasticity and pathology in the living mouse brain. *NeuroRx*. 2006; 3:489–496. [PubMed: 17012063]
- Grutzendler J, Kasthuri N, Gan WB. Long-term dendritic spine stability in the adult cortex. *Nature*. 2002; 420:812–816. [PubMed: 12490949]
- Grutzendler J, Yang G, Pan F, Parkhurst CN, Gan WB. Transcranial two-photon imaging of the living mouse brain. *Cold Spring Harb Protoc*. 2011; 2011
- Halpain S. Actin and the agile spine: how and why do dendritic spines dance? *Trends Neurosci*. 2000; 23:141–146. [PubMed: 10717670]
- Harris KM. Structure, development, and plasticity of dendritic spines. *Curr. Opin. Neurobiol*. 1999; 9:343–348. [PubMed: 10395574]
- Harris, KM., Spacek, J. Dendrite structure. In: Stuart, G., Spruston, N., Hausser, M., editors. *Dendrites*. Oxford University Press; Oxford: 2016. p. 1–46.
- Hayashi-Takagi A, Yagishita S, Nakamura M, Shirai F, Wu YI, Loshbaugh AL, Kuhlman B, Hahn KM, Kasai H. Labelling and optical erasure of synaptic memory traces in the motor cortex. *Nature*. 2015; 525:333–338. [PubMed: 26352471]
- He CX, Portera-Cailliau C. The trouble with spines in fragile X syndrome: density, maturity and plasticity. *Neuroscience*. 2013; 251:120–128. [PubMed: 22522472]
- Heinsbroek, JA., Kupchik, Scofield, MD., Kalivas, PW., Gipson, CD. Rapid transient plasticity in dopamine D1 and D2 receptor expressing medium spiny neurons in the addiction circuitry mediate relapse to cocaine seeking. Presented at the 2014 Annual Meeting of the Society for Neuroscience; Washington, DC. 2014.
- Holmes A, Spanagel R, Krystal JH. Glutamatergic targets for new alcohol medications. *Psychopharmacology (Berl)*. 2013; 229:539–554. [PubMed: 23995381]
- Holtmaat A, de PV, Wilbrecht L, Trachtenberg JT, Svoboda K, Portera-Cailliau C. Imaging neocortical neurons through a chronic cranial window. *Cold Spring Harb Protoc*. 2012; 2012:694–701. [PubMed: 22661440]
- Holtmaat AJ, Trachtenberg JT, Wilbrecht L, Shepherd GM, Zhang X, Knott GW, Svoboda K. Transient and persistent dendritic spines in the neocortex in vivo. *Neuron*. 2005; 45:279–291. [PubMed: 15664179]
- Hotulainen P, Hoogenraad CC. Actin in dendritic spines: connecting dynamics to function. *J. Cell Biol*. 2010; 189:619–629. [PubMed: 20457765]
- Hyman SE, Malenka RC, Nestler EJ. Neural mechanisms of addiction: the role of reward-related learning and memory. *Annual review of neuroscience*. 2006; 29:565–598.
- Isshiki M, Okabe S. Evaluation of cranial window types for in vivo two-photon imaging of brain microstructures. *Microscopy (Oxf)*. 2014; 63:53–63. [PubMed: 24212360]
- Kang HJ, Voleti B, Hajszan T, Rajkowska G, Stockmeier CA, Licznarski P, Lepack A, Majik MS, Jeong LS, Banasr M, Son H, Duman RS. Decreased expression of synapse-related genes and loss of synapses in major depressive disorder. *Nat. Med*. 2012; 18:1413–1417. [PubMed: 22885997]
- Kasai H, Hayama T, Ishikawa M, Watanabe S, Yagishita S, Noguchi J. Learning rules and persistence of dendritic spines. *Eur J Neurosci*. 2010; 32:241–249. [PubMed: 20646057]
- Kasai H, Matsuzaki M, Noguchi J, Yasumatsu N, Nakahara H. Structure-stability-function relationships of dendritic spines. *Trends Neurosci*. 2003; 26:360–368. [PubMed: 12850432]
- Kenny PJ. Common cellular and molecular mechanisms in obesity and drug addiction. *Nature reviews. Neuroscience*. 2011a; 12:638–651. [PubMed: 22011680]
- Kenny PJ. Reward mechanisms in obesity: new insights and future directions. *Neuron*. 2011b; 69:664–679. [PubMed: 21338878]
- Khibnik LA, Beaumont M, Doyle M, Heshmati M, Slesinger PA, Nestler EJ, Russo SJ. Stress and Cocaine Trigger Divergent and Cell Type-Specific Regulation of Synaptic Transmission at Single Spines in Nucleus Accumbens. *Biol Psychiatry*. 2015
- Knobloch M, Mansuy IM. Dendritic spine loss and synaptic alterations in Alzheimer's disease. *Mol. Neurobiol*. 2008; 37:73–82. [PubMed: 18438727]

- Knott GW, Holtmaat A, Wilbrecht L, Welker E, Svoboda K. Spine growth precedes synapse formation in the adult neocortex in vivo. *Nat. Neurosci.* 2006; 9:1117–1124. [PubMed: 16892056]
- Koch C, Zador A. The function of dendritic spines: devices subserving biochemical rather than electrical compartmentalization. *J. Neurosci.* 1993; 13:413–422. [PubMed: 8426220]
- Koleske AJ. Molecular mechanisms of dendrite stability. *Nat Rev Neurosci.* 2013; 14:536–550. [PubMed: 23839597]
- Kopeck C, Malinow R. Neuroscience. Matters of size. *Science.* 2006; 314:1554–1555. [PubMed: 17158312]
- Kopeck CD, Li B, Wei W, Boehm J, Malinow R. Glutamate receptor exocytosis and spine enlargement during chemically induced long-term potentiation. *J. Neurosci.* 2006; 26:2000–2009. [PubMed: 16481433]
- Koyama Y, Tohyama M. A novel, Golgi-Cox-based fluorescent staining method for visualizing full-length processes in primary rat neurons. *Neurochem. Int.* 2013; 63:35–41. [PubMed: 23619398]
- Krzyszczak M, Fulling C, Jabinet L, Armida J, Gebara E, Casse F, Habbas S, Volterra A, Hornung JP, Toni N. Synaptic Adhesion Molecules Regulate the Integration of New Granule Neurons in the Postnatal Mouse Hippocampus and their Impact on Spatial Memory. *Cerebral cortex.* 2016
- Kwon HB, Sabatini BL. Glutamate induces de novo growth of functional spines in developing cortex. *Nature.* 2011; 474:100–104. [PubMed: 21552280]
- Lamprecht R. The actin cytoskeleton in memory formation. *Prog. Neurobiol.* 2014; 117C:1–19.
- Lamprecht R, Farb CR, Rodrigues SM, LeDoux JE. Fear conditioning drives profilin into amygdala dendritic spines. *Nat Neurosci.* 2006; 9:481–483. [PubMed: 16547510]
- Lee BR, Dong Y. Cocaine-induced metaplasticity in the nucleus accumbens: silent synapse and beyond. *Neuropharmacology.* 2011; 61:1060–1069. [PubMed: 21232547]
- Lee BR, Ma YY, Huang YH, Wang X, Otaka M, Ishikawa M, Neumann PA, Graziane NM, Brown TE, Suska A, Guo C, Lobo MK, Sesack SR, Wolf ME, Nestler EJ, Shaham Y, Schluter OM, Dong Y. Maturation of silent synapses in amygdala-accumbens projection contributes to incubation of cocaine craving. *Nat. Neurosci.* 2013; 16:1644–1651. [PubMed: 24077564]
- Lee KF, Soares C, Beique JC. Examining form and function of dendritic spines. *Neural Plast.* 2012; 2012:704103. [PubMed: 22577585]
- Leuner B, Falduto J, Shors TJ. Associative memory formation increases the observation of dendritic spines in the hippocampus. *J Neurosci.* 2003; 23:659–665. [PubMed: 12533625]
- Leuner B, Shors TJ. Stress, anxiety, and dendritic spines: what are the connections? *Neuroscience.* 2013; 251:108–119. [PubMed: 22522470]
- Licznerski P, Duman RS. Remodeling of axo-spinous synapses in the pathophysiology and treatment of depression. *Neuroscience.* 2013; 251:33–50. [PubMed: 23036622]
- Long VA, Fanselow MS. Stress-enhanced fear learning in rats is resistant to the effects of immediate massed extinction. *Stress.* 2012; 15:627–636. [PubMed: 22176467]
- Lopes S, Teplytska L, Vaz-Silva J, Dioli C, Trindade R, Morais M, Webhofer C, Maccarrone G, Almeida OF, Turck CW, Sousa N, Sotiropoulos I, Filiou MD. Tau Deletion Prevents Stress-Induced Dendritic Atrophy in Prefrontal Cortex: Role of Synaptic Mitochondria. *Cereb Cortex.* 2016
- Ma YY, Wang X, Huang Y, Marie H, Nestler EJ, Schluter OM, Dong Y. Re-silencing of silent synapses unmasks anti-relapse effects of environmental enrichment. *Proc Natl Acad Sci U S A.* 2016
- Magarinos AM, McEwen BS. Stress-induced atrophy of apical dendrites of hippocampal CA3c neurons: involvement of glucocorticoid secretion and excitatory amino acid receptors. *Neuroscience.* 1995; 69:89–98. [PubMed: 8637636]
- Majewska AK, Newton JR, Sur M. Remodeling of synaptic structure in sensory cortical areas in vivo. *J. Neurosci.* 2006; 26:3021–3029. [PubMed: 16540580]
- Maletic-Savatic M, Malinow R, Svoboda K. Rapid dendritic morphogenesis in CA1 hippocampal dendrites induced by synaptic activity. *Science.* 1999; 283:1923–1927. [PubMed: 10082466]
- Mancuso JJ, Chen Y, Li X, Xue Z, Wong ST. Methods of dendritic spine detection: from Golgi to high-resolution optical imaging. *Neuroscience.* 2013; 251:129–140. [PubMed: 22522468]

- Marchant NJ, Kaganovsky K, Shaham Y, Bossert JM. Role of corticostriatal circuits in context-induced reinstatement of drug seeking. *Brain Res.* 2015; 1628:219–232. [PubMed: 25199590]
- Martin KC, Zukin RS. RNA trafficking and local protein synthesis in dendrites: an overview. *J. Neurosci.* 2006; 26:7131–7134. [PubMed: 16822966]
- Matsuo N, Reijmers L, Mayford M. Spine-type-specific recruitment of newly synthesized AMPA receptors with learning. *Science.* 2008; 319:1104–1107. [PubMed: 18292343]
- Matsuzaki M, Honkura N, Ellis-Davies GC, Kasai H. Structural basis of long-term potentiation in single dendritic spines. *Nature.* 2004; 429:761–766. [PubMed: 15190253]
- Mattison HA, Popovkina D, Kao JP, Thompson SM. The role of glutamate in the morphological and physiological development of dendritic spines. *Eur. J. Neurosci.* 2014; 39:1761–1770. [PubMed: 24661419]
- McEwen BS, Conrad CD, Kuroda Y, Frankfurt M, Magarinos AM, McKittrick C. Prevention of stress-induced morphological and cognitive consequences. *Eur Neuropsychopharmacol.* 1997; 7(Suppl 3):S323–328. [PubMed: 9405958]
- McEwen BS, Eiland L, Hunter RG, Miller MM. Stress and anxiety: structural plasticity and epigenetic regulation as a consequence of stress. *Neuropharmacology.* 2012; 62:3–12. [PubMed: 21807003]
- McEwen BS, Nasca C, Gray JD. Stress Effects on Neuronal Structure: Hippocampus, Amygdala, and Prefrontal Cortex. *Neuropsychopharmacology.* 2016; 41:3–23. [PubMed: 26076834]
- McFarland K, Lapish CC, Kalivas PW. Prefrontal glutamate release into the core of the nucleus accumbens mediates cocaine-induced reinstatement of drug-seeking behavior. *J. Neurosci.* 2003; 23:3531–3537. [PubMed: 12716962]
- McLaughlin KJ, Baran SE, Conrad CD. Chronic stress- and sex-specific neuromorphological and functional changes in limbic structures. *Mol Neurobiol.* 2009; 40:166–182. [PubMed: 19653136]
- McLaughlin KJ, Baran SE, Wright RL, Conrad CD. Chronic stress enhances spatial memory in ovariectomized female rats despite CA3 dendritic retraction: possible involvement of CA1 neurons. *Neuroscience.* 2005; 135:1045–1054. [PubMed: 16165283]
- Micheva KD, Busse B, Weiler NC, O'Rourke N, Smith SJ. Single-synapse analysis of a diverse synapse population: proteomic imaging methods and markers. *Neuron.* 2010; 68:639–653. [PubMed: 21092855]
- Morrison JH, Baxter MG. The ageing cortical synapse: hallmarks and implications for cognitive decline. *Nat Rev Neurosci.* 2012; 13:240–250. [PubMed: 22395804]
- Morrison JH, Hof PR. Life and death of neurons in the aging brain. *Science.* 1997; 278:412–419. [PubMed: 9334292]
- Mulholland PJ, Chandler LJ. The thorny side of addiction: adaptive plasticity and dendritic spines. *ScientificWorldJournal.* 2007; 7:9–21.
- Mulholland PJ, Chandler LJ, Kalivas PW. Signals from the Fourth Dimension Regulate Drug Relapse. *Trends in neurosciences.* 2016; 39:472–485. [PubMed: 27173064]
- Murakoshi H, Yasuda R. Postsynaptic signaling during plasticity of dendritic spines. *Trends Neurosci.* 2012; 35:135–143. [PubMed: 2222350]
- Murayama M, Larkum M. Fiber-optic calcium monitoring of dendritic activity in vivo. *Cold Spring Harb Protoc.* 2012; 2012:218–225. [PubMed: 22301652]
- Nagerl UV, Eberhorn N, Cambridge SB, Bonhoeffer T. Bidirectional activity-dependent morphological plasticity in hippocampal neurons. *Neuron.* 2004; 44:759–767. [PubMed: 15572108]
- Nagerl UV, Kostinger G, Anderson JC, Martin KA, Bonhoeffer T. Protracted synaptogenesis after activity-dependent spinogenesis in hippocampal neurons. *J. Neurosci.* 2007; 27:8149–8156. [PubMed: 17652605]
- Neisewander JL, O'Dell LE, Tran-Nguyen LT, Castaneda E, Fuchs RA. Dopamine overflow in the nucleus accumbens during extinction and reinstatement of cocaine self-administration behavior. *Neuropsychopharmacology.* 1996; 15:506–514. [PubMed: 8914124]
- Nestler EJ. Neurobiology. Total recall-the memory of addiction. *Science.* 2001; 292:2266–2267. [PubMed: 11423644]
- Nyberg F. Structural plasticity of the brain to psychostimulant use. *Neuropharmacology.* 2014; 87C: 115–124.

- O'Brien JA, Lummis SC. Diolistic labeling of neuronal cultures and intact tissue using a hand-held gene gun. *Nat Protoc.* 2006; 1:1517–1521.
- Oe Y, Tominaga-Yoshino K, Hasegawa S, Ogura A. Dendritic spine dynamics in synaptogenesis after repeated LTP inductions: dependence on pre-existing spine density. *Sci Rep.* 2013; 3:1957. [PubMed: 23739837]
- Oh WC, Hill TC, Zito K. Synapse-specific and size-dependent mechanisms of spine structural plasticity accompanying synaptic weakening. *Proc. Natl. Acad. Sci. U. S. A.* 2013; 110:E305–312. [PubMed: 23269840]
- Orlowski D, Bjarkam CR. A simple reproducible and time saving method of semi-automatic dendrite spine density estimation compared to manual spine counting. *J. Neurosci. Methods.* 2012; 208:128–133. [PubMed: 22595026]
- Pan F, Aldridge GM, Greenough WT, Gan WB. Dendritic spine instability and insensitivity to modulation by sensory experience in a mouse model of fragile X syndrome. *Proc Natl Acad Sci U S A.* 2010; 107:17768–17773. [PubMed: 20861447]
- Parekh R, Ascoli GA. Neuronal morphology goes digital: a research hub for cellular and system neuroscience. *Neuron.* 2013; 77:1017–1038. [PubMed: 23522039]
- Paulin JJ, Haslehurst P, Fellows AD, Liu W, Jackson JD, Joel Z, Cummings DM, Edwards FA. Large and Small Dendritic Spines Serve Different Interacting Functions in Hippocampal Synaptic Plasticity and Homeostasis. *Neural Plast.* 2016; 2016:6170509. [PubMed: 26881123]
- Pereira AC, Lambert HK, Grossman YS, Dumitriu D, Waldman R, Jannetty SK, Calakos K, Janssen WG, McEwen BS, Morrison JH. Glutamatergic regulation prevents hippocampal-dependent age-related cognitive decline through dendritic spine clustering. *Proc Natl Acad Sci U S A.* 2014; 111:18733–18738. [PubMed: 25512503]
- Pillai AG, de Jong D, Kanatsou S, Krugers H, Knapman A, Heinzmann JM, Holsboer F, Landgraf R, Joels M, Touma C. Dendritic morphology of hippocampal and amygdalar neurons in adolescent mice is resilient to genetic differences in stress reactivity. *PLoS One.* 2012; 7:e38971. [PubMed: 22701737]
- Pinto L, Mateus-Pinheiro A, Morais M, Bessa JM, Sousa N. Immuno-Golgi as a tool for analyzing neuronal 3D-dendritic structure in phenotypically characterized neurons. *PLoS One.* 2012; 7:e33114. [PubMed: 22427964]
- Platt JE, Stone EA. Chronic restraint stress elicits a positive antidepressant response on the forced swim test. *Eur J Pharmacol.* 1982; 82:179–181. [PubMed: 6889973]
- Qiao H, Li MX, Xu C, Chen HB, An SC, Ma XM. Dendritic Spines in Depression: What We Learned from Animal Models. *Neural Plast.* 2016; 2016:8056370. [PubMed: 26881133]
- Quirk GJ, Armony JL, LeDoux JE. Fear conditioning enhances different temporal components of tone-evoked spike trains in auditory cortex and lateral amygdala. *Neuron.* 1997; 19:613–624. [PubMed: 9331352]
- Radley JJ, Johnson LR, Janssen WG, Martino J, Lamprecht R, Hof PR, LeDoux JE, Morrison JH. Associative Pavlovian conditioning leads to an increase in spinophilin-immunoreactive dendritic spines in the lateral amygdala. *Eur J Neurosci.* 2006a; 24:876–884. [PubMed: 16930415]
- Radley JJ, Rocher AB, Miller M, Janssen WG, Liston C, Hof PR, McEwen BS, Morrison JH. Repeated stress induces dendritic spine loss in the rat medial prefrontal cortex. *Cereb. Cortex.* 2006b; 16:313–320. [PubMed: 15901656]
- Radley JJ, Rocher AB, Rodriguez A, Ehlenberger DB, Dammann M, McEwen BS, Morrison JH, Wearne SL, Hof PR. Repeated stress alters dendritic spine morphology in the rat medial prefrontal cortex. *J Comp Neurol.* 2008; 507:1141–1150. [PubMed: 18157834]
- Rall, W. Dendritic spines and synaptic potency. Cambridge University Press; Cambridge: 1978.
- Ramón y Cajal S. Estructura de los centros nerviosos de las aves. *Rev Trim Histol Norm Pat.* 1888; 1:1–10.
- Ramón y Cajal, S. La Textura del Sistema Nerviosa del Hombre y los Vertebrados. Moya; Madrid: 1899.
- Richards DA, Mateos JM, Hugel S, de Paola V, Caroni P, Gahwiler BH, McKinney RA. Glutamate induces the rapid formation of spine head protrusions in hippocampal slice cultures. *Proc. Natl. Acad. Sci. U. S. A.* 2005; 102:6166–6171. [PubMed: 15831587]

- Robinson TE, Kolb B. Alterations in the morphology of dendrites and dendritic spines in the nucleus accumbens and prefrontal cortex following repeated treatment with amphetamine or cocaine. *Eur J Neurosci.* 1999a; 11:1598–1604. [PubMed: 10215912]
- Robinson TE, Kolb B. Morphine alters the structure of neurons in the nucleus accumbens and neocortex of rats. *Synapse.* 1999b; 33:160–162. [PubMed: 10400894]
- Rojas JJ, Deniz BF, Miguel PM, Diaz R, Hermel Edo E, Achaval M, Netto CA, Pereira LO. Effects of daily environmental enrichment on behavior and dendritic spine density in hippocampus following neonatal hypoxia-ischemia in the rat. *Exp Neurol.* 2013; 241:25–33. [PubMed: 23219882]
- Romanski LM, Clugnet MC, Bordi F, LeDoux JE. Somatosensory and auditory convergence in the lateral nucleus of the amygdala. *Behav Neurosci.* 1993; 107:444–450. [PubMed: 8329134]
- Romeo RD, Bellani R, Karatsoreos IN, Chhua N, Vernov M, Conrad CD, McEwen BS. Stress history and pubertal development interact to shape hypothalamic-pituitary-adrenal axis plasticity. *Endocrinology.* 2006; 147:1664–1674. [PubMed: 16410296]
- Romeo RD, McEwen BS. Stress and the adolescent brain. *Ann N Y Acad Sci.* 2006; 1094:202–214. [PubMed: 17347352]
- Russo SJ, Dietz DM, Dumitriu D, Morrison JH, Malenka RC, Nestler EJ. The addicted synapse: mechanisms of synaptic and structural plasticity in nucleus accumbens. *Trends Neurosci.* 2010; 33:267–276. [PubMed: 20207024]
- Ryan TJ, Roy DS, Pignatelli M, Arons A, Tonegawa S. Engram cells retain memory under retrograde amnesia. *Science.* 2015; 348:1007–1013. [PubMed: 26023136]
- Sala C, Segal M. Dendritic spines: the locus of structural and functional plasticity. *Physiol. Rev.* 2014; 94:141–188. [PubMed: 24382885]
- Schafe GE, Nader K, Blair HT, LeDoux JE. Memory consolidation of Pavlovian fear conditioning: a cellular and molecular perspective. *Trends Neurosci.* 2001; 24:540–546. [PubMed: 11506888]
- Schenk S, Partridge B. Influence of a conditioned light stimulus on cocaine self-administration in rats. *Psychopharmacology.* 2001; 154:390–396. [PubMed: 11349392]
- Scofield MD, Heinsbroek JA, Gipson CD, Kupchik YM, Spencer S, Smith AC, Roberts-Wolfe D, Kalivas PW. The Nucleus Accumbens: Mechanisms of Addiction across Drug Classes Reflect the Importance of Glutamate Homeostasis. *Pharmacological reviews.* 2016; 68:816–871. [PubMed: 27363441]
- Seabold GK, Daunais JB, Rau A, Grant KA, Alvarez VA. DiOLISTIC labeling of neurons from rodent and non-human primate brain slices. *J Vis Exp.* 2010
- See RE. Neural substrates of conditioned-cued relapse to drug-seeking behavior. *Pharmacology, biochemistry, and behavior.* 2002; 71:517–529.
- Segal M. Dendritic spines and long-term plasticity. *Nat Rev Neurosci.* 2005; 6:277–284. [PubMed: 15803159]
- Segal M. Dendritic spines: Morphological building blocks of memory. *Neurobiology of learning and memory.* 2016
- Sesack SR, Carr DB, Omelchenko N, Pinto A. Anatomical substrates for glutamate-dopamine interactions: evidence for specificity of connections and extrasynaptic actions. *Ann. N. Y. Acad. Sci.* 2003; 1003:36–52. [PubMed: 14684434]
- Shaham Y, Shalev U, Lu L, De Wit H, Stewart J. The reinstatement model of drug relapse: history, methodology and major findings. *Psychopharmacology.* 2003; 168:3–20. [PubMed: 12402102]
- Shen H, Moussawi K, Zhou W, Toda S, Kalivas PW. Heroin relapse requires long-term potentiation-like plasticity mediated by NMDA2b-containing receptors. *Proc Natl Acad Sci U S A.* 2011; 108:19407–19412. [PubMed: 22084102]
- Shen H, Sesack SR, Toda S, Kalivas PW. Automated quantification of dendritic spine density and spine head diameter in medium spiny neurons of the nucleus accumbens. *Brain Struct Funct.* 2008; 213:149–157. [PubMed: 18535839]
- Shen HW, Gipson CD, Huits M, Kalivas PW. Prelimbic cortex and ventral tegmental area modulate synaptic plasticity differentially in nucleus accumbens during cocaine-reinstated drug seeking. *Neuropsychopharmacology.* 2014; 39:1169–1177. [PubMed: 24232172]

- Shen HW, Toda S, Moussawi K, Bouknight A, Zahm DS, Kalivas PW. Altered dendritic spine plasticity in cocaine-withdrawn rats. *J. Neurosci.* 2009; 29:2876–2884. [PubMed: 19261883]
- Shepherd GM. The dendritic spine: a multifunctional integrative unit. *J. Neurophysiol.* 1996; 75:2197–2210. [PubMed: 8793734]
- Shimamoto A, Debold JF, Holly EN, Miczek KA. Blunted accumbal dopamine response to cocaine following chronic social stress in female rats: exploring a link between depression and drug abuse. *Psychopharmacology (Berl)*. 2011; 218:271–279. [PubMed: 21638221]
- Shirao T, Gonzalez-Billault C. Actin filaments and microtubules in dendritic spines. *J. Neurochem.* 2013; 126:155–164. [PubMed: 23692384]
- Singer BF, Bubula N, Li D, Przybycien-Szymanska MM, Bindokas VP, Vezina P. Drug-Paired Contextual Stimuli Increase Dendritic Spine Dynamics in Select Nucleus Accumbens Neurons. *Neuropsychopharmacology*. 2016
- Sotiropoulos I, Sousa N. Tau as the converging protein between chronic stress and alzheimer's disease synaptic pathology. *Neurodegener Dis.* 2016; 16:22–25. [PubMed: 26551025]
- Spiga S, Acquas E, Puddu MC, Mulas G, Lintas A, Diana M. Simultaneous Golgi-Cox and immunofluorescence using confocal microscopy. *Brain Struct Funct.* 2011; 216:171–182. [PubMed: 21461741]
- Spiga S, Lintas A, Migliore M, Diana M. Altered architecture and functional consequences of the mesolimbic dopamine system in cannabis dependence. *Addict Biol.* 2010; 15:266–276. [PubMed: 20477755]
- Spiga S, Mulas G, Piras F, Diana M. The “addicted” spine. *Front Neuroanat.* 2014a; 8:110. [PubMed: 25324733]
- Spiga S, Talani G, Mulas G, Licheri V, Fois GR, Muggironi G, Masala N, Cannizzaro C, Biggio G, Sanna E, Diana M. Hampered long-term depression and thin spine loss in the nucleus accumbens of ethanol-dependent rats. *Proc. Natl. Acad. Sci. U. S. A.* 2014b; 111:E3745–3754. [PubMed: 25122682]
- Staffend NA, Meisel RL. DiOlistic labeling in fixed brain slices: phenotype, morphology, and dendritic spines. *Curr Prot Neurosci.* 2011a; Chapter 2(Unit 2):13.
- Staffend NA, Meisel RL. DiOlistic labeling of neurons in tissue slices: a qualitative and quantitative analysis of methodological variations. *Front Neuroanat.* 2011b; 5:14. [PubMed: 21427781]
- Stefanik MT, Kupchik YM, Kalivas PW. Optogenetic inhibition of cortical afferents in the nucleus accumbens simultaneously prevents cue-induced transient synaptic potentiation and cocaine-seeking behavior. *Brain Struct Funct.* 2015
- Stefanik MT, Kupchik YM, Kalivas PW. Optogenetic inhibition of cortical afferents in the nucleus accumbens simultaneously prevents cue-induced transient synaptic potentiation and cocaine-seeking behavior. *Brain Struct Funct.* 2016; 221:1681–1689. [PubMed: 25663648]
- Stewart MG, Davies HA, Sandi C, Kraev IV, Rogachevsky VV, Peddie CJ, Rodriguez JJ, Cordero MI, Donohue HS, Gabbott PL, Popov VI. Stress suppresses and learning induces plasticity in CA3 of rat hippocampus: a three-dimensional ultrastructural study of thorny excrescences and their postsynaptic densities. *Neuroscience.* 2005; 131:43–54. [PubMed: 15680690]
- Swanger SA, Yao X, Gross C, Bassell GJ. Automated 4D analysis of dendritic spine morphology: applications to stimulus-induced spine remodeling and pharmacological rescue in a disease model. *Mol Brain.* 2011; 4:38. [PubMed: 21982080]
- Tada T, Sheng M. Molecular mechanisms of dendritic spine morphogenesis. *Curr. Opin. Neurobiol.* 2006; 16:95–101. [PubMed: 16361095]
- Thiel KJ, Painter MR, Pentkowski NS, Mitroi D, Crawford CA, Neisewander JL. Environmental enrichment counters cocaine abstinence-induced stress and brain reactivity to cocaine cues but fails to prevent the incubation effect. *Addict Biol.* 2012; 17:365–377. [PubMed: 21812872]
- Thiel KJ, Sanabria F, Pentkowski NS, Neisewander JL. Anti-craving effects of environmental enrichment. *Int J Neuropsychopharmacol.* 2009; 12:1151–1156. [PubMed: 19691875]
- Toda S, Shen H, Kalivas PW. Inhibition of actin polymerization prevents cocaine-induced changes in spine morphology in the nucleus accumbens. *Neurotox. Res.* 2010; 18:410–415. [PubMed: 20401643]

- Toda S, Shen HW, Peters J, Cagle S, Kalivas PW. Cocaine increases actin cycling: effects in the reinstatement model of drug seeking. *J. Neurosci.* 2006; 26:1579–1587. [PubMed: 16452681]
- Toni N, Buchs PA, Nikonenko I, Bron CR, Muller D. LTP promotes formation of multiple spine synapses between a single axon terminal and a dendrite. *Nature.* 1999; 402:421–425. [PubMed: 10586883]
- Tonnesen J, Katona G, Rozsa B, Nagerl UV. Spine neck plasticity regulates compartmentalization of synapses. *Nat. Neurosci.* 2014; 17:678–685. [PubMed: 24657968]
- Trachtenberg JT, Chen BE, Knott GW, Feng G, Sanes JR, Welker E, Svoboda K. Long-term in vivo imaging of experience-dependent synaptic plasticity in adult cortex. *Nature.* 2002; 420:788–794. [PubMed: 12490942]
- Tredici G, Di FA, Miani A Jr, Pizzini G. Real complete three-dimensional reconstruction of Golgi-impregnated neurons by means of a confocal laser scanning microscope. *Neuroimage.* 1993; 1:87–93. [PubMed: 9343560]
- Turner CA, Lewis MH, King MA. Environmental enrichment: effects on stereotyped behavior and dendritic morphology. *Dev Psychobiol.* 2003; 43:20–27. [PubMed: 12794775]
- Uchoa ET, Aguilera G, Herman JP, Fiedler JL, Deak T, de Sousa MB. Novel aspects of glucocorticoid actions. *J. Neuroendocrinol.* 2014; 26:557–572. [PubMed: 24724595]
- Ultanir SK, Kim JE, Hall BJ, Deerinck T, Ellisman M, Ghosh A. Regulation of spine morphology and spine density by NMDA receptor signaling in vivo. *Proc. Natl. Acad. Sci. U. S. A.* 2007; 104:19553–19558. [PubMed: 18048342]
- Uys JD, McGuier NS, Gass JT, Griffin WC 3rd, Ball LE, Mulholland PJ. Chronic intermittent ethanol exposure and withdrawal leads to adaptations in nucleus accumbens core postsynaptic density proteome and dendritic spines. *Addict Biol.* 2016; 21:560–574. [PubMed: 25787124]
- van der Zee EA. Synapses, spines and kinases in mammalian learning and memory, and the impact of aging. *Neurosci. Biobehav. Rev.* 2015; 50:77–85. [PubMed: 24998408]
- Van Harrevelde A, Fifkova E. Swelling of dendritic spines in the fascia dentata after stimulation of the perforant fibers as a mechanism of post-tetanic potentiation. *Exp. Neurol.* 1975; 49:736–749. [PubMed: 173566]
- van Spronsen M, Hoogenraad CC. Synapse pathology in psychiatric and neurologic disease. *Curr Neurol Neurosci Rep.* 2010; 10:207–214. [PubMed: 20425036]
- von Bohlen und Halbach O, Zacher C, Gass P, Unsicker K. Age-related alterations in hippocampal spines and deficiencies in spatial memory in mice. *J Neurosci Res.* 2006; 83:525–531. [PubMed: 16447268]
- Wallace M, Frankfurt M, Arellanos A, Inagaki T, Luine V. Impaired recognition memory and decreased prefrontal cortex spine density in aged female rats. *Ann N Y Acad Sci.* 2007; 1097:54–57. [PubMed: 17413010]
- Wang XB, Yang Y, Zhou Q. Independent expression of synaptic and morphological plasticity associated with long-term depression. *J. Neurosci.* 2007; 27:12419–12429. [PubMed: 17989307]
- Wang XB, Zhou Q. Spine remodeling and synaptic modification. *Mol. Neurobiol.* 2010; 41:29–41. [PubMed: 20049655]
- Watson DJ, Ostroff L, Cao G, Parker PH, Smith H, Harris KM. LTP enhances synaptogenesis in the developing hippocampus. *Hippocampus.* 2016; 26:560–576. [PubMed: 26418237]
- Weinstock M. Sex-dependent changes induced by prenatal stress in cortical and hippocampal morphology and behaviour in rats: an update. *Stress.* 2011; 14:604–613. [PubMed: 21790452]
- Weiss F, Markou A, Lorang MT, Koob GF. Basal extracellular dopamine levels in the nucleus accumbens are decreased during cocaine withdrawal after unlimited-access self-administration. *Brain Res.* 1992; 593:314–318. [PubMed: 1450939]
- Yagishita S, Hayashi-Takagi A, Ellis-Davies GC, Urakubo H, Ishii S, Kasai H. A critical time window for dopamine actions on the structural plasticity of dendritic spines. *Science.* 2014; 345:1616–1620. [PubMed: 25258080]
- Yang G, Pan F, Gan WB. Stably maintained dendritic spines are associated with lifelong memories. *Nature.* 2009; 462:920–924. [PubMed: 19946265]
- Yasuda, R. Molecular signaling during plasticity of dendritic spines. In: Stuart, G. Spruston, N., Hausser, M., editors. *Dendrites*. Oxford University Press; Oxford: 2016. p. 581–602.

- Yasumatsu N, Matsuzaki M, Miyazaki T, Noguchi J, Kasai H. Principles of long-term dynamics of dendritic spines. *J. Neurosci.* 2008; 28:13592–13608. [PubMed: 19074033]
- Young EJ, Aceti M, Griggs EM, Fuchs RA, Zigmond Z, Rumbaugh G, Miller CA. Selective, retrieval-independent disruption of methamphetamine-associated memory by actin depolymerization. *Biol. Psychiatry.* 2014; 75:96–104. [PubMed: 24012327]
- Young EJ, Blouin AM, Briggs SB, Sullivan SE, Lin L, Cameron MD, Rumbaugh G, Miller CA. Nonmuscle myosin IIB as a therapeutic target for the prevention of relapse to methamphetamine use. *Mol. Psychiatry.* 2016; 21:615–623. [PubMed: 26239291]
- Yuste, R. *Dendritic Spines*. MIT Press; Cambridge, MA: 2010.
- Yuste R. The discovery of dendritic spines by Cajal. *Front Neuroanat.* 2015; 9:18. [PubMed: 25954162]
- Yuste R, Bonhoeffer T. Morphological changes in dendritic spines associated with long-term synaptic plasticity. *Annu. Rev. Neurosci.* 2001; 24:1071–1089. [PubMed: 11520928]
- Yuste R, Majewska A. On the function of dendritic spines. *Neuroscientist.* 2001; 7:387–395. [PubMed: 11597098]
- Zhou Q, Homma KJ, Poo MM. Shrinkage of dendritic spines associated with long-term depression of hippocampal synapses. *Neuron.* 2004; 44:749–757. [PubMed: 15572107]
- Zuo Y, Lin A, Chang P, Gan WB. Development of long-term dendritic spine stability in diverse regions of cerebral cortex. *Neuron.* 2005; 46:181–189. [PubMed: 15848798]

Figure 1.

(A) Illustration of a dendritic spine. Spines typically have neck and head compartments, whose structure is largely built on an actin cytoskeleton. G-actin polymerizes into F-actin to build the primary cytoskeletal structure, and thus remodels the spine during synaptic plasticity. This process continually reshapes brain circuitry and synaptic connections. At the PSD, membrane-bound proteins are continually inserted or retracted from the membrane, including ionotropic and metabotropic receptors. (B) Dendritic spines are often categorized into different subtypes, some of which are thought to be immature and filopodium-like, and others are thought to be mature with increased head diameters relative to that of the neck, as well as bifurcated and/or multi-branched morphologies. See text for a discussion of the limitations of spine categorization based on overall morphology. Abbreviations: PSD = post-synaptic density; G-actin = globular (monomeric) actin, F-actin = filamentous actin.

Figure 2.

(A) Image of a Golgi-Cox stained striatal medium spiny neuron and a higher magnification of the selected area showing dendritic spines. (B) A high resolution three-dimensional image of a Lucifer-Yellow filled neuron in the NA. The magnification of the boxed region is 63 $\times$, and a 3-D segment with clearly defined spines is reconstructed and analyzed. (C) Example of a DiI-filled MSN in the NA. The boxed region is magnified to 63 $\times$ and analyzed using Imaris software (Bitplane).