Published in final edited form as:

J Org Chem. 2016 May 20; 81(10): 4149–4157. doi:10.1021/acs.joc.6b00459.

Lithium Enolates Derived from Pyroglutaminol: Aggregation, Solvation, and Atropisomerism

Michael J. Houghton[¶], Naomi A. Biok[¶], Christopher J. Huck[¶], Russell F. Algera[¶], Ivan Keresztes[¶], Stephen W. Wright[‡], and David B. Collum^{*,¶}

[¶]Department of Chemistry and Chemical Biology, Baker Laboratory, Cornell University, Ithaca, New York 14853–1301

[‡]Worldwide Medicinal Chemistry, Pfizer Global Research and Development, 445 Eastern Point Road, Groton, CT 06340

Abstract

Lithium enolates derived from protected pyroglutaminols were characterized by using ⁶Li, ¹³C, and ¹⁹F NMR spectroscopies in conjunction with the method of continuous variation. A mixture of tetrasolvated dimers and tetrasolvated tetramers in proportions depend on the steric demands of the hemiaminal protecting group, tetrahydrofuran concentration, and the presence or absence of an α -fluoro moiety. The high steric demands of the substituted bicyclo[3.3.0] ring system promote dimers to an unusual extent and allow solvents and atropisomers in cubic tetramers to be observed in the slow-exchange limit. Pyridine used as a ⁶Li chemical shift reagent proved useful in assigning solvation numbers.

Graphical abstract

Introduction

As part of a program to develop anti-inflammatory agents, chemists at Pfizer examined functionalizations of readily available protected pyroglutaminols (eq 1).^{1,2} These stereochemically rich, nitrogen-based synthons contain critical features that allow for stereocontrolled functionalizations of both the α and the β carbons. Reactions that exploit the chirality of these compounds include alkylations,^{3,4} aldol and aza-aldol additions,^{5,6} and halogenations^{7,8} of enolates as well as 1,4-additions,^{9,10} cycloadditions,¹¹ dihydroxylations,¹² and cyclopropanations^{13,14} of the analogous α , β -unsaturated lactam.

*David B. Collum, dbc6@cornell.edu.

The authors declare no competing financial interests.

Supporting Information: Spectroscopic, kinetic, and computational data and authors for reference 21. This material is available free of charge via the Internet at <http://pubs.acs.org>.

The stereocontrol is often unremarkable; however, compared with the more popular benzylidene analog **1a**, the acetonide-protected variant **1b** enhances the preference for functionalization from the convex face.^{1,8,15} Given the potential importance of these synthons, surprisingly few variants of the hemiaminal linkage have been reported.¹

(1)

The Cornell contingent was drawn to these protected pyroglutaminols to expand studies of how the aggregation and solvation of lithium enolates influence their reactivities and selectivities.¹⁶ The appeal of the pyroglutaminol-derived skeleton is that the hemiaminal linkage can be varied without perturbing the core γ -lactam moiety. Moreover, the enolate geometries are necessarily *E*, and the chirality provides a means to correlate mechanism and reactivity with stereochemistry.^{17,18}

We describe herein the first of a two-part investigation of lithium enolates **2a–2k** and **3–5** (Chart 1) in tetrahydrofuran (THF)/toluene solutions. The deceptively high steric demands stemming from the substituted bicyclo[3.3.0] ring system promote tetramer-to-dimer deaggregation and offer insights into solvation and unexpected atropisomerism within the tetramers. In ongoing studies we are examining the mechanistic and stereochemical consequences of aggregation and solvation.

Results

Lithium enolates **2–5** were generated from their corresponding lactams¹ by using recrystallized lithium diisopropylamide (LDA, [⁶Li]LDA, or [⁶Li, ¹⁵N]LDA).¹⁹ Structures were determined with a combination of ⁶Li NMR spectroscopy, the method of continuous variation (MCV),²⁰ and density function theory (DFT) calculations at the B3LYP/6–31G(d) level with single-point calculations at the MP2 level of theory.²¹

As a brief reminder, using MCV to characterize aggregates involves the mixing two similar enolates, **A_n** and **B_n**, to generate an ensemble of homo- and heteroaggregates (eq 2).^{16,20} The number and symmetries of the heteroaggregates reveal the aggregation state, *n*, which is confirmed by plotting the relative concentrations versus measured²² mole fraction (*X_A* or *X_B*) to generate a Job plot (exemplified by Figures 2 and 4).²³

Enolate–LDA mixed aggregates in THF

We first addressed the role of LDA mixed aggregates observed when excess LDA is present. By example, generating enolate **3** using excess [^6Li , ^{15}N]LDA revealed mixed dimer **6** and confirmed that the structures assigned as enolate homoaggregates (**7** and **8**, below) contained no LDA fragments. The mixed dimer **6** displayed the expected ^6Li doublet and ^{15}N quintet ($J_{\text{Li-N}} = 5 \text{ Hz}$).²⁴ We observed an unusually large THF-concentration-dependent ^6Li chemical shift, which suggested that **6** might attain a solvation number >2 . DFT calculations showed some preference for trisolvation over disolvation (eq 3), which is logical given that LDA is a disolvate²⁴ in THF and homodimeric enolates **8** (below) are tetrasolvates.²⁵

(3)

Aggregation: dimer–tetramer mixtures in THF/toluene

^6Li NMR spectroscopy revealed that solutions of enolates **2–5** in THF/toluene solutions afforded two aggregates eventually shown using representative cases to be tetrasolvated tetramers (**7**) and tetrasolvated dimers (**8**; eq 4). The dimers were the only observable form in neat THF. Tetramers became observable typically below 5.0 M THF and were the exclusive form at $<2.0 \text{ M}$ THF. Assignment of the aggregation and solvation states is described below.

(4)

With MCV, mixtures of enolates **2b** and **3** in neat THF afforded the two homoaggregates along with a single heteroaggregate (Figure 1). Monitoring of their proportions versus measured²² mole fraction provided a Job plot showing a nearly statistical distribution (Figure 2). Dimers derived from **2a**, **2c**, **2e–j**, and **3** in neat THF were characterized similarly (supporting information).

The higher aggregates observed at low THF concentrations were shown to be tetramers by using MCV on a more select group of enolates. For example, mixtures of enolates **2b** and **3** in 0.13 M *total* THF concentration (coordinated and free) resulted in mole-fraction-

dependent ^6Li spectra consistent with an ensemble of homo- and heteroaggregated tetramers (Figure 3). The resulting Job plot shows a nearly statistical distribution of aggregates (Figure 4). Similar protocols confirmed the higher aggregates of **2i** and **2j** to be tetrameric at low THF concentration (supporting information). Enolates **2g**, **2k**, and **2i** also displayed clean ^6Li spectral data and concentration dependencies consistent with tetramer–dimer mixtures, but only the dimers were rigorously characterized. The dependencies on enolate and THF concentration, however, leave little doubt that the tetramers exist in those cases.

As a point of interest in light of efforts to develop MCV for studying non-lithium-based organometallics,¹⁶ mixtures of remotely fluorinated enolates **2b** and **2c** were characterized with MCV by monitoring well-resolved aggregates using ^{19}F NMR spectroscopy (supporting information.)^{16,26} The corresponding ^6Li resonances, by contrast, failed to resolve. α -Fluoro enolates **4** and **5** were also characterized with MCV and ^{19}F NMR spectroscopy. The fluorine tag on pairing partner **5** was used to characterize unfluorinated enolate **2j**.

Solvation: tetramers

It would be reasonable to infer from crystallographic¹⁸ and computational evidence²⁵ that the enolate tetramers are tetrasolvated, as in **8** (Scheme 1), but partial solvates have been observed in tetramers comprising hindered enolates.²⁷ To demonstrate qualitatively the *existence* of solvation, we turned to pyridine as a chemical shift reagent.¹⁶ Incremental additions of pyridine to THF/hydrocarbon solutions of lithium salts left unsolvated nuclei unchanged, whereas resonances corresponding to nuclei bearing substitutionally labile THF ligands shifted markedly downfield.^{16d,e,28}

Incremental replacement of THF by pyridine in 0.10 M solutions of enolate **2j** at $-30\text{ }^\circ\text{C}$ —while keeping the total ligand concentration constant (0.50 M)—resulted in marked downfield shifts of the observed tetramer. At $-95\text{ }^\circ\text{C}$, however, we detected additional spectral complexity (Figure 5) suggesting that the homo- and heterosolvated tetramers resolved (Scheme 1). Also readily apparent were 3:1, 2:2, and 1:3 mixed pyridine–THF solvates. Slow exchange of free and lithium-bound monodentate ligands on NMR time scales is rarely observed²⁹ [hexamethylphosphoramide (HMPA) excepted].^{18a,b} [^{15}N]Pyridine caused the downfield resonances to appear as doublets ($J_{\text{Li-N}} = 4.2\text{--}4.5\text{ Hz}$). Monitoring of the proportions of mixed-solvated tetramers versus the *measured* mole fraction²² (X_{p}) in THF–pyridine mixtures at constant donor ligand concentrations of 0.50 M afforded the relative proportions shown in the Job plot in Figure 6. Finally, varying the total solvent concentration caused no apparent changes in the coalescence temperature, which indicated that the exchange is dissociative rather than associative.

The Job plot in Figure 6 does *not* attest to the relative binding constants of THF and pyridine because we use the *measured* mole fraction—the mole fraction of only the bound ligands derived from the ^6Li resonances. Figure 6 shows deviations from a statistical distribution within the bound forms. Gaining insight into the relative binding affinities of THF and pyridine requires a plot of relative integration versus *intended* mole fraction—the actual or total mole fraction of THF and pyridine (Figure 7). A description of the parametric fits is

found in the supporting information. A discussion of the distortions in Figure 7 and the Job plot in Figure 6 appears below.

Table 1 illustrates the experimentally and computationally determined relative free energies of THF substitution by pyridine. The first row shows the measured free energies extracted directly from the Job plot in Figure 7. Embedded in these numbers are the statistical contributions³⁰ owing to the different numbers of available sites—ranging from four in the fully THF-solvated tetramer to one in the (ROLi)₄(pyr)₃(THF) tetramer. The second row comprises experimentally measured energies with the statistical factor omitted. The third row corresponds to the DFT-computed energies, which can be directly compared with the statistically corrected experimental numbers. Figure 8 shows a plot of theory versus experiment, which would be linear with a slope of unity in the event of a perfect correlation. Because of the skewed distribution of the data, the linearity is correct but unconvincing, and the slope is clearly not unity owing to discrepancies between theory and experiment. The trend, however, is clear.

We wondered whether the slow exchange in pyridine–THF mixtures arose from strong pyridine binding (akin to HMPA in HMPA–THF mixtures^{18a,b}) and whether THF alone could be observed in the slow-exchange limit. ¹³C NMR spectroscopy of 0.10 M **2j** with 2.25 equiv of THF (total) showed discrete resonances for the α-carbon corresponding to free and bound THF (δ66.3 and 66.8 ppm, respectively). The approximate 1.3:1 ratio confirmed the 1:1 THF–lithium proportion in the tetramer.

Solvation: dimers

Addition of pyridine caused a marked (>1.0 ppm) downfield shift of the ⁶Li resonance of the dimer, which confirmed that the dimer was solvated. Confident that the tetramers were tetrasolvates in THF/toluene, we monitored the tetramer–dimer proportions of enolate **3** as a function of THF concentration in toluene (Figure 9 and eq 5). The fit to eq 5 (supporting information) shows that the dimer is tetrasolvated (eq 6).¹⁸ Analogous THF dependencies were quantitated to show that the dimer of **2j** is also tetrasolvated.³¹

$$[A_2S_{2+n}]/[A_4S_4]^{1/2} = K_{eq}[S]^n \quad (6)$$

Atropisomerism: tetramers

Cooling solutions containing exclusively homotetrameric enolate to –120 °C elicited broadening and even multiple resonances in some cases. By example, after cooling, **2j** decoalesced to two broad resonances in a 1:1 proportion. With differential aggregation or solvation excluded as culprits, we were left with rotational isomerism (atropisomerism) arising from restricted rotation around the enolate C–O bonds. Calculations of enolate **2j**

show that the cyclohexyl substituents reside over faces of the cube (Scheme 2, top). There are, however, four substituents and *six* faces, which result in the three possible isomers depicted in Scheme 2. Each face of the cubes that is unencumbered by an enolate substituent is labeled an “open face.” We have also included parenthetically the number of magnetically inequivalent nuclei as well as the relative DFT computed energies. The computationally preferred atropisomer has an S_4 core structure that would result in the observed 1:1 resonance ratio. A simpler computational model using a methyl rather than a cyclohexyl moiety still showed a preference for the observed high-symmetry atropisomer (supporting information). The atropisomer exchange was THF-concentration-*independent*, consistent with either a rate-limiting THF dissociation or a mechanism requiring no THF dissociation.

Discussion

Enolates derived from hemiaminal-protected pyroglutaminol have captured the attention of synthetic organic chemists as nitrogen-containing chiral synthons. The concave and convex faces offer stereocontrol for further functionalizations. Lithium enolates **2–5** (see Chart 1) are also appealing templates with which to study enolate solvation and aggregation owing to variations available within the hemiaminal linkage that do not invade the inherent γ -lactam local structure.

Aggregation

The deceptively high steric demands of pyroglutaminol-derived enolates **2–5** have a striking impact on aggregation and solvation. Steric effects are notoriously difficult to quantitate because both size and shape are critical. Nonetheless, the hemiaminal substituents that we perceive to be large do indeed promote deaggregation of tetramer to dimer. *tert*-Butyl-substituted enolate **2k**, for example, gives no detectable tetramer with THF concentrations in toluene above 2.0 M, whereas the tetramer of **2a** persists at THF concentrations up to 6.0 M. Compared with their isostructural unfluorinated analogs, the α -fluoro groups on **4** and **5** also promote dimer by stabilizing the charge. (We observed no evidence of consequential F–Li interactions.)

Solvation

We routinely focus on how ligating solvents determine structure and reactivity. We assigned the tetramers and dimers as tetrasolvates **7** and **8**, respectively, by first showing that tetramer **7** is tetrasolvated and subsequently determining the relative solvation number of dimer **8**.

Assigning the solvation state of tetramer **4a** without merely relying on crystallographic analogy¹⁸ required some luck that began with a simple experiment. We are enamored with the probative role of pyridine as a ^6Li chemical shift reagent.^{16d,e,28} Adding pyridine to THF–toluene solutions of organolithiums results in marked downfield shifts for resonances corresponding to lithium nuclei bearing substitutionally labile THF ligands but has no effect on the chemical shifts of unsolvated ^6Li nuclei. Of course, in this case we were confident that the ^6Li nuclei bore at least one coordinated solvent, so it was intended as a control experiment. We were surprised, however, to observe pyridine–THF mixed solvates in the slow-exchange limit (see Scheme 1): the homo- and heterosolvated tetramers appear as

discrete species. The pyridine-solvated nuclei are distinguished by substantial downfield shifts and ^{15}N – ^6Li scalar coupling when ^{15}N pyridine is used. *Outside of reactions with HMPA, slow exchange of monodentate ligands on lithium is rarely observed.*²⁹ Slow pyridine exchange is *not* merely a consequence of strong coordination by pyridine: free and bound THF can be distinguished on tetramer **2j** with ^{13}C NMR spectroscopy, which allowed us to confirm the 1:1 THF–Li stoichiometry.

Bolstered by the results from ongoing studies of other enolates,¹⁶ a transiently formed five-coordinate lithium on a tetrameric cube seemed quite plausible.³² The steric hindrance of enolate **2** is critical to slowing the solvent exchange. Solvent-concentration-independent coalescence behavior confirmed what we suspected: solvent substitution is *dissociative*. Given that hindrance should *accelerate* such a dissociative substitution through steric relief,^{29,33} the bulk must suppress otherwise facile associative substitutions. Control experiments showed that less congested enolates such as that derived from indanone undergo rapid THF and pyridine exchange.

Atropisomerism

Cooling solutions of tetramer **7** to $-120\text{ }^{\circ}\text{C}$ revealed what we have assigned as atropisomerism. Several of these solutions decoalesce to complex mixtures, whereas others neatly decoalesce to a pair of resonances in a 1:1 ratio. The consequences of restricted rotation about the enolate C–O bonds are highlighted in Scheme 2. The most symmetric form that afforded two resonances is computed to be the most stable (albeit within computational error). We cannot help but see analogy between the shape of enolate **2** and the naphthyl groups in highly restricted binaphthyl moieties. Carlier³⁴ described a series of benzodiazepine-derived enolates in which conformational constraints within a ring had remarkable effects on reactivity. We are unaware of other reports on the atropisomerism of the type described herein,³⁵ but we also cannot currently imagine tangible consequences beyond merely spectroscopic effects.

Comments on MCV

We continue to find applications for MCV (Job plots) in determinations of organolithium structure. A recent review underscores our opinion that such applications reach beyond organolithium chemistry.²⁰ With that said, this study offers an excellent opportunity to emphasize a nuance of Job plots that may be unappreciated by many users, ideas and concerns that mirror those recently articulated by Jurczak and coworkers.³⁶ We use measured mole fraction—the mole fraction determined by integrating all species in the ensemble of aggregates or mixed solvates of a single aggregate—to determine solution structures even on impure samples or when multiple species coexisted.¹⁶ By example, Figure 6 shows how pyridine and THF distribute within the ensemble (eq 7) but *provides no insight whatsoever about the relative tendencies of THF and pyridine to bind lithium*. The presence of monomers, dimers, mixed aggregates, and even decomposition does not influence the outcome. Deviations from a statistical distribution reflect only deviations within the ensemble, which makes the method robust.

Contrast Figure 6 with Figure 7, in which absolute mole fraction—the mole fraction dictated by the quantities added—is used. Distortion of the maxima along the x axis toward the lower pyridine mole fractions reflects the relatively higher binding constants of pyridine compared with those of THF. *The analysis was more challenging but achievable provided that the composition and distribution of all species could be monitored.* It allowed us to extract the binding constants for the sequential substitution of pyridine for THF. The serial substitution by pyridine (see Scheme 1 and eq 8) is progressively less favorable with each substitution. A comparison with the values calculated with DFT (see Table 1 and Figure 8) shows a qualitative correlation that is credible given the inherent limitations of both experiment and theory.

Now we get to the ugly part: *all Job plots are susceptible to this error.* Imagine monitoring species **A** and **B** to form adduct **A_mB_n**. The position of the maximum along the x axis and the curvature reveal the composition of **A_mB_n** *provided that A_mB_n is the only associated form.* If other undetected equilibria or decomposition—any process by which the proportions of free **A** and **B** are distorted—are present, the answer will be quantitatively wrong. We escape this trap by using measured mole fraction—by monitoring all species—within ensembles, but this protocol seems to be rare. One can only imagine how many stoichiometries gleaned from the approximately 6500 studies using MCV²⁰ have been misassigned. It seems almost certain that the answer is not zero.

Conclusion

Controlling selectivity is certainly one of the paramount goals of the synthetic organic community. We are currently completing investigations of an aza-aldol addition by pyroglutaminol-derived enolate **2** in which changes in stereoselectivity correlate with mechanistic changes. We are also extending seminal structural studies from the laboratories of Jackman, Seebach, Streitwieser, Williard, Arnett, and Reich (just to name a few)¹⁸ to try to understand if and when tetrameric and dimeric enolates play a role beyond simply being precursors to monomers.³⁷ There is a nagging central question: can one control stereoselectivity by controlling enolate aggregation state? Flowers³⁸ appears to have achieved this goal, but it is a relatively untested hypothesis.

There is a more subtle but noteworthy consequence to the work described herein. The opportunity to observe lithium ion solvation in the slow-exchange limit is not trivial.²⁹

Reich's^{18a,b} studies of HMPA opened an enormous window into solvation as a molecular phenomenon rather than just a medium. Our own experiences with the solvation of lithium hexamethyldisilazide by ethereal solvents in the slow-exchange limit²⁹ were central to our efforts to understand lithium amide structure and reactivity. Gathering hard data on the mechanisms of solvent exchange, relative binding constants, and correlated solvation within mixed solvates is critical to nudging the chemistry of nondescript lithium salts toward a legitimate subdiscipline of coordination chemistry.

Experimental

Reagents and solvents

THF, toluene, and pyridine were distilled from solutions containing sodium benzophenone ketyl. The toluene stills contained approximately 1% tetraglyme to dissolve the ketyl. [⁶Li]LDA and [⁶Li,¹⁵N]LDA were prepared as described previously.¹⁹ Solutions of LDA were titrated for active base with a literature method.³⁹ Air- and moisture-sensitive materials were manipulated under argon with standard glovebox, vacuum line, and syringe techniques.

NMR spectroscopy

Individual stock solutions of substrates and LDA were prepared at room temperature, and samples of enolate were generated in an NMR tube and flame-sealed as described previously.¹⁶ Standard ⁶Li, ¹³C, ¹⁵N, and ¹⁹F NMR spectra were recorded on a 500 MHz spectrometer at 73.57, 125.79, 50.66, and 470.35 MHz, respectively. The ⁶Li, ¹³C, ¹⁵N, and ¹⁹F resonances were referenced to 0.30 M [⁶Li]LiCl/MeOH at –80 °C (0.0 ppm), the CH₂O resonance of THF at –90 °C (67.57 ppm), neat Me₂NEt at –90 °C (25.7 ppm), and C₆H₅F in neat THF at –80 °C (–112.0 ppm).

Known compounds were prepared according to the literature preparations.^{7a,40}

(3*R*,7*aS*)-3-(2-fluorophenyl)tetrahydro-3*H*,5*H*-pyrrolo[1,2-*c*]oxazol-5-one (1*c*)—L-pyrroglutaminol (500 mg, 4.34 mmol), *p*-toluenesulfonic acid (18.0 mg, 0.090 mmol) and 2-fluorobenzaldehyde (0.33 mL, 3.08 mmol) were added to a 25 mL round bottom flask containing toluene (8.3 mL). The round bottom flask was attached to a Dean–Stark trap containing toluene and freshly activated 3 Å molecular sieves and was brought to reflux for 5 hours. The resulting yellow solution was allowed to cool to rt, and then 10 mL of saturated brine solution were added. This mixture was extracted with 3×25 mL of Et₂O, and the organic layers were dried over Na₂SO₄ and dried in vacuo. The resulting yellow oil was purified using flash chromatography in Et₂O and rotatory evaporated to yield 470 mg (69%) of white solid. *R*_f = 0.34 in 60% ethyl acetate/hexanes. Melting point 62.9 – 68.3 °C. ¹H NMR (400 MHz, CDCl₃) δ 7.22 (td, ⁴*J*_{H-F} = 7.5 Hz, *J*_{H-H} = 7.5, 1.8 Hz, 1H), 7.15 (dddd, ⁴*J*_{H-F} = 5.2 Hz, *J*_{H-H} = 8.0, 7.2, 1.8 Hz, 1H), 6.96 (td, *J* = 7.6, 1.1 Hz, 1H), 6.90 (ddd, ³*J*_{H-F} = 10.5 Hz, *J*_{H-H} = 8.2, 1.1 Hz, 1H), 4.15 – 3.98 (m, 2H), 3.24 (t, *J* = 7.8 Hz, 1H), 2.59 (ddd, *J* = 17.3, 10.3, 8.8 Hz, 1H), 2.31 (ddd, *J* = 17.3, 10.0, 4.0 Hz, 1H), 2.16 (dddd, *J* = 14.2, 10.1, 7.4, 4.0 Hz, 1H), 1.73 (dddd, *J* = 13.6, 9.9, 8.8, 5.0 Hz, 1H). ¹³C NMR (101 MHz, Chloroform-*d*) δ 177.8, 160.5 (d, ¹*J*_{C-F} = 249.3 Hz), 130.4 (d, ³*J*_{C-F} = 8.3 Hz), 127.6 (d, ³*J*_{C-F} = 3.8 Hz), 125.9 (d, ²*J*_{C-F} = 12.7 Hz), 123.8 (d, ⁴*J*_{C-F} = 3.6 Hz), 115.5 (d, ²*J*_{C-F} =

21.0 Hz), 83.4 (d, $^3J_{\text{C-F}} = 2.7$ Hz), 71.4, 59.0 (d, $^5J_{\text{C-F}} = 1.0$ Hz), 32.9, 22.3. ^{19}F NMR (376 MHz, CDCl_3) δ -117.20 (ddd, $^3J_{\text{H-F}} = 10.4$ Hz, $^4J_{\text{H-F}} = 7.4$, 5.2 Hz). HRMS (DART ionization, orbitrap mass analyzer) calcd for $\text{C}_{12}\text{H}_{12}\text{FNO}_2$ [M+H] 222.09303, found 222.09248.

(3R,7aS)-3-(3-fluorophenyl)tetrahydro-3H,5H-pyrrolo[1,2-c]oxazol-5-one (1d)—Yield = 75% (721 mg of white solid). $R_f = 0.33$ in 60% ethyl acetate/hexanes. ^1H NMR (500 MHz, CDCl_3) δ 7.31 (td, $^4J_{\text{H-F}} = 5.7$ Hz, $J_{\text{H-H}} = 7.9$, 5.7 Hz, 1H), 7.23 (ddt, $^5J_{\text{H-F}} = 0.9$ Hz, $J_{\text{H-H}} = 7.7$, 1.7, 0.9 Hz, 1H), 7.14 (dt, $^3J_{\text{H-F}} = 9.6$ Hz, $J_{\text{H-H}} = 2.1$ Hz, 1H), 6.99 (tdd, $^3J_{\text{H-F}} = 8.4$, $J_{\text{H-H}} = 8.4$, 2.6, 1.0 Hz, 1H), 6.28 (s, 1H), 4.22 (dd, $J = 8.1$, 6.3 Hz, 1H), 4.14 – 4.08 (m, 1H), 3.48 (t, $J = 8.2$ Hz, 1H), 2.80 (ddd, $J = 17.3$, 10.2, 9.0 Hz, 1H), 2.54 (ddd, $J = 17.4$, 10.0, 3.8 Hz, 1H), 2.38 (dddd, $J = 14.1$, 10.2, 7.6, 3.8 Hz, 1H), 1.94 (dddd, $J = 13.3$, 9.9, 8.9, 5.4 Hz, 1H). ^{13}C NMR (126 MHz, Chloroform-*d*) δ 178.2, 162.8 (d, $^1J_{\text{C-F}} = 246.4$ Hz), 141.5 (d, $^3J_{\text{C-F}} = 6.8$ Hz), 130.1 (d, $^3J_{\text{C-F}} = 8.0$ Hz), 121.7 (d, $^4J_{\text{C-F}} = 3.0$ Hz), 115.4 (d, $^2J_{\text{C-F}} = 21.2$ Hz), 113.0 (d, $^2J_{\text{C-F}} = 22.7$ Hz), 86.3 (d, $^4J_{\text{C-F}} = 2.2$ Hz), 71.7, 58.7, 33.3, 23.0. ^{19}F NMR (376 MHz, CDCl_3) δ -112.79 (dddd, $^3J_{\text{H-F}} = 9.6$ Hz, $^3J_{\text{H-F}} = 8.6$ Hz, $^4J_{\text{H-F}} = 5.7$ Hz, $^5J_{\text{H-F}} = 0.9$ Hz). HRMS (DART ionization, orbitrap mass analyzer) calcd for $\text{C}_{12}\text{H}_{12}\text{FNO}_2$ [M+H] 222.09303, found 222.09248.

(3R,7aS)-3-(2-methoxyphenyl)tetrahydro-3H,5H-pyrrolo[1,2-c]oxazol-5-one (1e)—Yield = 24% (486 mg of white solid). $R_f = 0.21$ in 0.34 in 60 % ethyl acetate/hexanes. Melting point 94.5 – 101.6 °C. ^1H NMR (400 MHz, CDCl_3) δ 7.31 (td, $J = 7.8$, 1.7 Hz, 1H), 7.26 (dd, $J = 7.5$, 1.7 Hz, 1H), 4.32 (tt, $J = 7.9$, 5.7 Hz, 1H), 4.21 (dd, $J = 8.1$, 6.2 Hz, 1H), 3.88 (s, 3H), 3.46 (t, $J = 8.2$ Hz, 1H), 2.85 (dt, $J = 17.2$, 9.6 Hz, 1H), 2.57 (ddd, $J = 17.3$, 10.0, 3.7 Hz, 1H), 2.42 (dddd, $J = 13.8$, 10.0, 7.6, 3.9 Hz, 1H), 1.96 (dtd, $J = 18.8$, 9.5, 5.4 Hz, 1H). ^{13}C NMR (126 MHz, CDCl_3) δ 178.0, 157.7, 130.4, 126.6, 126.2, 120.4, 111.2, 84.6, 71.6, 59.9, 55.9, 33.6, 23.1. HRMS (DART ionization, orbitrap mass analyzer) calcd for $\text{C}_{13}\text{H}_{15}\text{NO}_3$ [M+H] 234.11302, found 234.11247.

(3R,7aS)-3-(2,6-dimethylphenyl)tetrahydro-3H,5H-pyrrolo[1,2-c]oxazol-5-one (1f)—Yield = 30% (174 mg of white solid). $R_f = 0.34$ in 60 % ethyl acetate/hexanes. Melting point 110.6 – 112.9 °C. ^1H NMR (400 MHz, CDCl_3) δ 7.10 (t, $J = 7.5$ Hz, 1H), 6.99 (d, $J = 7.5$ Hz, 2H), 6.44 (s, 1H), 4.52 – 4.43 (m, 1H), 4.40 (dd, $J = 8.3$, 5.7 Hz, 1H), 3.48 (t, $J = 8.7$ Hz, 1H), 2.97 – 2.83 (m, 1H), 2.49 (s, 8H), 2.38 (dddd, $J = 12.8$, 9.4, 6.9, 1.9 Hz, 1H), 1.95 (dddd, $J = 12.8$, 11.5, 9.5, 7.0 Hz, 1H). ^{13}C NMR (126 MHz, CDCl_3) δ 178.0, 157.7, 130.4, 126.6, 126.2, 120.4, 111.2, 84.6, 71.6, 59.9, 55.9, 33.6, 23.1. HRMS (DART ionization, orbitrap mass analyzer) calcd for $\text{C}_{14}\text{H}_{17}\text{NO}_2$ [M+H] 232.13375, found 232.13321.

(3R,7aS)-3-(naphthalen-1-yl)tetrahydro-3H,5H-pyrrolo[1,2-c]oxazol-5-one (1g)—Yield = 51% (561 mg of tan solid). $R_f = 0.31$ in 60 % ethyl acetate/hexanes. Melting point 154.6 – 161.3 °C. ^1H NMR (400 MHz, CDCl_3) δ 8.44 (d, $J = 8.4$ Hz, 1H), 7.85 (dd, $J = 12.0$, 8.1 Hz, 2H), 7.58 (td, $J = 6.8$, 6.1, 2.2 Hz, 2H), 7.51 (t, $J = 7.4$ Hz, 1H), 7.43 (t, $J = 7.7$ Hz, 1H), 7.06 (s, 1H), 4.29 (t, $J = 7.2$ Hz, 1H), 4.19 (qd, $J = 7.6$, 4.3 Hz, 1H), 3.56 (t, $J = 8.0$ Hz, 1H), 2.84 (ddd, $J = 17.9$, 10.5, 7.6 Hz, 1H), 2.62 (ddd, $J = 17.6$, 10.1, 4.9 Hz, 1H), 2.40

(dddd, $J = 12.9, 10.3, 7.7, 4.9$ Hz, 1H), 2.00 (dddd, $J = 14.0, 10.0, 7.7, 4.3$ Hz, 1H). ^{13}C NMR (126 MHz, CDCl_3) δ 179.0, 134.0, 133.7, 130.9, 129.6, 128.6, 126.7, 126.1, 125.0, 124.2, 122.8, 86.2, 71.3, 58.8, 32.8, 22.3. HRMS (DART ionization, orbitrap mass analyzer) calcd for $\text{C}_{16}\text{H}_{15}\text{NO}_2$ [M+H] 254.11810, found 254.11756.

(3*R*,7*aS*)-3-(anthracen-9-yl)tetrahydro-3*H*,5*H*-pyrrolo[1,2-*c*]oxazol-5-one (1h)—

Yield = 10% (26.4 mg of yellow solid). $R_f = 0.23$ in 60 % ethyl acetate/hexanes. ^1H NMR (500 MHz, CDCl_3) δ 8.63 (dd, $J = 9.1, 1.1$ Hz, 1H), 8.50 (s, 1H), 8.05 – 7.97 (m, 1H), 7.56 (s, 1H), 7.54 (ddd, $J = 9.1, 6.5, 1.5$ Hz, 1H), 7.46 (ddd, $J = 8.4, 6.5, 1.0$ Hz, 1H), 4.76 (dq, $J = 9.2, 6.5$ Hz, 1H), 4.58 (dd, $J = 8.6, 5.8$ Hz, 1H), 3.72 (t, $J = 8.9$ Hz, 1H), 2.93 (dt, $J = 17.0, 10.2$ Hz, 1H), 2.60 (ddd, $J = 17.0, 9.7, 2.7$ Hz, 1H), 2.50 (dddd, $J = 12.9, 9.9, 7.2, 2.7$ Hz, 1H), 2.09 (dtd, $J = 13.1, 10.1, 6.4$ Hz, 1H). ^{13}C NMR (126 MHz, CDCl_3) δ 178.4, 131.7, 130.5, 130.4, 129.5, 126.7, 126.5, 124.9, 124.1, 86.5, 73.2, 60.8, 34.9, 24.3. HRMS (DART ionization, orbitrap mass analyzer) calcd for $\text{C}_{13}\text{H}_{15}\text{NO}_3$ [M+H] 304.13375, found 304.13440.

(3*R*,7*aS*)-3-([1,1'-biphenyl]-2-yl)tetrahydro-3*H*,5*H*-pyrrolo[1,2-*c*]oxazol-5-one (1i)—

Yield = 62% (3.01 g of white solid). $R_f = 0.33$ in 60 % ethyl acetate/hexanes. Melting point 108.8 – 116.2 °C. ^1H NMR (400 MHz, CDCl_3) δ 7.52 – 7.46 (m, 2H), 7.45 – 7.38 (m, 3H), 7.38 – 7.32 (m, 3H), 7.30 – 7.23 (m, 1H), 4.29 (ddt, $J = 8.5, 7.2, 6.0$ Hz, 1H), 4.18 (dd, $J = 8.2, 5.9$ Hz, 1H), 3.32 (t, $J = 8.4$ Hz, 1H), 2.45 (ddd, $J = 17.2, 9.9, 3.1$ Hz, 1H), 2.34 (dddd, $J = 13.1, 10.3, 7.4, 3.1$ Hz, 1H), 1.86 (dtd, $J = 13.2, 9.9, 6.0$ Hz, 1H). ^{13}C NMR (126 MHz, CDCl_3) δ 177.3, 142.3, 140.2, 135.7, 130.8, 129.9, 128.8, 128.1, 127.7, 127.3, 126.4, 86.3, 71.9, 60.9, 34.2, 23.5. HRMS (DART ionization, orbitrap mass analyzer) calcd for $\text{C}_{18}\text{H}_{17}\text{NO}_2$ [M+H] 280.13375, found 280.13321.

(3*R*,6*R*,7*aS*)-3-cyclohexyl-6-fluorotetrahydro-3*H*,5*H*-pyrrolo[1,2-*c*]oxazol-5-one (1k)—

Lithium diisopropylamide (279 mg, 2.64 mmol) was dissolved in THF (11 mL) and cooled to –78 °C. To this mixture was added **1j** (500 mg, 2.40 mmol) dissolved in THF (10 mL) and was allowed to stir for 10 minutes to ensure complete enolization. *N*-fluorobenzenesulfonimide (983 mg, 3.12 mmol) dissolved in THF (4 mL) was added. After 2 h, the reaction was quenched with pH 7 phosphate buffer (6 mL) and allowed to warm. The mixture was extracted 3 × 20 mL EtOAc, dried over Na_2SO_4 , and rotary evaporated. The resulting yellow oil was purified using flash chromatography with a 70% diethyl Et_2O /pentane solvent system and rotary evaporated to give an oil that was further recrystallized from Et_2O /pentane to yield 136 mg (25%) of white solid as a single diastereomer. $R_f = 0.54$ in 60% ethyl acetate/hexanes. Melting point 68.4 – 69.7 °C. ^1H NMR (400 MHz, CDCl_3) δ 5.08 (ddd, $^2J_{\text{H-F}} = 51.9$ Hz, $J_{\text{H-H}} = 6.4, 1.5$ Hz, 1H), 4.99 (dd, $^5J_{\text{H-F}} = 2.4$ Hz, $J_{\text{H-H}} = 5.4$ Hz, 1H), 4.26 – 4.16 (m, 2H), 3.28 – 3.15 (m, 1H), 2.58 – 2.40 (m, 1H), 2.17 – 1.96 (m, 1H), 1.83 – 1.67 (m, 5H), 1.67 – 1.52 (m, 2H), 1.30 – 1.01 (m, 5H). ^{13}C NMR (126 MHz, Chloroform-*d*) δ 170.4 (d, $^2J_{\text{C-F}} = 17.0$ Hz), 94.1 (d, $^1J_{\text{C-F}} = 184.7$ Hz), 90.8, 71.9 (d, $^3J_{\text{C-F}} = 2.4$ Hz), 58.0, 42.5 (d, $^5J_{\text{C-F}} = 1.0$ Hz), 31.6 (d, $^2J_{\text{C-F}} = 23.7$ Hz), 27.47, 27.36, 26.19, 25.67, 25.61. ^{19}F NMR (376 MHz, CDCl_3) δ –180.90 – –181.29 (m). HRMS (DART ionization, orbitrap mass analyzer) calcd for $\text{C}_{12}\text{H}_{18}\text{FNO}_2$ [M+H] 228.13998, found 228.13943.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Institutes of Health (GM077167) for support.

References and Footnotes

1. (a) Wright SW, Choi C, Chung S, Boscoe BP, Drozda SE, Mousseau JJ, Trzupek JD. *Org Lett.* 2015; 17:5204. [PubMed: 26485576] (b) Anderson, DR., Bunnage, ME., Curran, KJ., Dehnhardt, CM., Gavrin, LK., Goldberg, JA., Han, S., Hepworth, D., Huang, H-C., Lee, A., Lee, KL., Lovering, FE., Lowe, MD., Mathias, JP., Papaioannou, N., Patny, A., Pierce, BS., Saiah, E., Strohbach, JW., Trzupek, JD., Vargas, R., Wang, X., Wright, SW., Zapf, CW. Bicyclic-fused heteroaryl or aryl compounds and their use as IRAK4 inhibitors. WO2015150995. 2015.
2. Najera, Carmen, Yus, Miguel. *Tetrahedron: Asymmetry.* 1999; 10:2245.
3. (a) Yoshioka S, Nagatomo M, Inoue M. *Org Lett.* 2015; 17:90. [PubMed: 25526222] (b) Kyle AF, Jakubec P, Cockfield DM, Cleator E, Skidmore J, Dixon DJ. *Chem Commun.* 2011; 47:10037.(c) Imai T, Nagasaka T. *Chem Pharm Bull.* 1995; 43:1081.(d) Zhang R, Brownnewell F, Madalengoitia JS. *Tetrahedron Lett.* 1999; 40:2707.(e) Clark AJ, Filik RP, Thomas GH, Sherringham J. *Tetrahedron Lett.* 2013; 54:4094.(f) Cowley AR, Hill TJ, Kocis P, Moloney MG, Stevenson RD, Thompson AL. *Org Biomol Chem.* 2011; 9:7042. [PubMed: 21858317] (g) Jao E, Bogen S, Saksena AK, Girijavallabhan V. *Synthesis.* 2003; 17:2643.(h) Makino K, Shintani K, Yamatake T, Hara O, Hatano K, Hamada Y. *Tetrahedron.* 2002; 58:9737.(i) Okamoto N, Hara O, Makino K, Hamada Y. *Tetrahedron: Asymmetry.* 2001; 12:1353.(j) Langlois N, Rakotonradany F. *Tetrahedron.* 2000; 56:2437.(k) Gu L, Yu M, Wu X, Zhang Y, Zhao G. *Adv Synth Catal.* 2006; 348:2223.(l) Sridhar J, Wei ZL, Nowak I, Lewin NE, Ayres JA, Pearce LV, Blumberg PM, Kozikowski AP. *J Med Chem.* 2003; 46:4196. [PubMed: 12954072] (m) Davies SG, Dixon DJ, Doisneau GJM, Prodger JC, Sanganee HJ. *Tetrahedron: Asymmetry.* 2002; 13:647.(n) Langlois N, Rakotonradany F. *Tetrahedron.* 2000; 56:2437.(o) Bailey JH, Byfield ATJ, Davis PJ, Foster AC, Leech M, Moloney MG, Muller M, Prout CK. *J Chem Soc Perkin Trans.* 2000; 1:1977.
4. a) Qiu, YL., Wang, C., Peng, X., Cao, H., Ying, L., Gao, X., Wang, B., Or, YS. Novel Benzimidazole Derivatives. WO 2013052369 A1. Apr 11. 2013 (b) Qiu, YL., Wang, C., Peng, X., Cao, H., Ying, L., Or, YS. Hepatitis c virus inhibitors. US 20110142798 A1. Jun 16. 2011 (c) Borriello, M., Pucci, S., Stasi, LP., Rovati, L. Cyclic amine derivatives as ep4 receptor antagonists. WO 2013004290 A1. Jan 10. 2013 (d) Dahmann, G., Fiegen, D., Fleck, M., Hoffmann, M., Klicic, J., East, SP., Napier, Spencer Charles R., Scott, John. PCT Int Appl. 2012101013. Aug 02. 2012 (e) Qian, X., Chuang, C., Lu, PP., Yao, B., Lu, Q., Jiang, H., Wang, W., Morgan, BP., Morgans, D, Jr. Certain chemical entities, compositions, and methods. WO 2009023193 A1. Feb 19. 2009 (f) Kozikowski, AP., Blumberg, PM., Hennings, H., Nowak, I. Protein kinase c modulators, and methods of use thereof. WO 2004007445 A3. Jan 22. 2004
5. (a) Beard MJ, Bailey JH, Cherry DT, Moloney MG, Shim SB, Statham KA, Bamford MJ, Lamont RB. *Tetrahedron.* 1996; 52:3719.(b) Baldwin JE, Moloney MG, Shim SB. *Tetrahedron Lett.* 1991; 32:1379.(c) Starkman BA, Young DW. *J Chem Soc, Perkin Trans.* 2002; 1:725.
6. (a) Bowler AN, Doyle PM, Hitchcock PB, Young DW. *Tetrahedron Lett.* 1991; 32:2679.(b) Avent AG, Bowler AN, Hitchcock PB, Young DW. *Tetrahedron Lett.* 1992; 33:1509.(c) Anwar M, Bailey JH, Dickinson LC, Edwards HJ, Goswami R, Moloney MG. *Org Biomol Chem.* 2003; 1:2364. [PubMed: 12945710]
7. (a) Konas DW, Coward JK. *J Org Chem.* 2001; 66:8831. [PubMed: 11749613] (b) Martinez-Montero S, Fernandez S, Sanghvi YS, Theodorakis EA, Detorio MA, Mcbrayer TR, Whitaker T, Schinazi RF, Gotor V, Ferrero M. *Bioorg Med Chem.* 2012; 20:6885. [PubMed: 23085031] (c) Doyle MP, Hu W, Phillips IM, Moody CJ, Pepper AG, Slawin AMZ. *Adv Synth Catal.* 2001; 343:112.(d) Konas DW, Coward JK. *Org Lett.* 1999; 1:2105. [PubMed: 10836063]

8. (a) Barrett, SD., Colombo, JM., Germain, BD., Kornilov, A., Kramer, JB., Uzieblo, A., Endres, GW., Ciske, FL., Owen, TA., O'Malley, JP. Preparation of prostaglandin analogs for methods, systems, and compositions for promoting bone growth. WO 2015009991 A2. Jan 22. 2015 (b) Barrett, SD., Colombo, JM., Germain, BD., Kornilov, A., Kramer, JB., Uzieblo, A. Synthesis of difluorolactam analogs. WO 2014144500 A2. Sep 18. 2014 (c) Barrett, SD., Ciske, FL., Colombo, JM., Endres, GW., Germain, BD., Kornilov, A., Kramer, JB., Uzieblo, A., Maxey, KM. Preparation of difluorolactams as selective EP4 receptor agonists and their use for in the treatment of EP4-mediated diseases and conditions. WO 2014015247 A1. Jan 23. 2014 (d) Doyle, MP., Hu, W., Phillips, I. Preparation and use of chiral rhodium complexes as Lewis acid catalysis for enantioselective hetero-Diels-Alder and addition reactions. WO 2002045853 A1. Jun 13. 2002 (e) Qian, X., Chuang, C., Lu, PP., Yao, B., Lu, Q., Jiang, H., Wang, W., Morgan, BP., Morgans, D, Jr. Preparation of isoquinolinylcarbamates as smooth muscle myosin or non-muscle myosin modulators. WO 2009023193 A1. Feb 19. 2009
9. (a) Hanessian S, Ratovelomanana V. Synlett. 1990; 9:501.(b) Hara S, Makino K, Hamada Y. Tetrahedron. 2004; 37:8031.(c) Okamoto N, Hara O, Makino K, Hamada Y. Tetrahedron: Asymmetry. 2001; 9:1353.(d) Endo A, Danishefsky SJ. J Am Chem Soc. 2005; 127:8298. [PubMed: 15941259]
10. Danishefsky, S., Endo, A. Synthesis of salinosporamide A and analogues thereof. PCT Int Appl. 2006124902. Nov 23. 2006
11. (a) Tsujishima H, Nakatani K, Shimamoto K, Shigeri Y, Yumoto N, Ohfuné Y. Tetrahedron Lett. 1998; 39:1193.(b) Bailey JH, Cherry DT, Crapnell KM, Moloney MG, Shim SB, Bamford MJ, Lamont RB. Tetrahedron. 1997; 53:11731.(c) Bamford MJ, Beard M, Cherry DT, Moloney MG. Tetrahedron: Asymmetry. 1995; 6:337.
12. (a) Cottrell IF, Davis PJ, Moloney MG. Tetrahedron: Asymmetry. 2004; 15:1239.(b) Kamath VP, Xue J, Juarez-Brambila JJ, Morris PE. Tetrahedron Lett. 2009; 50:5198.(c) Smith AB III, Friestad JB, Emmanuel B, Duan JJW, Hull KG, Iwashima M, Qiu Y, Spoors PG, Salvator BA. J Am Chem Soc. 1999; 121:10478.
13. Zhang R, Mamai A, Madalengoitia JS. J Org Chem. 1999; 64:547.
14. Aissaoui, A., Gude, M., Boss, C., Koberstein, R., Sifferlen, T., Trachsel, D., Lehmann, D. Trans-3-Aza-Bicyclo[3.1.0]Hexane Derivatives. WO 200916560 A2. Feb 5. 2009
15. Boyd DB, Foster BJ, Hatfield LD, Hornback WJ, Jones ND, Munroe EJ, Swartzendruber JK. Tetrahedron Lett. 1986; 27:3457.
16. (a) Liou LR, McNeil AJ, Ramírez A, Toombes GES, Gruver JM, Collum DB. J Am Chem Soc. 2008; 130:4859. [PubMed: 18336025] (b) De Vries TS, Goswami A, Liou LR, Gruver JM, Jayne E, Collum DB. J Am Chem Soc. 2009; 131:13142. [PubMed: 19702308] (c) Tomasevich LL, Collum DB. J Org Chem. 2013; 78:7498. [PubMed: 23806055] (d) Tomasevich LL, Collum DB. J Am Chem Soc. 2014; 136:9710. [PubMed: 24915602] (e) Tallmadge EH, Collum DB. J Am Chem Soc. 2015; 137:13087. [PubMed: 26437278]
17. (a) Green, JR. In Science of Synthesis. Vol. 8a. Georg Thieme Verlag; New York: 2005. p. 427-486.(b) Farina V, Reeves JT, Senanayake CH, Song JJ. Chem Rev. 2006; 106:2734. [PubMed: 16836298] (c) Wu G, Huang M. Chem Rev. 2006; 106:2596. [PubMed: 16836294]
18. (a) Reich HJ. Chem Rev. 2013; 113:7130. [PubMed: 23941648] (b) Reich HJ. J Org Chem. 2012; 77:5471. [PubMed: 22594379] (c) Seebach D. Angew Chem, Int Ed Engl. 1988; 27:1624.(d) Braun M. Helv Chim Acta. 2015; 98:1.(e) Williard, PG. Comprehensive Organic Synthesis. Trost, BM., Fleming, I., editors. Vol. 1. Pergamon; New York: 1991. Chapter 1.1(f) Kim YJ, Streitwieser A. Org Lett. 2002; 4:573. [PubMed: 11843594] (g) Jackman LM, Lange BC. Tetrahedron. 1977; 33:2737.(h) Li D, Keresztes I, Hopson R, Williard PG. Acc Chem Res. 2009; 42:270. [PubMed: 19105594] (i) Zune C, Jerome R. Prog Polymer Sci. 1999; 24:631.(k) Baskaran D. Prog Polym Sci. 2003; 28:521.(l) Harrison-Marchand A, Mongin F. Chem Rev. 2013; 113:7470. [PubMed: 23952819]
19. Ma Y, Hoepker AC, Gupta L, Faggini MF, Collum DB. J Am Chem Soc. 2010; 132:15610. [PubMed: 20961095]
20. Renny JS, Tomasevich LL, Tallmadge EH, Collum DB. Angew Chem, Int Ed. 2013; 52:11998.
21. Frisch, MJ., et al. GaussianVersion 3.09; revision A1. Gaussian, Inc; Wallingford, CT: 2009.

22. The measured mole fraction—the mole fraction within only the ensemble of interest—rather intended mole fraction of the enolates added to the samples—eliminates the distorting effects of impurities.
23. Job P. *Ann Chim.* 1928; 9:113.
24. Collum DB. *Acc Chem Res.* 1993; 26:227.
25. For leading references to theoretical studies of O-lithiated species, see: Khartabi HK, Gros PC, Fort Y, Ruiz-Lopez MF. *J Org Chem.* 2006; 73:9393. Streitwieser A. *J Mol Model.* 2006; 12:673. [PubMed: 16341718] Pratt LM, Streitwieser A. *J Org Chem.* 2003; 68:2830. [PubMed: 12662059] Pratt LM, Nguyen SC, Thanh BT. *J Org Chem.* 2008; 73:6086. [PubMed: 18646860]
26. (a) Gakh YG, Gakh AA, Gronenborn AM. *Magn Reson Chem.* 2000; 38:551. (b) McGill CA, Nordon A, Littlejohn D. *J Process Anal Chem.* 2001; 6:36. (c) Espinet P, Albeniz AC, Casares JA, Martinez-Ilarduya JM. *Coord Chem Rev.* 2008; 252:2180.
27. Pospisil PJ, Wilson SR, Jacobsen EN. *J Am Chem Soc.* 1992; 114:7585.
28. Reich HJ, Kulicke KJ. *J Am Chem Soc.* 1996; 118:273. Jackman LM, Chen X. *J Am Chem Soc.* 1997; 119:8681. Jackman LM, Petrei MM, Smith BD. *J Am Chem Soc.* 1991; 113:3451. Jackman LM, DeBrosse CW. *J Am Chem Soc.* 1983; 105:4177. Deana RK, Recklinga AM, Chena H, Daweab LN, Schneiderac CM, Kozak CM. *Dalton Trans.* 2013; 42:3504. [PubMed: 23280340] Boyle TJ, Pedrotty DM, Alam TM, Vick SC, Rodriguez MA. *Inorg Chem.* 2000; 39:5133. [PubMed: 11233212]. (g) See also ref 16c.
29. (a) Hilmersson G, Davidsson O. *J Org Chem.* 1995; 60:7660. (b) Hilmersson G, Ahlberg P, Davidsson O. *J Am Chem Soc.* 1996; 118:3539. (c) Boche G, Fraenkel G, Cabral J, Harms K, van Eikema Hommes NJR, Lohrenz J, Marsch M, Schleyer PvR. *J Am Chem Soc.* 1992; 114:1562. (d) Lucht BL, Collum DB. *Acc Chem Res.* 1999; 32:1035.
30. Benson SW. *J Am Chem Soc.* 1958; 80:5151. For additional leading references to statistical contributions in “redistribution reactions”, see: Fay RC, Lowry RN. *Inorg Chem.* 1974; 13:1309.
31. Mixtures of enolate **2b**, **2i**, and **3** solvated by TMEDA in toluene afford ensembles of homo- and heteroaggregates consistent with (Supporting Information).
32. (a) Lucht BL, Collum DB. *J Am Chem Soc.* 1995; 117:9863. (b) Scheschkewitz D. *Angew Chem, Int Ed Engl.* 2004; 43:2965. [PubMed: 15170317] (c) Niecke E, Nieger M, Schmidt O, Gudat D, Schoeller WW. *J Am Chem Soc.* 1999; 121:519. (d) Becker G, Eschbach B, Mundt O, Reti M, Niecke E, Issberner K, Nieger M, Thelen V, Noth H, Waldhor R, Schmidt M. *Z Anorg Allg Chem.* 1998; 624:469. (e) Becker G, Schwarz W, Seidler N, Westerhausen M. *Z Anorg Allg Chem.* 1992; 612:72. (f) Wang H, Wang H, Li HW, Xie Z. *Organometallics.* 2004; 23:875. (g) Xu X, Zhang Z, Yao Y, Zhang Y, Shen Q. *Inorg Chem.* 2007; 46:9379. [PubMed: 17880064] (h) Thiele K, Goris H, Imhof W, Seide W. *Z Anorg Allg Chem.* 2002; 628:107. (i) Ramirez A, Lobkovsky E, Collum DB. *J Am Chem Soc.* 2003; 125:15376. [PubMed: 14664582] (j) Buchalski P, Grabowska I, Kaminska E, Suwinska K. *Organometallics.* 2008; 27:2346.
33. Choi M-G, Brown TL. *Inorg Chem.* 1993; 32:1548.
34. Hsu DC, Lam PC-H, Slebodnick C, Carlier PR. *J Am Chem Soc.* 2009; 131:18168. [PubMed: 19921816]
35. For discussions of conformational equilibrations, see Clegg W, Liddle ST, Snaith R, Wheatley AEH. *New J Chem.* 1998; 22:1323. Boche G, Fraenkel G, Cabral J, Harms K, van Eikema Hommes NJR, Lohrenz J, Marsch M, Schleyer PvR. *J Am Chem Soc.* 1992; 114:1562. Remenar JF, Lucht BL, Kruglyak D, Romesberg FE, Gilchirst JH, Collum DB. *J Org Chem.* 1997; 62:5748.
36. Ulatowski F, D browa K, Bałakier T, Jurczak J. *J Org Chem.* 2016; 81:1746. [PubMed: 26866984]
37. (a) Yamataka, K., Yamada, H., Tomioka, H. *The Chemistry of Organolithium Compounds.* Rappoport, Z., Marek, I., editors. Vol. 2. Wiley; New York: 2004. p. 908. (b) Jackman LM, Lange BC. *J Am Chem Soc.* 1981; 103:4494. (c) Streitwieser A, Wang DZ. *J Am Chem Soc.* 1999; 121:6213. (d) Leung SSW, Streitwieser A. *J Org Chem.* 1999; 64:3390. [PubMed: 11674455] (e) Wang DZ, Kim YJ, Streitwieser A. *J Am Chem Soc.* 2000; 122:10754. (f) Kim YJ, Wang DZ. *Org Lett.* 2001; 3:2599. [PubMed: 11483070] (g) Streitwieser A, Leung SSW, Kim YJ. *Org Lett.* 1999; 1:145. [PubMed: 10822549] (h) Abbotto A, Leung SSW, Streitwieser A, Kilway KV. *J Am Chem Soc.* 1998; 120:10807. (i) Leung SSW, Streitwieser A. *J Am Chem Soc.* 1998; 120:10557. (j) Abu-Hasanayn F, Streitwieser A. *J Org Chem.* 1998; 63:2954. (k) Abu-Hasanayn F, Streitwieser A. *J Org Chem.* 1996; 118:8136. (l) Gareyev R, Ciula JC, Streitwieser A. *J Org Chem.* 1996; 61:4589.

[PubMed: 11667384] (m) Abu-Hasanayn F, Stratakis M, Streitwieser A. *J Org Chem.* 1995; 60:4688.(n) Dixon RE, Williams PG, Saljoughian M, Long MA, Streitwieser A. *Magn Reson Chem.* 1991; 29:509.

38. Casy BM, Flowers RA. *J Am Chem Soc.* 2011; 133:11492. [PubMed: 21721524]

39. Kofron WG, Baclawski LM. *J Org Chem.* 1976; 41:1879.

40. (a) Himmelsback, F., Austel, V., Pieper, H., Eisert, W., Mueller, T., Weisenberger, J., Linz, G., Krueger, GE. Preparation of 5-[(aminoaryloxy)methyl]-2-pyrrolidinoneacetates and analogs as drugs. EP 483667 A2 19920506. 1992. (b) Bodur C, Kutuk O, Karsli-Uzunbas G, Isimjan TT, Harrison P, Basaga H. *PloS one.* 2013; 8:e56369. [PubMed: 23441183]

Figure 1.

^6Li NMR spectra of 0.10 M solutions of **2b** and **3** in 12.3 M tetrahydrofuran (THF) at $-80\text{ }^\circ\text{C}$ with 0.13 M $[\text{}^6\text{Li}]\text{LDA}$ (LDA = lithium diisopropylamide). The measured²² mole fractions of **B** (X_{B}) are 0.00, 0.34, 0.54, 0.81, and 1.00 for (a) through (e), respectively.

Figure 2. Job plot showing relative integration of the ^6Li resonances versus the measured ^{22}Li mole fraction (X_B) of **3** for 0.10 M mixtures of **2b** (A_2) and **3** (B_2) in 12.3 M THF at -80°C .

Figure 3. ^6Li NMR spectra of 0.10 M solutions of **2b** and **3** in 0.13 M THF/toluene at $-95\text{ }^\circ\text{C}$ with 0.13 M $[^6\text{Li}]\text{LDA}$. The measured²² mole fractions of **2b** (X_{B}) are 0.00, 0.35, 0.52, 0.81, and 1.00 for (a) through (e), respectively.

Figure 4. Job plot showing the relative integrations of the ^6Li resonances versus the measured²² mole fractions (X_B) of **2b** for 0.10 M mixtures of **3** (**A₂**) and **2b** (**B₂**) in 0.13 M THF/toluene at -95°C .

Figure 5. ^6Li NMR spectra of 0.10 M solutions of **2j** in mixtures of THF and pyridine at 0.50 M total concentration with toluene as the cosolvent at -95°C with 0.105 M $[\text{}^6\text{Li}]\text{LDA}$. The measured mole fractions of pyridine (X_P) are 0.00, 0.50, 0.63, 0.75, and 1.00 for (a) through (e), respectively.

Figure 6. Job plot showing relative integration versus *measured* mole fraction (X_P) of pyridine for 0.50 M total solvent concentration for mixtures of pyridine (**P**) and THF (**T**) with 0.10 M **2j** at $-95\text{ }^{\circ}\text{C}$.

Figure 7. Job plot showing relative integration versus *intended* (or absolute) mole fraction (X_P) of pyridine for 0.50 M total solvent concentration in mixtures of pyridine (**P**) and THF (**T**) with 0.10 M **4** at -95°C .

Figure 8.

Plot of computed free energies of solvation (ΔG_{DFT} ; from Table 1, row 3) versus statistically corrected experimentally measured free energies of substitution (ΔG_{exp} ; from Table 1, row 2).

Figure 9.

Fit of aggregate relative dimer–tetramer concentrations versus THF concentration for enolate **3** at $-80\text{ }^{\circ}\text{C}$. The curve corresponds to a best fit to $y = k[\text{THF}]^n$ [eq 6; $k = (0.193 \pm 0.002)$, $n = 1.80 \pm 0.02$].

Scheme 1.

Scheme 2.

Chart 1.

Free energies of serial solvation by pyridine (see Scheme 1) as measured experimentally with the method of continuous variation (see Figure 7) and computationally with DFT.

Table 1

	G_1	G_2	G_3	G_4
Experimentally determined	Direct measurement	-0.86	0.03	0.48
	Statistically ^a corrected	-0.70	-0.13	-0.06
Computationally determined	-3.16	-2.77	0.16	0.01

^aStatistical correction removes the influence of the number of available sites for substitution, rendering the values directly comparable to the values computed with density functional theory.