Published in final edited form as:

ACS Chem Biol. 2016 November 18; 11(11): 2996–3001. doi:10.1021/acscchembio.6b00565.

Phenanthriplatin Acts as a Covalent Topoisomerase II Poison

Imogen A. Riddell^a, Ga Young Park^a, Keli Agama^b, Yves Pommier^{b,†}, and Stephen J. Lippard^{a,†}

^aDepartment of Chemistry, Massachusetts Institute of Technology, Cambridge, MA 02139

^bCenter for Cancer Research, National Cancer Institute, Bethesda, MD 20892

Abstract

Drugs capable of trapping topoisomerase II (Top2), an essential enzyme that cleaves double-stranded DNA in order to remove naturally occurring knots and tangles, can serve as potent anticancer agents. The monofunctional platinum anticancer agent phenanthriplatin, *cis*-[Pt(NH₃)₂(phenanthridine)Cl](NO₃), is shown here to act as a Top2 poison in addition to its known modes of action as a DNA and RNA polymerase inhibitor. Its potency therefore combines diverse modes of action by which phenanthriplatin can kill cancer cells. Covalent DNA-platinum lesions implicated in Top2 poisoning are distinctive from those generated by known topoisomerase poisons, which typically exert their action by intercalation at the topoisomerase cleavage site.

Graphical abstract

Introduction

Nuclear topoisomerases play an essential role in replication, transcription, repair, and chromosome organization. Specifically, topoisomerases resolve topological problems associated with DNA, unwinding supercoiled DNA and untangling inter- and intra-molecular DNA knots and catenanes. Two distinct classes of topoisomerase have been identified, type 1 and type 2, which include topoisomerase I (Top1)^{1–3} and topoisomerase II (Top2),^{4–7} respectively. These two classes of enzymes are distinguished by their differential

[†]Corresponding authors: lippard@mit.edu and pommier@nih.gov.

Supplemental Information

Detailed protocols for biochemical assays and additional experiments can be found in the Supplemental Information.

In vertebrates, Top2 exists as two closely related isoforms Top2 α and Top2 β .^{8,9} These two isoforms have greater than 70% amino acid similarity.^{4,7,9,10} Within the cell, they are differentially regulated and involved in dissimilar cellular processes. Top2 α , investigated in this study, is essential for cell growth and occurs in high levels in rapidly proliferating and cancer cells; its expression peaks at the G2/M phase of the cell cycle. By contrast, the concentration of the β isoform is more or less constant throughout the cell cycle.^{7,11,12}

Although this enzymatic action is essential to life, it is also an important target in cancer treatment. Under normal circumstances Top2cc is a transient intermediate, present at low steady state levels in the catalytic cycle, and the cleavage/religation equilibrium in which it is established is readily reversible, lying in favor of the ligation reaction.^{7,15} Top2 poisons exert their mode of action by binding to the Top2cc to generate a ternary complex^{13,18,19} that may either prevent religation or promote the forward cleavage reaction.¹⁵ The cytotoxic effects of this class of compounds do not result from stalling of the catalytic cycle at the Top2cc stage. Instead, the ternary complex blocks polymerases and downstream processing of the adducts, resulting in DNA fragmentation and ultimately cell death.^{10,19,20}

The early discovery that the Top2 catalytic cycle could be exploited in the development of anticancer drugs has resulted in significant research in this field, both in elucidating the mechanism of action of different Top2 poisons^{21,22} and inhibitors²³ and in designing novel complexes with better anticancer properties and reduced side effects.^{15,20,24} Topoisomerase poisons and inhibitors are used extensively in the clinics,^{4,7,25,26} with almost every form of cancer sensitive to chemotherapy being treated, at least partially, with drugs that target topoisomerases. The majority of Top2 drugs currently in the clinic do not distinguish between the two Top2 isoforms, but research indicates that the α form is the primary cytotoxic target.^{20,26–28}

ACS Chem Biol. Author manuscript; available in PMC 2017 January 20.

DNA³² polymerases along the DNA. In the present study we describe the results of experiments designed to test the hypothesis that a significant contribution to the potency of phenanthriplatin may be its ability to act as a Top2 poison.

Results and Discussion

Analysis of phenanthriplatin using the COMPARE algorithm of the NCI60 screening panel³³ and CellMiner³⁴ provided the initial indication that the monofunctional agent might be acting as a Top2 inhibitor.²⁹ The results, shown in Table 1, indicated that the effects of phenanthriplatin correlated well with those of known Top2 poisons and differed significantly from those of tubulin inhibitors and other N⁷-alkylating agents, including the commonly used platinum agents cisplatin, carboplatin and oxaliplatin.

Two independent methods were employed to test the formation of Top2cc in cancer cells. Both analyses indicated that treatment of HT-29 human colorectal adenocarcinoma cells with phenanthriplatin at increasing time intervals resulted in an increase in Top2cc, the transient, covalent DNA-Top2 complex normally observed at low levels in cells.

First, an in-vivo complex of enzyme (ICE) assay (Figure 1A) indicated that, after treatment with phenanthriplatin (50 μ M for 2 h), HT-29 cell extracts contained significantly more Top2cc than was observed in a control reaction that lacked a Top2 poison. Top2cc were observed down to 25 nM DNA concentrations, in contrast to the control reaction in which 100 nM DNA was required to observe Top2cc. In these experiments, etoposide (10 μ M) was employed as a positive control.

Alkaline elution assays were performed next to independently detect and quantify the formation of Top2cc as DNA-protein crosslinks (DPC).³⁵ It is well established that only DNA covalently bound to protein resists elution under the conditions of this experiment,^{36–38} therefore non-covalent polymerase-DNA interactions do not contribute to the amount of DNA retained by the protein-binding filter. Figure 1B clearly indicates that, as the concentration of phenanthriplatin in the HT-29 cell media increases from 1 – 100 μ M (constant 5 h), so too do the DPC.³⁵ This result reflects the total platinum concentration in the cell media and not the concentration involved in stabilization of the Top2-DNA complex. The effective concentration of phenanthriplatin available to stabilize crosslinks would be expected to be considerably lower than 50 μ M because a significant proportion of phenanthriplatin will bind covalently to DNA at sites not recognized by Top2.

The co-retention of protein and DNA in both the ICE assay and the alkaline elution experiments supports formation and persistence of Top2cc in the presence of phenanthriplatin. Research has shown that formation and stabilization of Top2cc is not toxic per se,³⁹ but it is the inability of polymerases to bypass this cleavage complex, combined with the downstream processing of the DPC, that results in fragmentation of genomic DNA and eventual cell death.^{4,19,40,41} Fluorescence activated cell sorting (FACS) analysis of the HT-29 cells treated with varying concentrations of phenanthriplatin support G1/S arrest (Supplemental Figure S2), as expected when DNA synthesis is impaired. Accordingly, one of the major modes of action by which phenanthriplatin exerts its anticancer activity is in

blocking RNA and DNA polymerases as a covalent Pt-DNA adduct. The FACS analysis most likely represents contributions from both of these modes of action.

Observation of Top2cc supports inhibition of Top2 by a conventional Top2 poisoning mechanism in which phenanthriplatin prevents religation during the cleavage/ligation equilibrium. In contrast, no covalent Top2-DNA intermediate is observed with cisplatin,⁴² and its activity profile is not correlated with known Top2 poisons. Platinum complexes that are currently established as Top2 poisons^{10,43} are known DNA intercalators, and for this reason their molecular mode of action has been postulated to be the same as traditional Top2 poisons such as etoposide. Whereas the inability of phenanthriplatin to intercalate into free DNA²⁹ does not preclude intercalation of phenanthriplatin in the presence of both Top2 and DNA,¹³ we believe the strong affinity of phenanthriplatin for N⁷-guanosine³² makes covalent bond formation more likely.

When the small molecule single-crystal structure of an alkylguanine-phenanthriplatin adduct³² was superimposed into the crystal structure of the etoposide-DNA-Top2 ternary complex¹³ (Figure 2), the phenanthridine ring was easily accommodated within the binding pocket known to house the glycosidic moiety of etoposide, indicating that the proposed platinated DNA-Top2 complex is sterically viable.

Having identified phenanthriplatin as a Top2 poison we were interested in determining the sequence requirements for DNA cleavage by Top2 α in the presence of phenanthriplatin. The nucleotide sequences for several different classes of drug-stimulated Top2 cleavage sites have been elucidated and shown to be strongly complex specific.^{21,22,44} A 117-bp DNA fragment derived from the 161-bp DNA fragment of pBluescript SK-phagemid was 5'-end-labeled with ³²P-ATP and incubated with phenanthriplatin at a range of concentrations and with recombinant Top2 for 3 h. Linear amplification was performed using standard conditions and the transcription stalling sites were visualized and compared to mapped binding sites of phenanthriplatin on the same 117-bp sequence (Figure 3; Supplemental Figures S3, S4 and S5).

The observation of common Top2 cleavage and phenanthriplatin binding sites supports the hypothesis that, when phenanthriplatin is bound at a guanosine within the active site of the Top2-DNA complex, it is capable of stabilizing the Top2cc intermediate and thereby exerting its action as a Top2 poison. That phenanthriplatin, and not cisplatin, forms adducts at nucleobases, which are also important with respect to Top2 cleavage is easily rationalized when considering that phenanthriplatin is a monofunctional platinum agent, whereas cisplatin generates bifunctional adducts, and within the region of interest there are few opportunities for bifunctional adduct formation.

Conclusions

The observation that phenanthriplatin shares more in common with Top2 poisons than other platinum anticancer agents led us to investigate the possibility that, in addition to acting as a polymerase roadblock,²⁹⁻³² phenanthriplatin may also inhibit the action of topoisomerases. Stabilization of the Top2cc intermediate was confirmed by using alkaline elution and ICE

bioassays, confirming that the catalytic cycle of the Top2 is stalled during the cleavage/religation inhibition, and that phenanthriplatin is indeed a Top2 poison. Furthermore, the observation of mutual Top2 cleavage and platination sites leads us to postulate that phenanthriplatin is bound covalently within the active site of the DNA:Top2 complex in such a way as to prevent religation of the cleaved DNA strands. The Top2 cleavage sites are also observed to be monofunctional platination sites explaining why phenanthriplatin, a monofunctional platination agent, and not cisplatin, a bifunctional agent, is observed to act as a Top2 poison.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work is supported by the NCI under grant CA034992 (S.J.L.). G.Y.P. received support from a Misrock Fellowship. KA and YP are supported by the Intramural Program of the National Cancer Institute, Center for Cancer Research (Z01-BC006161).

References

1. Champoux JJ, Dulbecco R. An activity from mammalian cells that untwists superhelical DNA—a possible swivel for DNA replication (polyoma-ethidium bromide-mouse-embryo cells-dye binding assay). *Proc Natl Acad Sci USA*. 1972; 69:143–146. [PubMed: 4333036]
2. Champoux JJ. DNA topoisomerases: Structure, function, and mechanism. *Annu Rev Biochem*. 2001; 70:369–413. [PubMed: 11395412]
3. Pommier Y. Topoisomerase I inhibitors: camptothecins and beyond. *Nat Rev Cancer*. 2006; 6:789–802. [PubMed: 16990856]
4. Pommier Y, Leo E, Zhang H, Marchand C. DNA topoisomerases and their poisoning by anticancer and antibacterial drugs. *Chem Biol*. 2010; 17:421–433. [PubMed: 20534341]
5. Wang JC. DNA topoisomerases: why so many? *J Biol Chem*. 1991; 266:6659–6662. [PubMed: 1849888]
6. McClendon AK, Osheroff N. DNA topoisomerase II, genotoxicity, and cancer. *Mutat Res*. 2007; 623:83–97. [PubMed: 17681352]
7. Nitiss JL. DNA topoisomerase II and its growing repertoire of biological functions. *Nat Rev Cancer*. 2009; 9:327–337. [PubMed: 19377505]
8. Chung TD, Drake FH, Tan KB, Per SR, Crooke ST, Mirabelli CK. Characterization and immunological identification of cDNA clones encoding two human DNA topoisomerase II isozymes. *Proc Natl Acad Sci U S A*. 1989; 86:9431–9435. [PubMed: 2556712]
9. Austin CA, Sng JH, Patel S, Fisher LM. Novel HeLa topoisomerase II is the II beta isoform: complete coding sequence and homology with other type II topoisomerases. *Biochimica et biophysica acta*. 1993; 1172:283–291. [PubMed: 8383537]
10. Deweese JE, Osheroff N. The DNA cleavage reaction of topoisomerase II: wolf in sheep's clothing. *Nucleic Acids Res*. 2009; 37:738–748. [PubMed: 19042970]
11. Woessner R, Mattern M, Mirabelli C, Johnson R, Drake F. Proliferation- and cell cycle-dependent differences in expression of the 170 kilodalton and 180 kilodalton forms of topoisomerase II in NIH-3T3 cells. *Cell Growth Differ*. 1991; 2:209–214. [PubMed: 1651102]
12. Li TK, Liu LF. Tumor cell death induced by topoisomerase-targeting drugs. *Annu Rev Pharmacol Toxicol*. 2001; 41:53–77.
13. Wu C-C, Li T-K, Farh L, Lin L-Y, Lin T-S, Yu Y-J, Yen T-J, Chiang C-W, Chan N-L. Structural Basis of Type II Topoisomerase Inhibition by the Anticancer Drug Etoposide. *Science*. 2011; 333:459–462. [PubMed: 21778401]

14. Berger JM, Fass D, Wang JC, Harrison SC. Structural similarities between topoisomerases that cleave one or both DNA strands. *Proc Natl Acad Sci USA*. 1998; 95:7876–7881. [PubMed: 9653108]
15. Burden DA, Osheroff N. Mechanism of action of eukaryotic topoisomerase II and drugs targeted to the enzyme. *BBA - Gene Struct Expr*. 1998; 1400:139–154.
16. Dong KC, Berger JM. Structural basis for gate-DNA recognition and bending by type IIA topoisomerases. *Nature*. 2007; 450:1201–1205. [PubMed: 18097402]
17. Mueller-Planitz F, Herschlag D. Coupling between ATP Binding and DNA cleavage by DNA topoisomerase II: a unifying kinetic and structural mechanism. *J Biol Chem*. 2008; 283:17463–17476. [PubMed: 18403371]
18. Pommier Y, Marchand C. Interfacial inhibitors: targeting macromolecular complexes. *Nat Rev Drug Discov*. 2012; 11:25–36.
19. Nitiss JL. Targeting DNA topoisomerase II in cancer chemotherapy. *Nat Rev Cancer*. 2009; 9:338–350. [PubMed: 19377506]
20. Pommier Y. Drugging topoisomerases: lessons and challenges. *ACS Chem Biol*. 2013; 8:82–95. [PubMed: 23259582]
21. Capranico G, Kohn KW, Pommier Y. Local sequence requirements for DNA cleavage by mammalian topoisomerase II in the presence of doxorubicin. *Nucleic Acids Res*. 1990; 18:6611–6619. [PubMed: 2174543]
22. Pommier Y, Capranico G, Orr A, Kohn KW. Local base sequence preferences for DNA cleavage by mammalian topoisomerase II in the presence of amsacrine or teniposide. *Nucleic Acids Res*. 1991; 19:5973–5980. [PubMed: 1658748]
23. Andoh T, Ishida R. Catalytic inhibitors of DNA topoisomerase II. *Biochimica et biophysica acta*. 1998; 1400:155–171. [PubMed: 9748552]
24. Wu C-C, Li Y-C, Wang Y-R, Li T-K, Chan N-L. On the structural basis and design guidelines for type II topoisomerase-targeting anticancer drugs. *Nucleic Acids Res*. 2013; 41:10630–10640. [PubMed: 24038465]
25. Hande KR. Clinical applications of anticancer drugs targeted to topoisomerase II. *BBA - Gene Struct Expr*. 1998; 1400:173–184.
26. Chen W, Qiu J, Shen YM. Topoisomerase II α , rather than II β , is a promising target in development of anti-cancer drugs. *Drug Discoveries Ther*. 2012; 6:230–237.
27. Azarova AM, Lyu YL, Lin CP, Tsai YC, Lau JY, Wang JC, Liu LF. Roles of DNA topoisomerase II isozymes in chemotherapy and secondary malignancies. *Proc Natl Acad Sci U S A*. 2007; 104:11014–11019. [PubMed: 17578914]
28. Cowell IG, Sondka Z, Smith K, Lee KC, Manville CM, Sidorczuk-Lesthuruge M, Rance HA, Padgett K, Jackson GH, Adachi N, Austin CA. Model for MLL translocations in therapy-related leukemia involving topoisomerase II β -mediated DNA strand breaks and gene proximity. *Proc Natl Acad Sci U S A*. 2012; 109:8989–8994. [PubMed: 22615413]
29. Park GY, Wilson JJ, Song Y, Lippard SJ. Phenanthriplatin, a monofunctional DNA-binding platinum anticancer drug candidate with unusual potency and cellular activity profile. *Proc Natl Acad Sci U S A*. 2012; 109:11987–11992. [PubMed: 22773807]
30. Wang D, Zhu G, Huang X, Lippard SJ. X-ray structure and mechanism of RNA polymerase II stalled at an antineoplastic monofunctional platinum-DNA adduct. *Proc Natl Acad Sci U S A*. 2010; 107:9584–9589. [PubMed: 20448203]
31. Kellinger MW, Park GY, Chong J, Lippard SJ, Wang D. Effect of a Monofunctional Phenanthriplatin-DNA Adduct on RNA Polymerase II Transcriptional Fidelity and Translesion Synthesis. *J Am Chem Soc*. 2013; 135:13054–13061. [PubMed: 23927577]
32. Gregory MT, Park GY, Johnstone TC, Lee YS, Yang W, Lippard SJ. Structural and mechanistic studies of polymerase η bypass of phenanthriplatin DNA damage. *Proc Natl Acad Sci U S A*. 2014; 111:9133–9138. [PubMed: 24927576]
33. Shoemaker RH. The NCI60 human tumour cell line anticancer drug screen. *Nat Rev Cancer*. 2006; 6:813–823. [PubMed: 16990858]

34. Reinhold WC, Sunshine M, Liu H, Varma S, Kohn KW, Morris J, Doroshow J, Pommier Y. CellMiner: a web-based suite of genomic and pharmacologic tools to explore transcript and drug patterns in the NCI-60 cell line set. *Cancer Res.* 2012; 72:3499–3511. [PubMed: 22802077]
35. Kohn KW. Beyond DNA cross-linking: history and prospects of DNA-targeted cancer treatment-fifteenth Bruce F. Cain Memorial Award Lecture. *Cancer Res.* 1996; 56:5533–5546. [PubMed: 8971150]
36. Zwelling LA, Michaels S, Erickson LC, Ungerleider RS, Nichols M, Kohn KW. Protein-associated deoxyribonucleic acid strand breaks in L1210 cells treated with the deoxyribonucleic acid intercalating agents 4'-(9-acridinylamino)methanesulfon-m-aniside and adriamycin. *Biochemistry.* 1981; 20:6553–6563. [PubMed: 6895473]
37. Oleinick NL, Chiu SM, Ramakrishnan N, Xue LY. The formation, identification, and significance of DNA-protein cross-links in mammalian cells. *Br J Cancer.* 1987; 8:135–140.
38. Zhitkovich A, Costa M. A simple, sensitive assay to detect DNA–protein crosslinks in intact cells and in vivo. *Carcinogenesis.* 1992; 13:1485–1489. [PubMed: 1499101]
39. Puc J, Kozbial P, Li W, Tan Y, Liu Z, Suter T, Ohgi KA, Zhang J, Aggarwal AK, Rosenfeld MG. Ligand-dependent enhancer activation regulated by topoisomerase-I activity. *Cell.* 2015; 160:367–380. [PubMed: 25619691]
40. Holm C, Covey JM, Kerrigan D, Pommier Y. Differential requirement of DNA replication for the cytotoxicity of DNA topoisomerase I and II inhibitors in Chinese hamster DC3F cells. *Cancer Res.* 1989; 49:6365–6368. [PubMed: 2553254]
41. D'Arpa P, Beardmore C, Liu LF. Involvement of nucleic acid synthesis in cell killing mechanisms of topoisomerase poisons. *Cancer Res.* 1990; 50:6919–6924. [PubMed: 1698546]
42. Hasinoff BB, Wu X, Krokhin OV, Ens W, Standing KG, Nitiss JL, Sivaram T, Giorgianni A, Yang S, Jiang Y, Yalowich JC. Biochemical and proteomics approaches to characterize topoisomerase II α cysteines and DNA as targets responsible for cisplatin-induced inhibition of topoisomerase II α . *Mol Pharmacol.* 2005; 67:937–947. [PubMed: 15602006]
43. Liu J, Leung CH, Chow ALF, Sun RWY, Yan SC, Che CM. Cyclometalated platinum(II) complexes as topoisomerase II α poisons. *Chem Commun.* 2011; 47:719–721.
44. Tewey K, Rowe T, Yang L, Halligan B, Liu L. Adriamycin-induced DNA damage mediated by mammalian DNA topoisomerase II. *Science.* 1984; 226:466–468. [PubMed: 6093249]

Figure 1.

Induction of Top2 cleavage complexes by phenanthriplatin in human colon HT-29 cells. **A)** Cells were treated with 10 μ M of etoposide (VP-16) for 3 h, or 50 μ M phenanthriplatin (Phen-Pt) for 1, 2, 3 or 5 h. Equal numbers of cells were lysed in DNAzol reagent and submitted to the ICE assay. Different concentrations of genomic DNA (0.2, 0.1, 0.05 and 0.025 μ g) were probed with an anti-Top2 antibody. Etoposide was used as a positive control (0.2, 0.1, 0.05 and 0.025 μ g) for Top2cc formation. **B)** Cells treated with phenanthriplatin for 5 h were detected by the alkaline elution assay. Quantification of DNA–Top2 cross-linking (in rad equivalents [rad-eq]). Data at each concentration of phenanthriplatin represent three independent determinations.

Figure 2.

Phenanthriplatin can be accommodated within the active site of the DNA:Top2cc when bound through the N⁷ position of the guanine adjacent to the site of DNA cleavage. Pymol images showing **A**) phenanthridine ring colored in yellow; platinum atom grey; **B**) the same view also showing etoposide (orange) bound within the active site of the DNA:Top2cc. Original structure with etoposide from PDB 3QX3.

Figure 3.

Chart showing the positions of mapped Top2-linked DNA breaks and platinum adducts. Covalently linked nucleobase colored red, phenanthriplatin adducts closed circles, and cisplatin adducts open squares on the 117-bp strand investigated.

Table 1

Correlation between phenanthriplatin and Top2 inhibitors (T2, red), N⁷-alkylating agents (A7, blue), and tubulin inhibitors (Tu, green) determined using the COMPARE algorithm of the NCI60 screening panel.³³ Significantly higher correlations were observed between the Top2 inhibitors than with the other platinum agents.

NSC	Name	MOA	Correlation
759703	Phenanthriplatin	-	1
123127	Doxorubicin	T2	0.519
301739	Mitoxantrone	T2	0.517
349174	Oxanthrazole	T2	0.494
82151	Daunorubicin	T2	0.446
164011	Rubidazole	T2	0.438
337766	Bisantrene hydrochloride	T2	0.433
246131	Valrubicin	T2	0.381
249992	M-AMSA	T2	0.358
122819	Teniposide	T2	0.336
256439	Idarubicin	T2	0.297
141540	Etoposide	T2	0.286
707389	Eribulin mesilate/mesylate	T2	0.282
178248	Chlorozotocin	A7	0.254
95466	PCNU	A7	0.232
750	Busulfan	A7	0.205
135758	Piperazinedione	A7	0.203
9706	Triethylenemelamine	A7	0.202
353451	Mitozolomide	A7	0.201
338947	Clomesone	A7	0.199
182986	AZQ	A7	0.195
119875	Cisplatin	A7	0.185
6396	Thiotepa	A7	0.178
241240	Carboplatin	A7	0.175
25154	Pipobroman	A7	0.172
762	Mechlorethamine	A7	0.143
34462	Uracil mustard	A7	0.14
3088	Chlorambucil	A7	0.138
8806	Melphalan	A7	0.108
138783	Bendamustine	A7	0.097
67574	Vincristine sulfate	Tu	0.095
77213	Procarbazine hydrochloride	A7	0.093
363812	Tetraplatin	A7	0.046
409962	Carmustine	A7	0.046

NSC	Name	MOA	Correlation
628503	Docetaxel, Docetaxol	Tu	0.041
109724	Ifosfamide	A7	0.013
266046	Oxaliplatin	A7	0.006
271674	Carboxyphthalatoplatinum	A7	-0.013
85998	Streptozocin	A7	-0.112
45388	Dacarbazine	A7	-0.301
26271	Cyclophosphamide	A7	-0.382