Received:
1 September 2015

Revised:
12 November 2015

Accepted:
9 March 2016

<http://dx.doi.org/10.1259/bjr.20150724>

Cite this article as:

Seo Y, Tamari K, Yoshioka Y, Isohashi F, Suzuki O, Hayashi K, et al. Characterization of *in vitro* radiosensitization in mammalian cells using biomathematical modelling: implications for hypofractionated radiotherapy with a combined modality approach. *Br J Radiol* 2016; **89**: 20150724.

FULL PAPER

Characterization of *in vitro* radiosensitization in mammalian cells using biomathematical modelling: implications for hypofractionated radiotherapy with a combined modality approach

YUJI SEO, MD, KEISUKE TAMARI, MD, PhD, YASUO YOSHIOKA, MD, PhD, FUMIAKI ISOHASHI, MD, PhD, OSAMU SUZUKI, MD, PhD, KAZUHIKO HAYASHI, MD, PhD, YUTAKA TAKAHASHI, PhD, SUNGJAE BAEK, MD, KEISUKE OTANI, MD and KAZUHIKO OGAWA, MD, PhD

Department of Radiation Oncology, Osaka University Graduate School of Medicine, Osaka, Japan

Address correspondence to: Dr Yuji Seo
E-mail: seo@radonc.med.osaka-u.ac.jp

Objective: It is unclear whether radiosensitization is beneficial when radiotherapy is administered at a high dose per fraction. The aim of this study was to assess the impact of radiation dose on the effectiveness of a broad range of radiosensitizers.

Methods: We analyzed 653 pairs of clonogenic survival curves in 285 published articles, in which modifications of radiosensitivity were studied using the colony-forming assay. The modifications of radiosensitivity were arbitrarily classified into 20 classes. The survival curves were fitted to two biomathematical models: the linear-quadratic model and the repair-misrepair (RMR) model.

Results: We found that radiosensitization was predominantly characterized by an increase of the α value (α -sensitization) without an increase of the β value (β -sensitization). A subset analysis revealed that all 20 classes showed significant α -sensitization. In contrast, only oxygen/hypoxic sensitizers (oxygen) and

poly(adenosine diphosphate-ribose) polymerase inhibition (PARPi) exhibited β -sensitization. An analysis using the RMR model revealed two major sources of radiosensitization: an increased residual DNA lesion through repair inhibition and a shift from linear repairs to quadratic misrepairs, leading to enhanced lethal chromosomal aberrations.

Conclusion: Oxygen and PARPi were found to show β -sensitization, which was favourable for eliciting a comparable degree of sensitization in the higher dose range. Reduced fidelity of the repair was suggested to be a possible mechanism of β -sensitization. Further study targeting β -sensitization is needed to develop a novel combined modality therapy with high-dose-per-fraction radiotherapy.

Advances in knowledge: Radiosensitization can be classified into two groups, α - and β -sensitizations. These two phenomena may stem from distinct underlying mechanisms.

INTRODUCTION

Oncologic outcomes have been shown to improve in various types of malignant tumours with the administration of chemotherapy and/or molecular-targeted therapy in addition to radiotherapy.¹ Numerous data in cell-based studies and pre-clinical animal studies have indicated that the enhanced efficacy of combined modality therapy can be attributed, at least in part, to synergistic cancer cell killings. Regarding radiation dose-fractionation schedules, conventional fractionation with 1.8–2 Gy per fraction once daily is the standard approach in definitive treatments, especially when systemic therapy is concurrently administered. Therefore, most research in radiosensitization has focused on

relatively low doses of radiation. On the other hand, a recent development in intensity-modulated radiotherapy and image-guided radiotherapy has enabled the safe delivery of higher doses per fraction, and hypofractionated treatments are becoming more and more common. High-precision radiotherapy improves the therapeutic ratio by conformal delivery of radiation to targets and physically spares the surrounding normal tissues. Combined therapy with such high-precision radiotherapy and systemic agents may further improve the therapeutic ratio. However, it is unclear whether combined treatment with a radiosensitizer is effective when radiotherapy is given at a high dose per fraction.

The linear-quadratic (LQ) model is based on the theory of dual radiation action, which refers to the formation of sublesions and subsequent interactions between pairs of sublesions to produce lethal lesions.² The LQ model is currently the *de facto* standard for analyzing radiation dose responses and is often utilized to convert one dose-fractionation schedule to another through the concept of biologically effective doses. A degree of radiosensitization can also be evaluated on the basis of changes in the LQ model parameters between the absence and presence of radiosensitizers.³

The repair–misrepair (RMR) model is another method used to fit acute radiation dose responses. The LQ and RMR models are conceptually similar in the sense that both models are based on chromosome aberration data and the lesion interaction hypothesis. However, unlike the LQ model, the RMR model assumes that the initial generation of lesions and subsequent molecular biological responses are distinct processes.⁴ Therefore, the biological responses following radiation-induced DNA damage can be quantitated by the RMR model. Although the RMR model is rarely used in clinical radiation oncology, we believe that it is useful for gaining insights into the underlying mechanisms of radiosensitization.

Here, we attempt to characterize radiosensitization as a function of radiation doses by analyzing a large clonogenic survival data set with various radiosensitizing agents in mammalian cells. Using the LQ and RMR models, we show that synergistic modifications of radiosensitivity can be classified into two groups, α - and β -sensitization, and that these two phenomena may stem from distinct underlying mechanisms.

METHODS AND MATERIALS

Data collection and review of published data

We systematically reviewed clonogenic survival curves in published articles in which pharmaceutical and/or genetic modification(s) of radiosensitivity were studied in mammalian cells. The data were collected by searching the PubMed database using the medical subject headings (MeSH) terms “radiation tolerance” or “radiation-sensitizing agents”. We selected studies using the following inclusion criteria: the standard colony-forming assay was used; the survival curves were plotted at multiple dose points; the dose range from 2 to 10 Gy was primarily focused on except under hypoxic conditions; and mechanism(s) of action appeared to be known or clearly claimed. All data points of the figures that fulfilled the criteria were digitized using ImageJ 1.48 for Windows (US National Institutes of Health, Bethesda, MD; <http://imagej.nih.gov/ij>) for statistical analysis.

Definition of additivity

Interactions between radiation and drugs were estimated relative to a definition of additivity widely accepted in cancer research.⁵ Bliss independence (survival probability independence) exists if the combined modality survival probability equals the product of the single-modality survival

probabilities; e.g. if the surviving fraction (SF) following radiation and treatment with drug A is:

$$SF^{R,A} = SF^R \times SF^A$$

or equivalent in terms of fraction affected (Fa) (using $SF = 1 - Fa$), if

$$Fa^{R,A} = Fa^R + (1 - Fa^R) \times Fa^A$$

Linear-quadratic model

The survival curves were fitted to the standard LQ model in the absence [Equation (1)] and presence [Equation (2)] of radiosensitizers.

$$SF^R = e^{-\alpha_0 D - \beta_0 D^2} \quad (1)$$

$$SF^{R,A} / SF^A = e^{-\alpha_1 D - \beta_1 D^2} \quad (2)$$

A relative comparison between the surviving fraction following radiation and treatment with drug A ($SF^{R,A}$)/drug A-induced cytotoxicity (SF^A) and the surviving fraction following radiation alone (SF^R) estimates radiosensitization. Since most investigators had normalized the data to drug-induced cytotoxicity (SF^A) in their survival plots, direct drug cytotoxicity was not available for analyses in this study.

To evaluate the degree of radiosensitization, the dose-modifying factor (DMF) was defined as the ratio of the isoeffective radiation dose in the absence and presence of radiosensitization. An isoeffective radiation dose that corresponded to each observed data point was estimated using the derived LQ model parameters in the range of survival fractions from the mathematical constant e^{-1} to e^{-8} . Thus, multiple DMFs were calculated from one pair of survival curves. The definition of the DMF was analogous to that of relative biologic effectiveness to characterize high linear energy transfer radiation effects. Variations of the DMFs may behave in a similar way to the relative biologic effectiveness. As opposed to point-to-point evaluations, the mean DMF (mDMF) was defined as the ratio of the mean inactivation dose in the absence ($MID_{control}$) and presence ($MID_{modified}$) of radiosensitization to evaluate the average radiosensitizing effect throughout the entire dose range [Equation (3)]. The mean inactivation dose was calculated using the area under the survival curve [Equation (4)].⁶

$$mDMF = MID_{control} / MID_{modified} \quad (3)$$

$$MID = \int_0^{\infty} e^{-\alpha x - \beta x^2} dx \quad (4)$$

Similar to DMF, we also defined the sensitizer enhancement ratio (SER) as the ratio of the log survival fraction in the presence and absence of radiosensitization. The SER was calculated at multiple isodose pairs of the observed data points in the range from 2 to 12 Gy.

Repair-misrepair model

We used the RMR model to elucidate the underlying mechanisms of radiosensitization in each class of radiosensitization. The RMR model assumes that the initial yield of lesions per cell [uncommitted lesion (U_0)] is proportional to the dose (D).

$$U_0 = \delta D \quad (5)$$

The quantity, $U(t)$, indicates the mean number of lesions per cell as a function of time.

$$\frac{dU}{dt} = -\lambda U - \kappa U^2$$

λ is the coefficient for linear self-repair, while κ is the coefficient for cooperative repairs through interactions of pairs of U lesions. After the time (t) required for the repair processes, the number of residual uncommitted lesions (rU) is:

$$U(t) = \frac{\delta(1-T)D}{1 + \frac{\delta TD}{\varepsilon}} \text{ where } T = 1 - e^{-\lambda t} \text{ and } \varepsilon = \lambda/\kappa \quad (6)$$

The total number of linear self-repairs, $R_L(t)$, is:

$$R_L(t) = \int_0^t \lambda U dt = \varepsilon \log \left\{ 1 + \frac{\delta TD}{\varepsilon} \right\}$$

The total number of quadratic misrepairs, $R_Q(t)$, is:

$$R_Q(t) = \int_0^t \kappa U^2 dt = \frac{\delta TD \left(1 + \frac{\delta D}{\varepsilon} \right)}{1 + \frac{\delta TD}{\varepsilon}} - \varepsilon \log \left\{ 1 + \frac{\delta TD}{\varepsilon} \right\}$$

Linear self-repair may not be a perfect process, and linear misrepair (mR_L) may occur, resulting in cell death. When the

Table 1. Mean inactivation doses (MID) and mean dose-modifying factors (mDMF) in 20 radiosensitization classes [median (interquartile range)]

Sensitization class	N	MID (Gy)		mDMF
		Control	Sensitizer	
Apoptosis	19	2.8 (2.32–3.51)	1.83 (1.58–2.58)	1.34 (1.21–1.49)
Cell cycle	29	2.65 (1.86–3.6)	1.79 (1.37–2.48)	1.38 (1.19–1.66)
COXi	32	2.99 (2.46–3.37)	2.15 (1.81–2.65)	1.37 (1.17–1.64)
DNA-alkylation	19	3.31 (2.72–3.42)	2.3 (2.15–2.91)	1.38 (1.08–1.48)
DNA-dsb-repair	12	2.58 (2.12–3.43)	1.42 (1.23–1.97)	1.7 (1.57–1.89)
DNA-dsb-signal	70	2.83 (2.48–3.22)	1.74 (1.35–2.19)	1.59 (1.37–1.97)
EGFR/HER2	55	2.85 (2.59–3.19)	2.19 (1.79–2.76)	1.29 (1.12–1.53)
Gemcitabine	28	3.24 (2.64–3.65)	2 (1.54–2.41)	1.59 (1.35–1.76)
HDACi	37	3.13 (2.75–4.56)	1.95 (1.41–2.44)	1.57 (1.43–1.96)
IUdR/BUdR	20	2.98 (2.32–3.67)	1.69 (1.18–2.17)	1.97 (1.52–2.4)
Oxygen	35	7.74 (5.34–9.97)	3.06 (2.46–3.72)	2.44 (2.26–2.95)
PARPi	32	2.86 (2.46–3.45)	2.1 (1.59–2.33)	1.35 (1.23–1.52)
PDGFR	11	3.59 (3.04–4.88)	2.6 (1.97–2.96)	1.47 (1.31–1.56)
PI3K/AKT/mTOR	73	3.34 (2.47–3.82)	2.24 (1.75–2.87)	1.36 (1.26–1.53)
Platinum	30	2.97 (2.51–3.61)	2.38 (1.9–3.83)	1.31 (1.16–1.38)
RAS/MAPK/MEK	31	3.02 (2.73–3.53)	2.25 (2.03–2.84)	1.23 (1.13–1.39)
RRi	17	3.07 (2.66–3.4)	1.77 (1.56–2.39)	1.62 (1.42–1.82)
Taxane	37	2.84 (2.26–3.38)	1.92 (1.51–2.57)	1.31 (1.18–1.67)
TOPOi	31	3.03 (2.24–4.23)	2.07 (1.62–2.69)	1.32 (1.17–1.57)
TSi	35	3.37 (2.73–3.86)	2.35 (1.7–2.95)	1.35 (1.19–1.79)

AKT, v-akt murine thymoma viral oncogene homolog; BUdR, bromodeoxyuridine; COXi, cyclooxygenase inhibition; dsb, double-strand break; EGFR, epidermal growth factor receptor; HDACi, histone deacetylase inhibition; HER2, human epidermal growth factor receptor 2; IUdR, iododeoxyuridine; MAPK, mitogen-activated protein kinase; MEK, MAPK kinase; mTOR, mammalian target of rapamycin; N, number of pairs of survival curves; PARPi, poly(adenosine diphosphate-ribose) polymerase inhibition; PDGFR, platelet-derived growth factor receptor; PI3K, phosphoinositide 3-kinase; RAS, rat sarcoma viral oncogene homolog; RRi, ribonucleotide reductase inhibition; TOPOi, topoisomerase I/II inhibition; TSi, thymidylate synthase inhibition.

probability of complete linear repair is φ , linear eurepair (eR_L) and mR_L are:

$$eR_L(t) = \varphi R_L(t) \quad (7)$$

$$mR_L(t) = (1 - \varphi) R_L(t) \quad (8)$$

Conversely, the surviving cells may contain chromosome aberrations caused by quadratic misrepair. When the probability of viable quadratic misrepair (vmR_Q) is ψ , vmR_Q and non-viable quadratic misrepair (nmR_Q) are:

$$vmR_Q(t) = \psi R_Q(t) \quad (9)$$

$$nmR_Q(t) = (1 - \psi) R_Q(t) \quad (10)$$

An RMR model with five parameters is derived as the probability that there are no lethal lesions in a cell.⁴

$$SF = e^{-\{U(t) + (1 - \varphi)R_L(t) + (1 - \psi)R_Q(t)\}} \\ = e^{-\{\delta D - \psi(\delta D - U(t))\}} \left\{ 1 + \frac{\delta TD}{\varepsilon} \right\}^{(\varphi - \psi)\varepsilon} \quad (11)$$

Since quadratic repair has a greater potential risk of causing asymmetric aberrations than linear repair, the coefficient φ was constrained to be greater than ψ .

Statistical analysis

Curve fitting to the models was performed by maximum likelihood estimation using the statistical software R 3.1.1 for Windows® (R Foundation for Statistical Computing, Vienna, Austria; <http://www.R-project.org>) with bbmle package. Statistical significances of differences between two groups were assessed by the Wilcoxon signed-rank test. Correlations between two variables were examined by Spearman's rank correlation test, and the resulting p -values were adjusted for multiple comparisons by the Benjamini–Hochberg procedure.

RESULTS

From our extensive search of publications, we collected and digitized experimental data from 653 pairs of survival curves in 285 articles (shown in [Supplemental file 1](#)). In total, there were 197 cell lines (163 human and 34 rodents). Of the 197 cell lines, 27 were derived from normal tissues, and the remaining 170 were cancer cells. We classified genetic or pharmacologic modifications of radiosensitivity into 20 groups ([Table 1](#)), which included major classes of conventional radiosensitizers and chemotherapeutic agents: halogenated pyrimidine analogues [iododeoxyuridine/bromodeoxyuridine (IUdR/BUdR)], hypoxic radiosensitizers/oxygen (oxygen), DNA-alkylating agents (DNA-alkylation), gemcitabine, platinum, ribonucleotide reductase inhibition (RRI), taxane, thymidylate synthase inhibition (TSi) and topoisomerase I/II inhibition (TOPOi). There were substantial amounts of experimental data with cyclooxygenase inhibition (COXi), histone deacetylase and poly[adenosine diphosphate (ADP)-ribose] polymerase inhibition (PARPi);

Figure 1. Representative pairs of survival curves fitted to the data points extracted from the 142nd reference in the [Supplemental file 1](#) (solid lines, without radiosensitization; dashed lines, with radiosensitization). The plots (a, b) show the fitted curves by the linear-quadratic (LQ) model and the repair-misrepair (RMR) model, respectively. Derived model parameters are shown in the plots. C, without radiosensitization; S, with sensitization; SF, surviving fraction.

therefore, we analyzed these three targets as specific classes of radiosensitization. The other molecular targets were grouped according to the Kyoto Encyclopedia of Genes and Genomes: apoptosis, homologous recombination/non-homologous end-joining [DNA-double-strand break (dsb)-repair], p53 signalling pathway (DNA-dsb-signal), cell cycle, avian erythroblastosis oncogene B (ErbB) signalling pathway [epidermal growth factor receptor/human epidermal growth factor receptor 2 (EGFR/HER2)], platelet-derived growth factor receptor (PDGFR), phosphoinositide 3-kinase (PI3K)-v-akt murine thymoma viral oncogene homolog (AKT) and mammalian target of rapamycin (mTOR) signalling pathway (PI3K/AKT/mTOR) and mitogen-activated protein kinase (MAPK) signaling pathway [rat sarcoma viral oncogene homolog (RAS)/MAPK/MAPK kinase (MEK)].

Analysis using linear-quadratic model

Radiosensitization can be characterized by the LQ model.³ The degree of radiosensitization was quantitated based on the changes in the α and β values in the 653 pairs of survival

Table 2. Derived parameters [(median (interquartile range)] of the linear-quadratic model

Sensitization class	α (Gy^{-1})			β (Gy^{-2})		
	Control	Sensitizer	<i>p</i> -value	Control	Sensitizer	<i>p</i> -value
Apoptosis	0.19 (0.094–0.34)	0.41 (0.21–0.57)	<0.001	0.034 (0.026–0.047)	0.030 (0.022–0.052)	NS
Cell cycle	0.21 (0.12–0.38)	0.40 (0.29–0.63)	<0.001	0.037 (0.03–0.054)	0.034 (0.02–0.051)	NS
COXi	0.16 (0.1–0.23)	0.38 (0.19–0.51)	<0.001	0.037 (0.027–0.054)	0.029 (0.011–0.046)	NS
DNA-alkylation	0.10 (0.061–0.17)	0.33 (0.19–0.37)	<0.001	0.051 (0.038–0.064)	0.035 (0.022–0.047)	0.017
DNA-dsb-repair	0.24 (0.053–0.31)	0.63 (0.35–0.75)	0.0015	0.027 (0.02–0.046)	0.025 (0.0096–0.045)	NS
DNA-dsb-signal	0.22 (0.1–0.3)	0.49 (0.34–0.65)	<0.001	0.030 (0.017–0.048)	0.028 (0.011–0.042)	NS
EGFR/HER2	0.20 (0.077–0.25)	0.34 (0.17–0.49)	<0.001	0.034 (0.021–0.047)	0.031 (0.015–0.055)	NS
Gemcitabine	0.14 (0.081–0.23)	0.38 (0.32–0.54)	<0.001	0.034 (0.026–0.048)	0.036 (0.011–0.048)	NS
HDACi	0.13 (0.066–0.19)	0.43 (0.23–0.62)	<0.001	0.031 (0.022–0.053)	0.030 (0.015–0.045)	NS
IUdR/BUdR	0.17 (0.075–0.29)	0.43 (0.35–0.78)	<0.001	0.039 (0.023–0.049)	0.042 (0.026–0.065)	NS
Oxygen	0.054 (0.044–0.12)	0.19 (0.13–0.27)	<0.001	0.0051 (0.003–0.0066)	0.027 (0.018–0.05)	<0.001
PARPi	0.19 (0.11–0.27)	0.33 (0.24–0.41)	<0.001	0.036 (0.022–0.049)	0.052 (0.041–0.071)	<0.001
PDGFR	0.083 (0.034–0.15)	0.29 (0.24–0.44)	0.002	0.039 (0.023–0.044)	0.022 (0.015–0.028)	NS
PI3K/AKT/mTOR	0.13 (0.038–0.25)	0.32 (0.13–0.51)	<0.001	0.034 (0.026–0.048)	0.033 (0.018–0.055)	NS
Platinum	0.12 (0.054–0.22)	0.20 (0.093–0.31)	0.002	0.036 (0.021–0.047)	0.041 (0.015–0.067)	NS
RAS/MAPK/MEK	0.091 (0.058–0.23)	0.28 (0.16–0.4)	<0.001	0.040 (0.029–0.057)	0.032 (0.022–0.052)	NS
RRi	0.12 (0.11–0.22)	0.36 (0.24–0.59)	<0.001	0.046 (0.034–0.059)	0.057 (0.017–0.07)	NS
Taxane	0.15 (0.039–0.32)	0.35 (0.26–0.51)	<0.001	0.042 (0.021–0.06)	0.038 (0.0085–0.071)	NS
TOPOi	0.22 (0.059–0.38)	0.39 (0.16–0.58)	<0.001	0.024 (0.012–0.042)	0.022 (0.0033–0.042)	NS
TSi	0.057 (0.021–0.21)	0.25 (0.12–0.36)	<0.001	0.040 (0.02–0.058)	0.052 (0.034–0.086)	NS

AKT, v-akt murine thymoma viral oncogene homolog; BUdR, bromodeoxyuridine; COXi, cyclooxygenase inhibition; dsb, double-strand break; EGFR, epidermal growth factor receptor; HDACi, histone deacetylase inhibition; HER2, human epidermal growth factor receptor 2; IUdR, iododeoxyuridine; MAPK, mitogen-activated protein kinase; MEK, MAPK kinase; mTOR, mammalian target of rapamycin; N, number of pairs of survival curves; NS, not significant; PARPi, poly(adenosine diphosphate-ribose) polymerase inhibition; PDGFR, platelet-derived growth factor receptor; PI3K, phosphoinositide 3-kinase; RAS, rat sarcoma viral oncogene homolog; RRi, ribonucleotide reductase inhibition; TOPOi, topoisomerase I/II inhibition; TSi, thymidylate synthase inhibition.

curves. A representative pair of survival curves is shown in [Figure 1a](#). We found that radiosensitization was predominantly characterized by an increase of the α value (α -sensitization) without an increase of the β value (β -sensitization). The median α and β values were 0.15 Gy^{-1} [interquartile range (IQR): 0.056–0.26] and 0.034 Gy^{-2} (IQR: 0.021–0.051), respectively, in the absence of radiosensitization. In the presence of radiosensitization, the median α and β values were 0.35 Gy^{-1} (IQR: 0.19 to 0.52) and 0.033 Gy^{-2} (IQR: 0.016 to 0.057), respectively. The α value increased by a median of 0.17 Gy^{-1} (IQR: 0.066 to 0.31) and the β value changed by a median of -0.00023 Gy^{-2} (IQR: -0.014 to 0.014). The subset analysis of the 20 radiosensitization classes revealed that all 20 classes showed significant α -sensitization ([Table 2](#)). We found that only oxygen and PARPi elicited significant β -sensitization, while DNA-alkylation slightly decreased β values. The remaining 17 classes showed no significant changes in the β values. The differential effects on the α and β values suggest that radiosensitization is dependent on the levels of cellular survival or radiation doses. Thus, we assessed the degree of radiosensitization using two different

approaches; DMFs evaluated correlations between the degree of radiosensitization and levels of survival fractions ([Figure 2](#)), while SERs evaluated correlations between the degree of radiosensitization and radiation doses ([Figure 3](#)). The mean DMFs are also shown in [Table 1](#) for reference. In general, the DMFs seemed to be larger at higher survival levels. Significant positive correlations were found between the DMFs and the log survival fractions in 14 of the radiosensitization classes. Similarly, greater SERs were elicited by lower radiation doses than higher doses in 16 classes. In both analyses, the sensitizers with stronger β -sensitization (oxygen and PARPi) showed relatively constant synergistic effects throughout the clinically relevant ranges. These results suggest that β -sensitization is more favourable than α -sensitization to gain a therapeutic index by combining systemic therapy with high-dose-per-fraction radiotherapy.

Analysis using repair-misrepair model

We utilized the RMR model to elucidate the underlying mechanisms of radiosensitization, paying special attention to the differences between α - and β -sensitization. The RMR

Figure 2. The scatter plots show dose-modifying factors (DMFs) as a function of log survival fractions in 20 classes of radiosensitization. Spearman's rank correlation coefficients (ρ) and p -values are shown in the plots. There were weak but statistically significant correlations between the DMFs and the log survival fractions in 14 of the radiosensitization classes; greater DMFs were elicited at higher levels of survival. There were no correlations in the other six classes, including poly(adenosine diphosphate-ribose) polymerase inhibition (PARPi) and hypoxic sensitizers/oxygen (oxygen). AKT, v-akt murine thymoma viral oncogene homolog; BUdR, bromodeoxyuridine; COXi, cyclooxygenase inhibition; dsb, double-strand break; EGFR, epidermal growth factor receptor; HDACi, histone deacetylase inhibition; HER2, human epidermal growth factor receptor 2; IUdR, iododeoxyuridine; MAPK, mitogen-activated protein kinase; MEK, MAPK kinase; mTOR, mammalian target of rapamycin; PDGFR, platelet-derived growth factor receptor; PI3K, phosphoinositide 3-kinase; RAS, rat sarcoma viral oncogene homolog; RRI, ribonucleotide reductase inhibition; SF, surviving fraction; TOPOi, topoisomerase I/II inhibition; TSi, thymidylate synthase inhibition.

model can estimate the number of initial DNA damage lesions and the subsequent biological processes,⁴ including U_0 , eR_L , mR_L , vmR_Q , nmR_Q and rU . A sum of the latter five quantities, *i.e.* an entire length of each bar in Figure 4, represents the U_0 . Cell death can be attributed to the three types of quantities: mR_L , nmR_Q and rU . The RMR model was built on the assumption that SFs were directly correlated with these quantities. To calculate the quantities, the 653 pairs of survival curves were fitted to the RMR model, estimating the five parameters (δ , ε , T , φ and ψ) in Equation (11). Figure 1b shows a representative pair of survival curves fitted by the RMR model. Once the model parameters were determined, one could calculate U_0 by plugging the parameter δ into Equation (5). Similarly, rU , eR_L , mR_L , vmR_Q and nmR_Q were calculated by plugging the parameters δ , ε , T , φ or ψ into Equation 6, 7, 8, 9 or 10. Since the five quantities except for U_0 were not linearly correlated with the radiation dose, the data are presented at a dose of 10 Gy (Figure 4). The five derived parameters are summarized in Table 3 according to the culture conditions (air vs hypoxia).

Under aerobic conditions without radiosensitization, the median U_0 was 2.8 Gy^{-1} per cell (IQR: 1.8–10.4). Approximately 60% of U_0 was linearly repaired, while the remaining 40% was repaired in a quadratic fashion. About one-half of the

quadratic misrepair was non-viable, leading to cell death. The median values of the coefficients φ and T were both 1. As a result, there were neither considerable mR_L nor rU . Therefore, the RMR model indicates that radiation-induced cell death was primarily attributable to quadratic misrepair. When combined with the radiosensitizing agents, there was no substantial change in U_0 , although eR_L decreased. Instead, rU increased significantly. There was a modest increase in mR_L ; however, this change was subtle and was not visible on the plot. While the linear repair was inhibited to a certain extent, both non-viable and viable quadratic repairs paradoxically increased. Collectively, the RMR model indicated that the mechanisms of radiosensitization were generally two-fold, with increased residual lesions through repair inhibition and shifts from linear repair to quadratic repair, leading to enhanced lethal chromosomal aberrations.

Under hypoxic conditions, U_0 was 1.3 Gy^{-1} per cell (IQR: 0.8–4.3), which is significantly lower than that under aerobic conditions. The majority of U_0 was linearly repaired. In the presence of hypoxic radiosensitizers/oxygen, the increased U_0 was repaired predominantly in a quadratic fashion. Therefore, radioresistance under hypoxia and radiosensitization by oxygen or hypoxic sensitizers seemed to be mediated not only by the increased initial yield of DNA damage but also

Figure 3. The scatterplots show sensitizer enhancement ratios (SERs) as a function of radiation doses in 20 classes of radiosensitization. Spearman's rank correlation coefficients (ρ) and p -values are shown in the plots. There were negative correlations between the SERs and radiation doses in 16 of the radiosensitization classes; greater SERs were elicited by lower doses of radiation. No correlations were found in the halogenated pyrimidine analogues (IUDr/BUDr), topoisomerase inhibition (TOPOi), poly(adenosine diphosphate-ribose) polymerase inhibition (PARPi) and hypoxic sensitizers/oxygen (oxygen). AKT, v-akt murine thymoma viral oncogene homolog; COXi, cyclooxygenase inhibition; dsb, double-strand break; EGFR, epidermal growth factor receptor; HDACi, histone deacetylase inhibition; HER2, human epidermal growth factor receptor 2; MAPK, mitogen-activated protein kinase; MEK, MAPK kinase; mTOR, mammalian target of rapamycin; PDGFR, platelet-derived growth factor receptor; PI3K, phosphoinositide 3-kinase; RAS, rat sarcoma viral oncogene homolog; RRI, ribonucleotide reductase inhibition; TOPOi, topoisomerase I/II inhibition; TSi, thymidylate synthase inhibition.

through an altered repair mechanism in a linear vs quadratic fashion.

To further characterize the mechanisms of radiosensitization, the biological quantities were summarized according to the 20 classes of radiosensitization. Since there were no substantial differences in U_0 or mR_L in this subset analysis, only rU and nmR_Q are shown in Figures 5 and 6, respectively. Six classes (COXi, DNA-dsb-signal, IUDr/BUDr, PDGFR, RAS/MAPK/MEK and taxane) showed increased rU without a significant change in nmR_Q , suggesting that repair inhibition is a possible mechanism in these radiosensitizations. In contrast, four classes (oxygen, PARPi, RRI and DNA-dsb-repair) showed increased nmR_Q without a significant change in rU . This suggests that more DNA lesions were repaired in a quadratic fashion rather than in a linear fashion without a substantial reduction of the total repair. Apoptosis, gemcitabine, HDACi, PI3K/AKT/mTOR, platinum and TSi showed mixed results, significantly affecting both rU and nmR_Q .

Relationship between the linear-quadratic and repair-misrepair models

We analyzed the correlations between the LQ model parameters and the biological quantities calculated in the RMR model for the 1306 survival curves (Figure 7). The α value

was negatively correlated with R_L and positively correlated with rU . In contrast, the β value showed little correlation with rU , but showed the greatest correlation with nmR_Q . Therefore, α -sensitization is likely to be mediated by an inhibition of the total repair activity. In contrast, β -sensitization can result from augmented quadratic misrepair, which may be due to reduced fidelity repair with similar or increased repair activity.

DISCUSSION

High-precision radiotherapy is able to deliver relatively high doses of radiation to the target while sparing the surrounding normal tissues. Ablative radiation doses may be enough for tumour control if targets are small and surrounded by parallel organs. However, there are multiple data indicating that radiation alone, even with high-dose-per-fraction radiotherapy, results in suboptimal outcomes for local tumour control as well as treatment-related toxicities.⁷⁻⁹ One possible way to further improve therapeutic indices, other than physical dose escalations, would be combined modality therapy with radiosensitizers. Accordingly, this study was initiated to find an optimal sensitizer and/or target of sensitization combined with high-dose radiation. A framework for drug-radiation combination therapy has been proposed, encompassing five mechanisms: spatial cooperation, cytotoxic enhancement, biological

Figure 4. The barplots represent biological responses to radiation-induced DNA damage at 10 Gy in the absence or presence of radiosensitization under aerobic conditions (air; the left two bars). The mean numbers of five types of DNA lesions per cell (linear eurepair, linear misrepair, residual lesion, viable quadratic misrepair and non-viable quadratic misrepair) were estimated using the derived parameters in the repair-misrepair model. In the absence of radiosensitization, almost all initial DNA damage lesions were repaired by either linear eurepair or quadratic misrepair, leaving no residual lesions or linear misrepair. This suggests that quadratic misrepair is a primary mode of cell death following the application of 10 Gy of radiation. In the presence of radiosensitization, the initial production of DNA damage lesions (the sum of the five quantities) did not change significantly. The linear eurepair significantly decreased. Instead, there were significant increases in viable/non-viable quadratic misrepair as well as the residual lesion. The right two barplots show a similar analysis under hypoxic conditions, where the initial production of DNA lesions was significantly reduced as compared with the aerobic conditions. The majority of the DNA damage was linearly repaired in the absence of radiosensitization. Hypoxic sensitizers/oxygen remarkably increased the initial production of DNA lesions to a comparable level with the data obtained under aerobic conditions. The amount of net increase was mostly repaired by quadratic misrepair without any significant change in linear eurepair.

cooperation, temporal modulation and normal tissue protection.¹⁰ Although the five components should be evaluated to develop a new combination of drug and radiotherapy, this study

focused on the aspect of only cytotoxic enhancement. The results of this study need to be further evaluated in future studies on the basis of other components of the drug–radiation combination framework.

Radiosensitization in high-dose-per-fraction radiotherapy

In this study, we attempted to evaluate the concept of radiosensitization when radiation is administered at a high dose per fraction using mathematical modelling. We found that synergistic effects were significantly dependent on radiation doses in many classes of radiosensitization we explored. In other words, there was differential sensitization in favour of lower radiation doses compared with higher doses. Many radiosensitizers showed only α -sensitization but not β -sensitization, which renders sensitization a challenging task in the setting of hypofractionated high-precision radiotherapy. The α -sensitizers may confer a greater sensitizing effect in the low-dose region and a smaller effect in the high-dose region, which may offset the preferential dose gradients in the target volume vs at-risk organ achieved by dose-painting techniques. A component of β -sensitization would be necessary to elicit comparable radiosensitization in high-dose regions. Significant β -sensitization was found for oxygen under hypoxic conditions and to a lesser extent for PARPi under aerobic conditions. These sensitizers may be suitable for combining with high-dose radiation. In agreement with our results, Brown et al¹¹ reported that hypoxic sensitizers are suitable candidates when administered with stereotactic radiotherapy because hypoxic cells are radio-resistant even at high doses of radiation. Using the LQ model, Franken et al¹² also examined the dose response aspect of radiosensitization in their clonogenic survival data with five different radiosensitizers, as well as hyperthermia. They found that α and β were separately influenced depending on the radiosensitivity of the cell lines and culture conditions used; however, consistent with our results, the radiosensitizers appeared to predominantly increase the α component. They concluded that radiosensitizing effects could be further exploited using smaller fractions or by lowering the dose rates. Interestingly, they showed that the β value was specifically increased by mild hyperthermia (41 °C), resulting in a lowered α/β ratio.

Radiosensitization is likely dependent on drug doses, timing of administration or duration of exposure. However, our data set

Table 3. Derived parameters [median (interquartile range)] of the repair-misrepair model under aerobic conditions (air) and hypoxic conditions (hypoxia)

Parameter	Air			Hypoxia		
	Control	Sensitizer	<i>p</i> -value	Control	Sensitizer	<i>p</i> -value
δ	2.8 (1.8–10)	2.9 (1.9–8.6)	NS	1.3 (0.83–4.3)	3.1 (1.7–13)	0.01
ϵ	47 (5.9–320)	25 (3.8–210)	<0.001	37 (7.3–400)	12 (5.1–280)	NS
φ	1 (1–1)	1 (1–1)	<0.001	1 (1–1)	1 (1–1)	NS
ψ	0.54 (0.24–0.79)	0.59 (0.25–0.79)	NS	0.61 (0.23–0.74)	0.56 (0.36–0.8)	NS
T	1 (0.97–1)	0.99 (0.92–1)	<0.001	1 (0.98–1)	1 (0.99–1)	NS

Figure 5. The number of residual lesions per cell following 10 Gy of radiation was compared between the absence (C) and presence (S) of radiosensitization in 20 classes of radiosensitization. The residual lesions were significantly increased in 12 of the 20 classes. The significance of the differences was tested using the Wilcoxon signed-rank test. Adjusted *p*-values by the Benjamini–Hochberg procedure are shown in the plots. AKT, v-akt murine thymoma viral oncogene homolog; BUdR, bromodeoxyuridine; COX*i*, cyclooxygenase inhibition; dsb, double-strand break; EGFR, epidermal growth factor receptor; HDAC*i*, histone deacetylase inhibition; HER2, human epidermal growth factor receptor 2; IUdR, iododeoxyuridine; MAPK, mitogen-activated protein kinase; MEK, MAPK kinase; mTOR, mammalian target of rapamycin; PDGFR, platelet-derived growth factor receptor; PI3K, phosphoinositide 3-kinase; RAS, rat sarcoma viral oncogene homolog; R*Ri*, ribonucleotide reductase inhibition; TOPO*i*, topoisomerase I/II inhibition; TS*i*, thymidylate synthase inhibition.

lacked such detailed information regarding drug administration. In addition, our data represented averaged effects of the arbitrarily classified 20 groups of sensitizers on the various types of cell lines. Therefore, caution needs to be exercised while interpreting the results of this study. It cannot be excluded that any agent may behave differently when administered in a specific way.

Mechanism(s) of action in α - and β -sensitizers

When DNA is considered as the primary molecular target of ionizing radiation, synergistic interactions between radiation and drugs may be conceptually attributed to several factors, including increased yield of DNA damage, impaired DNA repair processes and reduced tolerance to misrepaired lesions. Likewise, the RMR model formulates the processes of radiation dose responses into the initial DNA damage lesion and subsequent biological processes: eR_L , mR_L , rU , vmR_Q and nmR_Q . Thus, we found the RMR model to be useful for analyzing the mechanisms of action for radiosensitivity modification. The model-based analysis suggested that there were three mechanisms of radiosensitization: (1) increased yield of DNA damage; (2) increased residual DNA damage by inhibition of DNA repair; and (3) increased chromosomal translocation through enhanced quadratic misrepair. The initial yield of DNA damage was significantly increased by the oxygen/hypoxic radiosensitizers. Notably, oxygen molecules

play an important role in the physical and chemical phases of cellular responses to irradiation. It is necessary for oxygen molecules to physically fix DNA radicals following the interactions between DNA macromolecules and radiation-induced hydroxyl radicals; otherwise, they can be chemically restored within a short length of time.¹³ In addition, oxygen has been shown to modulate enzymatic DNA repair machineries by multiple mechanisms.¹³

The second and third mechanisms appear to contradict each other, since reduced DNA repair activity would decrease the generation of chromosomal translocations and *vice versa*. Therefore, if we inhibit the total activity of DNA repair, with residual DNA damage being increased, the α component would be enhanced without also enhancing the β component. This α -sensitization can be seen in many types of radiosensitizers/chemotherapeutic agents. On the contrary, if we can enhance quadratic misrepair, the β component would be increased, with a modest effect on the α component. At present, the underlying molecular mechanisms that modulate the amount of quadratic misrepair have not been well defined.

DNA dsbs can be repaired by three distinct repair machineries: DNA dependent protein kinase (DNA-PK)-dependent non-homologous end-joining (D-NHEJ), back up/alternative non-

Figure 6. The numbers of non-viable quadratic misrepairs per cell following 10Gy of radiation were compared between the absence (C) and presence (S) of radiosensitization in 20 classes of radiosensitization. Non-viable quadratic misrepair was significantly increased in 10 of the 20 classes. Significance of the difference was tested using the Wilcoxon signed-rank test. Adjusted *p*-values by the Benjamini-Hochberg procedure are shown in the plots. AKT, v-akt murine thymoma viral oncogene homolog; BUdR, bromodeoxyuridine; COX_i, cyclooxygenase inhibition; dsb, double-strand break; EGFR, epidermal growth factor receptor; HDAC_i, histone deacetylase inhibition; HER2, human epidermal growth factor receptor 2; IUdR, iododeoxyuridine; MAPK, mitogen-activated protein kinase; MEK, MAPK kinase; mTOR, mammalian target of rapamycin; PDGFR, platelet-derived growth factor receptor; PI3K, phosphoinositide 3-kinase; RAS, rat sarcoma viral oncogene homolog; RRI, ribonucleotide reductase inhibition; TOPO_i, topoisomerase I/II inhibition; TSi, thymidylate synthase inhibition.

homologous end-joining (B-NHEJ) and homologous recombination repair (HRR).¹⁴ The fidelity of DNA dsb repair decreases in the following order: HRR, D-NHEJ and B-NHEJ. It is well known that HRR is strictly controlled and operates primarily in the S-phase of the cell cycle when sister chromatids are available as templates for repair. In contrast, NHEJ is the primary repair machinery in the vertebrate¹⁵ and is active throughout the cell cycle. However, it has a potential risk for mR_L, *i.e.* sequence alterations in the vicinity of the dsb, to lose/alter genetic information. Moreover, NHEJ is thought to be a major source of chromosomal translocation.¹⁶ B-NHEJ is thought to function as a back-up pathway for D-NHEJ. B-NHEJ takes longer to complete than D-NHEJ, leaving more time for the DNA ends to drift apart. As a result, the DNA ends are more likely to join in an aberrant interchromosomal fashion.¹⁷ Although B-NHEJ is also active throughout the cell cycle, it is more enhanced in the G2 phase than in the G1 phase. Thus, B-NHEJ may also function as a back-up for HRR.¹⁸ However, how cells choose the various repair pathways when encountering DNA dsbs remains a subject of active research.^{15,19–22}

Several investigators have examined the role of HRR compared with NHEJ on cellular survival following fractionated irradiation. Using chicken B lymphoma cells (DT40) proficient or deficient in RAD54 and/or Ku80, HRR was shown to play

a critical role for split-dose recovery.²³ Similarly, it was reported that HRR-deficient Chinese hamster ovary cells (irs1SF) were sensitive to fractionation, while NHEJ-deficient Chinese hamster ovary cells (V3-3) showed reduced sensitivity to fraction sizes. The authors postulated that HRR, a high-fidelity repair process independent of DNA damage levels, mediates less fractionation sensitivity.²⁴ In fact, we found that the so-called S-phase sensitizers, which primarily elicit cytotoxicity and radiosensitization in the S-phase subpopulation, TSi, RRI and gemcitabine, increased quadratic misrepair (Figure 6).

Inhibition of PARP has been known to confer radiosensitization for several decades,²⁵ although the exact mechanisms of radiosensitization are poorly understood. PARP1 plays multiple important roles in DNA repair pathways, including base excision repair, single-strand break repair and dsb repair. PARP1-deficient cells have been shown to increase the levels of sister chromatid exchanges.²⁶ Following DNA damage, PARP1 is required to slow replication fork progression and promote HRR at stalled and/or collapsed replication forks.^{27,28} PARP1-deficient chicken lymphoma cells (DT40), which do not carry the PARP2 gene, showed reduced levels of HRR. This phenotype was strictly dependent on the presence of Ku, since PARP1/Ku70 double-mutant cells were proficient at performing HRR. PARP1 was proposed to

Figure 7. The scatterplot matrix from a total of 1306 survival curves shows correlations between the two linear-quadratic model parameters (α and β) and six quantities estimated in the repair-misrepair model (U_0 , R_L , R_Q , rU , mR_L and nmR_Q). The upper half of the matrix shows Spearman's rank correlation coefficients between each combination of the variables. The α value was negatively correlated with linear repair and positively correlated with rU . The β value was most highly correlated with nmR_Q . U_0 , uncommitted lesion; R_L , linear repair; R_Q , quadratic misrepair; rU , unrepaired residual lesion; mR_L , linear misrepair; nmR_Q , non-viable quadratic misrepair.

promote HRR by minimizing the suppressive effect of Ku on HRR.²⁹ Therefore, inhibition of PARP1 may reduce the levels of HRR and in turn increase error-prone NHEJ, resulting in increased quadratic misrepair. On the contrary, several subsequent studies have indicated that PARP1 competed with Ku for the free ends of DNA dsbs but actually promoted B-NHEJ,

which is a more mutagenic repair pathway than D-NHEJ.^{30–32} To further complicate matters, PARP1 facilitated alternative end-joining, while PARP2 suppressed chromosomal translocation during the class-switch recombination.³³ Nevertheless, these findings suggest that PARP1/2 is involved in an interface among the different repair machineries and plays an important role in the

choice of the pathways. We speculate that an altered choice to a dsb repair pathway with less fidelity would lead to enhanced quadratic misrepair, resulting in β -sensitization. Clearly, more data are necessary to elucidate how PARP and/or other molecules contribute to this type of radiosensitization by regulating the repair pathway choice.

In summary, a degree of synergistic effect was dependent on the levels of cellular survival or radiation doses in many classes of radiosensitization. Lower doses of radiation elicited greater synergism, namely α -sensitization, which was correlated with increased residual DNA lesions resulting from repair inhibition. In contrast, significant β -sensitization was found only in PARP inhibition and oxygen/hypoxic sensitizers.

Although β -sensitization has not been well characterized, the model-based analysis in this study suggests that increased chromosomal translocations mediated by a type of repair with reduced fidelity are a possible mechanism of β -sensitization. Pursuing targets of this type of radiosensitization may help in developing a novel combined modality therapy with high-dose-per-fraction radiotherapy. Since this is an exploratory study to generate hypotheses, further studies are required to validate our conclusion.

FUNDING

This work was supported by Grant-in-Aids for Scientific Research (C) 23591839 and 24591844 to YS from Japan Society for the Promotion of Science.

REFERENCES

- Lawrence YR, Vikram B, Dignam JJ, Chakravarti A, Machtay M, Freidlin B, et al. NCI-RTOG translational program strategic guidelines for the early-stage development of radiosensitizers. *J Natl Cancer Inst* 2013; **105**: 11–24. doi: <http://dx.doi.org/10.1093/jnci/djs472>
- Dale RG. The application of the linear-quadratic dose-effect equation to fractionated and protracted radiotherapy. *Br J Radiol* 1985; **58**: 515–28. doi: <http://dx.doi.org/10.1259/0007-1285-58-690-515>
- Miller EM, Fowler JF, Kinsella TJ. Linear-quadratic analysis of radiosensitization by halogenated pyrimidines. I. Radiosensitization of human colon cancer cells by iododeoxyuridine. *Radiat Res* 1992; **131**: 90–7. doi: <http://dx.doi.org/10.2307/3578321>
- Tobias CA. The repair-misrepair model in radiobiology: comparison to other models. *Radiat Res Suppl* 1985; **8**: S77–95. doi: <http://dx.doi.org/10.2307/3583515>
- Greco WR, Bravo G, Parsons JC. The search for synergy: a critical review from a response surface perspective. *Pharmacol Rev* 1995; **47**: 331–85.
- Fertil B, Dertinger H, Courdi A, Malaise EP. Mean inactivation dose: a useful concept for intercomparison of human cell survival curves. *Radiat Res* 1984; **99**: 73–84. doi: <http://dx.doi.org/10.2307/3576448>
- Rowe BP, Boffa DJ, Wilson LD, Kim AW, Dettmerbeck FC, Decker RH. Stereotactic body radiotherapy for central lung tumors. *J Thorac Oncol* 2012; **7**: 1394–9. doi: <http://dx.doi.org/10.1097/JTO.0b013e3182614bf3>
- Shibamoto Y, Hashizume C, Baba F, Ayakawa S, Miyakawa A, Murai T, et al. Stereotactic body radiotherapy using a radiobiology-based regimen for stage I non-small-cell lung cancer. *J Thorac Oncol* 2015; **10**: 960–4. doi: <http://dx.doi.org/10.1097/JTO.0000000000000525>
- Habermehl D, Herfarth KK, Bermejo JL, Hof H, Rieken S, Kuhn S, et al. Single-dose radiosurgical treatment for hepatic metastases—therapeutic outcome of 138 treated lesions from a single institution. *Radiat Oncol* 2013; **8**: 175. doi: <http://dx.doi.org/10.1186/1748-717X-8-175>
- Bentzen SM, Harari PM, Bernier J. Exploitable mechanisms for combining drugs with radiation: concepts, achievements and future directions. *Nat Clin Pract Oncol* 2007; **4**: 172–80. doi: <http://dx.doi.org/10.1038/ncponc0744>
- Brown JM, Diehn M, Loo BW. Stereotactic ablative radiotherapy should be combined with a hypoxic cell radiosensitizer. *Int J Radiat Oncol Biol Phys* 2010; **78**: 323–7. doi: <http://dx.doi.org/10.1016/j.ijrobp.2010.04.070>
- Franken NA, Oei AL, Kok HP, Rodermond HM, Sminia P, Crezee J, et al. Cell survival and radiosensitisation: Modulation of the linear and quadratic parameters of the LQ model (Review). *Int J Oncol* 2013; **42**: 1501–15. doi: <http://dx.doi.org/10.3892/ijo.2013.1857>
- Liu C, Lin Q, Yun Z. Cellular and molecular mechanisms underlying oxygen-dependent radiosensitivity. *Radiat Res* 2015; **183**: 487–96. doi: <http://dx.doi.org/10.1667/RR13959.1>
- Mladenov E, Magin S, Soni A, Iliakis G. DNA double-strand break repair as determinant of cellular radiosensitivity to killing and target in radiation therapy. *Front Oncol* 2013; **3**: 113. doi: <http://dx.doi.org/10.3389/fonc.2013.00113>
- Sonoda E, Hochegger H, Saberi A, Taniguchi Y, Takeda S. Differential usage of non-homologous end-joining and homologous recombination in double strand break repair. *DNA Repair (Amst)* 2006; **5**: 1021–9. doi: <http://dx.doi.org/10.1016/j.dnarep.2006.05.022>
- Povirk LF. Biochemical mechanisms of chromosomal translocations resulting from DNA double-strand breaks. *DNA Repair (Amst)* 2006; **5**: 1199–212. doi: <http://dx.doi.org/10.1016/j.dnarep.2006.05.016>
- Wray J, Williamson EA, Singh SB, Wu Y, Cogle CR, David M, et al. PARP1 is required for chromosomal translocations. *Blood* 2013; **121**: 4359–65. doi: <http://dx.doi.org/10.1182/blood-2012-10-460527>
- Dueva R, Iliakis G. Alternative pathways of non-homologous end joining (NHEJ) in genomic instability and cancer. *Transl Cancer Res* 2013; **2**: 163–77.
- Shrivastav M, De Haro LP, Nickoloff JA. Regulation of DNA double-strand break repair pathway choice. *Cell Res* 2008; **18**: 134–47. doi: <http://dx.doi.org/10.1038/cr.2007.111>
- Daley JM, Sung P. 53BP1, BRCA1, and the choice between recombination and end joining at DNA double-strand breaks. *Mol Cell Biol* 2014; **34**: 1380–8. doi: <http://dx.doi.org/10.1128/MCB.01639-13>
- Quennet V, Beucher A, Barton O, Takeda S, Löbrich M. CtIP and MRN promote

- non-homologous end-joining of etoposide-induced DNA double-strand breaks in G1. *Nucleic Acids Res* 2011; **39**: 2144–52. doi: <http://dx.doi.org/10.1093/nar/gkq1175>
22. Kato K, Nakajima K, Ui A, Muto-Terao Y, Ogiwara H, Nakada S. Fine-tuning of DNA damage-dependent ubiquitination by OTUB2 supports the DNA repair pathway choice. *Mol Cell* 2014; **53**: 617–30. doi: <http://dx.doi.org/10.1016/j.molcel.2014.01.030>
 23. Utsumi H, Elkind MM. Requirement for repair of DNA double-strand breaks by homologous recombination in split-dose recovery. *Radiat Res* 2001; **155**: 680–6. doi: [http://dx.doi.org/10.1667/0033-7587\(2001\)155\[0680:RFRODD\]2.0.CO;2](http://dx.doi.org/10.1667/0033-7587(2001)155[0680:RFRODD]2.0.CO;2)
 24. Somaiah N, Yarnold J, Lagerqvist A, Rothkamm K, Helleday T. Homologous recombination mediates cellular resistance and fraction size sensitivity to radiation therapy. *Radiother Oncol* 2013; **108**: 155–61. doi: <http://dx.doi.org/10.1016/j.radonc.2013.05.012>
 25. Oleinick NL, Evans HH. Poly(ADP-ribose) and the response of cells to ionizing radiation. *Radiat Res* 1985; **101**: 29–46. doi: <http://dx.doi.org/10.2307/3576301>
 26. Wang ZQ, Stingl L, Morrison C, Jantsch M, Los M, Schulze-Osthoff K, et al. PARP is important for genomic stability but dispensable in apoptosis. *Genes Dev* 1997; **11**: 2347–58. doi: <http://dx.doi.org/10.1101/gad.11.18.2347>
 27. Schultz N. Poly(ADP-ribose) polymerase (PARP-1) has a controlling role in homologous recombination. *Nucleic Acids Res* 2003; **31**: 4959–64. doi: <http://dx.doi.org/10.1093/nar/gkg703>
 28. Pears CJ, Couto CA, Wang H, Borer C, Kiely R, Lakin ND. The role of ADP-ribosylation in regulating DNA double-strand break repair. *Cell Cycle* 2012; **11**: 48–56. doi: <http://dx.doi.org/10.4161/cc.11.1.18793>
 29. Hohegger H, Dejsuphong D, Fukushima T, Morrison C, Sonoda E, Schreiber V, et al. Parp-1 protects homologous recombination from interference by Ku and Ligase IV in vertebrate cells. *EMBO J* 2006; **25**: 1305–14. doi: <http://dx.doi.org/10.1038/sj.emboj.7601015>
 30. Audebert M, Salles B, Calsou P. Involvement of poly(ADP-ribose) polymerase-1 and XRCC1/DNA ligase III in an alternative route for DNA double-strand breaks rejoining. *J Biol Chem* 2004; **279**: 55117–26. doi: <http://dx.doi.org/10.1074/jbc.M404524200>
 31. Wang M, Wu W, Wu W, Rosidi B, Zhang L, Wang H, et al. PARP-1 and Ku compete for repair of DNA double strand breaks by distinct NHEJ pathways. *Nucleic Acids Res* 2006; **34**: 6170–82. doi: <http://dx.doi.org/10.1093/nar/gkl840>
 32. Paddock MN, Bauman AT, Higdon R, Kolker E, Takeda S, Scharenberg AM. Competition between PARP-1 and Ku70 control the decision between high-fidelity and mutagenic DNA repair. *DNA Repair (Amst)* 2011; **10**: 338–43. doi: <http://dx.doi.org/10.1016/j.dnarep.2010.12.005>
 33. Robert I, Dantzer F, Reina-San-Martin B. Parp1 facilitates alternative NHEJ, whereas Parp2 suppresses IgH/c-myc translocations during immunoglobulin class switch recombination. *J Exp Med* 2009; **206**: 1047–56. doi: <http://dx.doi.org/10.1084/jem.20082468>