Published in final edited form as:

Curr Diab Rep. 2016 July ; 16(7): 66. doi:10.1007/s11892-016-0755-1.

Patterns of user engagement with mobile- and web-delivered self-care interventions for adults with T2DM: A review of the literature

Lyndsay A. Nelson, PhD^{1,2}, Taylor D. Coston, BS³, Andrea L. Cherrington, MD, MPH^{4,5}, and Chandra Y. Osborn, PhD, MPH^{1,2,6,7}

¹Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

²Center for Health Behavior and Health Education, Vanderbilt University Medical Center, Nashville, TN, USA

³School of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

⁴Department of Medicine, University of Alabama at Birmingham, Birmingham, AL, USA

⁵Division of Preventative Medicine, University of Alabama at Birmingham, Birmingham, AL, USA

⁶Center for Diabetes Translational Research, Vanderbilt University Medical Center, Nashville, TN, USA

⁷Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, TN, USA

Abstract

Technology-delivered interventions can improve the health behaviors and clinical outcomes of persons with diabetes, but only if end users engage with these interventions. To summarize the current knowledge on engagement with technology-based interventions, we conducted a review of recent mobile- and web-delivered intervention studies for adults with type 2 diabetes published from 2011–2015. Among 163 identified studies, 24 studies satisfied our inclusion criteria. There was substantial variation in how intervention engagement was reported across studies.

Engagement rates were lower among interventions with a longer duration, and engagement decreased over time. In several studies, older age and lower health literacy were associated with less engagement, and more engagement was associated with intervention improvement in at least one outcome, including glycemic control. Future technology-based intervention studies should report on engagement, examine and report on associations between user characteristics and engagement, and aim to standardize how this is reported, particularly in longer trials.

Corresponding Author: Chandra Y. Osborn, PhD, MPH, Vanderbilt University Medical Center, Division of General Internal Medicine & Public Health, Center for Health Behavior and Health Education, 2525 West End Ave., Suite 370, Nashville, TN 37203 United States, Phone: 1 615 936 8468, chandra.osborn@vanderbilt.edu.

Compliance with Ethical Standards

Conflict of Interest

Lyndsay A. Nelson, Taylor D. Coston, Andrea L. Cherrington, and Chandra Y. Osborn declare that they have no conflict of interest.

Human and Animal Rights and Informed Consent

This article does not contain any studies with human or animal subjects performed by any of the authors.

Keywords

Technology; Internet; Mobile; Intervention; Engagement; Diabetes; Review

Introduction

Suboptimal glycemic control can lead to a variety of complications including stroke, heart attacks, and death [1]. Adherence to recommended self-care behaviors (e.g., healthy eating, exercise, medication adherence, self-monitoring of blood glucose [SMBG]) can maintain optimal glycemic control [2, 3] and prevent complications [1], but self-care adherence is challenging, and, in turn, low among adults with type 2 diabetes (T2DM) [4, 5].

Technology-delivered interventions (e.g., mobile- and web-based) can effectively promote the self-care behaviors [6, 7], and improve the glycemic control of adults with T2DM [8, 9], but assume users will engage with interventions independently. The extent to which they do, however, is unclear [6, 10].

The degree to which users engage with an intervention can tell us why an intervention had an effect or not [6], yet technology-based interventions rarely report on users' engagement [11]. Among studies that do, engagement is operationalized inconsistently. Schubart et al.'s 2011 [12] review of web-delivered self-care interventions found differences in how studies report on user engagement (e.g., frequency of website visits, number of users logging in at least once, and pages viewed). Such reporting differences are also common among mobile-delivered interventions (e.g., text message response rate [13], number of users who responded to text messages at least three times per week [14]).

Users' level of intervention engagement varies widely across studies operationalizing engagement the same way (e.g., text message response rate, average number of website logins) [6]. In 2009, Fjeldsoe et al. [15] reviewed behavior change text message-based interventions and found substantial variability in users' compliance to the intervention. Studies report that, on average, users respond to as few as 17% [16] to as many as 84% [13] of text messages sent during a study period. Web-delivered intervention studies also report wide variability in user engagement, with some studies reporting an average of one to five logins and others reporting more than ten average logins [17]. Intervention engagement variability may be in large part due to the wide variability in mobile- and web-based intervention attributes (i.e. intervention duration, dosing, relevance, timing, user requirements), and/or user attributes (e.g., user age, socioeconomic or health literacy status).

Understanding the determinants of intervention engagement can help design interventions that optimize engagement; however, the factors influencing engagement in technology-delivered diabetes interventions are unclear. In their 2011 review of mobile interventions for patients with type 1 diabetes and T2DM, Mulvaney et al. [11] found insufficient reporting on engagement to associate intervention design features with engagement. This review only included studies published through 2010 and did not include web-delivered interventions. In 2014, Cotter et al. [10] conducted a review of web-delivered diabetes interventions published through January 2013, and called for more research to understand under which conditions and with what types of users, engagement changes.

Lastly, it is unclear whether engagement with a technology-delivered intervention is associated with changes in outcomes. In Fjeldsoe et al.'s [15] review, they reported on one study in which a subgroup of users who were more actively engaged in the intervention showed a trend toward greater glycemic control (A1c) improvement compared to users who were less engaged in the intervention. Moreover, in Cotter et al.'s [10] review, engagement in web-delivered interventions was inconsistently associated with diabetes outcomes.

In this review, we examine the recent literature to provide an updated account of studies reporting user engagement with both mobile- and web-delivered diabetes self-care interventions for adults with T2DM. The goals of our review are to describe the different ways engagement is operationalized, describe the extent to which T2DM users are engaging with technology-delivered interventions and how this varies by user characteristics, identify the design features that may facilitate or hinder engagement, and describe if and when engagement is associated with better outcomes.

Methods

Data Sources and Search Strategy

In November 2015, we searched PubMed for technology-delivered self-care interventions used by adults with T2DM. We searched for studies published between 2011 and 2015 to capture those studies not included in Mulvaney et al.'s [13] review. All searches included a term from each of three categories: (1) technology (e.g., mobile intervention, web program, cell phone program), (2) self-care (e.g., self-management, behavior), and (3) diabetes (i.e., type 2 diabetes adults). Terms were intralinked with "OR" and interlinked with "AND."

Study Eligibility

Eligible studies: (1) included adults with a diagnosis of T2DM (>50% of the study sample), (2) tested a technology-delivered intervention designed to improve a diabetes self-care behavior, (3) asked participants to use the intervention on their own time (i.e., outside the lab or clinic setting), (4) tested interventions using <50% telehealth or human interaction, and (5) reported on participants' engagement with the intervention. We excluded studies focused on the design and development phase of the intervention and studies testing interventions designed to improve a psychological (e.g., depressive symptoms, distress) or psychosocial (e.g., self-care knowledge, self-efficacy) outcome. Additionally, we excluded studies if they were a systematic review, a meta-analysis, editorial/commentary, or were not published in English.

Our search terms identified 163 published articles. Authors LAN and TDC independently reviewed each abstract of the identified articles. Studies were excluded at this stage if both reviewers agreed eligibility criteria were not met. Both authors independently reviewed the full text of the remaining articles, excluding those not meeting eligibility criteria, resulting in 24 articles being included in our review (see Figure 1).

Data Abstraction and Narrative Synthesis

Given variation in study designs, methodologies employed, and outcomes assessed between the 24 included studies, we used a narrative synthesis approach [18] to describe and identify patterns of engagement across included studies. We created two separate tables to report on both the general intervention characteristics (e.g., study design, sample characteristics, length of intervention, outcomes) (Table 1) and information specific to participants' engagement with an intervention (e.g., engagement rates, change in engagement across time, and associations between engagement and outcomes) (Table 2). Although all studies in our review tested an intervention designed to improve at least one diabetes self-care behavior, many still reported psychological and psychosocial outcomes and are therefore included in Table 1 and in our results. In both tables, we organized the interventions into two groups based on whether they were primarily a mobile- or web-based intervention. Although some mobile interventions contained web components, all interventions used either mobile phones or the Internet as their dominant platform. The subsequent review represents information available about engagement with mobile- or web-based self-care interventions for adults with T2DM published in PubMed from 2011–2015.

Results

Of the 24 articles included in our review, 15 tested a mobile-delivered intervention and nine tested a web-delivered intervention. The majority of studies ($n=10$) were formative research studies (i.e., pilot, feasibility, and usability studies), six were randomized controlled trials, four were non-experimental pre-post studies (i.e., no control group), three were studies reporting data from a larger trial, and one was a quasi-experimental study (i.e., used matched controls). Nearly two-thirds (58.3%) of these articles were published in the past three years (i.e., 2013–2015).

Operationalizing engagement

The studies in our review operationalized intervention engagement according to the types of technical tasks asked of users (e.g., visiting/logging in to a website, responding to text messages, and uploading blood glucose readings) and how engagement could be assessed based on these tasks (e.g., the average number of website logins or the number of users who logged in to a website). Interventions required users to perform a wide range of technical tasks, resulting in wide variability in how studies operationalized intervention engagement. In addition, there was wide variation in the extent to which studies described and reported on engagement. That is, some studies provided very basic information regarding the rate of engagement with one task, whereas others provided detailed information on engagement with different parts of the intervention and during specific periods of time.

Mobile-delivered interventions—The 15 mobile-delivered intervention studies in our review reported on engagement with text messages ($n=14$), phone calls ($n=5$), mobile apps (overall, $n=1$; feature-specific, $n=3$), and/or a web-based portion of the intervention ($n=2$). Six studies *only* reported the average number of tasks users engaged with (e.g., the rate of text message responses or interactive voice response [IVR] call completions), five studies *only* reported the number of users who engaged with a specific task (e.g. the number who

responded to a text message, the number who opened an app), and four studies reported on both. In most studies (79%), engagement was based on the full duration of the intervention study period, but in some cases engagement was based on a specific period of time within the study period. For example, Katz et al.'s [25] 12-month intervention reported the average number of blood glucose uploads during the first and last 10 weeks of the study period, but not the full 12 months. When detailed engagement information was included, studies reported on users' median time to respond to a text message [22, 26] and the number of users who met a minimum level of engagement (e.g., users who completed at least nine of 12 IVR calls, users who sent responses at least 3 times per week) [14, 28, 31]. Lastly, several studies categorized users by their level of engagement [21, 25, 29]. For example, Bell et al. [21] organized users into three categories based on how often they viewed daily videos sent via text messages (i.e., non-viewers, periodic viewers, and persistent viewers).

Web-delivered interventions—Of the nine web-delivered intervention studies in our review, most (67%) reported the number of times users logged into the website. Because many web-delivered interventions requested users interact with a website, studies also (44%) reported engagement with these specific tasks (e.g., number of blood glucose uploads and discussion posts, how often goals were set) [36–39]. In addition, three studies (33%) reported the frequency with which different sections of a webpage were viewed [32, 36, 40]. For example, Glasgow et al. [32] reported the exercise section of their website's "Action Plan" area was visited more than any other self-care section. Two studies (22%) reported the number of users who logged into a site based on a minimum frequency (i.e., at least monthly [34], and at least once during the study period [37]), and two studies reported the total average of time spent on the site [32, 35]. Similar to Bell et al. [21], Yu et al. [40] grouped users based on the number of times logged in per month (i.e., non-users, infrequent, frequent, and heavy users), and Jernigan and Lorig [38] separated users into two groups by their level of website participation (i.e., sporadically vs. regularly).

Levels of engagement

Due to wide variability in how engagement was operationalized across studies, we were unable to draw conclusions about how much users engage with mobile- or web-delivered interventions. We were, however, able to identify patterns for those studies using a common operationalization.

Mobile phone engagement—Among the four studies reporting the average response rate to daily, two-way text messages [13, 22, 23, 26], rates ranged from 52% to 84%. The types of daily text messages varied with some requesting adherence information (e.g., whether users took their medication) and others requesting blood glucose readings. Additionally, the intervention length (i.e. the period of time users received text messages) varied. For the two studies reporting lower rates of engagement (57% and 52%), the intervention length was three [23] and six months [26], respectively. Alternatively, for those reporting higher engagement rates (80% and 84%), the intervention length was one [22] and three months [13], respectively. In four studies [14, 21, 23, 28], authors reported the number of users who did not engage at all or did so very minimally; rates ranged from 5.8% to 24%. In the six studies reporting changes in engagement, engagement decreased in two studies

[13, 25], remained relatively stable in three studies [24, 26, 30], and varied by the individual user in one study [29].

Although not reported often enough to identify patterns, there were instances when articles reported associations between user characteristics and intervention engagement. For users around 60 years of age, the probability of completing IVR calls [28] and responding to text messages [13] decreased with increasing age. Additionally, Nelson et al. [13] found users who were non-White and users with lower health literacy were less likely to participate in IVR calls compared to White users and users with higher health literacy, respectively. Lastly, Piette et al. [28] found less education was associated with less IVR completion. The majority of mobile-delivered intervention studies ($n=11$) did not report on associations between user characteristics and engagement or the lack thereof.

Website engagement—For a web-delivered intervention length of at least three months, average logins per month ranged from 0.7 to 7 logins. Studies with fewer logins per month (0.7, 2.4, and 4) had intervention lengths of nine [40], 12 [37], and 13 months [39], respectively, whereas those with more logins per month (7 and 7) had intervention lengths of three [36] and four months [32].

Two studies reported the average time spent on a website [32, 35]. Glasgow et al. [32] reported a total average time of three hours spent on their website, or about seven minutes per visit in a 4-month intervention. Heinrich et al. [35] reported users spent an average of 58 total minutes on their site during a 2-week study, or about 16.3 minutes per visit. Of the four studies reporting on changes in engagement [32, 33, 37, 40], three described a decrease in usage over time [37] (two described different timeframes from the same intervention [32, 33]); in one study, website logins peaked during weeks 10 and 27 of the intervention, which the authors attribute to increased blog use [40].

Only two web-delivered interventions reported associations between user characteristics and engagement [34, 38]. Glasgow et al. [34] reported users with higher health literacy and numeracy skills, and baseline levels of physical activity, were the most engaged (i.e., visited the website at least monthly) during the 12-month intervention period. Additionally, they found users who were non-smokers were more likely to visit the site at least monthly compared to smokers. Jernigan and Lorig [38] did not find differences between American Indian (AI)/Alaskan Native (AN) users and non-AI/AN users in the level of website usage, but reported AI/AN users were more likely to use their website during day time hours compared to non-AI/AN users; the authors noted this may be because more AI/AN users had access to high-speed Internet in an office setting and dial-up Internet at home. Two studies reported no associations between user characteristics and engagement [32, 35], and five studies did not report whether or not there were associations [33, 36, 37, 39, 40].

Facilitators and barriers to intervention engagement

Our review also identified intervention design features used to facilitate user engagement. In addition, we identified barriers to user engagement.

The majority of interventions included in our review employed some form of tailoring ($n=22$); however, the extent to which tailoring was used varied widely. The most common form of tailoring was providing users with personalized feedback based on information they submitted to a system [13, 27, 36, 39]. For example, Ryan et al. [39] provided educational and motivational messages based on users' blood glucose uploads; on average, users uploaded data two times per week during a 13-month study period [39]. In Jennings et al. [36], users received tailored weekly feedback based on completion of an online logbook used to track exercise goals; over 12 weeks, users logged in to the website an average of 21 ± 34 times. Studies also provided tailored content based on users' responses to assessments. In Osborn and Mulvaney [27] and in Nelson et al. [13], daily text messages were tailored to address users' self-reported barriers to medication adherence; users responded to about 82% of messages in a 2-week study [27] and 84% of messages in a 3-month study [13]. In a web-delivered intervention, Jernigan and Lorig [38] provided content based on users' responses to a baseline questionnaire (e.g., feedback on the meaning of their A1c); 85% of users participated with the website regularly (i.e., about 3 times per week for 6 weeks). Other studies used very minimal tailoring. Bell et al. [21] tailored their 6-month intervention by sending text messages at a time specified by the user; over one-third of users did not view any video messages or only did so minimally at the start of the study. Similarly, Arora et al. [20] limited tailoring to users' preference of receiving content in either English or Spanish; only 35% and 9% of users responded to trivia and phone-linked messages, respectively, during a 3-week study. Fisher et al. [24] also allowed users the choice of English or Spanish content and tailored information in appointment reminder messages (appointment time, date, and location); users responded to 68% of messages during a 3-month study.

Additionally, there were vast differences in how interventions prompted, reminded, or incentivized users to engage with an intervention, which we are calling *engagement promotion*. In mobile-delivered interventions, intervention components can serve as reminders to use the intervention. For example, receipt of text messages and/or calls reminds users to engage. In two interventions using IVR, the IVR system made multiple attempts to reach users if they did not answer the call [19, 28]. Alternatively, in mobile interventions consisting of only an app and in web-delivered interventions consisting of only a website, users may be expected to engage without prompting. In order to foster continued use in these types of interventions, studies often sent weekly email reminders to visit a website [32, 36, 38, 40]. In one study, if engagement waned, staff members called users to inquire about their low responsiveness [26].

Studies also examined how support from a healthcare provider or peer impacted intervention engagement. In three studies (13%) users said a provider's involvement in the intervention (e.g., being able to communicate with a provider or knowing a provider was reviewing their messages) facilitated their engagement [19, 26, 37]. Further, two studies using an RCT design compared engagement between a condition receiving only the technology and a condition receiving both the technology and human support. When reporting the 4-month results of their web-delivered intervention, Glasgow et al. [32] reported greater use of the website among users assigned to the condition with the website plus social support compared to the condition with website access only. In contrast, Torbjørnsen et al. [30]

reported no difference in engagement between the condition with access to a mobile app versus the condition with access to the app plus health counseling. This suggests there may be inconsistencies in how providing human support as part of an automated technology impacts users' engagement.

Studies also relied on users' feedback to identify barriers to engaging with mobile- or web-delivered interventions. Three studies (13%), reported technical and usability problems interfered with engagement [13, 37, 40], and in three studies (13%) users said an inability to afford a required cell phone plan, or having limited access to a computer or the Internet hindered engagement [23, 25, 40]. To address cost barriers, Katz et al. [25] provided users with a reduction on their cell phone bill each month in exchange for continued engagement. At study end, 59% who remained active in the program said this discount was needed for their participation. Lastly, there may be unique individual barriers preventing engagement. For example, Yu et al. [40] reported users' sense of futility affected website use, while users in Jernigan and Lorig's [38] study said a lack of time, illness, family obligations, and depression were reasons for not completing website activities. Information on these barriers could inform intervention design choices, such as using a technology that can be more easily integrated into users' lives, requiring minimal deliberate effort. For example, if lack of time is a significant barrier, a text message-based intervention may encourage more engagement than one in which users must access a computer.

Associations between engagement and outcomes

Among studies included in this review, the majority ($n=16$) did not report whether there were associations between engagement and outcomes, but some ($n=3$) reported null associations and others ($n=7$) reported significant associations between engagement and outcomes. Heinrich et al. [35] reported a null association between engagement and diabetes knowledge; however, their web-delivered intervention was only 2 weeks long. Katz et al. [25] found users who remained active in a mobile intervention reduced their hospitalizations and emergency room visits during the 12-month study period compared to the 12 months before the study. Glasgow et al. [32] reported greater website usage was associated with improved eating patterns and physical activity, but not medication adherence at 4 months; notably, of the users whose outcomes improved and who visited the website monthly from months 5 to 12, 53.8% maintained these outcomes at 12 months [33]. Ryan et al. [39] found more frequent engagement in chat messages and interactive activities were each associated with reductions in LDL cholesterol levels. Yu et al. [40] reported users of a diabetes self-care website had greater reductions in diabetes distress than non-users; however, users and non-users did not differ in their self-efficacy or self-care. Finally, in two studies, greater use of an intervention was associated with improved A1c [21, 37]. Bell et al. [21], reported users who persistently viewed text message-delivered videos of diabetes-related tips and reminders experienced the greatest A1c improvement. Jethwani et al. [37] reported users with more blood glucose uploads had the greatest A1c improvement, with those taking less time to upload their first blood glucose reading experiencing the greatest A1c improvement.

Discussion

With the surge of technology-delivered interventions aimed at improving diabetes outcomes, more research is needed to understand how users engage with these interventions. Unlike most face-to-face approaches, users are typically expected to use mobile- and web-delivered interventions on their own time; that is, the intervention relies on the assumption users will engage with the technology on their own volition. The extent to which users actually engage is critical to understanding how interventions affect outcomes. We conducted a narrative review of studies published between 2011–2015 reporting engagement with technology-delivered self-care interventions for adults with T2DM, and found variability across studies in engagement reporting, lower rates of engagement in web-based interventions with a longer duration, older users and those with lower health literacy/less education being less engaged, and more engagement being associated with better outcomes in several studies. Based on our findings, we make recommendations for reporting and improving user engagement.

Our findings are consistent with Schubart et al.'s [12] review of web-delivered self-care interventions which also reported wide variation in how engagement is operationalized. In order to systematically combine and evaluate engagement findings, studies need to adopt uniform engagement reporting. Of the more common operationalizations used in our review, interventions using two-way text messaging should report users' average text message response rate and web-delivered interventions should report average website logins and average time spent on a website. Further, studies should report whether there is an expectation for how much users should engage with an intervention to help identify gaps between intended and actual use. Studies should also report on engagement change by reporting usage at different time intervals throughout the study period (e.g., website logins each month) to identify the point at which engagement wanes. Additionally, rather than only reporting users' engagement with one primary task, studies should break down engagement by reporting on usage with individual tasks. For example, reporting the number of different website tasks completed can reveal those parts of a website that were more or less engaging. Finally, dichotomizing users into high and low engagement groups and examining group differences in outcomes, can identify at which level of engagement users experience benefit.

Despite variability in operationalizing engagement, we did observe some general patterns across studies. First, it appears engagement may be impacted by intervention length, particularly for web-delivered interventions in which lower rates of website use were reported in longer interventions. Additionally, our results support past reviews indicating engagement decreases over the course of an intervention [7, 10]. It may be the longer a user participates in an intervention, the more they lose interest in it. However, because a longer intervention may be necessary to obtain effects, researchers should adopt strategies for boosting engagement throughout an intervention (e.g., tailoring and engagement promotion, discussed below). Lastly, although Cotter et al. [10] did not identify common user characteristics associated with engagement, our review suggests older adults may be less engaged with mobile interventions than younger adults, and individuals with lower health literacy/less education may be less engaged with both mobile- and web-based interventions compared to more literate/educated individuals.

Based on patterns of engagement reporting, it appears interventions with more tailoring may have higher levels of engagement than those with less tailoring. Technology-delivered interventions using tailored content are more effective at improving self-care than those using targeted or generic content [41, 42]. Most interventions in our review used some form of tailoring, although the type and extent of tailoring varied across interventions and was sometimes limited to users' preferred time to receive elements of the intervention. One way to improve user engagement is to adapt intervention content to users' individual needs [12]. Rather than only tailoring message delivery based on users' preferences (e.g., timing and frequency of texts), more interventions should tailor intervention content based on users' unique concerns for improving self-care (e.g., content addressing barriers to self-care adherence [13]). Similarly, just-in-time interventions which provide immediate, tailored support based on user data sent to a system (e.g., feedback on blood glucose uploads), may also improve engagement [43].

When comparing mobile- and web-delivered interventions in our review, we identified a need for more web-delivered interventions to use engagement promotion to remind and encourage users to engage. This finding aligns with both Connelly et al.'s [7] and Brouwer et al.'s [17] reviews that found additional components within the technology (e.g., emails) increased intervention engagement. Future research should explore additional forms of prompting in the form of rewards and/or new content to sustain users' interest in an intervention over time [17]. This may be particularly important in longer interventions.

Technology-delivered interventions should also anticipate possible barriers to engagement. If working with a low-income population, for example, adopting a low-cost technology (e.g., basic cell phone technology: text messaging and voice communications) and/or providing a stipend to cover costs may improve engagement [20]. Additionally, conducting usability studies can help identify and address technical issues that may discourage engagement in a longer trial [44].

Our review compiled evidence that increased use of a technology-delivered intervention is associated with improved outcomes among adults with T2DM. Among the eight studies reporting intervention engagement's association with outcomes, seven found at least one association between increased usage and outcome improvement, including two studies reporting improved glycemic control [21, 37]. Bell et al. [21] grouped users by their level of engagement to determine at what dose of the intervention users would benefit. Users who viewed >10 video messages per month experienced the greatest improvement in A1c, but none of the users viewed messages daily [21]. Considering that users still benefited despite not engaging every day, Bell et al. [21] suggest intermittent reinforcement (i.e., frequent, but not daily contact) may be an effective approach for sustaining users' engagement and improving A1c. More research needs to examine how dosing level may explain the association between engagement and outcomes. Additionally, studies should examine whether dosing should be tailored to different types of users; that is, some users may get annoyed with daily messages and engage more regularly with weekly messages, whereas others may appreciate daily contact.

There are several limitations to our review. The majority of studies included in our review were feasibility, usability, and pilot studies. Because these types of studies tend to have a shorter duration, engagement may be different in longer trials testing the same intervention. Further, we excluded studies from our review testing interventions with >50% telehealth because we were primarily interested in user engagement with automated interventions. Given the human-to-human element of telehealth, engagement with those interventions may be higher than with automated interventions. Several studies in our review reported a human's involvement in the intervention was important to users' engagement. Additionally, although we organized interventions by whether they were primarily mobile- or web-based, some mobile interventions used web components, which may have affected our results. As more interventions combine mobile and web platforms, research should explore how engagement differs between these combined interventions and those relying on just one platform. Despite our attempt to identify relevant articles, it is possible some were missed and publication bias led to an underreporting of data. Finally, the studies included in our review all had samples with >50% T2DM, limiting the generalizability of our findings to other patient populations.

Conclusions

Despite variability in engagement reporting and its associations, engagement remains difficult to sustain in technology-delivered self-care interventions for adults with T2DM and can be impacted by a variety of factors including how the intervention promotes engagement and the characteristics of users. In order to advance our understanding of how engagement varies and impacts outcomes, studies need to report on engagement in a standardized way, consistently report whether or not engagement is associated with user characteristics and outcomes, and report on engagement in longer-term RCTs. Exploring engagement as a mechanism by which an intervention affects outcomes and the optimal dose of engagement will inform how and when users experience benefit.

Acknowledgments

Drs. Nelson and Osborn are supported by NIH/NIDDK R01-DK100694, and Dr. Osborn is also supported by K01-DK087894. Dr. Cherrington is supported by NIH/NIDDK P30-DK07962.

References

Papers of particular interest, published recently, have been highlighted as:

- Of importance

1. Centers for Disease Control and Prevention. Reports to Congress: Diabetes Report Card 2014. 2014.
2. Osborn C, Mayberry L, Kim J. Medication adherence may be more important than other behaviors for optimizing glycemic control among low-income adults. *J Clin Pharm Ther.* in press;in press.
3. Gao J, Wang J, Zheng P, Haardörfer R, Kegler MC, Zhu Y, Fu H. Effects of self-care, self-efficacy, social support on glycemic control in adults with type 2 diabetes. *BMC family practice.* 2013; 14:1. [PubMed: 23281961]
4. Kirkman MS, Rowan-Martin MT, Levin R, Fonseca VA, Schmittiel JA, Herman WH, Aubert RE. Determinants of adherence to diabetes medications: findings from a large pharmacy claims database. *Diabetes care.* 2015; 38:604–609. [PubMed: 25573883]

5. García-Pérez L-E, Álvarez M, Dilla T, Gil-Guillén V, Orozco-Beltrán D. Adherence to therapies in patients with type 2 diabetes. *Diabetes Therapy*. 2013; 4:175–194. [PubMed: 23990497]
6. Holtz B, Lauckner C. Diabetes management via mobile phones: a systematic review. *Telemedicine journal and e-health : the official journal of the American Telemedicine Association*. 2012; 18:175–184. [PubMed: 22356525]
7. Connelly J, Kirk A, Masthoff J, MacRury S. The use of technology to promote physical activity in Type 2 diabetes management: a systematic review. *Diabetic medicine : a journal of the British Diabetic Association*. 2013; 30:1420–1432. [PubMed: 23870009]
8. Pal K, Eastwood SV, Michie S, Farmer A, Barnard ML, Peacock R, Wood B, Edwards P, Murray E. Computer-based interventions to improve self-management in adults with type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care*. 2014; 37:1759–1766. [PubMed: 24855158]
9. Liang X, Wang Q, Yang X, Cao J, Chen J, Mo X, Huang J, Wang L, Gu D. Effect of mobile phone intervention for diabetes on glycaemic control: a meta-analysis. *Diabetic medicine : a journal of the British Diabetic Association*. 2011; 28:455–463. [PubMed: 21392066]
10. Cotter AP, Durant N, Agne AA, Cherrington AL. Internet interventions to support lifestyle modification for diabetes management: a systematic review of the evidence. *Journal of Diabetes and its Complications*. 2014; 28:243–251. [PubMed: 24332469]
11. Mulvaney SA, Ritterband LM, Bosslet L. Mobile intervention design in diabetes: review and recommendations. *Curr Diab Rep*. 2011; 11:486–493. [PubMed: 21960031]
12. Schubart JR, Stuckey HL, Ganeshamoorthy A, Sciamanna CN. Chronic health conditions and internet behavioral interventions: a review of factors to enhance user engagement. *Computers, informatics, nursing : CIN*. 2011; 29:TC9–20.
13. Nelson LA, Mulvaney SA, Gebretsadik T, Ho YX, Johnson KB, Osborn CY. Disparities in the use of a mHealth medication adherence promotion intervention for low-income adults with type 2 diabetes. *Journal of the American Medical Informatics Association : JAMIA*. 2015
14. Capozza K, Woolsey S, Georgsson M, Black J, Bello N, Lence C, Oostema S, North C. Going mobile with diabetes support: a randomized study of a text message-based personalized behavioral intervention for type 2 diabetes self-care. *Diabetes spectrum : a publication of the American Diabetes Association*. 2015; 28:83–91. [PubMed: 25987806]
15. Fjeldsoe BS, Marshall AL, Miller YD. Behavior change interventions delivered by mobile telephone short-message service. *American journal of preventive medicine*. 2009; 36:165–173. [PubMed: 19135907]
16. Hanauer DA, Wentzell K, Laffel N, Laffel LM. Computerized Automated Reminder Diabetes System (CARDS): e-mail and SMS cell phone text messaging reminders to support diabetes management. *Diabetes Technol Ther*. 2009; 11:99–106. [PubMed: 19848576]
17. Brouwer W, Kroeze W, Crutzen R, de Nooijer J, de Vries NK, Brug J, Oenema A. Which intervention characteristics are related to more exposure to internet-delivered healthy lifestyle promotion interventions? A systematic review *Journal of medical Internet research*. 2011; 13:e2. [PubMed: 21212045]
18. Popay J, Roberts H, Sowden A, Petticrew M, Arai L, Rodgers M, Britten N, Roen K, Duffy S. Guidance on the conduct of narrative synthesis in systematic reviews. *ESRC methods programme*. 2006; 15:047–071.
19. Aikens JE, Rosland AM, Piette JD. Improvements in illness self-management and psychological distress associated with telemonitoring support for adults with diabetes. *Prim Care Diabetes*. 2015; 9:127–134. [PubMed: 25065270]
20. Arora S, Peters AL, Agy C, Menchine M. A mobile health intervention for inner city patients with poorly controlled diabetes: proof-of-concept of the TExT-MED program. *Diabetes Technol Ther*. 2012; 14:492–496. [PubMed: 22524591]
- 21. Bell AM, Fonda SJ, Walker MS, Schmidt V, Vigersky RA. Mobile phone-based video messages for diabetes self-care support. *J Diabetes Sci Technol*. 2012; 6:310–319. This study grouped users by their level of engagement in order to examine which group experienced the most improvement in A1c. [PubMed: 22538140]

22. Dick JJ, Nundy S, Solomon MC, Bishop KN, Chin MH, Peek ME. Feasibility and usability of a text message-based program for diabetes self-management in an urban African-American population. *J Diabetes Sci Technol*. 2011; 5:1246–1254. [PubMed: 22027326]
23. Dobson R, Carter K, Cutfield R, Hulme A, Hulme R, McNamara C, Maddison R, Murphy R, Shepherd M, Strydom J, Whittaker R. Diabetes Text-Message Self-Management Support Program (SMS4BG): A Pilot Study. *JMIR mHealth and uHealth*. 2015; 3:e32. [PubMed: 25830952]
24. Fischer HH, Moore SL, Ginosar D, Davidson AJ, Rice-Peterson CM, Durfee MJ, MacKenzie TD, Estacio RO, Steele AW. Care by cell phone: text messaging for chronic disease management. *Am J Manag Care*. 2012; 18:e42–47. [PubMed: 22435883]
25. Katz R, Mesfin T, Barr K. Lessons from a community-based mHealth diabetes self-management program: “it’s not just about the cell phone”. *J Health Commun*. 2012; 17(Suppl 1):67–72.
26. Nundy S, Dick JJ, Chou CH, Nocon RS, Chin MH, Peek ME. Mobile phone diabetes project led to improved glycemic control and net savings for Chicago plan participants. *Health Aff (Millwood)*. 2014; 33:265–272. [PubMed: 24493770]
27. Osborn CY, Mulvaney SA. Development and feasibility of a text messaging and interactive voice response intervention for low-income, diverse adults with type 2 diabetes mellitus. *J Diabetes Sci Technol*. 2013; 7:612–622. [PubMed: 23759393]
28. Piette JD, Valverde H, Marinec N, Jantz R, Kamis K, de la Vega CL, Woolley T, Pinto B. Establishing an independent mobile health program for chronic disease self-management support in bolivia. *Front Public Health*. 2014; 2:95. [PubMed: 25165687]
29. Tata N, Arsand E, Bratteteig T, Hartvigsen G. Usage and perceptions of a mobile self-management application for people with type 2 diabetes: qualitative study of a five-month trial. *Studies in health technology and informatics*. 2013; 192:127–131. [PubMed: 23920529]
30. Torbjornsen A, Jenum AK, Smastuen MC, Arsand E, Holmen H, Wahl AK, Ribu L. A Low-Intensity Mobile Health Intervention With and Without Health Counseling for Persons With Type 2 Diabetes, Part 1: Baseline and Short-Term Results From a Randomized Controlled Trial in the Norwegian Part of RENEWING HEALTH. *JMIR mHealth and uHealth*. 2014; 2:e52. [PubMed: 25499592]
31. Wood FG, Alley E, Baer S, Johnson R. Interactive Multimedia Tailored to Improve Diabetes Self-Management. *The Nursing clinics of North America*. 2015; 50:565–576. [PubMed: 26333610]
32. Glasgow RE, Christiansen SM, Kurz D, King DK, Woolley T, Faber AJ, Estabrooks PA, Strycker L, Toobert D, Dickman J. Engagement in a diabetes self-management website: usage patterns and generalizability of program use. *J Med Internet Res*. 2011; 13:e9. [PubMed: 21371992]
33. Glasgow RE, Kurz D, King D, Dickman JM, Faber AJ, Halterman E, Woolley T, Toobert DJ, Strycker LA, Estabrooks PA, Osuna D, Ritzwoller D. Twelve-month outcomes of an Internet-based diabetes self-management support program. *Patient Educ Couns*. 2012; 87:81–92. [PubMed: 21924576]
34. Glasgow RE, Strycker LA, King DK, Toobert DJ. Understanding who benefits at each step in an internet-based diabetes self-management program: application of a recursive partitioning approach. *Med Decis Making*. 2014; 34:180–191. [PubMed: 23913917]
35. Heinrich E, de Nooijer J, Schaper NC, Schoonus-Spit MH, Janssen MA, de Vries NK. Evaluation of the web-based Diabetes Interactive Education Programme (DIEP) for patients with type 2 diabetes. *Patient Educ Couns*. 2012; 86:172–178. [PubMed: 21616626]
36. Jennings CA, Vandelanotte C, Caperchione CM, Mummery WK. Effectiveness of a web-based physical activity intervention for adults with Type 2 diabetes-a randomised controlled trial. *Prev Med*. 2014; 60:33–40. [PubMed: 24345601]
- 37. Jethwani K, Ling E, Mohammed M, Myint UK, Pelletier A, Kvedar JC. Diabetes connect: an evaluation of patient adoption and engagement in a web-based remote glucose monitoring program. *J Diabetes Sci Technol*. 2012; 6:1328–1336. This study provided a thorough description of website engagement patterns and compared differences in outcomes between users who were more or less engaged. [PubMed: 23294777]
38. Jernigan VB, Lorig K. The internet diabetes self-management workshop for American Indians and Alaska Natives. *Health Promot Pract*. 2011; 12:261–270. [PubMed: 20534807]

39. Ryan JG, Schwartz R, Jennings T, Fedders M, Vittoria I. Feasibility of an internet-based intervention for improving diabetes outcomes among low-income patients with a high risk for poor diabetes outcomes followed in a community clinic. *Diabetes Educ.* 2013; 39:365–375. [PubMed: 23610181]
40. Yu CH, Parsons JA, Mamdani M, Lebovic G, Hall S, Newton D, Shah BR, Bhattacharyya O, Laupacis A, Straus SE. A web-based intervention to support self-management of patients with type 2 diabetes mellitus: effect on self-efficacy, self-care and diabetes distress. *BMC Med Inform Decis Mak.* 2014; 14:117. [PubMed: 25495847]
41. Lustria ML, Noar SM, Cortese J, Van Stee SK, Glueckauf RL, Lee J. A meta-analysis of web-delivered tailored health behavior change interventions. *J HEALTH COMMUN.* 2013; 18:1039–1069. [PubMed: 23750972]
42. Head KJ, Noar SM, Iannarino NT, Grant Harrington N. Efficacy of text messaging-based interventions for health promotion: A meta-analysis. *Social Science & Medicine.* 2013; 97:41–48. [PubMed: 24161087]
43. Klasnja P, Pratt W. Healthcare in the pocket: mapping the space of mobile-phone health interventions. *Journal of biomedical informatics.* 2012; 45:184–198. [PubMed: 21925288]
44. Nelson LA, Bethune MC, Lagotte A, Osborn CY. The usability of Diabetes MAP: A web-delivered intervention for improving medication adherence. *JMIR Human Factors.* in press.

Figure 1.

Table 1

Characteristics of interventions included in this review

Mobile-delivered interventions						
Authors [reference]	Study Design	Sample size, mean age, majority ethnicity, mean A1c; % T2DM	Intervention characteristics	Intervention Tailoring	Duration	Outcomes
Capozza et al. [14]	2-arm RCT	N=93; mean age 53.25 years; 76% NHW; A1c 9.0%; 100% T2DM	Care4Life sends 1–7 daily texts with content on diabetes education, health improvement, and self-care behaviors. Users have access to a web-based portal for accessing response trends to two-way texts.	Type, timing, and frequency of text messages.	6 months	A1c improved in the intervention group and in the control group. There were no between differences at 3 or 6 months.
Aikens et al. [19]	Non-experimental pre-post	N=301; mean age 66.7 years; 92.8% NHW; A1c not reported; 100% T2DM	mHealth service makes weekly automated IVR calls to users asking for responses to questions about experiences with diabetes and providing immediate verbal feedback.	Verbal feedback based on user experience. System provides immediate problem-tailored support if user reports difficulty with self-care.	6 months	Self-reported medication adherence, physical functioning, depressive symptoms, and diabetes-related distress improved among users at 3 weeks.
Arora et al. [20]	Proof-of-concept study	N=23; mean age 45.4 years; 70% Latino; Median A1c 8.9% ± 2.2%; 70% T2DM	TEXT-MED sends users 3 text messages daily for 3 weeks providing educational and motivational content, medication reminders, healthy living challenges, trivia, or phone links.	Choice of receiving content in either English or Spanish	3 weeks	Self-reported healthy eating, physical activity, foot checks, and self-efficacy improved (non-statistical) among users at 3 weeks.
Bell et al. [21]	2-arm RCT	N=64; mean age 58 years; 58% AA; A1c 9.3% ± 1.3%; 92% T2DM	Intervention sends daily, asynchronous one-way videos of diabetes-related tips and reminders delivered via cell phones. Users provided with cell phones and service.	Time of day for receiving video messages.	6 months (tracked change for 1 year)	A1c improved more in the intervention group compared to the control group over 12 months, especially in the first 3 months
Dick et al. [22]	Feasibility/pilot study	N=19; mean age 55 years; 100% AA; A1c not reported; 100% T2DM	SMS-DMCare sends texts with personalized reminders for medication, SMBG, foot care, and appointments. Users send texts reporting on their self-care adherence.	Text timing and frequency; option to receive additional texts related to self-care.	4 weeks	Self-reported medication adherence and foot checks improved among users at 1 month and 2 months.
Dobson et al. [23]	Pilot study	N=42; mean age 45.7 years; 38% NZE; A1c 10.3%; 71% T2DM	Self-Management Support for Blood Glucose sends modules via texts on diabetes education, encouragement, and illness perceptions. Users and clinicians have access to a website to view users' BG responses trends.	Timing and frequency of BG messages; user's name included in messages.	3 months	A1c improved among users with complete data (62 %) at 3 months.
Fischer [24]	Feasibility study	N=47; 35.4% were 50–59 years of age; 77% Hispanic/Latino;	Intervention sends 3 text messages per week requesting BG readings and 3 text messages with appointment reminders in advance of	Choice of English or Spanish; reminders include time, date, and location of user's appointment.	3 months	BG result availability improved among users at 3 months. Cancellation and no-

Mobile-delivered interventions						
Authors [reference]	Study Design	Sample size, mean age, majority ethnicity, mean A1c; % T2DM	Intervention characteristics	Intervention Tailoring	Duration	Outcomes
Katz et al. [25]	Feasibility/pilot study	A1c not reported; >90% T2DM N=32; mean age 39 years; 100% AA; A1c 8%; 100% T2DM	each scheduled appointment (requests "Y" or "N" response). The WellDoc Diabetes Manager system is an interactive platform for users, case managers, and providers to track users' BG, and receive real-time feedback. Users offered web-enabled handsets and a low-cost monthly prepaid phone contract.	Case managers sent personalized messages weekly with feedback based on BG readings.	12 months	show rates did not change at 3 months. Emergency department visits and hospitalizations decreased among active users during the 12-month study compared to the 12-month period before the study.
Nelson et al. [13]	Used data from quasi experimental study	N=80; mean age 50 years; 69% Non-white; A1c 8.3%; 100% T2DM (patients recruited from FQHC)	MEssaging for Diabetes (MED) sends daily one-way text messages addressing user-specific barriers to adherence, daily two-way text messages inquiring about med adherence, and weekly IVR calls providing adherence feedback, encouragement, and problem-solving questions.	Timing of texts and IVR calls; text content tailored to users' self-reported adherence barriers. IVR call feedback tailored to adherence text responses.	3 months	Engagement results reported in Table 2.
Nundy et al. [26]	Quasi-experimental/pre-post	N=74; mean age 53 years; 65.6% AA; A1c 7.6%; users had T1DM or T2DM	CareSmarts is a mHealth diabetes program providing self-care support and team-based care via texts. Users receive self-care texts that either prompt a specific action or ask questions. Users follow a curriculum of modules on different self-care topics.	Text content modified every 2 weeks; Users have option of receiving more modules or only self-care texts.	6 months	Self-reported SMBG, foot care, and med adherence improved among the intervention group at 6 months. A1c improved in the intervention group did not change in the control group at 6 months.
Osborn and Mulvaney [27]	Feasibility/usability study	N=20; mean age 51.6 years; 45% AA; A1c 7.6%; 100% T2DM (patients recruited from FQHC)	(See description for MED above – Nelson et al. [13])	See description of tailoring above – Nelson et al. [13].	3 rounds of 2 week-testing	Engagement results reported in Table 2.
Piette et al. [28]	Non-experimental pre-post	N=165; mean age 65 years; Patients recruited in Bolivia; Did not report A1c; 55.2% T2DM	mHealth platform sends 12 weekly IVR calls assessing users' self-care behaviors, perceived health status, and diabetes symptoms. After providing responses, users receive self-care education.	Self-care education tailored to users' responses to IVR calls	12 weeks	Likelihood of reporting good health increased during the 12-month study, and was associated with improved self-reported med adherence.
Tatara et al. [29]	Usability study	N=10; mean age 57.2 years; Did not report ethnicity; Did not	The Few Touch application (FTA) is a mobile app with self-care systems for BG, food habits, and physical activity, a goal-setting system, and a diabetes information look-up	Goal setting and feedback tailored to users' input.	5 months	Engagement results reported in Table 2.

Mobile-delivered interventions						
Authors [reference]	Study Design	Sample size, mean age, majority ethnicity, mean A1c; % T2DM	Intervention characteristics	Intervention Tailoring	Duration	Outcomes
Torbjornsen et al. [30]	3-arm RCT	report A1c; 90% T2DM N=151; mean age 57.7 years; 100% Norwegian; A1c 8.2%; 100% T2DM	system. Users were provided a mobile phone and BG meter. In both intervention arms, users interacted with FTA (see Tataru et al. [29]). In one intervention arm, users also received health counseling. All users were provided with a mobile phone.	See description of tailoring above – Tataru et al. [29].	4- month results of 12- month study	A1c improved in both intervention groups and in the control group at 4 months. There were no between group differences.
Wood et al. [31]	Pilot study	N=7; mean age 50.1 years; 83% NHW; did not report A1c; 100% T2DM (patients recruited at free clinic)	Diabetes 101 is a native app including videos of T2DM self-care activities and reminders, and a dictionary with 50 common diabetes terms defined in lay terminology and an audio recording of pronunciation. Users were given iPod Touch and asked to use the app as frequently as desired.	Option to view a video was based on users' knowledge about the topic (assessed with a brief quiz)	2 weeks	Health literacy and self-efficacy improved among users at 2 weeks.
Web-delivered interventions						
Glasgow et al. [32]	Used data from RCT [33] (baseline to 4 months)	N=270; mean age 57.8 years; 67.4% NHW; 22.3% Hispanic/Latino; A1c 8.2%±1.8%; 100% T2DM	My Path to Healthy Life is a web-based diabetes self-care intervention. It consists of interactive and multimedia features including quiz questions and tips. Action plans list users' goal and tailored information for reaching goal. Users have access to forums, SM displays, and test results. Half of intervention arm received social/peer support.	Choice of English or Spanish; Optional audio voice-overs, choice in setting behavioral goals, and choice among various features.	4 months	Engagement reported in Table 2.
Glasgow et al. [33]	3-arm RCT	N=463; mean age 58.4 years; 72% NHW; 21.8% Hispanic; 100% T2DM	See My Path to Healthy Life described above (Glasgow et al. [32]). Participants were randomly assigned to either 1) My Path to Healthy Life, 2) My Path to Healthy Life plus enhanced social support, or 3) usual care.	See description above (Glasgow et al. [32]); also, system re-tailors three self-care goals at 6 months	12 months	Self-reported health behaviors improved more in both of the intervention groups than in the control group at 12 months. Biological and psychosocial outcomes improved moderately in all groups.
Glasgow et al. [34]	Used data from RCT [33] (baseline to 12 months)	N=462; mean age 58.4 years; 72% NHW; 21.8% Hispanic; 100% T2DM	See My Path to Healthy Life described above (Glasgow et al. [32]). Data from participants in either of two intervention conditions (see Glasgow [33] described above) were combined in analyses to identify subgroups associated with success at each step of the program.	See description above (Glasgow et al. [32])	12 months	Engagement reported in Table 2.

Mobile-delivered interventions

Authors [reference]	Study Design	Sample size, mean age, majority ethnicity, mean A1c, % T2DM	Intervention characteristics	Intervention Tailoring	Duration	Outcomes
Heinrich et al. [35]	3-arm RCT	N=135; mean age 57 years; Race/ethnicity not reported; A1c not reported; 100% T2DM	Diabetes Interactive Education Programme (DIEP) is a web-based diabetes education program. The website provides an overview of T2DM in seven chapters with both basic and detailed information. Information is mostly presented in spoken language, supported by headlines, images, video and real patient experiences on different topics.	None	2 weeks	Diabetes knowledge scores improved in the intervention group at 2 weeks. No changes in knowledge were observed in the control group.
Jennings et al. [36]	2-arm RCT	N=397; mean age 58 years; participants recruited in Australia; A1c not reported; 100% T2DM	The intervention consisted of a website aiming to improve diabetes self-care based on the Theory of Planned Behavior. Components included educational modules, social support, positive reinforcement, personalized feedback and activities such as goal setting and planning.	Personalized goals and personalized feedback	12 weeks (tracked change for 36 weeks)	Self-reported physical activity improved more among those who completed the intervention compared to the control group at 12 weeks (the intent-to-treat analysis was not significant).
Jethwani et al. [37]	Non-experimental pre-post	N=75; mean age 61 years; race/ethnicity not reported; A1c 9.2%; 100% T2DM	Diabetes Connect (DC) is a web-based glucose home monitoring program facilitating diabetes self-care that allows patients and providers to communicate. DC enables users to collect BG readings and transmit data to a web-based platform. Users can monitor trends and share data with providers in real time. Users have access to educational material.	Users can monitor their trends in BG readings.	12 months	Both the number of high and low out-of-range BG events declined among users during the 12-month study. A1c improved among users from pre-intervention to post-intervention.
Jernigan and Lorig [30]	Feasibility/Pilot study	Two pilots: 1) AI/ANs; N=27; mean age: 42.9 years; A1c 6.8%; 2) non-AI/ANs; N=27; Mean age: 54.2 years; A1c: 6.9%	The Stanford Internet Diabetes Self-management Workshop is based on self-efficacy theory and offers online activities and resources for managing diabetes. Users are asked to log in at least 3 times/week for about 2 hours and participate in weekly activities.	Intervention content is based on users' responses to baseline questions. Users are asked to form a personal action plan each week.	6 weeks	Engagement reported in Table 2.
Ryan et al. [39]	Feasibility study	N=21; mean age 54.4 years; 77.8% AA; A1c 7.54%; 100% T2DM (patients were all low-income, majority not insured)	HEAT-IT-UP is a web-based suite of applications, designed to improve diabetes outcomes via improvements in working alliance, health-related quality of life, and self-efficacy. Users can communicate with providers and network with peers. Users were given a computer, glucometer and test strips, internet connectivity, and user support.	Users given feedback based on uploaded blood glucose values.	13 months	Triglyceride levels increased, and LDL cholesterol and total cholesterol both decreased (non-significant) among users at 13 months. A1c improved among users at 13 months.
Yu et al. [40]	Non-experimental pre-post	N=81; mean age 57.2 years; 62% White, 30%	The Diabetes Online Companion is a diabetes self-care website based on self-efficacy theory; it provides general static information,	Interactive tailored diabetes-related information	9 months	Self-reported self-care improved among users at 9 months. Self-

Mobile-delivered interventions						
Authors [reference]	Study Design	Sample size, mean age, majority ethnicity, mean A1c; % T2DM	Intervention characteristics	Intervention Tailoring	Duration	Outcomes
		Asian; A1c 7.6%; 100% T2DM	tailored interactive information, SM logs, and a community blog.			efficacy, diabetes distress, and clinical outcomes (i.e., A1c, blood pressure, LDL-C, and weight) did not change from baseline to 9 months.
<i>RCT</i> randomized controlled trial, <i>NHW</i> non-Hispanic White, <i>A1c</i> glycemic control, <i>T2DM</i> type 2 diabetes, <i>mHealth</i> mobile health, <i>IVR</i> interactive voice response, <i>med</i> medication, <i>AA</i> African-American, <i>SMBG</i> self-monitoring of blood glucose, <i>NZE</i> New Zealand European, <i>BG</i> blood glucose, <i>FQHC</i> federally qualified health center, <i>T1DM</i> type 1 diabetes, <i>AI/AN</i> American Indian/Alaskan Native						

Table 2

Engagement descriptions for studies included in the review

Authors [reference]; duration	Engagement promotion	Engagement results	Engagement change	Engagement associations with user characteristics and outcomes	User satisfaction and concerns
Mobile-delivered interventions					
Capozza et al. [14]; 6 months	Users received messages daily.	Among those enrolled, 29% used the intervention for at least 90 days and sent responses to texts at least three times/week. 11% of users never replied to a text.	Not reported	Not reported	Most (94%) would recommend the intervention; 45% said the intervention improved diabetes knowledge.
Aikens et al. [19]; 3 and 6 months	Users received weekly IVR calls; if they did not answer, the program made multiple call attempts; Users received \$20 for each of two interviews.	Users completed an IVR call during 84% of the weeks they participated.	Not reported	Not reported	89% wanted the intervention to become part of their usual care.
Arora et al. [20]; 3 weeks	Users received two texts/week; the phone-linked text included a number users could call to receive a free gift. Users were paid \$50 for phone costs.	Among users, 35% responded to trivia text and 9% called in responses to phone linked text.	Not reported	Not reported	90% wanted to continue the intervention; 100% would recommend it to others.
Bell et al. [21]; 6 months	Users received video texts daily; each text could be viewed multiple times for 24h.	Eleven users either did not view videos at all or only did briefly at the start; 10 users viewed videos throughout, but sometimes missed whole weeks; 10 users viewed videos persistently.	Not reported	Persistent viewers experienced the greatest A1c improvement.	Not reported
Dick et al. [22]; 4 weeks	Users received at least one text/day requesting a response.	Users responded to 80% of text messages requiring a response; median response time was 6.1 minutes. It was common for patients to respond "Ok" or "Thank you" to messages not requiring a response.	Not reported	Not reported	Overall, users were very satisfied with the intervention and agreed it was easy to use.
Dobson et al. [23]; 3 months	Users received up to four SMBG texts/day and were given a voucher (NZD \$20) at study end to cover phone costs.	24% of users never replied to a text. Of those who responded to at least one text, the average response rate was 57% (range 1 to 99%). Only 12% of users reported accessing and viewing their BG results online.	Not reported	Not reported	Cost was a leading barrier for users who did not reply to a text. Not having access to a computer/Internet was a common barrier to using the website component.
Fischer et al. [24]; 3 months	Users received three texts/week requesting BG readings three text reminders before each appointment.	Users responded to 68.2% of text message prompts. Most users (66%) provided BG data upon request.	79% of users responded regularly to >50% of message prompts.	Not reported	Users rated texts favorably. They wanted the intervention to continue and were

Authors [reference]; duration	Engagement promotion	Engagement results	Engagement change	Engagement associations with user characteristics and outcomes	User satisfaction and concerns
Mobile-delivered interventions					
Katz et al. [25]; 12 months	Users who routinely entered BG readings received a \$20 discount on their cell phone bill each month.	Active users averaged 9.1 BG entries/week in the first 10 weeks and 3.3 BG entries/week in the last 10 weeks. Inactive users averaged 5.5 BG entries/week.	BG entries/week for active group decreased from 9.1 in the first 10 weeks to 3.3 in last 10 weeks.	Hospitalizations and emergency department visits reduced for active users.	willing to recommend it to others. Nine users discontinued using the intervention because cell phone costs. Active users satisfied with the intervention; 59% said discounts were needed to participate.
Nelson et al. [13]; 3 months	Users received a daily tailored text addressing med adherence barriers, a daily text assessing adherence, and a weekly IVR call with med adherence feedback and problems-solving questions.	Users responded to 84% of assessment text messages and participated in 57% of the IVR calls.	Participation in IVR calls appeared to decrease following the 5 th week.	After age 50, the probability of text responses decreased with increasing age. Whites and users with higher health literacy were more likely to complete IVR calls than non-Whites and users with lower health literacy.	There were technical issues with IVR calls that may have limited the opportunities for engagement.
Nundy et al. [26]; 6 months	Users received daily texts with prompts to respond. If users had low responsiveness, research staff contacted users by phone.	Users responded to an average of 52% of self-assessment questions (range, 0–91%). The average median response time was 19 minutes (range, 1–192 minutes).	Response rate remained relatively constant during the study period (between 48% and 55% each month).	Not reported	Most users (73%) were satisfied with the intervention; knowing a health provider was reviewing messages was important for engagement (88%).
Osborn and Mulvaney [27]; three rounds of 2-week testing	See Nelson et al. [13] above.	Out of 14 opportunities to respond to texts, users responded to an average of 10.3±2.4 texts in Round 1, 13.5±1.0 texts in Round 2, and 9.8±2.4 texts in Round 3. IVR calls were answered by 80% of users in Round 1, 67% of users in Round 2, and 65% of users in Round 3.	Not reported	Not reported	Users in all three rounds said they would recommend the intervention to others with diabetes. Users rated the helpfulness of the tailored text 8.3 out of 10 and the helpfulness of the IVR call 7.5 out of 10.
Piette et al. [28]; 12 weeks	IVR calls sent weekly; if user did not respond, multiple call attempts were made at times indicated by the user.	Users completed 51% of calls. Among all users, 35% completed > 9 calls and 15% completed calls for 11 or 12 weeks in which one was attempted. 19% of users never participated in an IVR call.	Not reported	More education associated with more IVR call completion. IVR call completion declined after age 65.	95% of users would recommend the intervention; 84% of users would use it again. Several users had problems hearing and responding to questions on the call. Some users

Authors [reference]; duration	Engagement promotion	Engagement results	Engagement change	Engagement with user characteristics and outcomes	User satisfaction and concerns
Mobile-delivered interventions					
Tatara et al. [29]; 5 months	None/Not reported	Four users were categorized as having frequent use with a positive experience; three users had moderate use with a neutral experience; four users had little use with a mixed experience.	Five users increased usage with at least one of the three self-care systems, and four users decreased usage.	Greater usage was associated with more perceived usefulness of the intervention	wanted to speak with a provider. Many usability problems were reported regardless of age. However, most design features were perceived positively.
Torbjornsen et al. [30]; 4 months	None/Not reported	Engagement was assessed with the number of BG measurements and keystrokes across the study period. Trend of app use did not differ between the intervention groups.	Usage was lowest in the first month, increased slightly in the second month, and remained stable in the third and fourth months.	Not reported	Not reported
Wood et al. [31]; 2 weeks	None/not reported	All seven users opened the app at least twice, and some opened it as many as 16 times.	Not reported	Not reported	Users liked the flexibility of using the app without Internet and appreciated videos showing "real" people living with T2DM.
Web-delivered interventions					
Glasgow et al. [32]; 4 months	Users received periodic prompts to return to the site via email and IVR; Users were reimbursed \$25 for follow-ups.	Website usage varied widely (1 to 119 visits) (mean, 28 visits). Total time spent on site averaged about 3 h (7 min/visit). Most frequently visited section was self-monitoring. Users in the website plus social support condition had higher website use.	Usage decreased over time; 70% of users visited at least weekly during the first 6 weeks and 47% visited during weeks 7 to 16.	Website use not related to baseline variables. More website use was associated with greater improvement in eating and physical activity, but not med adherence.	Not reported
Glasgow et al. [33]; 12 months	Users assigned to social support condition received follow-up calls at 2 and 8 weeks to answer questions and discuss self-care goals.	Not reported	Website use high during the first 4 months; afterward, site visits declined from an average of 11/month to an average of about 3/month in month 12.	Not reported	Not reported
Glasgow et al. [34]; 12 months	See Glasgow et al. [33] described above.	Of those who remained in the study from baseline to 4 months, 219 (81.1%) visited the website at least monthly.	Not reported (see above; Glasgow et al. [33]).	The most engaged group had higher numeracy, health literacy, and baseline levels of physical activity. Of users who visited the website at least monthly, 58%	Not reported

Authors [reference]; duration	Engagement promotion	Engagement results	Engagement change	Engagement associations with user characteristics and outcomes	User satisfaction and concerns
Mobile-delivered interventions					
Heinrich et al. [35]; 2 weeks	Not reported	The average number of website visits was 3.5 ± 2.7 ; average total time spent on the website was 58.0 ± 56.1 min.	Not reported.	improved at least one self-care behavior. Time spent on the website was not related to baseline variables. There was no association between the time spent on the website and an increase in knowledge.	Not reported
Jennings et al. [36]; 12 weeks	All users were mailed a pedometer as a motivational tool. Users received weekly email reminders with updated content and a link to the website.	Users logged in an average of 21.1 ± 33.6 times. Within the online logbook section, at least 50% logged one type of entry (see article for details). 65% of users set at least one weekly exercise goal (average of 4.6 ± 3.7). There were a total of 2961 discussion post views.	Not reported.	Not reported.	Overall, the site was rated favorably: users said the website was friendly (91%), useful (92%), and the information was easy to understand (95%); users also liked accessing a personalized site (88%).
Jethwani et al. [37]; 12 months	Not reported	About 37% of users did not upload any BG readings, 32% had 1–15 uploads and 31% had >15 uploads. Of those who uploaded, 83% uploaded within first month, 40% logged into website at least once during the study period. Users averaged 29 ± 46 logins.	Both the frequency of BG readings and uploads decreased over time.	Users with >15 uploads had the greatest A1c improvement. Users with better outcomes typically took <10 days to upload their first BG reading.	Users reported several reasons for discontinued use (e.g., technical issues). Many said their providers' interest in program was the strongest predictor of whether or not they used it.
Jernigan and Lorig [38]; 6 weeks	Users received weekly prompts to post to one or more of the website's bulletin boards.	Of the 27 AI/AN pilot users, 15% participated in the workshop sporadically and 85% participated regularly. Among AI/AN users, 4% never made an action plan and 56% made action plans every week. There were a total of 354 bulletin board posts.	Not reported	No differences in website logins and participations rates between AI/AN and non-AI/AN users. AI/AN users more likely to log in during daytime hours.	User satisfaction was high. Reasons for not making weekly action plans included lack of time, illness, family obligations, and depression.
Ryan et al. [39]; 13 months	Not reported	On average, users uploaded BG data two times/week and logged into the app one time/week. Users generated an average of 59.2 ± 69.4 chat events and 11.5 ± 14.9 mail messages, and 56.3 ± 68.6 activities involving a nurse coordinator.	Not reported	Engaging more frequently in chat messages and interactive activities were each associated with reductions in LDL cholesterol levels.	Users valued chatting with nurses and doctors through the HEAT-IT-UP package intervention platform.

Authors [reference]; duration	Engagement promotion	Engagement results	Engagement change	Engagement associations with user characteristics and outcomes	User satisfaction and concerns
Mobile-delivered interventions					
Yu et al. [40]; 9 months	Users received weekly email reminders to visit the website and email prompts when new blog posts were posted.	Users logged in an average of 8.2 days and 0.7 times/month. Among those enrolled, 14% were non-users, 75% were infrequent users, 9% were frequent users, and 2% were heavy users. 67%, 90%, and 60% of users visited static, interactive, and log pages, respectively at least once.	Website use across all users ranged from 4 to 50 logins/week with peaks of 50 logins in week 10 and 37 logins in week 27.	No differences between users and non-users in self-efficacy or self-care. Users had a greater reduction in diabetes distress than non-users.	Patient factors and website factors affected use of website. Users were motivated to access the website based on current needs/concerns and new blog posts.

T2DM type 2 diabetes; *SM* self-monitoring; *BG* blood glucose; *IVR* interactive voice response; *med* medication; *app* application; *AI/AN* American Indian/Alaskan Native