the death certificates, but rather as to the prevalence of septic fevers of any degree in the practice of one or other doctor, nurse, or midwife, and this knowledge can only be obtained by all sorts of pyrexias after labour, except those of a merely evanescent or nervous type, being included in the definition. Much was made of the objections raised to notification on the ground that blame would be likely to attach in popular esteem to a medical practitioner in whose practice such cases occurred, and that the doctors would, in many cases, fail to notify except when cases were likely to terminate fatally; but we may doubt whether these objections are quite as real as some of the speakers seemed to imagine. At any rate, it was shown by Dr. Berry Hart that in large towns such as Sheffield, London, Leeds, &c., the notified cases considerably exceeded in number the fatal ones, which would suggest that where notification is in operation it is done conscientiously. It has so long been customary among our critics to blame the medical practitioner for these unfortunate cases, that people tend to forget that what a public official like the medical officer of health to a large district can do in searching out the origin of puerperal fever is by no means confined to tracking the medical attendant. There are many possibilities of infection in other directions. Both nurse and laundry come under suspicion, and it is only by notification that the matter can be traced back to its root. Dr. John Campbell, of Belfast, probably expressed the truth when he said that the nurse was responsible for a large proportion of the cases among the poor. Knowledge on the subject of antiseptics has become so widely spread throughout the whole body of the medical profession during the past quarter of a century, and the means by which a medical practitioner can free himself from infection are now so effectual, that even putting the matter on the lowest basis, namely, that of self-interest, it is inconceivable that medical practitioners should at all largely allow themselves to become the means of spreading puerperal fever among their patients. But the ordinary monthly nurse, who, so far as concerns the poor, is very often quite untrained in all that pertains to antiseptics, while even in regard to cleanliness her ideas are vague, remains, as she always has been, the probable source of much of the mischief. What we have to recognise is that, while puerperal fever has practically vanished from lying-in hospitals, the loss of life which it causes outside these hospitals has hardly diminished at all, and in seeking for the cause of this state of affairs it is only reasonable to attribute the continuance of this mortality to the continuance of the old custom of employing untrained women as nurses. We do not here refer to the midwife, but to the ordinary neighbourly nurse. The medical practitioner now possesses ample means of ridding himself of infection, and, so far as he himself is concerned, of putting a stop to the spread of this terrible disease, and in view of the fact that the disease still persists, we are forced to the conclusion that it has some other origin. The inquiry, then, must be carried further, and by aid of notification the medical officer of health must be put in a position in which he may trace any prevalence of puerperal sepsis to its origin, just as he would a milk outbreak of typhoid fever, and for this purpose the mild cases, as well as the serious ones, should be reported.

TUBERCULOUS JOINT LESIONS.

IN addressing the Canadian Medical Association at Ottawa on the Clinical Aspects of Tuberculous Lesions, Mr. Edmund Owen made some interesting observations regarding tuberculous joint disease, which are especially worthy of note at the present day when so strong a tendency is shown by many surgeons to regard such tuberculous diseases as always requiring operative treatment. He drew attention to the frequency with which tuberculous disease of the spine, which has gone so far as to produce angular curvature, severe and destructive as it may have been, gets well in the end. Everyone, he says, knows some example of this, or if he does not know him personally he has seen him in the street. "He is rather a short man, with peculiarly high square shoulders, and with a boss between them. And not only has he long since outgrown his tuberculous lesion without any operative assistance whatever, but could we see him in his own home we might not improbably find him—and I say it with some regret—surrounded by a number of apparently healthy sons and daughters." Such a case then is of considerable clinical importance, showing, as it does, that a man with an undoubted tuberculous lesion of the first magnitude can completely recover without having undergone any operative procedure. Another instance of the favourable course which severe tuberculous disease may run in the absence of active surgical interference is seen in the case of old-standing hip-joint disease, the boy actually "growing out of his trouble." When he becomes a young man he may walk lame, and may be precluded from taking an active part in athletics, yet he may be vigorous, and it may be evident that he has quite triumphed over his disease. Turning now to the nature of the tuberculous joint disease and the character of the operation which may be required in certain cases, Mr. Owen said (a) that chronic inflammation of a joint in a child or young person was always tuberculous—except in those very rare cases in which it was due to hereditary syphilis or osteo-arthritis; (b) that tuberculous inflammation might completely destroy a joint, and then leave it solidly and soundly synostosed without the surrounding tissues of the skin having been implicated; (c) that if tuberculous granulation tissue broke down into a fluid, that fluid was not pus, and the collection was not, properly speaking, an abscess—unless by bad fortune or by worse surgery it had become infected by septic micro organisms; and (d) that such a fluid collection was not to be treated as an abscess—by incision and drainage—but was to be opened and emptied, and scraped and cleansed of its unhealthy lining of granulation tissue. The wound in the skin was to be completely closed by sutures, firm pressure was to be applied, and the part was to be kept absolutely at rest. Lastly, he said that he had failed to discover that iodoform was of any peculiar value in the treatment of tuberculous lesions.

FORCIBLE STRAIGHTENING OF CARIOUS SPINES.

WITH his usual vigour of thought and of expression, Mr. Edmund Owen went on to state his views on the practice of forcibly straightening the spine in cases of tuberculous disease with deformity. "Though I should be grieved to stand in the way of surgical advance-

ment," he said, "I do not mind getting in the road and temporarily impeding traffic while we are taking time to consider the route, and are assuring ourselves that the stream of surgical progress is going in the right direction. My opinion is that the deformity of Pott's disease does not lend itself to operative treatment; that forcibly to interfere with it is to thwart Nature in her good attempts at effecting a curative consolidation in her own way. . . . I think, further, that in a short time we shall hear very little about the method. This is what I *think*; but I am absolutely *sure* of this—that if a child of my own had an angular deformity of its spine no person on earth should be allowed roughly to meddle with it. This is the only trustworthy way of testing one's opinion concerning the therapeutic value of speculative methods of treatment." There is, however, a small group of cases for which Mr. Owen thinks that forcible rectification of the angular deformity may perhaps eventually be found suitable, namely, a certain few of those in which pressure by bone, or by organising inflammatory deposits, has taken place upon the anterior surface of the cord, so that the patient has lost the power of voluntary movements in the lower extremities.

THE PREVENTION OF THE HUMPED BACK OF SPINAL DISEASE.

THE humped back of spinal disease is, of course, an opprobrium, and it is small wonder that the surgeon is anxious to efface it. But, if he had given proofs of such laudable efforts and anxiety at the beginning of his treatment, says Mr. Owen, he would probably have had no hump to deal with. This is a matter of much importance, serving as it does to illustrate the feeble treatment which sometimes results from lack of self-confidence in the practitioner. Mr. Owen says: "I have no hesitation in saying that, even at the present time, the treatment of spinal disease in its early stages is too often half-hearted, and sometimes actually blameworthy. It may be urged by way of excuse that, at the very beginning of spinal disease, the symptoms are so equivocal that the practitioner hesitates even to whisper his opinion lest the disappearance of the symptoms should suggest that after all he is an alarmist. He knew that the girl had symmetrical pains in her chest, belly, and legs; he knew that she got easily tired at play, or that she was inclined to loll and lie about when others were full of activity, and that, regardless of nursery manners, she persistently sat at meals with her elbows on the table. He suspected spinal disease; he even told the parents that the girl should be kept quiet. He may actually have gone so far as to sketch out a plan of treatment which was designed to secure a certain amount of rest, but he was slack in seeing that even this small measure was carried out. In short, he had not the courage of his opinion, and the case was allowed to drift. Oh! for the spirit of Lady Macbeth, who called out to her weak-kneed spouse and fellow-practitioner:

'Infirm of purpose! Give me the daggers!'

I am a great admirer of Lady Macbeth. . . . I am fully aware that her character was not faultless . . . but how splendid she would have been in the treatment of early spinal disease! There would have been no half-measures with her!" And the treatment required at

this the very earliest stage is rest, absolute rest. To secure this in the case of a child there is only one way, and that is by making him lie flat in bed. Whatever may be done later on, the treatment must at any rate be inaugurated by the patient being imprisoned in a pillowless bed, and the medical man must not merely recommend but must carefully consider all the details, and make it his business to see that they are loyally and faithfully carried out. This is strong teaching, but it is undoubtedly correct. In every direction it is becoming manifest that early diagnosis is the key to success. But when the diagnosis is made the surgeon must have the courage of his opinion, and we are afraid that he must be prepared in many cases, even when his success is greatest, to hear it said that if he had but let things alone they would have come all right. The reward of success does not always take the form of gratitude.

RAILWAY CARRIAGES AND INFECTION.

AMONG the many so-called causes which conspire to produce a disease it is often by no means easy to assign to each its due importance. In the case of an infectious disease it is but natural that the place of honour should be given to the presence of the germ. Truly, its presence is a *causa causans*, but while willing to pay to the microbe all the honour which is its due, no greater mistake could be made than to forget that without other contributory causes the germ is impotent. Dr. Noble Jones,¹ of Chicago, has lately been making observations in regard to the presence of tubercle bacilli in the nasal secretion of a number of apparently healthy persons, and he concludes from the results of thirty-one inoculations that virulent tubercle bacilli are found in the nasal cavities of healthy persons in the ordinary walks of life. Strauss long ago showed that the presence of virulent tubercle bacilli could be constantly demonstrated in the nasal cavities of those who, being non-tuberculous themselves, were in attendance on tuberculous persons. Turning to another disease, in the July number of the *Revue des Maladies de l'Enfance* there is an article describing two series of investigations by Kober in relation to the presence of virulent diphtheria bacilli in the throats of healthy persons. In the first of these he shows, as has long been known, that in a considerable proportion of those who, although healthy, were known to have been recently in contact with patients suffering from diphtheria, the bacilli were found in the throat. In the second series he further showed that, even among those who were not known to have recently come in contact with any diphtheria case, a diphtheria-like bacillus could now and again be isolated. There is nothing very new in this, but its interest is revived by a question brought up at the recent Congress of the Royal Institute of Public Health in Aberdeen in regard to the dangers to health arising from the dirty condition of many railway carriages. There is every reason to believe that these are a danger, and that people must now and again become infected with various diseases from being shut up in their stuffy atmosphere. But we must not jump to the conclusion that they are a very great danger, for here we have the fact before us that people may carry about sources of infection actually