Depression and obesity are major causes of maternal death in Britain

Depression and obesity are now the major causes of deaths related to pregnancy among women in the United Kingdom, a report published by the Royal College of Obstetricians and Gynaecologists said last week. Although numbers of deaths from complications such as thromboembolism have fallen, indirect psychological and lifestyle factors continue to pose a significant risk.

The Confidential Enquiry into Maternal Death assessed 391 deaths over a period of three years, from 2000 to 2002. It found that 40% of maternal deaths are now attributed to indirect causes with depression and suicide a particular concern.

Obesity is also a problem. Thirty five per cent of the women studied were obese, 50% more than among the general population. No single complication was associated with these deaths, but there was thought to be a link between obesity and broader lifestyle factors such as smoking and social exclusion.

Madeleine Brettingham *London*

Why mothers die can be purchased from the Royal College of Obstetricians and Gynaecologists' bookshop on 020 7772 6275.

Quality of UK consumer health tests to be scrutinised

Concerns that consumer health tests offered by UK retail outlets fail to meet quality standards routinely used in NHS laboratories have prompted calls for regulations to be tightened to guarantee public safety.

The move follows a small amount of anecdotal evidence on cholesterol testing submitted to the Royal College of Pathologists and the National Patient Safety Agency amid fears that this might signal a wider problem.

Cholesterol testing has joined the range of point of sale

tests, since the government made the cholesterol lowering drug simvastatin available over the counter in England and Wales in July 2004. But, unlike in the NHS, retail outlets are not required to comply with internal and external quality controls to assure standards.

Dr Graham Beastall, college vice president and chairman of its chemistry advisory committee, said that point of sale testing was likely to grow, in view of the government's agenda to improve access to diagnostics. "But there are not enough checks and balances in the system."

Caroline White *London*

Ranbaxy withdraws all its AIDS drugs from WHO list

Ranbaxy, an Indian generic drug company, has withdrawn all of its AIDS medicines from the World Health Organization's list of recommended drugs, not because they are unsafe or of poor quality, but because they may not be as effective as they should be, a spokeswoman for WHO said.

WHO announced Ranbaxy's move on 9 November, six months after the health agency removed these and certain other AIDS medicines from its so called prequalification list of essential medicines. WHO had told its member countries that patients should stop using these delisted medicines unless they were unable to get alternative products.

Ranbaxy, however, has not explained why it took the step of voluntarily withdrawing the drugs. A spokeswoman for WHO, Daniela Bagozzi, said that Ranbaxy had found discrepancies in data intended to prove that the generic or copied AIDS medicines were bioequivalent (had the same effect as the patented originals).

Ms Bagozzi denied suggestions that the move came under direct pressure from the United States, which supports the big drug companies that produce patented products.

Fiona Fleck *Geneva*

Use of drug eluting stents increases in the US but only for some groups

Janice Hopkins Tanne *New York*

In the nine months after the first drug eluting stent was approved in the United States, the new style stent was used in 35% of all angioplasties. Stents—metal scaffold tubes that support arteries after balloon angioplasty—can be coated in a slow release drug to reduce restenosis, in which the artery becomes blocked again.

Use of drug eluting stents rose rapidly, from 20% in the first quarter after its introduction to 45% in the third quarter. Plain metal stents, however, were more likely to be used in patients without health insurance, Native Americans, African-Americans, and patients who had had a coronary artery bypass graft.

Sunil Rao, assistant professor of medicine at Duke University Medical Center in Durham, North Carolina, reported that both drug eluting and plain metal stents were used in 83% of the 162 969 percutaneous coronary interventions done at 259 US sites from April 2003 (when the Cypher stent became available) to December 2003.

In an abstract presented at the meeting of the American Heart Association in New Orleans, Dr Rao and colleagues used multivariable logistic regression, including hospital, patient, and lesion characteristics, to determine predictors of drug eluting stents compared with plain metal stents.

The data were collected in a large multicentre national quality improvement initiative for cardiac catheterisation laboratories run by the American Col-

lege of Cardiology's national cardiovascular disease registry. The college uses the data to generate a quality report that tells each centre how it compares with practices in the rest of the country, Dr Rao told the *BMJ*.

Dr Rao's study also found that low volume hospitals and government hospitals (mostly veterans' hospitals) were more likely to use plain metal stents. Bare metal stents were also likely to be used on patients who had received a stent as an emergency procedure, were in shock at presentation, or who had had an acute myocardial infarction in the 24 hours before a stent was fitted.

Elderly patients were more likely to have diffuse disease and to have arteries that were too small to use a stent and might receive bypass surgery, he said.

The drug eluting stent was almost twice as expensive as the plain metal one, although costs have come down, he said. Patients receiving a plain stent were usually prescribed aspirin and an expensive antiplatelet drug for one month afterwards; those receiving a drug eluting stent required the expensive drug for three to six months.

In a second study, Dr Rao found that the stents were used almost equally in approved clinical situations and in those not included in the initial indication—unapproved or "off-label" use. Unapproved uses included patients with a myocardial infarction, with complex lesions, and who had had restenosis with a previous stent. □

The sirolimus-eluting Cypher stent, the first to be approved in the US, was used in 35% of angioplasties in the nine months after its approval