Published in final edited form as:

Drug Alcohol Rev. 2017 January ; 36(1): 88–96. doi:10.1111/dar.12454.

Deconstructing the Associations Between Executive Functioning, Problematic Alcohol Use, and Intimate Partner Aggression: A Dyadic Analysis

Dominic J. Parrott, PhD¹ [Professor], Kevin M. Swartout, PhD² [Assistant Professor], Christopher I. Eckhardt, PhD³ [Professor], and Olivia S. Subramani, BS⁴ [Doctoral Student]

¹Department of Psychology, Georgia State University

²Department of Psychology, Georgia State University

³Department of Psychological Sciences, Purdue University

⁴Department of Psychology, Georgia State University

Abstract

Introduction and Aims—Problematic drinking and executive functioning deficits are two known risk factors for intimate partner aggression (IPA). However, executive functioning is a multifaceted construct, and it is not clear whether deficits in specific components of executive functioning are differentially associated with IPA perpetration generally and within the context of problematic alcohol use. To address this question, the present study investigated the effects of problematic drinking and components of executive functioning on physical IPA perpetration within a dyadic framework.

Design and Methods—Participants were 582 heavy drinking couples (total $N = 1,164$) with a recent history of psychological and/or physical IPA recruited from two metropolitan cities in the United States. Multilevel models were used to examine effects within an Actor-Partner Interdependence framework.

Results—The highest levels of physical IPA were observed among Actors who reported everyday consequences of executive functioning deficits related to emotional dysregulation whose partners were problematic drinkers. However, the association between executive functioning deficits related to emotional dysregulation and IPA was stronger toward partners who were non-problematic drinkers relative to partners who were problematic drinkers. No such effect was found for executive functioning deficits related to behavioral regulation.

Discussion and Conclusions—Results provide insight into how problematic drinking and specific executive functioning deficits interact dyadically in relation to physical IPA perpetration.

Keywords

intimate partner violence; heavy drinking; domestic violence; emotion regulation; actor-partner interdependence model

Intimate partner aggression (IPA) is a significant public health problem, with almost 7 million women and 5.5 million men experiencing physical aggression, stalking, or rape from an intimate partner each year [1]. Although alcohol use is a robust contributing cause of IPA perpetration [2], this relation is not ubiquitous. Thus, development of interactional, etiological models of alcohol-related IPA is critical [3]. The present study addressed this need by testing an interactional model of physical IPA perpetration that considers problematic drinking and components of executive functioning within an actor-partner framework.

Executive Functioning and Alcohol-Related IPA

Executive functioning comprises numerous neurocognitive domains critical to planning and execution of adaptive behavioral responses, including attentional control, response inhibition, working memory, cognitive flexibility, and self-monitoring [4, 5]. Narrative and meta-analytic reviews have established that executive functioning deficits are associated with a propensity for aggressive behavior [6-8]. In addition, both executive functioning deficits and alcohol intoxication are associated with behavioral disinhibition [9, 10]. Consistent with this view, numerous laboratory studies indicate that acute alcohol consumption facilitates aggression to a greater extent among persons with low, relative to high, premorbid executive functioning [11-13].

Extant research also suggests that executive functioning deficits are associated with an increased risk for IPA [for reviews, see 14, 15]. However, two critical questions remain untested. First, it is not clear how executive functioning deficits portend risk within a relationship context, particularly when one or both partners drink heavily. Relationship conflict can lead to aggression for many reasons. One empirically-supported explanation is that IPA results from an interaction between aggression-impelling forces (e.g., trait anger, antisocial traits) and aggression inhibiting/disinhibiting processes (e.g., self-regulation) in response to real or perceived partner provocation [16, 17]. It is well-established that heavy drinking compromises one's ability to adaptively regulate negative emotions elicited by relationship conflict [18, 19]. In addition, executive functioning deficits are thought to underlie maladaptive coping efforts and related inhibitory failures that contribute to IPA perpetration [20, 21]. In the presence of relationship conflict, the concomitant strong emotions and behavioral tendencies that impel aggression must be regulated to prevent IPA perpetration. Individuals who drink heavily and/or possess executive functioning deficits related to behavioral and emotional self-regulation are less capable of modulating these internal states and, as a result, are at greater risk for IPA. Second, it is not clear which components of executive functioning best predict IPA perpetration [14]. To this end, executive functions can be deconstructed into distinct subcomponents and thus afford the potential to examine various facets of self-regulation.

Deconstructing the Associations Between Executive Functioning and Alcohol-Related IPA

Within the alcohol-aggression literature, executive functioning is often conceptualized as a unidimensional construct [11]. However, executive functioning can be broken down into meaningful components, which may evidence differential associations with IPA. Unfortunately, when specific, theoretically-based components of executive functioning are targeted, studies are often limited by the use of single task measures that do not precisely assess the intended construct [22, 23]. This approach increases the risk of overlooking nuanced associations between specific components and aggression, inclusive of IPA. Thus, moving forward, studies that examine associations between executive functioning deficits, alcohol use, and IPA must assess subcomponents of executive functioning. To the extent that specific executive functions are identified as particularly relevant to the alcohol-IPA relation, intervention-based research and practice can be advanced.

Only one study to date has attempted to address this problem, although not specifically in relation to IPA. Giancola, Godlaski, and Roth [24] assessed self-reported everyday consequences of component deficits in executive functioning in social drinkers who consumed an alcoholic or placebo beverage prior to engaging in a laboratory aggression task. Consistent with prior work [11], alcohol intoxication predicted higher levels of aggression among participants who self-reported more frequent everyday consequences associated with greater deficits in general executive functioning. However, a specific component of executive functioning termed *behavioral regulation*, which represents an individual's ability to regulate behavioral and emotional responses, was a significantly stronger moderator of intoxicated aggression than general executive functioning or other subcomponents. This finding suggests that pre-morbid deficits in the regulation of behavioral and emotional responses significantly increase the likelihood that alcohol will facilitate aggression in response to provocation.

Pertinently, research indicates that deficits in emotion regulation are associated with a heightened risk for IPA perpetration [25, 26]. Similarly, a recent daily diary study found that men and women's capacity to regulate state negative affect is central to their risk for IPA perpetration [27]. While informative, this line of research has not examined the relative influence of emotional and behavioral regulation on IPA risk.

The Present Study

The reviewed literature suggests that problematic alcohol use and components of executive functioning related to behavioral and emotional regulation are relevant to the prediction of aggression. However, emotion regulation deficits may be a particularly critical predictor of aggression within intimate relationships [26, 27]. In addition, prior research also suggests the importance of considering the characteristics of both partners when understanding the association between alcohol use and IPA, given findings demonstrating that partner alcohol use predicts actor's IPA perpetration [28]. As such, the present study (1) examined the independent and interactive associations between various subcomponents of executive functioning as manifested in everyday life and problematic alcohol use and the perpetration

of physical IPA in a sample of men and women with a history of relationship conflict, (2) decomposed components of executive functioning as manifested in everyday life to examine directly the relative contributions of behavioral and emotional regulation, and (3) tested these effects within a dyadic Actor Partner Interdependence Model framework (APIM) [29]. The APIM allowed for simultaneous analysis of individual and partner predictors of IPA perpetration while accounting for the interdependence of actor and partner variables.

Although no prior research has examined associations among problematic drinking, executive functioning, and IPA within an APIM framework, several hypotheses were examined in light of available non-dyadic findings. It was hypothesized that actor problematic drinking and actor executive functioning deficits would be positively related to actor physical IPA perpetration. Additionally, it was hypothesized that actor executive functioning deficits would moderate the association between actor problematic drinking and physical IPA, such that the relation between problematic drinking and physical IPA perpetration would be stronger among individuals with greater deficits in executive functioning. Based on prior literature [26, 27], it was hypothesized that executive functions related to emotional, but not behavioral, regulation would moderate this association. Finally, though actor-partner interactions have only recently been examined in studies on alcohol-related IPA [28, 30-33], research clearly indicates that a partner's heavy and/or problematic drinking can elicit greater relationship conflict and facilitate an actor's perpetration of more frequent and intense IPA [31, 33]. Based on these data, it was hypothesized that partner problematic drinking would be positively related to actor physical IPA perpetration. In addition, two actor-partner interactions were advanced. First, it was expected that actor executive functioning deficits related to emotional, but not behavioral, regulation would be more positively associated with IPA perpetration toward partners who were problematic, relative to non-problematic, drinkers. Second, it was expected that actor problematic drinking would be more positively associated with actor physical IPA perpetration toward partners who were problematic, relative to non-problematic, drinkers.

Method

The distinct set of hypotheses tested herein used data that were drawn from a larger investigation on the effects of acute alcohol intoxication and IPA. Thus, couples were required to meet eligibility criteria for an alcohol administration study (see below). The present hypotheses are novel, and the analytic plan was developed specifically to address these aims.

Participants

Participants were 1,224 individuals nested within 612 intimate couples recruited from two U.S. cities through online and print advertisements. To be eligible, couples had to be dating for at least one month, be at least 21 years of age, and identify English as their native language. Couples were excluded if either partner reported serious head injuries, a condition in which alcohol is medically contraindicated, or a desire to seek treatment for alcohol use. At least one partner was required to meet two additional eligibility criteria. First, this individual had to report consumption of an average of at least five (for men) or four (for

women) alcoholic beverages per occasion at least twice per month during the past year. Second, this individual had to be identified as perpetrating psychological or physical IPA toward their current partner via self- or-partner report on the Revised Conflict Tactics Scale [34]. Site 1 enrolled 708 (57.8%) of the individuals in the study, with the remaining 516 (42.2%) individuals enrolled at Site 2.

Couples were screened separately by telephone to assess eligibility, which was then verified in a more comprehensive in-person laboratory assessment. Four couples ($n = 8$) were same-sex and at least one member of 26 couples failed to provide either alcohol use or executive functioning data and were excluded from the current analyses, resulting in a total sample of 582 heterosexual couples ($N = 1,164$). Table 1 contains sample demographics. The average relationship length was 52.15 ($SD = 56.88$) months. Most participants were not married and either non-cohabitating (40.8%) or cohabitating (35.4%), whereas only 15.1% were married. Most participants self-identified as African American (63.6%) or Caucasian (27.0%); 5.5% identified as Hispanic or Latino/a. Participants had a median income range of \$10,000 to \$20,000 per year. This study was approved by each university's Institutional Review Board.

Measures

Revised Conflict Tactics Scale (CTS-2)—The CTS-2 [34] is a 78-item self-report instrument that measures a range of events that occur during disagreements within intimate relationships. Participants are instructed to indicate on a seven-point scale how many times they have perpetrated or experienced the listed behaviors over the past 12 months. Responses may range from 0 (never) to 6 (more than 20 times). Frequency scores are calculated by adding the midpoints of the score range for each item to form a total score. For example, if a participant indicates a response of “3-5” times in the past year, a score of “4” would be assigned. Physical IPA perpetration was operationalized as the total number of acts reported on the respondent's Physical Assault Perpetration subscale or the partner's Physical Assault Victimization subscale; thus, for each participant, we operationalized that individual's physical IPA perpetration as the higher of these two scores [35, 36]. Sample items include “I have thrown something at my partner that could hurt” and “I choked my partner.”

Alcohol Use Disorder Identification Test (AUDIT)—The AUDIT [37] is a 10-item Likert-type scale developed as a screening measure for hazardous and harmful patterns of alcohol consumption. Participants rate items on a 0 to 4 scale, with higher scores indicative of greater problematic drinking. Both members of the dyad reported on their own alcohol use. Sample items include “how often during the past year have you failed to do what was normally expected of you because of drinking,” and “How often do you have a drink containing alcohol.” The AUDIT has high internal consistency across a range of samples [37], which is consistent with the current sample ($\alpha = .85$).

Behavior Rating Inventory of Executive Function—Adult Version (BRIEF-A)—This 75-item self-report inventory assesses a variety of executive functions used in everyday life [38]. Participants are instructed to indicate on a 3-point Likert-type scale (1 = never; 2 = sometimes; 3 = often) the extent to which they have experienced problems with various

behaviors related to executive functioning in the past month. Higher scores indicate poorer executive functioning. The original BRIEF-A yielded two primary factors: the Behavioral Regulation Index and the Metacognition Index. However, consistent with the BRIEF designed for children and adolescents [39, 40], recent exploratory and confirmatory factor analyses with adult samples support a three factor structure that retains the metacognition factor but separates the original behavioral regulation index into a *Behavioral Regulation factor* and an *Emotion Regulation factor* [41, 42]. The Behavioral Regulation factor reflects one's ability to inhibit inappropriate behaviors and judge the appropriateness of one's behavior in social situations. The Emotion Regulation factor reflects one's ability to regulate emotional responses, think flexibly, and adjust to changes in the social environment. Past research indicates that these factors, relative to the Metacognition Index, are more robust and proximally antecedent predictors of aggression [24]. Thus, only the Behavioral and Emotional Regulation factors were examined. The validity of the BRIEF-A has been well-documented in healthy [42, 43] and clinical populations [41, 44]. Internal consistency for the major scales ranges from .93 to .96. In the present sample, reliability for the Behavioral and Emotional Regulation factors exceeded .86.

Procedures

Upon the couple's arrival to the laboratory, each participant was led to a private testing room. After providing informed consent, participants completed the questionnaire battery on a computer using MediaLab 2006 software [45]. The experimenter provided instructions on how to operate the computer program and answered any questions during the session. After completion of the questionnaire battery, participants were compensated and thanked for their time.

Results

Analysis Strategy

Data were analyzed using a multilevel approach to modeling actor-partner interdependence [29, 46] within Mplus version 7.11. The count outcome, physical IPA, was significantly skewed (skewness = 2.28, *S.E.* = .07) with a preponderance of zeros; therefore, we tested a variety of analytic techniques specifically suited for count data with a high proportion of zeros. Negative binomial regression was ultimately selected because it modeled the data best compared with other count-based models based upon the Akaike Information Criterion [47, 48]. Specifically, actor physical IPA was modeled as a function of both actor and partner AUDIT scores, emotional regulation, behavioral regulation, and five empirically-based actor-partner interaction effects (see below). Dyads were distinguishable by gender in regard to the main effects ($\chi^2[6] = 39.23, p < .05$) and the intercepts ($\chi^2[1] = 670.57, p < .05$). Gender was therefore effect coded and entered as a model covariate to account for intercept distinguishability, and interactions between gender and all other predictors and interactions were estimated to account for effect distinguishability [49]. All predictors were standardized with regard to the respective grand means prior to computing interactions and analysis; standard errors were similar across standardized and unstandardized effects and therefore produced consistent significance levels [50]. All effects presented are standardized with respect to the predictor to aid interpretation – once exponentiated, the coefficients represent

the multiplicative effect that a one standard deviation change in the predictor has on the rate of physical IPA, an incidence rate ratio.

Two separate negative binomial regression models were initially estimated: the first model contained main effects of the predictors detailed above on physical IPA with additional effects to control for gender and site; the second model contained all of the model 1 effects plus 5 two-way interactions derived from the literature: (1) actor emotional regulation*actor problematic drinking, (2) actor behavioral regulation*actor problematic drinking, (3) actor emotional regulation*partner problematic drinking, (4) actor behavioral regulation*partner problematic drinking, and (5) actor problematic drinking*partner problematic drinking. This initial interaction model was subsequently trimmed, one hypothesized interactive effect at a time, to strengthen model fit and parsimony. The resulting model contained one significant interaction effect: actor emotional regulation*partner problematic drinking; trimming the other 4 non-significant interaction effects did not hinder model fit ($\chi^2[4] = 1.15, p > .75$). To rule out an alternative interpretation that the interaction between actor and partner emotion regulation accounts for the effect of emotional regulation*partner problematic drinking, we added as covariates to the model the two-way actor-partner emotion regulation interaction as well as the three-way interaction between actor and partner emotion regulation and partner AUDIT.

Actor-Partner Model Results

Descriptive statistics are presented in Table 2. There was significant covariation within couples on problematic drinking, emotional regulation, and behavioral regulation. In the preliminary, main-effects model, the average rate of physical IPA significantly differed across the couples ($u_0 = 1.77, 95\% CI = 1.49-2.04, p < .001$). All of the actor and partner physical IPA predictors were statistically significant predictors of IPA, with the exception of the marginally-significant effect of partner problematic drinking (see Table 3). A one standard deviation (*SD*) increase in actor problematic drinking corresponded with a 13% increase in the physical IPA incidence rate, and partner problematic drinking corresponded with a 7.5% increase. For actors, a one *SD* increase in emotional regulation and behavioral regulation deficits corresponded with increases of 44% and 17% in the physical IPA rate, respectively. A one *SD* increase in partner emotional and behavioral regulation deficits corresponded with 40.1% and 18.2% increases in actor physical IPA rate, respectively. Men had a 5% lower rate of physical IPA perpetration.

As noted above, the hypothesized actor-partner interaction – between actor emotional regulation and partner problematic drinking – significantly predicted physical IPA, even while accounting for numerous control variables (Table 4). Figure 1 illustrates how increases in emotion regulation deficits are associated with IPA rates among individuals whose partners had either high (1 *SD* above the mean) or low (1 *SD* below the mean) AUDIT scores. This interactive effect can be visually detected by attending to the curvature of the accelerating lines; compared with individuals whose partners are problematic drinkers (Partner AUDIT +1 *SD*, $b = 0.327$), the acceleration is more pronounced among individuals whose partners are non-problematic drinkers (Partner AUDIT –1 *SD*, $b = 0.413$), indicating a stronger association between emotional regulation and IPA rates within this group.

Discussion

The present study was the first to investigate the effects of problematic drinking and subcomponents of executive functioning as manifested in everyday life on physical IPA perpetration. The highest levels of physical IPA were observed among Actors who reported everyday consequences of executive functioning deficits related to emotional dysregulation whose partners were problematic drinkers. Studies indicate that a partner's problematic drinking contributes to greater relationship conflict and facilitates their partner's (i.e., actor) perpetration of more frequent and intense IPA [31, 33]. In essence, individuals with problematic drinking partners are exposed to more conflict and concomitant negative affect relative to individuals with non-problematic drinking partners. As evident in the present findings, individuals who self-report negative consequences in everyday life that are associated with executive functioning deficits related to emotional regulation are at particularly high risk to perpetrate physical IPA toward problematic drinking partners. However, the cross-over interaction between actor emotional regulation and partner problematic drinking was driven primarily by a stronger effect of emotional regulation on IPA toward non-problematic, relative to problematic, drinking partners. While further work is needed to fully reveal the nature of this finding, these results may align with the General Aggression Model, which posits that an aggressive response is influenced initially by the interaction of individual differences and situational factors [51]. Thus, the strength of the association between deficits in Actor emotion regulation (an individual difference factor) and physical IPA may depend upon a partner's level of problematic drinking (a situational factor). Collectively, these findings are consistent with recent research which found that proximal negative affect was associated with an increased likelihood of IPA among persons with poorer emotion regulation skills [27] as well as numerous studies that suggest affect regulation is a critical predictor of IPA perpetration [52]. Together, these data support prior calls for interventions that promote perpetrators' adaptive emotion regulation skills [53].

Although qualified by the aforementioned interaction, it is worth noting that actor and partner main effects of problematic alcohol use and everyday consequences of executive functioning deficits on physical IPA perpetration were found. First, actor and partner problematic alcohol use were associated with 13% and 7% increases, respectively, in the rate of actor physical IPA perpetration, although the latter finding was marginally significant. Of course, it is well-established that alcohol use is a contributing cause of IPA perpetration [2] and victimization [54]. However, these findings are noteworthy because they remained robust within the actor-partner framework. Second, actor and partner executive functioning deficits related to emotional regulation as manifested in everyday life were associated with 44% and 40% increases, respectively, in the rate of actor physical IPA perpetration. Also, actor and partner executive functioning deficits related to behavioral regulation as manifested in everyday life were associated with a 17% and 18% increases, respectively, in the rate of actor physical IPA perpetration. These are the first data to indicate that both partners' emotional and behavioral regulation difficulties contribute to the rate of actors' physical IPA perpetration. And, consistent with extant literature [52], results highlight the substantially larger impact of actor deficits in emotional, relative to behavioral, regulation on actor physical IPA.

Two null findings merit discussion. First, contrary to our hypothesis, actor problematic drinking did not interact with actor executive functioning deficits. Though contrary to extant literature [10], the dyadic nature of the present data suggests that associations between problematic alcohol use and executive functioning deficits may differ based on whether findings are perpetrator-based or are derived from dyadic data sources. Second, it merits attention that everyday consequences of executive functioning deficits related to behavioral regulation were not differentially associated with physical IPA perpetration toward problematic and non-problematic drinking partners. As indicated by the aforementioned significant actor main effect, actor behavioral regulation is an important correlate of actor physical IPA perpetration. However, it seems that the additional conflict and concomitant negative affect associated with problematic drinking partners does not exacerbate this effect. Perhaps this is due to the fact that the self-regulatory abilities captured by the Behavioral Regulation factor require one to judge the appropriateness of behavior within the content of relevant social norms. Thus, whereas behavioral regulation may indicate one's receptivity to strong social norms which prohibit physical IPA, it does not appear to differentiate one's reaction to heightened conflict and/or negative affect within the relationship.

Several limitations merit attention. First, due to the cross-sectional design, the temporal relationships among problematic alcohol use, executive functioning, and physical IPA perpetration cannot be unequivocally determined. Moreover, physical IPA during the past year was assessed via retrospective self-report and is thus vulnerable to inaccurate reporting and social desirability bias. Assessment of IPA within a relational context via laboratory-[3] or event-based designs [28] can elucidate temporal relationships and minimize these limitations of self-report. Second, the decision to use the BRIEF-A was based on its ability to assess the functional consequences of executive dysfunction [55] and its high ecological validity [56]. However, these advantages come at a cost, as this measurement approach is not designed to identify the neurobiological substrates of those consequences [55]. Moreover, unlike behavioral measures, the validity of the BRIEF-A is more dependent on the respondent's language and reading processing skills [57]. Clearly, the administration of both behavioral and self-report measures of executive functioning would address these weaknesses.

In conclusion, the present study is the first to disentangle the various subcomponents of executive functioning as manifested in everyday life that best predict IPA perpetration generally and within the context of problematic alcohol use. Results provide insight into how problematic drinking and specific, everyday consequences of executive functioning deficits interact dyadically in relation to physical IPA perpetration. Findings are consistent with the broader literature that considers IPA as a result of aggression-impelling forces (e.g., partner's drinking) and aggression disinhibiting processes (e.g., actor's executive functioning), and thus highlight the importance of considering etiological determinants of IPA within a dyadic context.

Acknowledgements

Dominic J. Parrott, Kevin M. Swartout, & Olivia S. Subramani, Department of Psychology, Georgia State University. Christopher I. Eckhardt, Department of Psychological Sciences, Purdue University.

This research was supported by grant R01-AA-020578 from the National Institute on Alcohol Abuse and Alcoholism awarded to the 1st and 3rd authors.

References

1. Black, MC., Basile, KC., Breiding, MJ., Smith, SG., Walter, ML., Merrick, MT., et al. National intimate partner and sexual violence survey (NISVS): 2010 summary report. National Center for Injury Prevention and Control, Centers for Disease Control and Prevention; Atlanta: 2011.
2. Leonard KE. Alcohol and intimate partner violence: When can we say that heavy drinking is a contributing cause of violence? *Addiction*. 2005; 100:422–425. [PubMed: 15784050]
3. Eckhardt CI, Parrott DJ, Sprunger JG. Mechanisms of alcohol-facilitated intimate partner violence. *Violence Against Women*. 2015; 21:939–957. [PubMed: 26059921]
4. Anderson, V., Jacobs, R., Anderson, PJ. Executive functions and the frontal lobes: A lifespan perspective. Philadelphia: Taylor & Francis: 2008.
5. Goldberg, E. The executive brain: Frontal lobes and the civilized mind. Oxford University Press; New York: 2002.
6. Moffitt TE. The neuropsychology of conduct disorder. *Dev Psychopathol*. 1993; 5:135–151.
7. Morgan AB, Lilienfeld SO. A meta-analytic review of the relation between antisocial behavior and neuropsychological measures of executive function. *Clin Psychol Rev*. 2000; 20:113–136. [PubMed: 10660831]
8. Ogilvie JM, Stewart AL, Chan RCK, Shum DHK. Neuropsychological measures of executive function and antisocial behavior: A meta-analysis. *Criminology*. 2011; 49:1063–1107.
9. Fishbein D. Neuropsychological function, drug abuse, and violence: A conceptual framework. *Crim Justice Behav*. 2000; 27:139–159.
10. Giancola PR. Executive functioning: a conceptual framework for alcohol-related aggression. *Exp Clin Psychopharmacol*. 2000; 8:576–597. [PubMed: 11127429]
11. Giancola PR. Executive functioning and alcohol-related aggression. *J Abnorm Psychol*. 2004; 113:541–555. [PubMed: 15535787]
12. Giancola PR, Parrott DJ, Roth RM. The influence of difficult temperament on alcohol-related aggression: Better accounted for by executive functioning? *Addict Behav*. 2006; 31:2169–2187. [PubMed: 16563644]
13. Hoaken P, Assaad J, Pihl RO. Cognitive functioning and the inhibition of alcohol-induced aggression. *J Stud Alcohol Drugs*. 1998; 59:599–607.
14. Bueso-Izquierdo N, Hart SD, Hidalgo-Ruzzante N, Kropp PR, Pérez-García M. The mind of the male batterer: A neuroscience perspective. *Aggress Violent Behav*. 2015:243–251. Part B.
15. Pinto LA, Sullivan EL, Rosenbaum A, Wyngarden N, Umhau JCM, Taft CT. Biological correlates of intimate partner violence perpetration. *Aggress Violent Behav*. 2010; 15:387–398. [PubMed: 23393423]
16. Finkel EJ. Impelling and inhibiting forces in the perpetration of intimate partner violence. *Rev Gen Psychol*. 2007; 11:193–207.
17. Finkel, EJ., Eckhardt, CI. Intimate partner violence. In: Simpson, JA., Campbell, L., editors. *The Oxford handbook of close relationships*. Oxford; New York: 2013. p. 452–474.
18. Eckhardt CI. Effects of alcohol intoxication on anger experience and expression among partner assaultive men. *J Consult Clin Psychol*. 2007; 75:61–71. [PubMed: 17295564]
19. Watkins LE, DiLillo D, Maldonado RC. The interactive effects of emotion regulation and alcohol intoxication on lab-based intimate partner aggression. *Psychol Addict Behav*. 2015; 29:653–663. [PubMed: 25844831]
20. Corvo K. The role of executive function deficits in domestic violence perpetration. *Partner Abuse*. 2014; 5:342–355.
21. Corvo K, Johnson P. Sharpening Ockham's Razor: The role of psychopathology and neuropsychopathology in the perpetration of domestic violence. *Aggress Violent Behav*. 2013; 18:175–182.

22. Ellis ML, Weiss B, Lochman JE. Executive functions in children: Associations with aggressive behavior and appraisal processing. *J Abnorm Child Psychol*. 2009; 37:945–956. [PubMed: 19408113]
23. Stanford MS, Greve KW, Gerstle JE. Neuropsychological correlates of self-reported impulsive aggression in a college sample. *Pers Individ Dif*. 1997; 23:961–965.
24. Giancola PR, Godlaski AJ, Roth RM. Identifying component-processes of executive functioning that serve as risk factors for the alcohol-aggression relation. *Psychol Addict Behav*. 2012; 26:201–211. [PubMed: 21875167]
25. Harper FWK, Austin AG, Cercone JL, Arias I. The role of shame, anger, and affect regulation in men's perpetration of psychological abuse in dating relationships. *J Interpers Violence*. 2005; 20:1648–1662. [PubMed: 16246922]
26. Shorey RC, Brasfield H, Febres J, Stuart GL. The association between impulsivity, trait anger, and the perpetration of intimate partner violence and general violence among women arrested for domestic violence. *J Interpers Violence*. 2011; 26:2681–2697. [PubMed: 21118862]
27. Shorey RC, McNulty JK, Moore TM, Stuart GL. Emotion regulation moderates the association between proximal negative affect and intimate partner violence perpetration. *Prev Sci*. 2015; 6:873–880.
28. Testa M, Derrick JL. A daily process examination of the temporal association between alcohol use and verbal and physical aggression in community couples. *Psychol Addict Behav*. 2014; 28:127–138. [PubMed: 24341618]
29. Cook WL, Kenny DA. The actor–partner interdependence model: A model of bidirectional effects in developmental studies. *Int J Behav Dev*. 2005; 29:101–109.
30. Crane CA, Testa M, Derrick JL, Leonard KE. Daily associations among self-control, heavy episodic drinking, and relationship functioning: An examination of actor and partner effects. *Aggress Behav*. 2014; 40:440–450. [PubMed: 24700558]
31. Testa M, Kubiak A, Quigley BM, Houston RJ, Derrick JL, Levitt A, et al. Husband and wife alcohol use as independent or interactive predictors of intimate partner violence. *J Stud Alcohol Drugs*. 2012; 73:268–276. [PubMed: 22333334]
32. Watkins LE, Maldonado RC, DiLillo D. Hazardous alcohol use and intimate partner aggression among dating couples: The role of impulse control difficulties. *Aggress Behav*. 2014; 40:369–381. [PubMed: 24464403]
33. Woodin EM, Caldeira V, Sotskova A, Galaugher T, Lu M. Harmful alcohol use as a predictor of intimate partner violence during the transition to parenthood: Interdependent and interaction effects. *Addict Behav*. 2014; 39:1890–1897. [PubMed: 25150656]
34. Straus MA, Hamby SL, Boney-McCoy S, Sugarman DB. The revised conflict tactics scales (CTS2): Development and preliminary psychometric data. *J Fam Issues*. 1996; 17:283–316.
35. Foran HM, O'Leary KD. Problem drinking, jealousy, and anger control: Variables predicting physical aggression against a partner. *J Fam Violence*. 2008; 23:141–148.
36. Slep AS, O'Leary SG. Parent and partner violence in families with young children: Rates, patterns, and connections. *J Consult Clin Psychol*. 2005; 73:435–444. [PubMed: 15982141]
37. Babor, TF., Biddle-Higgins, JC., Saunders, JB., Monteiro, MG. AUDIT: The Alcohol Use Disorders Identification Test: Guidelines for Use in Primary Health Care. World Health Organization; Geneva, Switzerland: 2001.
38. Roth, RM., Isquith, PK., Gioia, GA. Behavior Rating Inventory of Executive Function-Adult Version (BRIEF-A). Psychological Assessment Resources; Lutz, FL: 2005.
39. Egeland J, Fallmyr Ø. Confirmatory factor analysis of the Behavior Rating Inventory of Executive Function (BRIEF): Support for a distinction between emotional and behavioral regulation. *Child Neuropsychol*. 2010; 16:326–337. [PubMed: 20209415]
40. Gioia GA, Isquith PK. Two faces of monitor: They self and thy task. *J Int Neuropsychol Soc*. 2002; 8:229.
41. Donders J, Strong CH. Latent structure of the Behavior Rating Inventory of Executive Function – Adult Version (BRIEF-A) after a mild traumatic brain injury. *Arch Clin Neuropsychol* in press.
42. Roth RM, Lance CE, Isquith PK, Fischer AS, Giancola PR. Confirmatory factor analysis of the behavior rating inventory of executive function-adult version in healthy adults and application to

- attention-deficit/hyperactivity disorder. *Arch Clin Neuropsychol*. 2013; 28:425–434. [PubMed: 23676185]
43. Rabin LA, Fogel J, Nutter-Upham KE. Academic procrastination in college students: The role of self-reported executive function. *J Clin Exp Neuropsychol*. 2010; 33:344–357. [PubMed: 21113838]
 44. Chang WP, Davies PL, Gavin WJ. Error monitoring in college students with attention-deficit/hyperactivity disorder. *J Psychophysiol*. 2009; 23:113–125.
 45. Jarvis, BG. DirectRT (Version 2006) [Computer software]. Empirisoft Corporation; New York: 2006.
 46. Kenny DA, Cook WL. Partner effects in relationship research: Conceptual issues, analytic difficulties, and illustrations. *Pers Relatsh*. 1999; 6:433–448.
 47. Akaike H. A new look at the statistical model identification. *IEEE Trans Automat Contr*. 1974; 19:716–723.
 48. Swartout KM, Thompson MP, Koss MP, Su N. What is the best way to analyze less frequent forms of violence? The case of sexual aggression. *Psychol Violence*. 2015; 5:305–313. [PubMed: 26925298]
 49. Kashy, DA., Donnellan, MB. Conceptual and methodological issues in the analysis of data from dyads and groups. In: Deaux, K., Snyder, M., editors. *The Oxford handbook of personality and social psychology*. Oxford; New York: 2012. p. 209-238.
 50. Aiken, L., West, S. *Multiple regression: Testing and interpreting interactions*. Sage Publications; Thousand Oaks: 1991.
 51. DeWall CN, Anderson CA, Bushman BJ. The general aggression model: Theoretical extensions to violence. *Psychol Violence*. 2011; 1:245–258.
 52. Birkley EL, Eckhardt CI. Anger, hostility, internalizing negative emotions, and intimate partner violence perpetration: A meta-analytic review. *Clin Psychol Rev*. 2015; 37:40–56. [PubMed: 25752947]
 53. Shorey RC, Zucosky H, Brasfield H, Febres J, Cornelius TL, Sage C, et al. Dating violence prevention programming: Directions for future interventions. *Aggress Violent Behav*. 2012; 17:289–296. [PubMed: 22773916]
 54. Devries KM, Child JC, Bacchus LJ, Mak J, Falder G, Graham K, et al. Intimate partner violence victimization and alcohol consumption in women: A systematic review and meta-analysis. *Addiction*. 2014; 109:379–391. [PubMed: 24329907]
 55. McAuley T, Chen S, Goos L, Schachar R, Crosbie J. Is the behavior rating of executive function strongly associated with measures of impairment in executive function? *J Int Neuropsychol Soc*. 2010; 16:495–505. [PubMed: 20188014]
 56. Gioia GA, Isquith PK. Ecological assessment of executive function in traumatic brain injury. *Dev Neuropsychol*. 2004; 25:135–158. [PubMed: 14984332]
 57. Denckla M. The behavior rating inventory of executive function: Commentary. *Child Neuropsychol*. 2002; 8:304–306. [PubMed: 12759827]

Figure 1.

Interaction plot: Partner AUDIT score moderates the association between actor emotion regulation deficits and actor IPA.

Note: Point estimates reflect the natural log of the Physical IPA rate; AUDIT: Alcohol Use Disorders Identification Test score; ER: Emotional Regulation; IPA: Intimate Partner Aggression; Higher ER scores reflect greater deficits in emotional regulation.

Table 1

Sample demographics.

	Gender	
	Female	Male
No. (%)	582 (50)	582 (50)
Age, Mean (<i>SD</i>)	31.52 (9.98)	33.26 (10.54)
Years of education, Mean (<i>SD</i>)	14.27 (2.77)	13.83 (2.72)
Drinks per day, Mean (<i>SD</i>)	4.62 (3.45)	5.74 (4.00)
Drinking days per week, Mean (<i>SD</i>)	2.20 (1.89)	2.04 (1.99)

Table 2

Descriptive statistics for unstandardized model variables.

Variable	Mean	SD	<i>r</i> with Partner's Score
AUDIT	8.47	6.20	.381 [*]
ER	23.18	6.20	.14 [*]
BR	21.00	5.00	.20 [*]
IPA	2.35	2.35	--- ^a

Note:

^{*} < .001. AUDIT: Alcohol Use Disorders Identification Test score; ER: Emotional Regulation; BR: Behavioral Regulation; IPA: Intimate Partner Aggression.

^a Not applicable because IPA is count.

Table 3

Main effects model results.

	Predictor	Estimate	SE	95% CI	IRR	p value
Actor	AUDIT	.12	.04	.04, .20	1.13	.002
	ER	.36	.07	.22, .50	1.44	< .001
	BR	.16	.07	.02, .29	1.17	.025
Partner	AUDIT	.07	.04	-.005, .15	1.07	.068
	ER	.34	.07	.19, .48	1.40	< .001
	BR	.17	.07	.03, .31	1.18	.019
Control	Actor Gender	-.11	.02	-.15, -.07	.90	< .001
	Site	-.10	.12	-.34, .14	.90	.412
	Actor AUDIT*Gender	.06	.06	-.05, .16	1.06	.309
	Actor ER*Gender	.01	.07	-.12, .15	1.01	.844
	Actor BR*Gender	-.001	.06	-.13, .13	1.00	.993
	Partner AUDIT*Gender	-.07	.06	-.18, .04	0.93	.213
	Partner ER*Gender	.001	.07	-.14, .14	1.00	.986
	Partner BR*Gender	.02	.07	-.11, .14	1.02	.815

Note: AUDIT: Alcohol Use Disorders Identification Test score; ER: Emotional Regulation; BR: Behavioral Regulation.

Table 4

Final actor-partner interaction model results.

	Predictor	Est.	SE	95% CI	IRR	p value
Actor	AUDIT	.13	.04	.05, .20	1.14	.001
	ER	.37	.07	.23, .51	1.45	<.001
	BR	.15	.07	.01, .29	1.16	.03
Partner	AUDIT	.08	.04	.00, .16	1.08	.05
	ER	.36	.08	.21, .50	1.43	<.001
	BR	.15	.07	.01, .29	1.16	.033
Interaction	Actor ER*Partner AUDIT	-.04	.02	-.23, -.02	0.96	.046
Controls	Site ^a	-.08	.12	-.32, .16	0.92	.498
	Actor Gender ^b	-.11	.02	-.15, -.07	0.90	<.001
	Actor ER*Partner ER	-.13	.05	-.08, -.01	0.88	.015
	Actor ER*Partner ER*Partner AUDIT	.04	.02	-.01, .09	1.04	.089
	Actor ER*Partner ER*Gender	.02	.02	-.02, .05	1.02	.312
	Actor AUDIT*Gender	.05	.05	-.05, .16	1.05	.331
	Actor ER*Gender	.04	.07	-.10, .17	1.04	.582
	Actor BR*Gender	-.02	.06	-.14, .11	0.98	.796
	Partner AUDIT*Gender	-.08	.06	-.19, .03	0.92	.153
	Partner ER*Gender	-.03	.07	-.17, .11	0.97	.663
	Partner BR*Gender	.03	.07	-.10, .15	1.03	.688

Note: AUDIT: Alcohol Use Disorders Identification Test score; ER: Emotional Regulation; BR: Behavioral Regulation.

^aSite effect coded;

^bGender effect coded with Male = 1, Female = -1.