Published in final edited form as:

J Rheumatol. 2017 February ; 44(2): 170–173. doi:10.3899/jrheum.160026.

Explanatory Style in Patients with Rheumatoid Arthritis: An Unrecognized Predictor of Mortality

Aaron D. Crowson, BA¹, Robert C. Colligan, PhD², Eric L. Matteson, MD MPH^{1,3}, John M. Davis III, MD MS³, and Cynthia S. Crowson, MS^{1,3}

¹Department of Health Sciences Research, Mayo Clinic, Rochester, MN, USA

²Department of Psychiatry and Psychology, Mayo Clinic, Rochester, MN, USA

³Division of Rheumatology, Department of Medicine, Mayo Clinic, Rochester, MN, USA

Abstract

Objective—To determine whether pessimistic explanatory style altered the risk for and mortality of rheumatoid arthritis (RA) patients.

Methods—The study included subjects from a population-based cohort with incident RA and non-RA comparison cohort who completed the Minnesota Multiphasic Personality Inventory (MMPI).

Results—Among 148 RA and 135 non-RA subjects, pessimism was associated with development of rheumatoid factor positive (RF+) RA. Pessimism was associated with an increased risk of mortality (hazard ratio [HR]:2.88 with similar magnitude to RF+ (HR:2.28).

Conclusion—Pessimistic explanatory style was associated with an increased risk of developing RA and increased mortality rate in patients with RA.

Keywords

Rheumatoid arthritis; mortality; psychosocial; pessimism; Minnesota Multiphasic Personality Inventory

INTRODUCTION

The contribution of psychosocial factors to healthcare outcomes is widely appreciated and likely to play an even greater role as efforts towards integrated healthcare are pursued. Evidence suggests that personality traits can affect disease course and even mortality.

Explanatory style is a personality trait known to impact health and result in poor health outcomes (1, 2). For example, a pessimistic explanatory was associated with poorer survival in lung cancer patients (3). Pessimists had significantly more pain and lower physical activity after knee replacement surgery compared to non-pessimists (4).

Rheumatoid arthritis (RA) is a debilitating disease characterized by chronic pain and poorer outcomes, including increased risk of mortality (5). Psychosocial factors, including depression and anxiety, are associated with adverse outcomes of RA (6, 7), but little is known about the association between personality traits and RA. The aim of this study was to examine the relationship between pessimism and both the risk for developing RA and mortality rates among patients with RA.

METHODS

Study Subjects

This is a retrospective study of Olmsted County, Minnesota residents with incident diagnosis of RA in 1955–2007 who completed the Minnesota Multiphasic Personality Inventory (MMPI). The study cohorts were assembled, following approval by Institutional Review Boards of Mayo Clinic (15-007631) and Olmsted Medical Center (001-OMC-16), using the Rochester Epidemiology Project (REP), a unique medical record linkage system which makes population-based epidemiologic research possible by providing access to comprehensive (inpatient as well as outpatient) medical records for all Olmsted County residents, at any local provider (8).

The RA cohort included all residents of Rochester (the central city of Olmsted County) who were 18 years of age when fulfilling American College of Rheumatology 1987 classification criteria for RA in 1955–1979 and all Olmsted County residents fulfilling criteria in 1980–2007(9, 10). RA incidence was defined as the date of first fulfillment of the criteria. A comparison cohort of residents without RA was obtained by randomly selecting a subject of similar sex and birth year for each RA patient, who was assigned an index date corresponding to the RA incidence date of their matched pair. All study subjects were followed until death, migration, or 1/1/2015.

Smoking status and obesity (i.e., body mass index ≥ 30 kg/m²) at incidence/index date were collected previously. History of depression prior to RA incidence/index date was obtained using diagnostic codes. The Charlson comorbidity index was calculated without the rheumatologic component (11).

This study utilized patients who completed the MMPI at Mayo Clinic at least once prior to the RA incidence/index date. The MMPIs included those completed for medical evaluation and research purposes. The original MMPI was comprised of 550 unique true/false items regarding physical and emotional symptoms, attitudes, feelings, thoughts, and life experiences (12). MMPIs completed prior to age 18 years and those with >100 missing responses were excluded. For patients with multiple MMPIs, the earliest MMPI was used.

The Optimism-Pessimism (PSM) scale of the MMPI was developed from Seligman's theory of explanatory style. This theory postulates that people who attribute the cause of negative life events internally (to themselves), globally (affecting other aspects of their life), and with stability (continually happening), are at risk of disturbing their cognitive and emotional function, along with their physical health (3, 13). This validated scale, derived from 298 items, yields normalized T-scores with mean of 50 and standard deviation (SD) of 10 with

higher scores indicating pessimism. PSM scores were categorized using a cutpoint of 1 SD above the mean (i.e., PSM ≥ 60 defines pessimism)(1–3). The MMPI Depression Scale 2 and item 51 (“I am in just as good physical health as most of my friends”) were also examined (12).

Statistical Analysis

Cohorts were compared using chi-square and rank sum tests. Logistic regression was used to examine the association between pessimism and RA with adjustment for age, sex, smoking and obesity. Cox models, adjusted for age, sex, smoking, obesity and Charlson index, were used to examine the association between pessimism and mortality in RA. The interaction between RF positivity (RF+) and pessimism was also examined. Kaplan-Meier methods were used to estimate mortality. Survival curves adjusted to the whole RA population were computed by averaging individual predictions from the Cox models. Analyses were performed using SAS version 9.4 (SAS Institute, Cary, NC, USA) and R 3.1.1 (R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

The study population included 148 patients with RA and 135 non-RA subjects who completed the MMPI before incidence/index date. Among 86 RA and non-RA subjects who completed multiple MMPIs, only 12(14%) changed from pessimistic to non-pessimistic and 5(6%) changed from non-pessimistic to pessimistic. In addition, no difference was found in mean PSM scores between those who completed the MMPI before RA diagnosis (N=148; mean PSM score=55.8) and those who took the MMPI after RA diagnosis (N=53; mean PSM score=54.9; $p=0.56$).

The MMPIs were completed on average 13.7 years before incidence date for patients with RA and 14.2 years before index date for the non-RA subjects (Table 1). Only 9 patients with RA completed the MMPI between symptom onset and RA diagnosis. Patients with RA had higher Charlson index values at RA incidence/index date than non-RA subjects.

Pessimism was somewhat more common in RA (39%) than non-RA (31%; $p=0.19$; Table 1). Among patients with RA, pessimism was more frequent in those with RF+RA (44%) than in non-RA subjects (31%; $p=0.049$). There was no difference in pessimism between RF-RA and non-RA groups (29% vs 31%; $p=0.76$). After adjustment for known RA risk factors (age, sex, smoking and obesity), the association between pessimism and RF+RA persisted (odds ratio [OR]:1.74; 95% confidence interval [CI]:0.99–3.04; $p=0.053$).

Among the 148 patients with RA, 64 died during a median of 13.5 years of follow-up. In a multivariable model adjusted for age, sex, calendar year, smoking, obesity and Charlson index, RF+ patients had twice the risk of mortality as RF– (hazard ratio [HR]:2.28; 95%CI: 1.10–4.73; Table 2) and pessimism increased the risk of mortality nearly 3-fold (HR:2.88; 95%CI:1.02–8.14). A significant interaction between RF+ and pessimism ($p=0.037$) demonstrated patients with both RF+ and pessimism did not have a higher risk of mortality than those with either. Ten-year survival was highest for RF-RA non-pessimists (93%) and were similar for the other 3 groups (73% RF+RA non-pessimists, 77% RF-RA pessimists

and 84% RF+RA pessimists; Figure 1). The MMPI depression scale ($p=0.88$) and prior history of depression diagnosis ($p=0.32$) were not significantly associated with mortality and did not change the association between pessimism and mortality (HR for pessimism: 2.97 and 3.00, respectively).

DISCUSSION

Pessimism is associated with a higher likelihood of developing RF+RA. Among patients with RA, pessimism was associated with a substantial increased risk of mortality. This increased mortality in patients with RA is consistent with that from studies of other chronic diseases, demonstrating that pessimism is associated with an increase in all- cause mortality (3, 14). Similarly, a study on lung cancer showed that pessimistic patients lived on average 6 months less than non-pessimists (3).

Pessimism has also been associated with other poor health outcomes. A study of knee replacement surgery reported that patients who were classified as pessimistic had higher levels of reported pain and less physical activity two years after the surgery (4). Similarly, after heart transplant, patients who were pessimistic showed more depressive symptoms, while those considered optimistic reported better overall quality of life (15).

The effect and biology of pessimism in RA is poorly studied. An association between poor positive affect (i.e., less optimism) and elevated IL-6 levels, a biomarker of inflammation, may partially explain why pessimism could affect RA (16). A negative outlook (conceptualized by anxiety, depression, and pessimism) increased the chances of developing cardiovascular disease (CVD), also known to be influenced by inflammation, in patients with RA (6). Others reported depressive, anxious or pessimistic outlooks led to increased rates of CVD and mortality in the general population (17, 18).

Strengths of this study include its population-based design with extensive follow-up and that most MMPIs were completed years before the patients developed RA. The PSM scale has repeatedly shown reliability and validity (1, 14), and explanatory style has demonstrated stability across many years of adult life(19). The MMPI-2 and PSM-R are contemporary version with the same properties that reduce respondent burden(20). Some limitations are the study is retrospective, and the MMPIs were administered for a variety of reasons and not to all possible candidates, potentially resulting in a selection bias. Also the small sample size limited statistical power to detect differences. Finally, the Olmsted County population is predominately white; therefore our results may not be generalizable to more diverse populations.

In conclusion, a pessimistic explanatory style increased the risk of developing RA and rates of mortality among patients with existing RA. Our results suggest that personality traits may influence how patients perceive and manage illness, as well as their global outcomes. Patients who learn and have techniques to counter consequences of pessimistic explanatory style may be able to achieve better outcomes of RA. Therefore, understanding personality traits is an important step in the holistic management of disease and a vital component of integrated care.

Acknowledgments

Funding: This work was supported by a research grant R01 AR06849 and made possible by the Rochester Epidemiology Project (R01-AG034676) from the National Institutes of Health. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

References

1. Maruta T, Colligan RC, Malinchoc M, Offord KP. Optimists vs pessimists: survival rate among medical patients over a 30-year period. *Mayo Clin Proc.* 2000; 75:140–3. [PubMed: 10683651]
2. Maruta T, Colligan RC, Malinchoc M, Offord KP. Optimism-pessimism assessed in the 1960s and self-reported health status 30 years later. *Mayo Clin Proc.* 2002; 77:748–53. [PubMed: 12173709]
3. Novotny P, Colligan RC, Szydio DW, Clark MM, Rausch S, Wampfler J, et al. A pessimistic explanatory style is prognostic for poor lung cancer survival. *J Thorac Oncol.* 2010; 5:326–32. [PubMed: 20139778]
4. Singh JA, O’Byrne MM, Colligan RC, Lewallen DG. Pessimistic explanatory style: a psychological risk factor for poor pain and functional outcomes two years after knee replacement. *J Bone Joint Surg Br.* 2010; 92:799–806. [PubMed: 20513876]
5. Gonzalez A, Icen M, Kremers HM, Crowson CS, Davis JM 3rd, Thernau TM, et al. Mortality trends in rheumatoid arthritis: the role of rheumatoid factor. *J Rheumatol.* 2008; 35:1009–14. [PubMed: 18412312]
6. Liu YL, Szklo M, Davidson KW, Bathon JM, Giles JT. Differential association of psychosocial comorbidities with subclinical atherosclerosis in rheumatoid arthritis. *Arthritis Care Res.* 2015; 67:1335–44.
7. Dickens C, McGowan L, Clark-Carter D, Creed F. Depression in rheumatoid arthritis: a systematic review of the literature with meta-analysis. *Psychosom Med.* 2002; 64:52–60. [PubMed: 11818586]
8. St Sauver JL, Grossardt BR, Yawn BP, Melton LJ 3rd, Pankratz JJ, Brue SM, et al. Data resource profile: the Rochester Epidemiology Project (REP) medical records-linkage system. *Int J Epidemiol.* 2012; 41:1614–24. [PubMed: 23159830]
9. Arnett FC, Edworthy SM, Bloch DA, McShane DJ, Fries JF, Cooper NS, et al. The American Rheumatism Association 1987 revised criteria for the classification of rheumatoid arthritis. *Arth Rheum.* 1988; 31:315–24. [PubMed: 3358796]
10. Myasoedova E, Crowson CS, Kremers HM, Thernau TM, Gabriel SE. Is the incidence of rheumatoid arthritis rising?: Results from Olmsted County, Minnesota, 1955–2007. *Arth Rheum.* 2010; 62:1576–82. [PubMed: 20191579]
11. Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol.* 1992; 45:613–9. [PubMed: 1607900]
12. Hathaway, SR., McKinley, JC. The Minnesota Multiphasic Personality Inventory. Minneapolis: University of Minnesota; 1943.
13. Colligan RC, Offord KP, Malinchoc M, Schulman P, Seligman ME. CAVEing the MMPI for an Optimism-Pessimism Scale: Seligman’s attributional model and the assessment of explanatory style. *J Clin Psychol.* 1994; 50:71–95. [PubMed: 8150997]
14. Grossardt BR, Bower JH, Geda YE, Colligan RC, Rocca WA. Pessimistic, anxious, and depressive personality traits predict all-cause mortality: the Mayo Clinic cohort study of personality and aging. *Psychosom Med.* 2009; 71:491–500. [PubMed: 19321849]
15. Jowsey SG, Cutshall SM, Colligan RC, Stevens SR, Kremers WK, Vasquez AR, et al. Seligman’s theory of attributional style: optimism, pessimism, and quality of life after heart transplant. *Prog Transplant.* 2012; 22:49–55. [PubMed: 22489443]
16. Sin NL, Graham-Engeland JE, Ong AD, Almeida DM. Affective reactivity to daily stressors is associated with elevated inflammation. *Health Psychol.* 2015; 34:1154–65. [PubMed: 26030309]
17. Sin NL, Moskowitz JT, Whooley MA. Positive affect and health behaviors across 5 years in patients with coronary heart disease: the heart and soul study. *Psychosom Med.* 2015; 77:1058–66. [PubMed: 26428445]

18. Chamberlain AM, Vickers KS, Colligan RC, Weston SA, Rummans TA, Roger VL. Associations of preexisting depression and anxiety with hospitalization in patients with cardiovascular disease. *Mayo Clin Proc.* 2011; 86:1056–62. [PubMed: 22033250]
19. Burns MO, Seligman ME. Explanatory style across the life span: evidence for stability over 52 years. *J Pers Soc Psychol.* 1989; 56:471–7. [PubMed: 2926642]
20. Malinchoc M, Offord KP, Colligan RC. PSM-R: Revised Optimism-Pessimism Scale for the MMPI-2 and MMPI. *J Clin Psychol.* 1995; 51:205–14. [PubMed: 7797644]

Figure 1.

Survival curves for patients with rheumatoid arthritis (RA) with Minnesota Multiphasic Personality Inventory prior to RA diagnosis according to rheumatoid factor status (RF positive/negative) and pessimistic explanatory style (no/yes) adjusted for age, sex, calendar year of RA diagnosis, smoking status and Charlson comorbidity index

Table 1

Characteristics of patients with and without rheumatoid arthritis (RA) who completed the Minnesota Multiphasic Personality Inventory (MMPI)

	Non-RA (N=135)	RA (N=148)	p value
Age, years, mean ($\pm$SD)	55.4 ($\pm$ 12.9)	56.5 ($\pm$ 12.9)	0.33
Sex, female	92 (68%)	112 (76%)	0.16
Year of index date, mean ($\pm$SD)	1995.5 ($\pm$ 9.1)	1994.5 ($\pm$ 9.8)	0.42
Time from MMPI to index date, years, mean ($\pm$SD)	14.2 ($\pm$ 9.0)	13.7 ($\pm$ 9.1)	0.71
MMPI Item 51: Health similar to peers (% true)	102 (77%)	106 (74%)	0.55
Pessimistic Explanatory Style *	42 (31%)	57 (39%)	0.25
MMPI Depression scale, mean ($\pm$SD)	58.4 ($\pm$ 10.7)	58.7 ($\pm$ 11.2)	0.83
Rheumatoid factor positivity	–	96 (65%)	
Sedimentation rate, median (25th percentile, 75th percentile)	–	23.0 (10, 40)	
Obesity (body mass index $\geq$ 30 kg/m²)	46 (34%)	56 (38%)	0.51
Smoking status			0.13
Never	68 (50%)	60 (41%)	
Current	20 (15%)	34 (23%)	
Former	47 (35%)	54 (36%)	
Charlson comorbidity index, mean ($\pm$SD)	0.9 ($\pm$ 2.2)	1.3 ($\pm$ 1.9)	<0.001
History of depression diagnosis	76 (58%)	84 (57%)	0.88

SD = standard deviation; MMPI = Minnesota Multiphasic Personality Inventory

* defined using the MMPI PSM scale (PSM $\geq$ 60 classified as pessimistic)

Table 2

Multivariable model of mortality among patients with rheumatoid arthritis (RA) who completed the Minnesota Multiphasic Personality Inventory prior to RA incidence

Characteristic	Hazard Ratio (95% Confidence Interval)
Age, years	1.09(1.06, 1.12)
Sex, male	0.89(0.50, 1.60)
Calendar year of RA incidence	1.00(0.97, 1.03)
Current smoker	2.34(1.17, 4.67)
Former smoker	1.08(0.55, 2.12)
Charlson comorbidity index	1.10(0.97, 1.24)
RF and pessimism	
RF– and not pessimistic	1.0 (reference)
RF+ and not pessimistic	2.28(1.10, 4.73)
RF– and pessimistic	2.88(1.02, 8.14)
RF+ and pessimistic	1.67 (0.77, 3.62)

Abbreviations: RF+: rheumatoid factor positive; RF– rheumatoid factor negative