Published in final edited form as:

Int Rev Neurobiol. 2016 ; 126: 293–355. doi:10.1016/bs.irm.2016.02.014.

Genes and Alcohol Consumption: Studies with Mutant Mice

Jody Mayfield¹, Michael A. Arends², R. Adron Harris¹, and Yuri A. Blednov¹

¹Waggoner Center for Alcohol and Addiction Research, The University of Texas at Austin, Austin, TX 78712

²Committee on the Neurobiology of Addictive Disorders, The Scripps Research Institute, La Jolla, CA 92037

Abstract

In this chapter, we review the effects of global null mutant and overexpressing transgenic mouse lines on voluntary self-administration of alcohol. We examine approximately 200 publications pertaining to the effects of 155 mouse genes on alcohol consumption in different drinking models. The targeted genes vary in function and include neurotransmitter, ion channel, neuroimmune, and neuropeptide signaling systems. The alcohol self-administration models include operant conditioning, two- and four-bottle choice continuous and intermittent access, drinking in the dark limited access, chronic intermittent ethanol, and scheduled high alcohol consumption tests. Comparisons of different drinking models using the same mutant mice are potentially the most informative, and we will highlight those examples. More mutants have been tested for continuous two-bottle choice consumption than any other test; of the 137 mouse genes examined using this model, 97 (72%) altered drinking in at least one sex. Overall, the effects of genetic manipulations on alcohol drinking often depend on the sex of the mice, alcohol concentration and time of access, genetic background, as well as the drinking test.

Keywords

global homozygous knockout; knockin; transgenic overexpression; voluntary self-administration; operant; continuous and intermittent two-bottle choice; drinking in the dark; scheduled high alcohol consumption

Introduction

Alcohol use disorder (AUD) is a multifactorial disease and its risk factors are determined by the interplay of genetic and environmental factors, combined with neuroadaptations following acute and repeated alcohol exposure. Alcohol targets ion channels and signaling cascades, producing intoxication, anxiolysis, and a sense of reward. After prolonged, repeated exposure, alcohol-induced changes in gene expression and synaptic function are thought to contribute to the development tolerance, sensitization, and compulsive consumption and drug seeking.

More than a hundred genes have been shown to affect alcohol consumption and other alcohol-related behaviors in mouse models (Crabbe, Phillips, Harris, Arends, & Koob, 2006). Excessive alcohol consumption is a common model of addictive behavior, and animal models of voluntary self-administration are valuable for profiling genetic determinants of AUD (Green & Grahame, 2008). The preference to drink alcohol is a reliable measure that depends upon mouse genotype and has been consistent across laboratories despite variations of the drinking protocol used. Pre-clinical models, in conjunction with human genetic studies, may expose overlapping target genes and identify the most relevant drinking models and biological systems associated with AUD.

In this review, we focus on a single phenotype, voluntary alcohol self-administration, and summarize the global genetic manipulations in mice published to date on this behavior. Most of the studies used two-bottle choice (2BC) tests, where mice had a choice between water and ethanol and access was usually measured in continuous 24-hour periods. In some cases, 2BC access to ethanol was intermittent (e.g., every other day), which typically (Hwa et al., 2011; Melendez, 2011; Rosenwasser, Fixaris, Crabbe, Brooks, & Ascheid, 2013), but not always (Crabbe, Harkness, Spence, Huang, & Metten, 2012), results in higher ethanol intake compared to continuous access. A few studies used four-bottle choice (4BC) access, where mice have simultaneous access to water and three different concentrations of ethanol. Because rodents distribute their drinking across the circadian cycle and because limited access to ethanol tends to increase intake, restricted access during the dark cycle is often used to study periods of high consumption and to model binge-like drinking in humans (Thiele & Navarro, 2014). In the classic mouse drinking in the dark (DID) test, drinking session times begin a few hours after the start of the dark cycle and usually last 2-4 hours over a few days. High levels of ethanol drinking and pharmacologically relevant blood ethanol concentrations (BECs) are achieved using this model (Thiele & Navarro, 2014). The scheduled high access consumption (SHAC) test uses fluid restriction to promote drinking of a low ethanol concentration (Finn et al., 2005). This chronic drinking model can also produce high BECs. Fluid access is first restricted and then gradually relaxed until the effects of fluid limitation are minimized. In operant self-administration tests, mice are trained to self-administer quantities of ethanol that produce moderate to high BECs. Removal of access to alcohol followed by restored access transiently increases consumption in dependent mice, and the effects of mutant genes on this alcohol deprivation effect (ADE) (Rodd, Bell, Sable, Murphy, & McBride, 2004; Vengeliene, Bilbao, & Spanagel, 2014), a model of relapse drinking, are also presented. As shown in the tables, the effects of some mutants can depend on the drinking test used as well as the ethanol concentration, time of access, genetic background, and sex of the mice (Vanderlinden, Saba, Bennett, Hoffman, & Tabakoff, 2015).

There are important considerations regarding genetic engineering methods, including the potential alteration of genes other than the mutated gene and the influence of the background strain carrying the genotype. C57BL/6J (B6) mice are a high alcohol drinking strain, and as the tables in this chapter demonstrate, occupy a central role in voluntary drinking studies. This review focuses on global homozygous knockouts, although a few studies used hypofunctional or overexpressing transgenic lines. Strategies to reduce confounding and compensatory effects of null mutations include the use of knockin mice and brain regional or

cell-specific knockouts, and a few of these studies are noted. In this review, we do not debate the genetic engineering methods used; instead, our aim is to provide a summary of homozygous null or overexpressing mutants and their role (or lack thereof) in alcohol consumption in mice. Current mouse gene and protein names from Uniprot (<http://www.uniprot.org>) are listed in the tables, which in some cases differ from the nomenclature used in the published studies. We first review the effects of mutant neurotransmitter receptor subunits on alcohol drinking in mice and then examine mutations in other ion channel receptors, cannabinoid and opioid receptors, neuropeptides, kinases/enzymes, and immune-related genes.

Neurotransmitter Systems

γ -aminobutyric acid (GABA)

Alcohol potentiates GABA_A receptor-mediated responses and enhances inhibitory neurotransmission, and some of the top candidate genes implicated in alcohol consumption code for specific GABA_A receptor subunits (Trudell, Messing, Mayfield, & Harris, 2014). Deletion of the $\alpha 1$ subunit decreased ethanol consumption in operant and 2BC tests (Blednov et al., 2003; June et al., 2007), and knockdown of $\alpha 5$ reduced drinking in male (Boehm, Ponomarev, et al., 2004) but not female mice (Stephens, Pistovcakova, Worthing, Atack, & Dawson, 2005) (Table 1). However, loss of $\alpha 2$ or $\beta 2$ subunits had no effect on drinking (Blednov et al., 2003; Boehm, Ponomarev, et al., 2004). Decreased drinking was also observed in mice lacking δ or $\rho 1$ subunits of GABA_A receptors (Blednov et al., 2014; Mihalek et al., 2001). In rats, local knockdown of the δ or $\alpha 4$ subunit in select regions of the nucleus accumbens reduced ethanol intake (Nie, Rewal, Gill, Ron, & Janak, 2011; Rewal et al., 2009). Deletion of glutamic acid decarboxylase, the principal enzyme for synthesis of GABA, had no effect on ethanol consumption if the mutation was expressed on a B6 background; however, consumption increased in *Gad2* knockout mice on a B6 $\times$ 129/SvJ (N2) background (Blednov, Walker, Iyer, Homanics, & Harris, 2010). Knockin mice with ethanol-insensitive $\alpha 1$ subunits did not differ from control in ethanol intake (D. F. Werner et al., 2006), whereas $\alpha 2$ subunits were linked with alcohol consumption in some drinking models using a knockin strategy (Blednov, Borghese, et al., 2011). Furthermore, human genetic association studies have nominated the GABA_A receptor $\alpha 2$ subunit as a top candidate in alcohol-dependent individuals, reviewed in (Enoch, 2008). A recent meta-analysis also identified a strong association of $\alpha 2$ with alcohol dependence along with evidence for $\alpha 6$ and $\gamma 2$ subunits (D. Li et al., 2014).

Glutamate

Ionotropic and some metabotropic glutamate receptors are known targets for alcohol action (D'Souza, 2015). Mice lacking *Gria1* (glutamate receptor ionotropic AMPA 1), i.e., the GluR1 subunit, did not differ from wild-type in voluntary ethanol consumption, stress-induced drinking, or in the expression of ADE (Cowen, Schroff, Gass, Sprengel, & Spanagel, 2003) (Table 2). Mice lacking *Gria3* (encoding the GluR3 subunit) did not differ from wild-type in the number of reinforced lever presses during operant ethanol self-administration, although cue-induced reinstatement was reduced (Sanchis-Segura et al., 2006). There were no genotype differences in voluntary ethanol consumption or preference

at any concentration tested (2-16%) in a 2BC test. Both genotypes showed increased ethanol consumption after a deprivation period following 14 weeks of 16% ethanol exposure, but the increase was initially blunted in the knockout mice. Although self-administration was not different between wild-type and mice lacking GluR1 or GluR3, ethanol seeking behavior after operant administration and resumption of self-administration after alcohol deprivation were reduced in GluR3 knockout mice, suggesting a role for the GluR3 subunit in alcohol seeking and relapse.

Mice lacking *Grm2*, which encodes the metabotropic glutamate receptor 2, increased consumption and preference for high ethanol concentrations in a 2BC model using an escalation procedure in which the ethanol concentration was increased from 3-17% over 80 days (Zhou et al., 2013) (Table 2). However, *Grm4* KO mice did not differ from wild-type in ethanol intake or preference in a 2BC test using 3-9% ethanol (Blednov, Walker, Osterndorf-Kahanek, & Harris, 2004). *Grm5* knockout mice differed from wild-type in some drinking models (Blednov & Harris, 2008). There were no differences in ethanol (3-12%) intake and preference in male or female mice in a 2BC test. Because the effects of deleting *Grm5* were somewhat greater in females, only females were tested in 4BC, 1-bottle and 2BC DID, and SHAC tests. Female *Grm5* knockout mice showed lowered ethanol intake and preference in the 4BC and 2BC DID tests, but there were no genotype differences in the other tests (Blednov & Harris, 2008). Decreased ethanol intake and preference were also reported in male *Grm5* KO mice consuming either 5 or 10% ethanol in a 2BC model (Bird, Kirchhoff, Djouma, & Lawrence, 2008).

The Homer family of postsynaptic scaffolding proteins modulates metabotropic glutamate receptors and is involved in addiction-related neuroplasticity. *Homer2* knockout mice drank less ethanol than wild-type (Szumlinski, Toda, Middaugh, Worley, & Kalivas, 2003) and showed aversion to 12% ethanol, with total ethanol intake less than half of wild-type mice in a 2BC test (Szumlinski et al., 2005) (Table 2). Infusion of Homer 2 into the nucleus accumbens reversed the effects on ethanol preference and intake in knockout mice. Furthermore, Homer 2b overexpression in the nucleus accumbens increased operant self-administration of 6 and 12% alcohol in B6 mice (Szumlinski, Ary, Lominac, Klugmann, & Kippin, 2008). In contrast to continuous access, there was no difference in intake between *Homer2* knockout and wild-type mice in a limited-access DID model using 20% ethanol (Lum, Campbell, Rostock, & Szumlinski, 2014). These authors reported that infusion of a metabotropic glutamate receptor 1 negative allosteric modulator into the nucleus accumbens reduced drinking in the DID test in control but not *Homer2* knockout mice, indicating a role for metabotropic glutamate receptors scaffolded by Homer 2 proteins in alcohol consumption.

A family of neuronal pentraxins regulates AMPA receptor aggregation and synaptogenesis at the postsynaptic density. Neuronal activity regulated pentraxin (*Narp*), also known as *Nptx2*, is an immediate early gene that is induced by synaptic activity and produces an AMPA receptor binding protein that enhances AMPA receptor clustering and function. *Narp* knockout mice did not escalate their daily ethanol intake after repeated exposure, and total ethanol intake in these mice was significantly lower compared to wild-type after 10 days of testing (Ary et al., 2012) (Table 2). Further study showed that *Narp* deletion shifted intake

and preference away from the high alcohol concentration (12%) with repeated alcohol exposure. *Narp* induction may thus be important for escalating alcohol consumption under free-choice conditions.

Chronic alcohol exposure and withdrawal produce a hyperexcitable or hyperglutamatergic state, and restoring the balance of glutamate responses could be relevant for treating dependence (Rao, Bell, Engleman, & Sari, 2015). Acamprosate (FDA approved to treat AUD) may work by inhibiting a hyperglutamatergic state in the brain and helping to balance inhibitory and excitatory transmission. In mouse models, acamprosate was effective in preventing the development of withdrawal (Blednov & Harris, 2008) and reducing free-choice ethanol intake (A. J. Brager, R. A. Prosser, & J. D. Glass, 2011; A. Brager, R. A. Prosser, & J. D. Glass, 2011; Spanagel et al., 2005). The anticonvulsant gabapentin is a calcium channel blocker that can also modulate glutamatergic (and GABAergic) activity and showed some promise in treating AUD in a clinical trial (Mason et al., 2014).

Dopamine

The mesolimbic dopamine pathway and its role in the rewarding properties of drugs of abuse has been a long-term focus for addiction research (van Huijstee & Mansvelder, 2014). Studies consistently show that ethanol self-administration in operant and 2BC tests is decreased in D₁- and D₂- receptor deficient mice (Delis et al., 2013; El-Ghundi et al., 1998; Palmer, Low, Grandy, & Phillips, 2003; Phillips et al., 1998; Risinger, Freeman, Rubinstein, Low, & Grandy, 2000; Thanos, Rivera, et al., 2005); however, deletion of the D₂ long receptor increased DID (Bulwa et al., 2011) (Table 3). Overexpression of D₂ transiently increased intake in D₂ receptor knockout mice and decreased intake in wild-type mice (Thanos, Rivera, et al., 2005). Effects of D₃ and D₄ dopamine receptor deletion are less well studied, but there is a report of substantially decreased alcohol consumption in 2BC and DID tests in D₃ knockout mice (Leggio et al., 2014). An example of the complex role of sex and genetic background is illustrated by the DAT knockout, which decreased drinking in females but not males on a B6 background and increased drinking in males but not females on a mixed genetic background (Table 3).

Serotonin

The serotonergic system has important roles in mood and impulse control and is implicated in the development and maintenance of alcohol and other drug abuse disorders (Bauer, Graham, Soares, & Nielsen, 2015). Ethanol intake and preference increased in a knockin mouse line (Tph2KI) expressing a hypofunctional variant of the rate-limiting serotonin (5-HT) synthesis enzyme, tryptophan hydroxylase 2 (Sachs, Salahi, & Caron, 2014) (Table 4). Increased ethanol self-administration has also been reported in mice lacking 5-HT_{1B} receptors (Bouwknicht et al., 2000; Crabbe et al., 1996; Risinger, Doan, & Vickrey, 1999). Knockout of 5-HT₆ receptors had no effect on 2BC drinking (Bonasera, Chu, Brennan, & Tecott, 2006). However, decreased intake was observed in B6 mice overexpressing 5-HT_{3A} receptors (Metz, Chynoweth, & Allan, 2006). Mice overexpressing 5-HT₃ receptors in the forebrain also drank less alcohol than non-transgenic mice in a 2BC test (Engel, Lyons, & Allan, 1998). The 5-HT transporter is the main means for regulating 5-HT levels in brain,

and deletion of the transporter decreased ethanol intake in B6 mice (Boyce-Rustay et al., 2006; Kelai et al., 2003; Lamb & Daws, 2013).

Although selective 5-HT reuptake inhibitors and the 5-HT₃ antagonist ondansetron were not promising as first-line treatments for AUD, they may provide some benefit in treating select populations of alcoholics. A review of pharmacotherapies for AUD in preclinical and clinical models can be found in (Zindel & Kranzler, 2014).

Adenosine

Adenosine signaling is implicated in drug addiction and ethanol-related behaviors in preclinical models (Nam, Bruner, & Choi, 2013). Deletion of the adenosine A₁ receptor did not alter operant responding for ethanol in B6 × 129 mice (Houchi, Persyn, Legastelois, & Naassila, 2013) (Table 5). Although male and female adenosine A_{2A} receptor (A2AR) knockout mice generated on a CD1 background drank more ethanol compared to wild-type in 2BC tests (Houchi et al., 2008; Naassila, Ledent, & Daoust, 2002), no differences were found between A2AR knockout and wild-type mice on a B6 background (Houchi et al., 2008).

Type 1 equilibrative nucleoside transporter (ENT1) is an ethanol-sensitive adenosine transporter and a main regulator of adenosine levels in the brain. Table 5 shows that ENT1 knockout mice generated on a B6 × 129 background consume more ethanol compared to wild-type littermates in operant and 2BC drinking tests (J. Chen et al., 2010; Choi et al., 2004; M. R. Lee et al., 2013; Nam, Hinton, et al., 2013; Nam et al., 2011; Ruby et al., 2014; Ruby, Walker, An, Kim, & Choi, 2011). Thus, studies from multiple labs support a role for adenosine in controlling alcohol-seeking behavior.

Cannabinoids and Opioids

The endocannabinoid system is involved in brain reward signaling and drug-seeking behavior (Panagis, Mackey, & Vlachou, 2014). In several studies of different genetic backgrounds, male and female cannabinoid 1 receptor (CB1R) knockout mice showed reduced ethanol intake and/or preference for ethanol than wild-type mice (Hungund, Szakall, Adam, Basavarajappa, & Vadasz, 2003; Lallemand & de Witte, 2005; Naassila, Pierrefiche, Ledent, & Daoust, 2004; Poncelet, Maruani, Calassi, & Soubrie, 2003; Racz et al., 2003; Thanos, Dimitrakakis, Rice, Gifford, & Volkow, 2005; Vinod, Yalamanchili, et al., 2008) (Table 6). In contrast, CB2R knockout mice showed increased ethanol consumption in two different drinking models (Ortega-Alvaro et al., 2015). Fatty acid amide hydrolase (FAAH) metabolizes endocannabinoids, and FAAH inhibition increases anandamide levels in brain. Increased ethanol intake and preference were reported in FAAH knockout mice on B6 and B6×129/SvJ backgrounds (Basavarajappa, Yalamanchili, Cravatt, Cooper, & Hungund, 2006; Blednov, Cravatt, Boehm, Walker, & Harris, 2007; Vinod, Sanguino, Yalamanchili, Manzanares, & Hungund, 2008).

Endogenous opioids are expressed throughout the reinforcement circuitry in the brain, and cannabinoid and opioid systems may interact to regulate the effects of drugs of abuse (Charbogne, Kieffer, & Befort, 2014). Table 6 shows the effects of opioid receptor mutations

on voluntary alcohol drinking. Mice lacking the mu-opioid receptor (MOR) did not self-administer ethanol (Roberts et al., 2000). This was demonstrated in two different operant procedures (nose-poke and lever press) and in 2BC drinking with or without prior ethanol experience. Other studies using MOR knockout mice reported decreased drinking compared to wild-type mice (Becker et al., 2002; Contet et al., 2014; Hall, Sora, & Uhl, 2001). Decreased consumption was also observed in kappa-opioid receptor (KOR) knockout mice (Kovacs et al., 2005; van Rijn & Whistler, 2009). In contrast, mice lacking delta-type opioid receptors (DORs) showed increased alcohol drinking behavior (Roberts et al., 2001; van Rijn & Whistler, 2009). The effects of deleting β -endorphin, prodynorphin, or preproenkephalin varied depending on the ethanol concentration, time of exposure, sex, and genotype (Table 6).

Consistent with the preclinical research showing that knockout of select opioid receptors reduces drinking in mice, human genetic studies further support a role for polymorphisms in genes coding for MOR1, DOR1, KOR1, and other opioid receptors in clinical populations of alcoholics (Levrán, Yuferov, & Kreek, 2012). In addition, opioids are used in the treatment of AUD. The pan-opioid antagonist naltrexone is already FDA approved to treat AUD and may reduce craving and relapse to heavy drinking in humans. Nalmefene, another pan-opioid receptor antagonist structurally similar to naltrexone, is currently used to treat opioid overdose in the U.S. and is approved in Europe for as-needed use to reduce heavy drinking. It has several advantages over naltrexone, including longer duration of action, lack of dose-dependent liver toxicity, and higher affinity of binding to MORs and KORs. Nalmefene reduced the total amount of alcohol consumed and the number of heavy drinking days in alcohol-dependent patients (van den Brink et al., 2014).

Immune-related Genes

The interplay between brain, behavior, and immune responses in the etiology and progression of drug abuse is a current area of interest in addiction research (http://www.arcr.niaaa.nih.gov/arcr/arcr372/toc37_2.htm). The neuroimmune system, encompassing innate immune signaling within the peripheral and central nervous systems, is important in the pathophysiology and potential treatment of alcohol abuse and dependence (Crews & Vetreno, 2015; Mayfield, Ferguson, & Harris, 2013; Robinson et al., 2014). Alcohol may increase neuroimmune gene and protein expression through both peripheral-central signaling molecules and direct actions in the brain (Crews & Vetreno, 2015).

Deletion of many different immune-related genes lowered ethanol intake and preference in male and female mice on a B6 background (Table 7). In some cases, the effect depended on the drinking test used. Deletion of certain chemokine and chemokine receptor genes reduced ethanol preference and intake (Blednov et al., 2005). Knockout mice were also used to study other immune-associated genes previously linked to alcohol consumption in a gene expression analysis of mouse brain (Mulligan et al., 2006). The following null mutations were examined based on this transcriptome meta-analysis: *B2m* (beta-2 microglobulin), *Cd14* (cluster of differentiation 14), *Il1rn* (interleukin 1 receptor antagonist), *Il6* (interleukin 6), *Ctss* (cathepsin S), and *Ctsf* (cathepsin F) (Blednov et al., 2012). These candidate targets were subsequently validated in behavioral tests because knockout of these individual genes

reduced ethanol consumption and preference in 2BC tests (Blednov et al., 2012). In contrast, a transgenic mouse line overexpressing *Il6* showed increased alcohol preference in females but not males (Harris & Blednov, 2013). C3H/HeJ mice are naturally TLR4-deficient and showed decreased operant self-administration of ethanol compared to the control strain (Harris & Blednov, 2013); however, ethanol consumption was not changed in TLR4 knockout mice in a 2BC test (Pascual, Balino, Alfonso-Loeches, Aragon, & Guerri, 2011).

Pharmacological inhibition of immune signaling reduces alcohol drinking in mice, whereas activation of the innate immune system using lipopolysaccharide produces prolonged increases in drinking (Blednov, Benavidez, et al., 2011). Immune molecules, acting peripherally and/or centrally, appear to be involved in preference for alcohol and the behavioral evidence agrees with expression profiling of mouse brain (Robinson et al., 2014). Unique gene expression patterns in human post-mortem brain provide additional support for immune gene and related network effects in alcohol-dependent individuals (Liu et al., 2006; Ponomarev, Wang, Zhang, Harris, & Mayfield, 2012). Furthermore, genetic association studies in humans link immune-associated genes with alcohol dependence (Blednov, Benavidez, Black, Ferguson, et al., 2015). The use of agents that regulate immune responses as potential therapeutics for AUD is discussed in the Medication Development section later in the chapter.

Ion Channels

Some of the rapid-onset actions of alcohol are likely mediated by direct action on ion channels (Howard, Trudell, & Harris, 2014; Trudell et al., 2014). In addition to the prominent and well-studied neurotransmitter systems in alcohol dependence discussed previously (e.g., GABA, glutamate, dopamine, 5-HT), other ion channels that have been implicated in alcohol intake and preference are described below.

Transient receptor potential (TRP) cation channels

TRP channels are a group of non-selective ion channels that mediate sensations, tastes, temperature, pain, and other chemical and physical stimuli. There is a positive correlation between ethanol intake and consumption of sweet solutions in rodents and humans (Kampov-Polevoy et al., 2014). Deletion of TRP cation channel subfamily M, member 5 substantially decreased ethanol intake and preference (Blednov et al., 2008) (Table 8). Altered taste perception may explain the reduced preference for both saccharin and ethanol. Mice lacking TRP cation channel subfamily V, member 1 (TRPV1), also known as the vanilloid or capsaicin receptor, showed increased ethanol preference but no change in saccharin or quinine preference (Blednov & Harris, 2009). The endocannabinoid, anandamide, is an endogenous activator of this channel, and ethanol also directly activates TRPV1 at high concentrations. The increased ethanol consumption in *Trpv1* knockout mice may simply be due to a reduction in the ethanol-induced burning sensation, although some central actions, including endocannabinoid signaling, may be involved.

N-type calcium channels

Voltage-dependent calcium channels regulate neuronal excitability, neurotransmitter release, and gene expression. Several types exist with unique pharmacological and electrophysiological characteristics. L-, T-, and N-type voltage-gated calcium channels have all been implicated in the behavioral effects of ethanol. Reduced ethanol intake and preference were reported in mice lacking functional α_{1B} subunits of N-type calcium channels (Newton et al., 2004) (Table 8), suggesting these subunits as a potential target to control drinking. The calcium channel blocker gabapentin targets channels with $\alpha_2\delta$ subunits (Geisler, Schopf, & Obermair, 2015) and, as previously mentioned, offers some promise as a therapeutic for AUD (Mason et al., 2014).

GIRK channels

G protein-coupled inwardly rectifying K^+ (GIRK) channels are widely expressed in brain and are activated by ligand-stimulated G protein-coupled receptors, resulting in an outward K^+ current that hyperpolarizes neuronal membranes and decreases neuronal excitability. Genetic evidence from mouse models has provided extensive insight into the significance of GIRK channels in drug addiction, and their role in responses to ethanol and other drugs of abuse is reviewed in (Bodhinathan & Slesinger, 2014) and (Mayfield, Blednov, & Harris, 2015).

As shown in Table 8, ethanol consumption and preference did not differ in wild-type and *Girk2* knockout mice in the standard 2BC test, where the bottle positions were alternated daily to control for position preferences. However, when the ethanol bottles were always available in the preferred location, *Girk2* knockout mice consumed more ethanol compared to wild-type (Blednov, Stoffel, Chang, & Harris, 2001). Moreover, single nucleotide polymorphisms of the *KCJN6* gene (GIRK2) have been associated with alcohol dependence in humans (Clarke et al., 2011; Kang et al., 2012). Deletion of *Girk3* increased limited- but not continuous-access voluntary drinking, and overexpression of *Girk3* in the VTA reversed the binge-drinking phenotype and reduced drinking in wild-type mice (Herman et al., 2015). These results point to a role for GIRK subunits in the rewarding properties of ethanol and as a potential target for regulating binge-like drinking.

BK channels

BK channels are calcium-activated potassium channels that are characterized by their large conductance of potassium ions. The β subunits influence BK channel responses to acute and chronic ethanol, and ethanol action on one system cannot easily be extrapolated to another. In 2BC continuous and intermittent access tests, non-dependent mice lacking BK $\beta 1$ or $\beta 4$ subunits did not alter their drinking compared to wild-type (Kreifeldt, Le, Treistman, Koob, & Contet, 2013) (Table 8). Weeks of voluntary 2BC drinking alternated with weeks of CIE vapor escalated drinking in dependent wild-type mice to a greater extent than BK $\beta 4$ knockout mice. In contrast, $\beta 1$ knockout mice drank more after fewer CIE cycles than did wild-type mice. Thus loss of $\beta 4$ decreased, while loss of $\beta 1$ increased, the escalation of drinking during withdrawal from CIE (Kreifeldt et al., 2013).

Nicotinic acetylcholine receptors

Neuronal nicotinic acetylcholine receptors (nAChRs) are involved in several alcohol-mediated behaviors (Rahman, Engleman, & Bell, 2014). Varenicline (which has partial agonist activity at nAChRs and is used in smoking cessation programs) reduced drinking in both preclinical and clinical studies, and human genetic studies showed a strong association of polymorphisms in *CHRNA5* (which encodes the $\alpha 5$ subunit) with risk of developing alcohol dependence, reviewed in (Rahman et al., 2014). Mice lacking *Chrna5* did not differ from wild-type in ethanol consumption (Santos, Chatterjee, Henry, Holgate, & Bartlett, 2013), but transgenic mice overexpressing $\alpha 5\alpha 3\beta 4$ showed reduced intake (Gallego et al., 2012) (Table 8). Knockout of $\alpha 6$ nACh subunits had no effect on 2BC drinking (Kamens, Hoft, Cox, Miyamoto, & Ehringer, 2012). However, deletion of $\alpha 4$ decreased DID of a high ethanol concentration (Hendrickson, Zhao-Shea, Pang, Gardner, & Tapper, 2010), and loss of $\alpha 7$ decreased 2BC drinking in female mice (Kamens, Andersen, & Picciotto, 2010). Knockout of $\beta 2$ (Dawson, Miles, & Damaj, 2013; Kamens et al., 2010) or $\beta 3$ (Kamens et al., 2012) nAChR subunits had no effect in 2BC tests in male or female mice.

Glycine receptors

Supraspinal glycine receptors are found in reward pathways and play a key role in alcohol inhibitory responses in some neurons (Perkins, Trudell, Crawford, Alkana, & Davies, 2010). The glycine receptor $\alpha 1$ subunit has proven difficult to engineer and studies are complicated by the reduced glycinergic function and lethality of homozygous mutants. Deletion of glycine receptor subunits $\alpha 2$ or $\alpha 3$ had no effect on limited-access DID, while 2BC continuous-access drinking decreased in $\alpha 2$ but not $\alpha 3$ null mutants (Blednov, Benavidez, Black, Leiter, et al., 2015) (Table 8). However, intermittent-access drinking increased in $\alpha 3$ but not $\alpha 2$ null mutants. A linkage analysis in humans also supports a role for the $\alpha 3$ subunit in alcohol dependence (Han, Gelernter, Kranzler, & Yang, 2013).

Protein Kinases

In addition to direct effects on ion channels, ethanol indirectly modulates channel function via phosphorylation and other post-translational processing mechanisms (Trudell et al., 2014). Mice lacking protein kinase C type ϵ (PKC ϵ) drank less ethanol than wild-type, and this effect has been observed in different drinking tests across multiple labs (Besheer, Lepoutre, Mole, & Hodge, 2006; Choi, Wang, Dadgar, Chang, & Messing, 2002; Hodge et al., 1999; Olive et al., 2005; Olive, Mehmert, Messing, & Hodge, 2000; Wallace et al., 2007) (Table 9). Selective knockdown of PKC ϵ in the amygdala also reduced 2-hour ethanol consumption in a DID model (Lesscher et al., 2009). PKC ϵ is known to reduce the positive allosteric effects of benzodiazepines and ethanol at GABA $_A$ receptors, which may in turn regulate reward signaling. The effects of deleting other PKC types on alcohol intake and preference have not been studied as extensively, although drinking in 2BC tests increased in mice lacking PKC γ (Bowers & Wehner, 2001) or ζ (A. M. Lee et al., 2014).

Enzymes

Table 10 shows the effects of other assorted enzymes on ethanol drinking in mice. In particular, aldehyde dehydrogenase (ALDH) is one of the few known genes to affect risk of

developing AUD in humans (C. H. Chen, Ferreira, Gross, & Mochly-Rosen, 2014). ALDH2 plays a major role in the detoxification of ethanol-derived acetaldehyde, and inhibition of ALDH is the mechanism of action of disulfiram, an FDA-approved drug for AUD. A mutation in *ALDH2* produces an enzyme incapable of metabolizing ethanol, resulting in severe adverse reactions to even small amounts of ingested alcohol in individuals who are homozygous for the mutation. UChA and UChB rat lines show similar *Aldh2* polymorphisms that affect alcohol consumption (Quintanilla, Israel, Sapag, & Tampier, 2006). Knockout of the *Aldh2* gene in mice was also shown to limit drinking in 2BC tests (Fernandez et al., 2006; Isse et al., 2002). Thus, genetic polymorphisms that alter ethanol metabolism and susceptibility to its effects may decrease ethanol-induced metabolic tolerance and risk for developing AUD.

Neuropeptides/Hormones

Table 11 summarizes the effects of deletion or overexpression of classical or putative neuropeptides and hormones and their receptors on voluntary ethanol administration in mice. Of the genes represented here, the corticotropin releasing factor/urocortin family and other stress-related neuromodulators are promising for future studies of genetic determinants of AUD (Schank, Ryabinin, Giardino, Ciccocioppo, & Heilig, 2012).

Other Gene Targets

Table 12 represents an assortment of genes that did not specifically fit into the previous categories. These genes are associated with synaptic function, development, circadian regulation, and other cellular regulatory functions but are not discussed individually herein. One of the most pronounced phenotypes among mutant mice is the almost complete blockade of alcohol consumption in mice lacking any one of three taste genes (*Gnat3*, *Tas1r3*, *Trpm5*) (Blednov et al., 2008).

Concluding Remarks

Mouse models of voluntary ethanol administration have been instrumental for profiling putative behavioral and genetic determinants in human alcoholics, who exhibit excessive consumption as a hallmark of the disease. The impact of more than 150 genes on alcohol consumption has been evaluated by construction of mutant mice. The global knockout strategy has been used extensively in addiction research to link proteins with behavior, and most studies presented in this chapter used this approach. Collectively speaking, the studies suggest the potential involvement of a large number of different mouse genes in voluntary alcohol drinking. However, many of the effects are modest and global knockout or overexpression of a specific gene does not determine if the effect on drinking is directly linked to effects of decreased or increased expression of the respective protein. Other technologies such as knockin animal models and conditional and cell-specific knockouts will provide more discriminating tools for linking ethanol-sensitive sites on proteins to behaviors (Blednov, Borghese, et al., 2011). The goal of the knockin approach is to remove ethanol action on a specific protein without otherwise altering its function, but this strategy (like the knockout model) can still be complicated by compensatory actions from other proteins. Selectively targeting an ethanol site could also affect other modulatory sites of the

protein despite the appearance of normal function (Harris, Osterndorff-Kahanek, Ponomarev, Homanics, & Blednov, 2011).

When focusing on genes that have more profound effects on consumption and have consistently been implicated across multiple drinking tests, the number of targets diminishes. For example, null mutant mice for specific GABA_A (Table 1), dopamine (Table 3), cannabinoid and opioid receptors (Table 6), immune-related genes (Table 7), and PKC ϵ (Table 9) have demonstrated decreased drinking and preference in several drinking tests. These systems either have established roles in addiction research or represent emerging targets, as in the case of immune pathways. Further consideration should be given to the impact of the drinking paradigms on peripheral and central genes. For example, even when similar amounts of total ethanol were consumed in the continuous and intermittent access tests by B6 mice, brain and liver transcriptomes were differentially affected (Osterndorff-Kahanek, Ponomarev, Blednov, & Harris, 2013). The distinct genomic effects induced by the alcohol exposure protocol may explain the differing effects observed in some drinking tests. The peripheral effects of alcohol may be particularly important for neuroimmune modulation of drinking because production of cytokines outside the brain can influence brain function. This peripheral-central neuroimmune communication may also be important in alcohol craving and dependence in humans (Leclercq, De Saeger, Delzenne, de Timary, & Starkel, 2014).

Medication Development

FDA-approved drugs for AUD have provided only modest benefit and are not routinely prescribed, and so the search continues for more effective drugs. Identifying existing drugs that could be repurposed to treat AUD is a current goal for researchers and, if successful, would fast track therapeutic options for the disease. A strategy for prioritizing relevant genes from a large list of potential targets is to examine the preclinical evidence in combination with genetic association studies in human alcoholics. Overlapping evidence from animal and human studies will help pinpoint genes and their biological networks that could be most relevant in the treatment of AUD. Mouse drinking models could then be used to test viable targets using approved drugs known to modulate the target network.

Pro-inflammatory neuroimmune signaling pathways have been increasingly implicated in the etiology and progression of AUD and offer unexplored avenues in its treatment. Currently, this is an emphasis area of the Integrative Neuroscience Initiative on Alcoholism (INIA) sponsored by the National Institutes of Health, National Institute on Alcohol Abuse and Alcoholism. Several FDA-approved drugs with anti-inflammatory and immune inhibitory effects regulate alcohol responses in animal models, lending support that some of these could potentially be repurposed to treat AUD. For example, the peroxisome proliferator-activated receptor (PPAR) agonists, fenofibrate and tesaglitazar, reduced drinking in a PPAR α -dependent manner (Blednov, Black, Benavidez, Stamatakis, & Harris, 2016), altered expression of immune genes in the liver, and altered neuronal gene expression in mouse brain (Ferguson, Most, Blednov, & Harris, 2014). There is overlapping evidence in mice and humans for specific PPAR genes in alcohol consumption and dependence (Blednov, Benavidez, Black, Ferguson, et al., 2015). In addition, minocycline reduced DID

in adult mice, and a bioinformatics pathway analysis revealed an overexpression of neuroimmune-related pathways in these mice (Agrawal et al., 2014). Mouse models will continue to be essential for identifying candidate genes and screening medications to modulate the affected pathways.

Future Directions

While the study of individual genes is informative, combining gene network and systems biology approaches to identify inter-related networks and pathways is critical in the future treatment of AUD. Because complex trait diseases involve coordinated expression changes in multiple gene families, examining gene clusters is an important research direction, as supported by the Integrative Neuroscience Initiative on Alcoholism (INIA) studies showing that co-expression patterns can distinguish gene modules related to alcohol consumption in animal models (Iancu et al., 2013; Nunez et al., 2013; Saba et al., 2011; Saba et al., 2015). Another INIA study reported that RNA-Seq profiling of postmortem human prefrontal cortex also revealed disrupted gene networks in alcohol-dependent individuals compared to matched controls (Farris, Arasappan, Hunicke-Smith, Harris, & Mayfield, 2015). The subnetworks related to lifetime alcohol consumption in humans contained known alcohol targets, from the preclinical literature, and were overrepresented for genes involved in synaptic function. These genes were also enriched in a transcriptome meta-analysis of mouse drinking behavior (Mulligan et al., 2006), suggesting an overlapping set of alcohol-related genes across species. Comparison of human and animal model networks, combined with single gene-based approaches, will be important in understanding the neurobiology of AUD (Farris, Pietrzykowski, et al., 2015).

Genomic and proteomic techniques are capable of analyzing complex data sets into related biological processes (Gorini, Bell, & Mayfield, 2011; Gorini, Harris, & Mayfield, 2014; Gorini, Roberts, & Mayfield, 2013; Sikela et al., 2006). Ideally, integration of genetics and transcriptomics with convergent biological processes and phenotypic behaviors will reveal prospective therapeutic targets for the disease (Gorini et al., 2011; Gorini et al., 2014). Drug addiction is mediated by both genetic and environmental determinants and neuroadaptations in reward circuitry are related to disease progression. Therefore, in addition to individual candidate genes, future research must consider the gene and protein networks and associated biological system changes observed at different stages of the disease. These network-centric approaches can summarize complex gene lists into interrelated components and provide a systems-level picture of the disease. This biological systems framework will be key in determining causal factors of AUD and providing a more integrated, functional approach to its treatment.

Acknowledgments

The authors acknowledge funding from NIAAA grants AA006399, AA013520, AA020926, and AA012404. The authors have no conflicts of interest with this material. The views expressed herein are solely those of the authors and do not necessarily represent those of the funding agencies.

References

- Abramov U, Raud S, Innos J, Koks S, Matsui T, Vasar E. Gender specific effects of ethanol in mice, lacking CCK2 receptors. *Behav Brain Res.* 2006; 175(1):149–156. DOI: 10.1016/j.bbr.2006.08.015 [PubMed: 16970998]
- Agrawal RG, Owen JA, Levin PS, Hewetson A, Berman AE, Franklin SR, et al. Bergeson SE. Bioinformatics analyses reveal age-specific neuroimmune modulation as a target for treatment of high ethanol drinking. *Alcohol Clin Exp Res.* 2014; 38(2):428–437. DOI: 10.1111/acer.12288 [PubMed: 24125126]
- Ary AW, Cozzoli DK, Finn DA, Crabbe JC, Dehoff MH, Worley PF, Szumlinski KK. Ethanol up-regulates nucleus accumbens neuronal activity dependent pentraxin (Narp): implications for alcohol-induced behavioral plasticity. *Alcohol.* 2012; 46(4):377–387. DOI: 10.1016/j.alcohol.2011.10.003 [PubMed: 22444953]
- Bahi A, Tolle V, Fehrentz JA, Brunel L, Martinez J, Tomasetto CL, Karam SM. Ghrelin knockout mice show decreased voluntary alcohol consumption and reduced ethanol-induced conditioned place preference. *Peptides.* 2013; 43:48–55. DOI: 10.1016/j.peptides.2013.02.008 [PubMed: 23428971]
- Basavarajappa BS, Yalamanchili R, Cravatt BF, Cooper TB, Hungund BL. Increased ethanol consumption and preference and decreased ethanol sensitivity in female FAAH knockout mice. *Neuropharmacology.* 2006; 50(7):834–844. DOI: 10.1016/j.neuropharm.2005.12.005 [PubMed: 16448676]
- Bauer IE, Graham DP, Soares JC, Nielsen DA. Serotonergic gene variation in substance use pharmacotherapy: a systematic review. *Pharmacogenomics.* 2015; 16(11):1307–1314. DOI: 10.2217/pgs.15.72 [PubMed: 26265436]
- Bechtholt AJ, Smith R, Raber J, Cunningham CL. Enhanced ethanol-, but not cocaine-induced, conditioned place preference in Apoe(-/-) mice. *Pharmacol Biochem Behav.* 2004; 77(4):783–792. DOI: 10.1016/j.pbb.2004.02.002 [PubMed: 15099924]
- Becker A, Grecksch G, Kraus J, Loh HH, Schroeder H, Holtt V. Rewarding effects of ethanol and cocaine in mu opioid receptor-deficient mice. *Naunyn Schmiedeberg's Arch Pharmacol.* 2002; 365(4):296–302. DOI: 10.1007/s00210-002-0533-2 [PubMed: 11919654]
- Besheer J, Lepoutre V, Mole B, Hodge CW. GABAA receptor regulation of voluntary ethanol drinking requires PKCepsilon. *Synapse.* 2006; 60(6):411–419. DOI: 10.1002/syn.20314 [PubMed: 16881070]
- Bird MK, Kirchhoff J, Djouma E, Lawrence AJ. Metabotropic glutamate 5 receptors regulate sensitivity to ethanol in mice. *Int J Neuropsychopharmacol.* 2008; 11(6):765–774. DOI: 10.1017/S1461145708008572 [PubMed: 18400131]
- Bjork K, Rimondini R, Hansson AC, Terasmaa A, Hyytia P, Heilig M, Sommer WH. Modulation of voluntary ethanol consumption by beta-arrestin 2. *FASEB J.* 2008; 22(7):2552–2560. DOI: 10.1096/fj.07-102442 [PubMed: 18367649]
- Blednov YA, Benavidez JM, Black M, Ferguson LB, Schoenhard GL, Goate AM, et al. Harris RA. Peroxisome proliferator-activated receptors alpha and gamma are linked with alcohol consumption in mice and withdrawal and dependence in humans. *Alcohol Clin Exp Res.* 2015; 39(1):136–145. DOI: 10.1111/acer.12610 [PubMed: 25516156]
- Blednov YA, Benavidez JM, Black M, Leiter CR, Osterndorff-Kahanek E, Harris RA. Glycine receptors containing alpha2 or alpha3 subunits regulate specific ethanol-mediated behaviors. *J Pharmacol Exp Ther.* 2015; 353(1):181–191. DOI: 10.1124/jpet.114.221895 [PubMed: 25678534]
- Blednov YA, Benavidez JM, Black M, Leiter CR, Osterndorff-Kahanek E, Johnson D, et al. Harris RA. GABAA receptors containing rho1 subunits contribute to in vivo effects of ethanol in mice. *PLoS One.* 2014; 9(1):e85525.doi: 10.1371/journal.pone.0085525 [PubMed: 24454882]
- Blednov YA, Benavidez JM, Geil C, Perra S, Morikawa H, Harris RA. Activation of inflammatory signaling by lipopolysaccharide produces a prolonged increase of voluntary alcohol intake in mice. *Brain Behav Immun.* 2011; 25 Suppl 1:S92–S105. DOI: 10.1016/j.bbi.2011.01.008 [PubMed: 21266194]

- Blednov YA, Bergeson SE, Walker D, Ferreira VM, Kuziel WA, Harris RA. Perturbation of chemokine networks by gene deletion alters the reinforcing actions of ethanol. *Behav Brain Res.* 2005; 165(1):110–125. DOI: 10.1016/j.bbr.2005.06.026 [PubMed: 16105698]
- Blednov YA, Black M, Benavidez JM, Stamatakis EE, Harris RA. PPAR Agonists: I. Role of receptor subunits in alcohol consumption in male and female mice. *Alcohol Clin Exp Res.* 2016; doi: 10.1111/acer.12976
- Blednov YA, Borghese CM, McCracken ML, Benavidez JM, Geil CR, Osterndorff-Kahanek E, et al. Harris RA. Loss of ethanol conditioned taste aversion and motor stimulation in knockin mice with ethanol-insensitive alpha2-containing GABA(A) receptors. *J Pharmacol Exp Ther.* 2011; 336(1): 145–154. DOI: 10.1124/jpet.110.171645 [PubMed: 20876231]
- Blednov YA, Cravatt BF, Boehm SL 2nd, Walker D, Harris RA. Role of endocannabinoids in alcohol consumption and intoxication: studies of mice lacking fatty acid amide hydrolase. *Neuropsychopharmacology.* 2007; 32(7):1570–1582. DOI: 10.1038/sj.npp.1301274 [PubMed: 17164820]
- Blednov YA, Harris RA. Metabotropic glutamate receptor 5 (mGluR5) regulation of ethanol sedation, dependence and consumption: relationship to acamprosate actions. *Int J Neuropsychopharmacol.* 2008; 11(6):775–793. DOI: 10.1017/S1461145708008584 [PubMed: 18377703]
- Blednov YA, Harris RA. Deletion of vanilloid receptor (TRPV1) in mice alters behavioral effects of ethanol. *Neuropharmacology.* 2009; 56(4):814–820. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19705551>. [PubMed: 19705551]
- Blednov YA, Ponomarev I, Geil C, Bergeson S, Koob GF, Harris RA. Neuroimmune regulation of alcohol consumption: behavioral validation of genes obtained from genomic studies. *Addict Biol.* 2012; 17(1):108–120. DOI: 10.1111/j.1369-1600.2010.00284.x [PubMed: 21309947]
- Blednov YA, Stoffel M, Chang SR, Harris RA. Potassium channels as targets for ethanol: studies of G-protein-coupled inwardly rectifying potassium channel 2 (GIRK2) null mutant mice. *J Pharmacol Exp Ther.* 2001; 298(2):521–530. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11454913>. [PubMed: 11454913]
- Blednov YA, Walker D, Alva H, Creech K, Findlay G, Harris RA. GABAA receptor alpha 1 and beta 2 subunit null mutant mice: behavioral responses to ethanol. *J Pharmacol Exp Ther.* 2003; 305(3): 854–863. DOI: 10.1124/jpet.103.049478 [PubMed: 12626647]
- Blednov YA, Walker D, Harris RA. Blockade of the leptin-sensitive pathway markedly reduces alcohol consumption in mice. *Alcohol Clin Exp Res.* 2004; 28(11):1683–1692. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15547455>. [PubMed: 15547455]
- Blednov YA, Walker DL, Iyer SV, Homanics G, Harris AR. Mice lacking Gad2 show altered behavioral effects of ethanol, flurazepam and gabaxadol. *Addict Biol.* 2010; 15(1):45–61. DOI: 10.1111/j.1369-1600.2009.00186.x [PubMed: 20002022]
- Blednov YA, Walker D, Martinez M, Harris RA. Reduced alcohol consumption in mice lacking preprodynorphin. *Alcohol.* 2006; 40(2):73–86. DOI: 10.1016/j.alcohol.2006.12.002 [PubMed: 17307643]
- Blednov YA, Walker D, Martinez M, Levine M, Damak S, Margolskee RF. Perception of sweet taste is important for voluntary alcohol consumption in mice. *Genes Brain Behav.* 2008; 7(1):1–13. DOI: 10.1111/j.1601-183X.2007.00309.x
- Blednov YA, Walker D, Osterndorf-Kahanek E, Harris RA. Mice lacking metabotropic glutamate receptor 4 do not show the motor stimulatory effect of ethanol. *Alcohol.* 2004; 34(2-3):251–259. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15902920>. [PubMed: 15902920]
- Bodhinathan K, Slesinger PA. Alcohol modulation of G-protein-gated inwardly rectifying potassium channels: from binding to therapeutics. *Front Physiol.* 2014; 5:76.doi: 10.3389/fphys.2014.00076 [PubMed: 24611054]
- Boehm SL 2nd, Peden L, Chang R, Harris RA, Blednov YA. Deletion of the fyn-kinase gene alters behavioral sensitivity to ethanol. *Alcohol Clin Exp Res.* 2003; 27(7):1033–1040. DOI: 10.1097/01.ALC.0000075822.80583.71 [PubMed: 12878908]
- Boehm SL 2nd, Peden L, Jennings AW, Kojima N, Harris RA, Blednov YA. Over-expression of the fyn-kinase gene reduces hypnotic sensitivity to ethanol in mice. *Neurosci Lett.* 2004; 372(1-2):6–11. DOI: 10.1016/j.neulet.2004.08.028 [PubMed: 15531078]

- Boehm SL 2nd, Ponomarev I, Jennings AW, Whiting PJ, Rosahl TW, Garrett EM, et al. Harris RA. gamma-Aminobutyric acid A receptor subunit mutant mice: new perspectives on alcohol actions. *Biochem Pharmacol.* 2004; 68(8):1581–1602. DOI: 10.1016/j.bcp.2004.07.023 [PubMed: 15451402]
- Bonasera SJ, Chu HM, Brennan TJ, Tecott LH. A null mutation of the serotonin 6 receptor alters acute responses to ethanol. *Neuropsychopharmacology.* 2006; 31(8):1801–1813. DOI: 10.1038/sj.npp.1301030 [PubMed: 16452990]
- Bouwknicht JA, Hijzen TH, van der Gugten J, Maes RA, Hen R, Olivier B. Ethanol intake is not elevated in male 5-HT(1B) receptor knockout mice. *Eur J Pharmacol.* 2000; 403(1-2):95–98. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10969149>. [PubMed: 10969149]
- Bowers BJ, Wehner JM. Ethanol consumption and behavioral impulsivity are increased in protein kinase Cgamma null mutant mice. *J Neurosci.* 2001; 21(21):RC180. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11606660>. [PubMed: 11606660]
- Boyce-Rustay JM, Holmes A. Ethanol-related behaviors in mice lacking the NMDA receptor NR2A subunit. *Psychopharmacology (Berl).* 2006; 187(4):455–466. DOI: 10.1007/s00213-006-0448-6 [PubMed: 16835771]
- Boyce-Rustay JM, Risinger FO. Dopamine D3 receptor knockout mice and the motivational effects of ethanol. *Pharmacol Biochem Behav.* 2003; 75(2):373–379. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12873629>. [PubMed: 12873629]
- Boyce-Rustay JM, Wiedholz LM, Millstein RA, Carroll J, Murphy DL, Daws LC, Holmes A. Ethanol-related behaviors in serotonin transporter knockout mice. *Alcohol Clin Exp Res.* 2006; 30(12):1957–1965. DOI: 10.1111/j.1530-0277.2006.00241.x [PubMed: 17117959]
- Brager AJ, Prosser RA, Glass JD. Circadian and acamprosate modulation of elevated ethanol drinking in mPer2 clock gene mutant mice. *Chronobiol Int.* 2011; 28(8):664–672. DOI: 10.3109/07420528.2011.601968 [PubMed: 21929298]
- Brager A, Prosser RA, Glass JD. Acamprosate-responsive brain sites for suppression of ethanol intake and preference. *Am J Physiol Regul Integr Comp Physiol.* 2011; 301(4):R1032–1043. DOI: 10.1152/ajpregu.00179.2011 [PubMed: 21697518]
- Brasser SM, Norman MB, Lemon CH. Trp3 taste receptor involvement in gustatory neural responses to ethanol and oral ethanol preference. *Physiol Genomics.* 2010; 41(3):232–243. DOI: 10.1152/physiolgenomics.00113.2009 [PubMed: 20145204]
- Bulwa ZB, Sharlin JA, Clark PJ, Bhattacharya TK, Kilby CN, Wang Y, Rhodes JS. Increased consumption of ethanol and sugar water in mice lacking the dopamine D2 long receptor. *Alcohol.* 2011; 45(7):631–639. DOI: 10.1016/j.alcohol.2011.06.004 [PubMed: 21803530]
- Cai YQ, Cai GQ, Liu GX, Cai Q, Shi JH, Shi J, et al. Fei J. Mice with genetically altered GABA transporter subtype I (GAT1) expression show altered behavioral responses to ethanol. *J Neurosci Res.* 2006; 84(2):255–267. DOI: 10.1002/jnr.20884 [PubMed: 16683252]
- Caldwell HK, Stewart J, Wiedholz LM, Millstein RA, Iacangelo A, Holmes A, et al. Wersinger SR. The acute intoxicating effects of ethanol are not dependent on the vasopressin 1a or 1b receptors. *Neuropeptides.* 2006; 40(5):325–337. DOI: 10.1016/j.npep.2006.08.001 [PubMed: 17049983]
- Camp MC, Feyder M, Ihne J, Palachick B, Hurd B, Karlsson RM, et al. Holmes A. A novel role for PSD-95 in mediating ethanol intoxication, drinking and place preference. *Addict Biol.* 2011; 16(3):428–439. DOI: 10.1111/j.1369-1600.2010.00282.x [PubMed: 21309945]
- Charbogne P, Kieffer BL, Befort K. 15 years of genetic approaches in vivo for addiction research: Opioid receptor and peptide gene knockout in mouse models of drug abuse. *Neuropharmacology.* 2014; 76 Pt B:204–217. DOI: 10.1016/j.neuropharm.2013.08.028 [PubMed: 24035914]
- Chen CH, Ferreira JC, Gross ER, Mochly-Rosen D. Targeting aldehyde dehydrogenase 2: new therapeutic opportunities. *Physiol Rev.* 2014; 94(1):1–34. DOI: 10.1152/physrev.00017.2013 [PubMed: 24382882]
- Chen J, Nam HW, Lee MR, Hinton DJ, Choi S, Kim T, et al. Choi DS. Altered glutamatergic neurotransmission in the striatum regulates ethanol sensitivity and intake in mice lacking ENT1. *Behav Brain Res.* 2010; 208(2):636–642. DOI: 10.1016/j.bbr.2010.01.011 [PubMed: 20085785]

- Choi DS, Cascini MG, Mailliard W, Young H, Paredes P, McMahon T, et al. Messing RO. The type 1 equilibrative nucleoside transporter regulates ethanol intoxication and preference. *Nat Neurosci.* 2004; 7(8):855–861. DOI: 10.1038/nn1288 [PubMed: 15258586]
- Choi DS, Wang D, Dadgar J, Chang WS, Messing RO. Conditional rescue of protein kinase C epsilon regulates ethanol preference and hypnotic sensitivity in adult mice. *J Neurosci.* 2002; 22(22): 9905–9911. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12427847>. [PubMed: 12427847]
- Chu K, Koob GF, Cole M, Zorrilla EP, Roberts AJ. Dependence-induced increases in ethanol self-administration in mice are blocked by the CRF1 receptor antagonist antalarmin and by CRF1 receptor knockout. *Pharmacol Biochem Behav.* 2007; 86(4):813–821. DOI: 10.1016/j.pbb.2007.03.009 [PubMed: 17482248]
- Clarke TK, Laucht M, Ridinger M, Wodarz N, Rietschel M, Maier W, et al. Schumann G. KCNJ6 is associated with adult alcohol dependence and involved in gene x early life stress interactions in adolescent alcohol drinking. *Neuropsychopharmacology.* 2011; 36(6):1142–1148. DOI: 10.1038/npp.2010.247 [PubMed: 21307845]
- Contet C, Kim A, Le D, Iyengar SK, Kotzebue RW, Yuan CJ, et al. Mandyam CD. mu-Opioid receptors mediate the effects of chronic ethanol binge drinking on the hippocampal neurogenic niche. *Addict Biol.* 2014; 19(5):770–780. DOI: 10.1111/adb.12040 [PubMed: 23461397]
- Correa M, Viaggi C, Escrig MA, Pascual M, Guerri C, Vaglini F, et al. Corsini GU. Ethanol intake and ethanol-induced locomotion and locomotor sensitization in Cyp2e1 knockout mice. *Pharmacogenet Genomics.* 2009; 19(3):217–225. DOI: 10.1097/FPC.0b013e328324e726 [PubMed: 19177030]
- Cowen MS, Schroff KC, Gass P, Sprengel R, Spanagel R. Neurobehavioral effects of alcohol in AMPA receptor subunit (GluR1) deficient mice. *Neuropharmacology.* 2003; 45(3):325–333. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12871650>. [PubMed: 12871650]
- Cowen MS, Schumann G, Yagi T, Spanagel R. Role of Fyn tyrosine kinase in ethanol consumption by mice. *Alcohol Clin Exp Res.* 2003; 27(8):1213–1219. DOI: 10.1097/01.ALC.0000081630.14159.02 [PubMed: 12966312]
- Crabbe JC, Harkness JH, Spence SE, Huang LC, Metten P. Intermittent availability of ethanol does not always lead to elevated drinking in mice. *Alcohol Alcohol.* 2012; 47(5):509–517. DOI: 10.1093/alcalc/ags067 [PubMed: 22717273]
- Crabbe JC, Phillips TJ, Feller DJ, Hen R, Wenger CD, Lessov CN, Schafer GL. Elevated alcohol consumption in null mutant mice lacking 5-HT1B serotonin receptors. *Nat Genet.* 1996; 14(1):98–101. DOI: 10.1038/ng0996-98 [PubMed: 8782828]
- Crabbe JC, Phillips TJ, Harris RA, Arends MA, Koob GF. Alcohol-related genes: contributions from studies with genetically engineered mice. *Addict Biol.* 2006; 11(3-4):195–269. DOI: 10.1111/j.1369-1600.2006.00038.x [PubMed: 16961758]
- Crews FT, Vetreno RP. Mechanisms of neuroimmune gene induction in alcoholism. *Psychopharmacology (Berl).* 2015; doi: 10.1007/s00213-015-3906-1
- D'Souza MS. Glutamatergic transmission in drug reward: implications for drug addiction. *Front Neurosci.* 2015; 9:404.doi: 10.3389/fnins.2015.00404 [PubMed: 26594139]
- Dawson A, Miles MF, Damaj MI. The beta2 nicotinic acetylcholine receptor subunit differentially influences ethanol behavioral effects in the mouse. *Alcohol.* 2013; 47(2):85–94. DOI: 10.1016/j.alcohol.2012.12.004 [PubMed: 23419392]
- Delis F, Thanos PK, Rombola C, Rosko L, Grandy D, Wang GJ, Volkow ND. Chronic mild stress increases alcohol intake in mice with low dopamine D2 receptor levels. *Behav Neurosci.* 2013; 127(1):95–105. DOI: 10.1037/a0030750 [PubMed: 23148856]
- Dong L, Bilbao A, Laucht M, Henriksson R, Yakovleva T, Ridinger M, et al. Schumann G. Effects of the circadian rhythm gene period 1 (per1) on psychosocial stress-induced alcohol drinking. *Am J Psychiatry.* 2011; 168(10):1090–1098. DOI: 10.1176/appi.ajp.2011.10111579 [PubMed: 21828288]
- Duncan EA, Sorrell JE, Adamantidis A, Rider T, Jandacek RJ, Seeley RJ, et al. Woods SC. Alcohol drinking in MCH receptor-1-deficient mice. *Alcohol Clin Exp Res.* 2007; 31(8):1325–1337. DOI: 10.1111/j.1530-0277.2007.00427.x [PubMed: 17550369]

- El-Ghundi M, George SR, Drago J, Fletcher PJ, Fan T, Nguyen T, et al. O'Dowd BF. Disruption of dopamine D1 receptor gene expression attenuates alcohol-seeking behavior. *Eur J Pharmacol.* 1998; 353(2-3):149–158. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9726645>. [PubMed: 9726645]
- Engel SR, Lyons CR, Allan AM. 5-HT3 receptor over-expression decreases ethanol self administration in transgenic mice. *Psychopharmacology (Berl).* 1998; 140(2):243–248. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/9860116>. [PubMed: 9860116]
- Enoch MA. The role of GABA(A) receptors in the development of alcoholism. *Pharmacol Biochem Behav.* 2008; 90(1):95–104. DOI: 10.1016/j.pbb.2008.03.007 [PubMed: 18440057]
- Faber F, Gembardt F, Sun X, Mizutani S, Siems WE, Walther T. Lack of angiotensin II conversion to angiotensin III increases water but not alcohol consumption in aminopeptidase A-deficient mice. *Regul Pept.* 2006; 136(1-3):130–137. DOI: 10.1016/j.regpep.2006.06.001 [PubMed: 16889841]
- Falzone TL, Gelman DM, Young JJ, Grandy DK, Low MJ, Rubinstein M. Absence of dopamine D4 receptors results in enhanced reactivity to unconditioned, but not conditioned, fear. *Eur J Neurosci.* 2002; 15(1):158–164. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11860516>. [PubMed: 11860516]
- Farris SP, Arasappan D, Hunicke-Smith S, Harris RA, Mayfield RD. Transcriptome organization for chronic alcohol abuse in human brain. *Mol Psychiatry.* 2015; 20(11):1438–1447. DOI: 10.1038/mp.2014.159 [PubMed: 25450227]
- Farris SP, Pietrzykowski AZ, Miles MF, O'Brien MA, Sanna PP, Zakhari S, et al. Harris RA. Applying the new genomics to alcohol dependence. *Alcohol.* 2015; doi: 10.1016/j.alcohol.2015.03.001
- Fee JR, Sparta DR, Knapp DJ, Breese GR, Picker MJ, Thiele TE. Predictors of high ethanol consumption in RIIbeta knock-out mice: assessment of anxiety and ethanol-induced sedation. *Alcohol Clin Exp Res.* 2004; 28(10):1459–1468. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15597077>. [PubMed: 15597077]
- Femenia T, Manzanares J. Increased ethanol intake in prodynorphin knockout mice is associated to changes in opioid receptor function and dopamine transmission. *Addict Biol.* 2012; 17(2):322–337. DOI: 10.1111/j.1369-1600.2011.00378.x [PubMed: 21966993]
- Ferguson LB, Most D, Blednov YA, Harris RA. PPAR agonists regulate brain gene expression: relationship to their effects on ethanol consumption. *Neuropharmacology.* 2014; 86:397–407. DOI: 10.1016/j.neuropharm.2014.06.024 [PubMed: 25036611]
- Fernandez E, Koek W, Ran Q, Gerhardt GA, France CP, Strong R. Monoamine metabolism and behavioral responses to ethanol in mitochondrial aldehyde dehydrogenase knockout mice. *Alcohol Clin Exp Res.* 2006; 30(10):1650–1658. DOI: 10.1111/j.1530-0277.2006.00200.x [PubMed: 17010132]
- Ferraro FM 3rd, Sparta DR, Knapp DJ, Breese GR, Thiele TE. Increased consumption but not operant self-administration of ethanol in mice lacking the RIIbeta subunit of protein kinase A. *Alcohol Clin Exp Res.* 2006; 30(5):825–835. DOI: 10.1111/j.1530-0277.2006.00096.x [PubMed: 16634851]
- Finn DA, Belknap JK, Cronise K, Yoneyama N, Murillo A, Crabbe JC. A procedure to produce high alcohol intake in mice. *Psychopharmacology (Berl).* 2005; 178(4):471–480. DOI: 10.1007/s00213-004-2039-8 [PubMed: 15765261]
- Gallego X, Ruiz-Medina J, Valverde O, Molas S, Robles N, Sabria J, et al. Dierssen M. Transgenic over expression of nicotinic receptor alpha 5, alpha 3, and beta 4 subunit genes reduces ethanol intake in mice. *Alcohol.* 2012; 46(3):205–215. DOI: 10.1016/j.alcohol.2011.11.005 [PubMed: 22459873]
- Geisler S, Schopf CL, Obermair GJ. Emerging evidence for specific neuronal functions of auxiliary calcium channel alpha(2)delta subunits. *Gen Physiol Biophys.* 2015; 34(2):105–118. DOI: 10.4149/gpb_2014037 [PubMed: 25504062]
- Giardino WJ, Cocking DL, Kaur S, Cunningham CL, Ryabinin AE. Urocortin-1 within the centrally-projecting Edinger-Westphal nucleus is critical for ethanol preference. *PLoS One.* 2011; 6(10):e26997.doi: 10.1371/journal.pone.0026997 [PubMed: 22046429]

- Giardino WJ, Ryabinin AE. CRF1 receptor signaling regulates food and fluid intake in the drinking-in-the-dark model of binge alcohol consumption. *Alcohol Clin Exp Res*. 2013; 37(7):1161–1170. DOI: 10.1111/acer.12076 [PubMed: 23398267]
- Gorini G, Bell RL, Mayfield RD. Molecular targets of alcohol action: Translational research for pharmacotherapy development and screening. *Prog Mol Biol Transl Sci*. 2011; 98:293–347. DOI: 10.1016/B978-0-12-385506-0.00007-7 [PubMed: 21199775]
- Gorini G, Harris RA, Mayfield RD. Proteomic approaches and identification of novel therapeutic targets for alcoholism. *Neuropsychopharmacology*. 2014; 39(1):104–130. DOI: 10.1038/npp.2013.182 [PubMed: 23900301]
- Gorini G, Roberts AJ, Mayfield RD. Neurobiological signatures of alcohol dependence revealed by protein profiling. *PLoS One*. 2013; 8(12):e82656.doi: 10.1371/journal.pone.0082656 [PubMed: 24358215]
- Grahame NJ, Mosemiller AK, Low MJ, Froehlich JC. Naltrexone and alcohol drinking in mice lacking beta-endorphin by site-directed mutagenesis. *Pharmacol Biochem Behav*. 2000; 67(4):759–766. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11166066>. [PubMed: 11166066]
- Green AS, Grahame NJ. Ethanol drinking in rodents: is free-choice drinking related to the reinforcing effects of ethanol? *Alcohol*. 2008; 42(1):1–11. DOI: 10.1016/j.alcohol.2007.10.005 [PubMed: 18164576]
- Grisel JE, Mogil JS, Grahame NJ, Rubinstein M, Belknap JK, Crabbe JC, Low MJ. Ethanol oral self-administration is increased in mutant mice with decreased beta-endorphin expression. *Brain Res*. 1999; 835(1):62–67. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10448196>. [PubMed: 10448196]
- Hall FS, Sora I, Uhl GR. Ethanol consumption and reward are decreased in mu-opiate receptor knockout mice. *Psychopharmacology (Berl)*. 2001; 154(1):43–49. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11292005>. [PubMed: 11292005]
- Hall FS, Sora I, Uhl GR. Sex-dependent modulation of ethanol consumption in vesicular monoamine transporter 2 (VMAT2) and dopamine transporter (DAT) knockout mice. *Neuropsychopharmacology*. 2003; 28(4):620–628. DOI: 10.1038/sj.npp.1300070 [PubMed: 12655306]
- Han S, Gelernter J, Kranzler HR, Yang BZ. Ordered subset linkage analysis based on admixture proportion identifies new linkage evidence for alcohol dependence in African-Americans. *Hum Genet*. 2013; 132(4):397–403. DOI: 10.1007/s00439-012-1255-2 [PubMed: 23239122]
- Harris RA, Blednov YA. Neuroimmune genes and alcohol drinking behavior. In: Cui, C, Grandison, L., Noronha, A., editors. *Neural-Immune Interactions in Brain Function and Alcohol Related Disorders*. New York: Springer; 2013. p. 425–440.
- Harris RA, Osterndorff-Kahanek E, Ponomarev I, Homanics GE, Blednov YA. Testing the silence of mutations: Transcriptomic and behavioral studies of GABA(A) receptor alpha1 and alpha2 subunit knock-in mice. *Neurosci Lett*. 2011; 488(1):31–35. DOI: 10.1016/j.neulet.2010.10.075 [PubMed: 21056629]
- Hendrickson LM, Zhao-Shea R, Pang X, Gardner PD, Tapper AR. Activation of alpha4* nAChRs is necessary and sufficient for varenicline-induced reduction of alcohol consumption. *J Neurosci*. 2010; 30(30):10169–10176. DOI: 10.1523/JNEUROSCI.2601-10.2010 [PubMed: 20668200]
- Herman MA, Sidhu H, Stouffer DG, Kreifeldt M, Le D, Cates-Gatto C, et al. Contet C. GIRK3 gates activation of the mesolimbic dopaminergic pathway by ethanol. *Proc Natl Acad Sci U S A*. 2015; 112(22):7091–7096. DOI: 10.1073/pnas.1416146112 [PubMed: 25964320]
- Hilakivi-Clarke L, Goldberg R. Gonadal hormones and aggression-maintaining effect of alcohol in male transgenic transforming growth factor-alpha mice. *Alcohol Clin Exp Res*. 1995; 19(3):708–713. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/7573797>. [PubMed: 7573797]
- Hodge CW, Mehmert KK, Kelley SP, McMahon T, Haywood A, Olive MF, et al. Messing RO. Supersensitivity to allosteric GABA(A) receptor modulators and alcohol in mice lacking PKCepsilon. *Nat Neurosci*. 1999; 2(11):997–1002. DOI: 10.1038/14795 [PubMed: 10526339]
- Houchi H, Persyn W, Legastelois R, Naassila M. The adenosine A2A receptor agonist CGS 21680 decreases ethanol self-administration in both non-dependent and dependent animals. *Addict Biol*. 2013; 18(5):812–825. DOI: 10.1111/adb.12032 [PubMed: 23301633]

- Houchi H, Warnault V, Barbier E, Dubois C, Pierrefiche O, Ledent C, et al. Naassila M. Involvement of A2A receptors in anxiolytic, locomotor and motivational properties of ethanol in mice. *Genes Brain Behav.* 2008; 7(8):887–898. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/19097273>. [PubMed: 19097273]
- Howard RJ, Trudell JR, Harris RA. Seeking structural specificity: direct modulation of pentameric ligand-gated ion channels by alcohols and general anesthetics. *Pharmacol Rev.* 2014; 66(2):396–412. DOI: 10.1124/pr.113.007468 [PubMed: 24515646]
- Hungund BL, Szakall I, Adam A, Basavarajappa BS, Vadasz C. Cannabinoid CB1 receptor knockout mice exhibit markedly reduced voluntary alcohol consumption and lack alcohol-induced dopamine release in the nucleus accumbens. *J Neurochem.* 2003; 84(4):698–704. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12562514>. [PubMed: 12562514]
- Hwa LS, Chu A, Levinson SA, Kayyali TM, DeBold JF, Miczek KA. Persistent escalation of alcohol drinking in C57BL/6J mice with intermittent access to 20% ethanol. *Alcohol Clin Exp Res.* 2011; 35(11):1938–1947. DOI: 10.1111/j.1530-0277.2011.01545.x [PubMed: 21631540]
- Iancu OD, Oberbeck D, Darakjian P, Metten P, McWeeney S, Crabbe JC, Hitzemann R. Selection for drinking in the dark alters brain gene coexpression networks. *Alcohol Clin Exp Res.* 2013; 37(8):1295–1303. DOI: 10.1111/acer.12100 [PubMed: 23550792]
- Imai S, Kano M, Nonoyama K, Ebihara S. Behavioral characteristics of ubiquitin-specific peptidase 46-deficient mice. *PLoS One.* 2013; 8(3):e58566.doi: 10.1371/journal.pone.0058566 [PubMed: 23472206]
- Isse T, Oyama T, Kitagawa K, Matsuno K, Matsumoto A, Yoshida A, et al. Kawamoto T. Diminished alcohol preference in transgenic mice lacking aldehyde dehydrogenase activity. *Pharmacogenetics.* 2002; 12(8):621–626. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12439222>. [PubMed: 12439222]
- Jaholkowski P, Mierzejewski P, Zatorski P, Scinska A, Sienkiewicz-Jarosz H, Kaczmarek L, et al. Bienkowski P. Increased ethanol intake and preference in cyclin D2 knockout mice. *Genes Brain Behav.* 2011; 10(5):551–556. DOI: 10.1111/j.1601-183X.2011.00692.x [PubMed: 21429093]
- Jerlhag E, Landgren S, Egecioglu E, Dickson SL, Engel JA. The alcohol-induced locomotor stimulation and accumbal dopamine release is suppressed in ghrelin knockout mice. *Alcohol.* 2011; 45(4):341–347. DOI: 10.1016/j.alcohol.2010.10.002 [PubMed: 21145690]
- June HL Sr, Foster KL, Eiler WJ 2nd, Goergen J, Cook JB, Johnson N, et al. Homanics GE. Dopamine and benzodiazepine-dependent mechanisms regulate the EtOH-enhanced locomotor stimulation in the GABAA alpha1 subunit null mutant mice. *Neuropsychopharmacology.* 2007; 32(1):137–152. DOI: 10.1038/sj.npp.1301097 [PubMed: 16710315]
- Kamens HM, Andersen J, Picciotto MR. Modulation of ethanol consumption by genetic and pharmacological manipulation of nicotinic acetylcholine receptors in mice. *Psychopharmacology (Berl).* 2010; 208(4):613–626. DOI: 10.1007/s00213-009-1759-1 [PubMed: 20072781]
- Kamens HM, Hoft NR, Cox RJ, Miyamoto JH, Ehringer MA. The alpha6 nicotinic acetylcholine receptor subunit influences ethanol-induced sedation. *Alcohol.* 2012; 46(5):463–471. DOI: 10.1016/j.alcohol.2012.03.001 [PubMed: 22572056]
- Kampov-Polevoy A, Lange L, Bobashev G, Eggleston B, Root T, Garbutt JC. Sweet-liking is associated with transformation of heavy drinking into alcohol-related problems in young adults with high novelty seeking. *Alcohol Clin Exp Res.* 2014; 38(7):2119–2126. DOI: 10.1111/acer.12458 [PubMed: 24962796]
- Kang SJ, Rangaswamy M, Manz N, Wang JC, Wetherill L, Hinrichs T, et al. Porjesz B. Family-based genome-wide association study of frontal theta oscillations identifies potassium channel gene KCNJ6. *Genes Brain Behav.* 2012; 11(6):712–719. DOI: 10.1111/j.1601-183X.2012.00803.x [PubMed: 22554406]
- Kapfhamer D, Bettinger JC, Davies AG, Eastman CL, Smail EA, Heberlein U, McIntire SL. Loss of RAB-3/A in *Caenorhabditis elegans* and the mouse affects behavioral response to ethanol. *Genes Brain Behav.* 2008; 7(6):669–676. DOI: 10.1111/j.1601-183X.2008.00404.x [PubMed: 18397381]
- Kapfhamer D, Taylor S, Zou ME, Lim JP, Kharazia V, Heberlein U. Taok2 controls behavioral response to ethanol in mice. *Genes Brain Behav.* 2013; 12(1):87–97. DOI: 10.1111/j.1601-183X.2012.00834.x [PubMed: 22883308]

- Karatayev O, Baylan J, Leibowitz SF. Increased intake of ethanol and dietary fat in galanin overexpressing mice. *Alcohol*. 2009; 43(8):571–580. DOI: 10.1016/j.alcohol.2009.09.025 [PubMed: 20004335]
- Karatayev O, Baylan J, Weed V, Chang S, Wynick D, Leibowitz SF. Galanin knockout mice show disturbances in ethanol consumption and expression of hypothalamic peptides that stimulate ethanol intake. *Alcohol Clin Exp Res*. 2010; 34(1):72–80. DOI: 10.1111/j.1530-0277.2009.01068.x [PubMed: 19860804]
- Karlsson RM, Adermark L, Molander A, Perreau-Lenz S, Singley E, Solomon M, et al. Heilig M. Reduced alcohol intake and reward associated with impaired endocannabinoid signaling in mice with a deletion of the glutamate transporter GLAST. *Neuropharmacology*. 2012; 63(2):181–189. DOI: 10.1016/j.neuropharm.2012.01.027 [PubMed: 22342743]
- Kaur S, Li J, Stenzel-Poore MP, Ryabinin AE. Corticotropin-releasing factor acting on corticotropin-releasing factor receptor type 1 is critical for binge alcohol drinking in mice. *Alcohol Clin Exp Res*. 2012; 36(2):369–376. DOI: 10.1111/j.1530-0277.2011.01610.x [PubMed: 21895713]
- Kelai S, Aissi F, Lesch KP, Cohen-Salmon C, Hamon M, Lanfumey L. Alcohol intake after serotonin transporter inactivation in mice. *Alcohol Alcohol*. 2003; 38(4):386–389. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12814910>. [PubMed: 12814910]
- Kim KS, Kim H, Baek IS, Lee KW, Han PL. Mice lacking adenylyl cyclase type 5 (AC5) show increased ethanol consumption and reduced ethanol sensitivity. *Psychopharmacology (Berl)*. 2011; 215(2):391–398. DOI: 10.1007/s00213-010-2143-x [PubMed: 21193983]
- Koenig HN, Olive MF. Ethanol consumption patterns and conditioned place preference in mice lacking preproenkephalin. *Neurosci Lett*. 2002; 325(2):75–78. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12044625>. [PubMed: 12044625]
- Korkosz A, Kolomanska P, Kowalska K, Rogowski A, Radwanska K, Kaczmarek L, et al. Bienkowski P. Dissociation of ethanol and saccharin preference in fosB knockout mice. *Physiol Behav*. 2004; 82(2-3):391–395. DOI: 10.1016/j.physbeh.2004.04.054 [PubMed: 15276803]
- Kovacs KM, Szakall I, O'Brien D, Wang R, Vinod KY, Saito M, et al. Vadasz C. Decreased oral self-administration of alcohol in kappa-opioid receptor knock-out mice. *Alcohol Clin Exp Res*. 2005; 29(5):730–738. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15897716>. [PubMed: 15897716]
- Kreifeldt M, Le D, Treistman SN, Koob GF, Contet C. BK channel beta1 and beta4 auxiliary subunits exert opposite influences on escalated ethanol drinking in dependent mice. *Front Integr Neurosci*. 2013; 7:105doi: 10.3389/fnint.2013.00105
- Lallemant F, de Witte P. Ethanol induces higher BEC in CB1 cannabinoid receptor knockout mice while decreasing ethanol preference. *Alcohol Alcohol*. 2005; 40(1):54–62. DOI: 10.1093/alcalc/agh115 [PubMed: 15550449]
- Lamb RJ, Daws LC. Ethanol self-administration in serotonin transporter knockout mice: unconstrained demand and elasticity. *Genes Brain Behav*. 2013; 12(7):741–747. DOI: 10.1111/gbb.12068 [PubMed: 23927813]
- Lasek AW, Lim J, Kliethermes CL, Berger KH, Joslyn G, Brush G, et al. Heberlein U. An evolutionary conserved role for anaplastic lymphoma kinase in behavioral responses to ethanol. *PLoS One*. 2011; 6(7):e22636.doi: 10.1371/journal.pone.0022636 [PubMed: 21799923]
- Leclercq S, De Saeger C, Delzenne N, de Timary P, Starkel P. Role of inflammatory pathways, blood mononuclear cells, and gut-derived bacterial products in alcohol dependence. *Biol Psychiatry*. 2014; 76(9):725–733. DOI: 10.1016/j.biopsych.2014.02.003 [PubMed: 24629538]
- Lee AM, Zou ME, Lim JP, Stecher J, McMahon T, Messing RO. Deletion of *Prkcz* increases intermittent ethanol consumption in mice. *Alcohol Clin Exp Res*. 2014; 38(1):170–178. DOI: 10.1111/acer.12211 [PubMed: 23905844]
- Lee MR, Hinton DJ, Song JY, Lee KW, Choo C, Johng H, et al. Choi DS. Neurotensin receptor type 1 regulates ethanol intoxication and consumption in mice. *Pharmacol Biochem Behav*. 2010; 95(2):235–241. DOI: 10.1016/j.pbb.2010.01.012 [PubMed: 20122953]
- Lee MR, Hinton DJ, Unal SS, Richelson E, Choi DS. Increased ethanol consumption and preference in mice lacking neurotensin receptor type 2. *Alcohol Clin Exp Res*. 2011; 35(1):99–107. DOI: 10.1111/j.1530-0277.2010.01326.x [PubMed: 21039631]

- Lee MR, Ruby CL, Hinton DJ, Choi S, Adams CA, Young Kang N, Choi DS. Striatal adenosine signaling regulates EAAT2 and astrocytic AQP4 expression and alcohol drinking in mice. *Neuropsychopharmacology*. 2013; 38(3):437–445. DOI: 10.1038/npp.2012.198 [PubMed: 23032072]
- Legastelois R, Darcq E, Wegner SA, Lombroso PJ, Ron D. Striatal-enriched protein tyrosine phosphatase controls responses to aversive stimuli: implication for ethanol drinking. *PLoS One*. 2015; 10(5):e0127408.doi: 10.1371/journal.pone.0127408 [PubMed: 25992601]
- Leggio GM, Camillieri G, Platania CB, Castorina A, Marrazzo G, Torrisi SA, et al. Salomone S. Dopamine D3 receptor is necessary for ethanol consumption: an approach with buspirone. *Neuropsychopharmacology*. 2014; 39(8):2017–2028. DOI: 10.1038/npp.2014.51 [PubMed: 24584330]
- Lesscher HM, Wallace MJ, Zeng L, Wang V, Deitchman JK, McMahon T, et al. Newton PM. Amygdala protein kinase C epsilon controls alcohol consumption. *Genes Brain Behav*. 2009; 8(5):493–499. DOI: 10.1111/j.1601-183X.2009.00485.x [PubMed: 19243450]
- Levrin O, Yuferov V, Kreek MJ. The genetics of the opioid system and specific drug addictions. *Hum Genet*. 2012; 131(6):823–842. DOI: 10.1007/s00439-012-1172-4 [PubMed: 22547174]
- Li D, Sulovari A, Cheng C, Zhao H, Kranzler HR, Gelernter J. Association of gamma-aminobutyric acid A receptor alpha2 gene (GABRA2) with alcohol use disorder. *Neuropsychopharmacology*. 2014; 39(4):907–918. DOI: 10.1038/npp.2013.291 [PubMed: 24136292]
- Li H, Tao Y, Ma L, Liu X, Ma L. beta-Arrestin-2 inhibits preference for alcohol in mice and suppresses Akt signaling in the dorsal striatum. *Neurosci Bull*. 2013; 29(5):531–540. DOI: 10.1007/s12264-013-1350-y [PubMed: 23839051]
- Liu J, Lewohl JM, Harris RA, Iyer VR, Dodd PR, Randall PK, Mayfield RD. Patterns of gene expression in the frontal cortex discriminate alcoholic from nonalcoholic individuals. *Neuropsychopharmacology*. 2006; 31(7):1574–1582. DOI: 10.1038/sj.npp.1300947 [PubMed: 16292326]
- Loney KD, Uddin RK, Singh SM. Analysis of metallothionein brain gene expression in relation to ethanol preference in mice using cosegregation and gene knockouts. *Alcohol Clin Exp Res*. 2006; 30(1):15–25. DOI: 10.1111/j.1530-0277.2006.00015.x [PubMed: 16433728]
- Lopez-Jimenez A, Walter NA, Gine E, Santos A, Echeverry-Alzate V, Buhler KM, et al. Lopez-Moreno JA. A spontaneous deletion of alpha-synuclein is associated with an increase in CB1 mRNA transcript and receptor expression in the hippocampus and amygdala: effects on alcohol consumption. *Synapse*. 2013; 67(6):280–289. DOI: 10.1002/syn.21639 [PubMed: 23345080]
- Lum EN, Campbell RR, Rostock C, Szumlinski KK. mGluR1 within the nucleus accumbens regulates alcohol intake in mice under limited-access conditions. *Neuropharmacology*. 2014; 79:679–687. DOI: 10.1016/j.neuropharm.2014.01.024 [PubMed: 24467847]
- Lynch LJ, Sullivan KA, Vallender EJ, Rowlett JK, Platt DM, Miller GM. Trace amine associated receptor 1 modulates behavioral effects of ethanol. *Subst Abuse*. 2013; 7:117–126. DOI: 10.4137/SART.S12110 [PubMed: 23861588]
- Maas JW Jr, Vogt SK, Chan GC, Pineda VV, Storm DR, Muglia LJ. Calcium-stimulated adenylyl cyclases are critical modulators of neuronal ethanol sensitivity. *J Neurosci*. 2005; 25(16):4118–4126. DOI: 10.1523/JNEUROSCI.4273-04.2005 [PubMed: 15843614]
- Martin GE, Hendrickson LM, Penta KL, Friesen RM, Pietrzykowski AZ, Tapper AR, Treistman SN. Identification of a BK channel auxiliary protein controlling molecular and behavioral tolerance to alcohol. *Proc Natl Acad Sci U S A*. 2008; 105(45):17543–17548. DOI: 10.1073/pnas.0801068105 [PubMed: 18981408]
- Mason BJ, Quello S, Goodell V, Shadan F, Kyle M, Begovic A. Gabapentin treatment for alcohol dependence: a randomized clinical trial. *JAMA Intern Med*. 2014; 174(1):70–77. DOI: 10.1001/jamainternmed.2013.11950 [PubMed: 24190578]
- Maul B, Becker M, Gembardt F, Becker A, Schultheiss HP, Siems WE, Walther T. Genetic deficiency in neprilysin or its pharmacological inhibition initiate excessive stress-induced alcohol consumption in mice. *PLoS One*. 2012; 7(11):e50187.doi: 10.1371/journal.pone.0050187 [PubMed: 23185571]

- Maul B, Krause W, Pankow K, Becker M, Gembardt F, Alenina N, et al. Siems WE. Central angiotensin II controls alcohol consumption via its AT1 receptor. *FASEB J.* 2005; 19(11):1474–1481. DOI: 10.1096/fj.05-3742com [PubMed: 16126915]
- Maul B, Siems WE, Hoehe MR, Grecksch G, Bader M, Walther T. Alcohol consumption is controlled by angiotensin II. *FASEB J.* 2001; 15(9)(1640) Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11427512>.
- Mayfield J, Blednov YA, Harris RA. Behavioral and Genetic Evidence for GIRK Channels in the CNS: Role in Physiology, Pathophysiology, and Drug Addiction. *Int Rev Neurobiol.* 2015; 123:279–313. DOI: 10.1016/bs.irm.2015.05.016 [PubMed: 26422988]
- Mayfield J, Ferguson L, Harris RA. Neuroimmune signaling: a key component of alcohol abuse. *Curr Opin Neurobiol.* 2013; 23(4):513–520. DOI: 10.1016/j.conb.2013.01.024 [PubMed: 23434064]
- Melendez RI. Intermittent (every-other-day) drinking induces rapid escalation of ethanol intake and preference in adolescent and adult C57BL/6J mice. *Alcohol Clin Exp Res.* 2011; 35(4):652–658. DOI: 10.1111/j.1530-0277.2010.01383.x [PubMed: 21223302]
- Metz AV, Chynoweth J, Allan AM. Influence of genetic background on alcohol drinking and behavioral phenotypes of 5-HT3 receptor over-expressing mice. *Pharmacol Biochem Behav.* 2006; 84(1):120–127. DOI: 10.1016/j.pbb.2006.04.016 [PubMed: 16765427]
- Mihalek RM, Bowers BJ, Wehner JM, Kralic JE, VanDoren MJ, Morrow AL, Homanics GE. GABA(A)-receptor delta subunit knockout mice have multiple defects in behavioral responses to ethanol. *Alcohol Clin Exp Res.* 2001; 25(12):1708–1718. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11781502>. [PubMed: 11781502]
- Mittleman G, Call SB, Cockroft JL, Goldowitz D, Matthews DB, Blaha CD. Dopamine dynamics associated with, and resulting from, schedule-induced alcohol self-administration: analyses in dopamine transporter knockout mice. *Alcohol.* 2011; 45(4):325–339. DOI: 10.1016/j.alcohol.2010.12.006 [PubMed: 21354763]
- Miyasaka K, Hosoya H, Takano S, Ohta M, Sekime A, Kanai S, et al. Funakoshi A. Differences in ethanol ingestion between cholecystokinin-A receptor deficient and -B receptor deficient mice. *Alcohol Alcohol.* 2005; 40(3):176–180. DOI: 10.1093/alcalc/agh143 [PubMed: 15767271]
- Molander A, Vengeliene V, Heilig M, Wurst W, Deussing JM, Spanagel R. Brain-specific inactivation of the Crhr1 gene inhibits post-dependent and stress-induced alcohol intake, but does not affect relapse-like drinking. *Neuropsychopharmacology.* 2012; 37(4):1047–1056. DOI: 10.1038/npp.2011.297 [PubMed: 22113086]
- Moore R, Krstew EV, Kirchhoff J, Davisson RL, Lawrence AJ. Central overexpression of angiotensin AT(1A) receptors prevents dopamine D(2) receptor regulation of alcohol consumption in mice. *Alcohol Clin Exp Res.* 2007; 31(7):1128–1137. DOI: 10.1111/j.1530-0277.2007.00399.x [PubMed: 17451402]
- Mulligan MK, Ponomarev I, Hitzemann RJ, Belknap JK, Tabakoff B, Harris RA, et al. Bergeson SE. Toward understanding the genetics of alcohol drinking through transcriptome meta-analysis. *Proc Natl Acad Sci U S A.* 2006; 103(16):6368–6373. DOI: 10.1073/pnas.0510188103 [PubMed: 16618939]
- Mutschler J, Bilbao A, von der Goltz C, Demiralay C, Jahn H, Wiedemann K, et al. Kiefer F. Augmented stress-induced alcohol drinking and withdrawal in mice lacking functional natriuretic peptide-A receptors. *Alcohol Alcohol.* 2010; 45(1):13–16. DOI: 10.1093/alcalc/agg065 [PubMed: 19828462]
- Naassila M, Ledent C, Daoust M. Low ethanol sensitivity and increased ethanol consumption in mice lacking adenosine A2A receptors. *J Neurosci.* 2002; 22(23):10487–10493. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12451148>. [PubMed: 12451148]
- Naassila M, Pierrefiche O, Ledent C, Daoust M. Decreased alcohol self-administration and increased alcohol sensitivity and withdrawal in CB1 receptor knockout mice. *Neuropharmacology.* 2004; 46(2):243–253. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14680762>. [PubMed: 14680762]
- Nam HW, Bruner RC, Choi DS. Adenosine signaling in striatal circuits and alcohol use disorders. *Mol Cells.* 2013; 36(3):195–202. DOI: 10.1007/s10059-013-0192-9 [PubMed: 23912595]

- Nam HW, Hinton DJ, Kang NY, Kim T, Lee MR, Oliveros A, et al. Choi DS. Adenosine transporter ENT1 regulates the acquisition of goal-directed behavior and ethanol drinking through A2A receptor in the dorsomedial striatum. *J Neurosci*. 2013; 33(10):4329–4338. DOI: 10.1523/JNEUROSCI.3094-12.2013 [PubMed: 23467349]
- Nam HW, Lee MR, Zhu Y, Wu J, Hinton DJ, Choi S, et al. Choi DS. Type 1 equilibrative nucleoside transporter regulates ethanol drinking through accumbal N-methyl-D-aspartate receptor signaling. *Biol Psychiatry*. 2011; 69(11):1043–1051. DOI: 10.1016/j.biopsych.2011.02.013 [PubMed: 21489406]
- Navarro M, Cubero I, Chen AS, Chen HY, Knapp DJ, Breese GR, et al. Thiele TE. Effects of melanocortin receptor activation and blockade on ethanol intake: a possible role for the melanocortin-4 receptor. *Alcohol Clin Exp Res*. 2005; 29(86):949–957. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/15976520>. [PubMed: 15976520]
- Navarro M, Cubero I, Ko L, Thiele TE. Deletion of agouti-related protein blunts ethanol self-administration and binge-like drinking in mice. *Genes Brain Behav*. 2009; 8(4):450–458. DOI: 10.1111/j.1601-183X.2009.00493.x [PubMed: 19566712]
- Navarro M, Lerma-Cabrera JM, Carvajal F, Lowery EG, Cubero I, Thiele TE. Assessment of voluntary ethanol consumption and the effects of a melanocortin (MC) receptor agonist on ethanol intake in mutant C57BL/6J mice lacking the MC-4 receptor. *Alcohol Clin Exp Res*. 2011; 35(6):1058–1066. DOI: 10.1111/j.1530-0277.2011.01438.x [PubMed: 21332528]
- Newton PM, Orr CJ, Wallace MJ, Kim C, Shin HS, Messing RO. Deletion of N-type calcium channels alters ethanol reward and reduces ethanol consumption in mice. *J Neurosci*. 2004; 24(44):9862–9869. DOI: 10.1523/JNEUROSCI.3446-04.2004 [PubMed: 15525770]
- Nie H, Rewal M, Gill TM, Ron D, Janak PH. Extrasynaptic delta-containing GABAA receptors in the nucleus accumbens dorsomedial shell contribute to alcohol intake. *Proc Natl Acad Sci U S A*. 2011; 108(11):4459–4464. DOI: 10.1073/pnas.1016156108 [PubMed: 21368141]
- Nunez YO, Truitt JM, Gorini G, Ponomareva ON, Blednov YA, Harris RA, Mayfield RD. Positively correlated miRNA-mRNA regulatory networks in mouse frontal cortex during early stages of alcohol dependence. *BMC Genomics*. 2013; 14725doi: 10.1186/1471-2164-14-725
- Nuutinen S, Lintunen M, Vanhanen J, Ojala T, Rozov S, Panula P. Evidence for the role of histamine H3 receptor in alcohol consumption and alcohol reward in mice. *Neuropsychopharmacology*. 2011; 36(10):2030–2040. DOI: 10.1038/npp.2011.90 [PubMed: 21654737]
- Offenhauser N, Castelletti D, Mapelli L, Soppo BE, Regondi MC, Rossi P, et al. Di Fiore PP. Increased ethanol resistance and consumption in Eps8 knockout mice correlates with altered actin dynamics. *Cell*. 2006; 127(1):213–226. DOI: 10.1016/j.cell.2006.09.011 [PubMed: 17018287]
- Olive MF, McGeehan AJ, Kinder JR, McMahon T, Hodge CW, Janak PH, Messing RO. The mGluR5 antagonist 6-methyl-2-(phenylethynyl)pyridine decreases ethanol consumption via a protein kinase C epsilon-dependent mechanism. *Mol Pharmacol*. 2005; 67(2):349–355. DOI: 10.1124/mol.104.003319 [PubMed: 15548766]
- Olive MF, Mehmert KK, Koenig HN, Camarini R, Kim JA, Nannini MA, et al. Hodge CW. A role for corticotropin releasing factor (CRF) in ethanol consumption, sensitivity, and reward as revealed by CRF-deficient mice. *Psychopharmacology (Berl)*. 2003; 165(2):181–187. DOI: 10.1007/s00213-002-1248-2 [PubMed: 12397512]
- Olive MF, Mehmert KK, Messing RO, Hodge CW. Reduced operant ethanol self-administration and in vivo mesolimbic dopamine responses to ethanol in PKCepsilon-deficient mice. *Eur J Neurosci*. 2000; 12(11):4131–4140. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11069609>. [PubMed: 11069609]
- Olney JJ, Sprow GM, Navarro M, Thiele TE. The protective effects of the melanocortin receptor (MCR) agonist, melanotan-II (MTII), against binge-like ethanol drinking are facilitated by deletion of the MC3 receptor in mice. *Neuropeptides*. 2014; 48(1):47–51. DOI: 10.1016/j.npep.2013.11.001 [PubMed: 24290566]
- Ortega-Alvaro A, Ternianov A, Aracil-Fernandez A, Navarrete F, Garcia-Gutierrez MS, Manzanares J. Role of cannabinoid CB2 receptor in the reinforcing actions of ethanol. *Addict Biol*. 2015; 20(1):43–55. DOI: 10.1111/adb.12076 [PubMed: 23855434]
- Osterndorff-Kahanek E, Ponomarev I, Blednov YA, Harris RA. Gene expression in brain and liver produced by three different regimens of alcohol consumption in mice: comparison with immune

activation. PLoS One. 2013; 8(3):e59870.doi: 10.1371/journal.pone.0059870 [PubMed: 23555817]

- Palmer AA, Low MJ, Grandy DK, Phillips TJ. Effects of a Drd2 deletion mutation on ethanol-induced locomotor stimulation and sensitization suggest a role for epistasis. Behav Genet. 2003; 33(3): 311–324. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12837020>. [PubMed: 12837020]
- Panagis G, Mackey B, Vlachou S. Cannabinoid Regulation of Brain Reward Processing with an Emphasis on the Role of CB1 Receptors: A Step Back into the Future. Front Psychiatry. 2014; 592doi: 10.3389/fpsy.2014.00092
- Pascual M, Balino P, Alfonso-Loeches S, Aragon CM, Guerri C. Impact of TLR4 on behavioral and cognitive dysfunctions associated with alcohol-induced neuroinflammatory damage. Brain Behav Immun. 2011; 25 Suppl 1:S80–91. DOI: 10.1016/j.bbi.2011.02.012 [PubMed: 21352907]
- Pastor R, Reed C, Burkhart-Kasch S, Li N, Sharpe AL, Coste SC, et al. Phillips TJ. Ethanol concentration-dependent effects and the role of stress on ethanol drinking in corticotropin-releasing factor type 1 and double type 1 and 2 receptor knockout mice. Psychopharmacology (Berl). 2011; 218(1):169–177. DOI: 10.1007/s00213-011-2284-6 [PubMed: 21487655]
- Perkins DI, Trudell JR, Crawford DK, Alkana RL, Davies DL. Molecular targets and mechanisms for ethanol action in glycine receptors. Pharmacol Ther. 2010; 127(1):53–65. DOI: 10.1016/j.pharmthera.2010.03.003 [PubMed: 20399807]
- Phillips TJ, Brown KJ, Burkhart-Kasch S, Wenger CD, Kelly MA, Rubinstein M, et al. Low MJ. Alcohol preference and sensitivity are markedly reduced in mice lacking dopamine D2 receptors. Nat Neurosci. 1998; 1(7):610–615. DOI: 10.1038/2843 [PubMed: 10196569]
- Poncelet M, Maruani J, Calassi R, Soubrie P. Overeating, alcohol and sucrose consumption decrease in CB1 receptor deleted mice. Neurosci Lett. 2003; 343(3):216–218. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12770700>. [PubMed: 12770700]
- Ponomarev I, Wang S, Zhang L, Harris RA, Mayfield RD. Gene coexpression networks in human brain identify epigenetic modifications in alcohol dependence. J Neurosci. 2012; 32(5):1884–1897. DOI: 10.1523/JNEUROSCI.3136-11.2012 [PubMed: 22302827]
- Popova NK, Vishnivetskaya GB, Ivanova EA, Skrinkaya JA, Seif I. Altered behavior and alcohol tolerance in transgenic mice lacking MAO A: a comparison with effects of MAO A inhibitor clorgyline. Pharmacol Biochem Behav. 2000; 67(4):719–727. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11166062>. [PubMed: 11166062]
- Quintanilla ME, Israel Y, Sapag A, Tampier L. The UChA and UChB rat lines: metabolic and genetic differences influencing ethanol intake. Addict Biol. 2006; 11(3-4):310–323. DOI: 10.1111/j.1369-1600.2006.00030.x [PubMed: 16961761]
- Racz I, Bilkei-Gorzo A, Toth ZE, Michel K, Palkovits M, Zimmer A. A critical role for the cannabinoid CB1 receptors in alcohol dependence and stress-stimulated ethanol drinking. J Neurosci. 2003; 23(6):2453–2458. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12657705>. [PubMed: 12657705]
- Racz I, Markert A, Mauer D, Stoffel-Wagner B, Zimmer A. Long-term ethanol effects on acute stress responses: modulation by dynorphin. Addict Biol. 2013; 18(4):678–688. DOI: 10.1111/j.1369-1600.2012.00494.x [PubMed: 22994904]
- Racz I, Schurmann B, Karpushova A, Reuter M, Cichon S, Montag C, et al. Zimmer A. The opioid peptides enkephalin and beta-endorphin in alcohol dependence. Biol Psychiatry. 2008; 64(11): 989–997. DOI: 10.1016/j.biopsych.2008.05.008 [PubMed: 18589403]
- Rahman S, Engleman EA, Bell RL. Nicotinic receptor modulation to treat alcohol and drug dependence. Front Neurosci. 2014; 8426doi: 10.3389/fnins.2014.00426
- Rao PS, Bell RL, Engleman EA, Sari Y. Targeting glutamate uptake to treat alcohol use disorders. Front Neurosci. 2015; 9144doi: 10.3389/fnins.2015.00144
- Repunte-Canonigo V, Berton F, Cottone P, Reifel-Miller A, Roberts AJ, Morales M, et al. Sanna PP. A potential role for adiponectin receptor 2 (AdipoR2) in the regulation of alcohol intake. Brain Res. 2010; 1339:11–17. DOI: 10.1016/j.brainres.2010.03.060 [PubMed: 20380822]
- Rewal M, Jurd R, Gill TM, He DY, Ron D, Janak PH. Alpha4-containing GABAA receptors in the nucleus accumbens mediate moderate intake of alcohol. J Neurosci. 2009; 29(2):543–549. DOI: 10.1523/JNEUROSCI.3199-08.2009 [PubMed: 19144854]

- Risinger FO, Doan AM, Vickrey AC. Oral operant ethanol self-administration in 5-HT1b knockout mice. *Behav Brain Res.* 1999; 102(1-2):211–215. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10403028>. [PubMed: 10403028]
- Risinger FO, Freeman PA, Greengard P, Fienberg AA. Motivational effects of ethanol in DARPP-32 knock-out mice. *J Neurosci.* 2001; 21(1):340–348. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11150351>. [PubMed: 11150351]
- Risinger FO, Freeman PA, Rubinstein M, Low MJ, Grandy DK. Lack of operant ethanol self-administration in dopamine D2 receptor knockout mice. *Psychopharmacology (Berl).* 2000; 152(3):343–350. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11105945>. [PubMed: 11105945]
- Roberts AJ, Gold LH, Polis I, McDonald JS, Filliol D, Kieffer BL, Koob GF. Increased ethanol self-administration in delta-opioid receptor knockout mice. *Alcohol Clin Exp Res.* 2001; 25(9):1249–1256. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11584142>. [PubMed: 11584142]
- Roberts AJ, McDonald JS, Heyser CJ, Kieffer BL, Matthes HW, Koob GF, Gold LH. mu-Opioid receptor knockout mice do not self-administer alcohol. *J Pharmacol Exp Ther.* 2000; 293(3):1002–1008. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10869404>. [PubMed: 10869404]
- Robinson G, Most D, Ferguson LB, Mayfield J, Harris RA, Blednov YA. Neuroimmune pathways in alcohol consumption: evidence from behavioral and genetic studies in rodents and humans. *Int Rev Neurobiol.* 2014; 118:13–39. DOI: 10.1016/B978-0-12-801284-0.00002-6 [PubMed: 25175860]
- Rodd ZA, Bell RL, Sable HJ, Murphy JM, McBride WJ. Recent advances in animal models of alcohol craving and relapse. *Pharmacol Biochem Behav.* 2004; 79(3):439–450. DOI: 10.1016/j.pbb.2004.08.018 [PubMed: 15582015]
- Rosenwasser AM, Fixaris MC, Crabbe JC, Brooks PC, Ascheid S. Escalation of intake under intermittent ethanol access in diverse mouse genotypes. *Addict Biol.* 2013; 18(3):496–507. DOI: 10.1111/j.1369-1600.2012.00481.x [PubMed: 22862671]
- Ruby CL, Vадnie CA, Hinton DJ, Abulseoud OA, Walker DL, O'Connor KM, et al. Choi DS. Adenosinergic regulation of striatal clock gene expression and ethanol intake during constant light. *Neuropsychopharmacology.* 2014; 39(10):2432–2440. DOI: 10.1038/npp.2014.94 [PubMed: 24755889]
- Ruby CL, Walker DL, An J, Kim J, Choi DS. Sex-Specific Regulation of Depression, Anxiety-Like Behaviors and Alcohol Drinking in Mice Lacking ENT1. *J Addict Res Ther.* 2011; S4doi: 10.4172/2155-6105.S4-004
- Saba LM, Bennett B, Hoffman PL, Barcomb K, Ishii T, Kechris K, Tabakoff B. A systems genetic analysis of alcohol drinking by mice, rats and men: influence of brain GABAergic transmission. *Neuropharmacology.* 2011; 60(7-8):1269–1280. DOI: 10.1016/j.neuropharm.2010.12.019 [PubMed: 21185315]
- Saba LM, Flink SC, Vanderlinden LA, Israel Y, Tampier L, Colombo G, et al. Tabakoff B. The sequenced rat brain transcriptome--its use in identifying networks predisposing alcohol consumption. *FEBS J.* 2015; 282(18):3556–3578. DOI: 10.1111/febs.13358 [PubMed: 26183165]
- Sachs BD, Salahi AA, Caron MG. Congenital brain serotonin deficiency leads to reduced ethanol sensitivity and increased ethanol consumption in mice. *Neuropharmacology.* 2014; 77:177–184. DOI: 10.1016/j.neuropharm.2013.09.010 [PubMed: 24067926]
- Sakoori K, Murphy NP. Endogenous nociceptin (orphanin FQ) suppresses basal hedonic state and acute reward responses to methamphetamine and ethanol, but facilitates chronic responses. *Neuropsychopharmacology.* 2008; 33(4):877–891. DOI: 10.1038/sj.npp.1301459 [PubMed: 17522627]
- Salinas AG, Nguyen CT, Ahmadi-Tehrani D, Morrisett RA. Reduced ethanol consumption and preference in cocaine- and amphetamine-regulated transcript (CART) knockout mice. *Addict Biol.* 2014; 19(2):175–184. DOI: 10.1111/j.1369-1600.2012.00475.x [PubMed: 22823101]
- Sanbe A, Takagi N, Fujiwara Y, Yamauchi J, Endo T, Mizutani R, et al. Tanoue A. Alcohol preference in mice lacking the Avpr1a vasopressin receptor. *Am J Physiol Regul Integr Comp Physiol.* 2008; 294(5):R1482–1490. DOI: 10.1152/ajpregu.00708.2007 [PubMed: 18305023]

- Sanchis-Segura C, Borchardt T, Vengeliene V, Zghoul T, Bachteler D, Gass P, et al. Spanagel R. Involvement of the AMPA receptor GluR-C subunit in alcohol-seeking behavior and relapse. *J Neurosci*. 2006; 26(4):1231–1238. DOI: 10.1523/JNEUROSCI.4237-05.2006 [PubMed: 16436610]
- Santos N, Chatterjee S, Henry A, Holgate J, Bartlett SE. The alpha5 neuronal nicotinic acetylcholine receptor subunit plays an important role in the sedative effects of ethanol but does not modulate consumption in mice. *Alcohol Clin Exp Res*. 2013; 37(4):655–662. DOI: 10.1111/acer.12009 [PubMed: 23164049]
- Savarese A, Zou ME, Kharazia V, Maiya R, Lasek AW. Increased behavioral responses to ethanol in Lmo3 knockout mice. *Genes Brain Behav*. 2014; 13(8):777–783. DOI: 10.1111/gbb.12176 [PubMed: 25176312]
- Savelieva KV, Caudle WM, Findlay GS, Caron MG, Miller GW. Decreased ethanol preference and consumption in dopamine transporter female knock-out mice. *Alcohol Clin Exp Res*. 2002; 26(6):758–764. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12068242>. [PubMed: 12068242]
- Schank JR, Ryabinin AE, Giardino WJ, Ciccocioppo R, Heilig M. Stress-related neuropeptides and addictive behaviors: beyond the usual suspects. *Neuron*. 2012; 76(1):192–208. DOI: 10.1016/j.neuron.2012.09.026 [PubMed: 23040815]
- Sharpe AL, Coste SC, Burkhardt-Kasch S, Li N, Stenzel-Poore MP, Phillips TJ. Mice deficient in corticotropin-releasing factor receptor type 2 exhibit normal ethanol-associated behaviors. *Alcohol Clin Exp Res*. 2005; 29(9):1601–1609. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/16205360>. [PubMed: 16205360]
- Siems W, Maul B, Krause W, Gerard C, Hauser KF, Hersch LB, et al. Saria A. Neutral endopeptidase and alcohol consumption, experiments in neutral endopeptidase-deficient mice. *Eur J Pharmacol*. 2000; 397(2-3):327–334. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10844131>. [PubMed: 10844131]
- Sikela JM, Maclaren EJ, Kim Y, Karimpour-Fard A, Cai WW, Pollack J, et al. Miles MF. DNA microarray and proteomic strategies for understanding alcohol action. *Alcohol Clin Exp Res*. 2006; 30(4):700–708. DOI: 10.1111/j.1530-0277.2006.00081.x [PubMed: 16573589]
- Sillaber I, Rammes G, Zimmermann S, Mahal B, Zieglerberger W, Wurst W, et al. Spanagel R. Enhanced and delayed stress-induced alcohol drinking in mice lacking functional CRH1 receptors. *Science*. 2002; 296(5569):931–933. DOI: 10.1126/science.1069836 [PubMed: 11988580]
- Spanagel R, Pendyala G, Abarca C, Zghoul T, Sanchis-Segura C, Magnone MC, et al. Albrecht U. The clock gene *Per2* influences the glutamatergic system and modulates alcohol consumption. *Nat Med*. 2005; 11(1):35–42. DOI: 10.1038/nm1163 [PubMed: 15608650]
- Spanagel R, Siegmund S, Cowen M, Schrott KC, Schumann G, Fiserova M, et al. Putzke J. The neuronal nitric oxide synthase gene is critically involved in neurobehavioral effects of alcohol. *J Neurosci*. 2002; 22(19):8676–8683. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12351742>. [PubMed: 12351742]
- Specia DJ, Chihara D, Ashique AM, Bowers MS, Pierce-Shimomura JT, Lee J, et al. McIntire SL. Conserved role of *unc-79* in ethanol responses in lightweight mutant mice. *PLoS Genet*. 2010; 6(8):doi: 10.1371/journal.pgen.1001057
- Sperling RE, Gomes SM, Sypek EI, Carey AN, McLaughlin JP. Endogenous kappa-opioid mediation of stress-induced potentiation of ethanol-conditioned place preference and self-administration. *Psychopharmacology (Berl)*. 2010; 210(2):199–209. DOI: 10.1007/s00213-010-1844-5 [PubMed: 20401606]
- Stacey D, Bilbao A, Maroteaux M, Jia T, Easton AC, Longueville S, et al. Consortium, Imagen. RASGRF2 regulates alcohol-induced reinforcement by influencing mesolimbic dopamine neuron activity and dopamine release. *Proc Natl Acad Sci U S A*. 2012; 109(51):21128–21133. DOI: 10.1073/pnas.1211844110 [PubMed: 23223532]
- Steffensen SC, Bradley KD, Hansen DM, Wilcox JD, Wilcox RS, Allison DW, et al. Edwards JG. The role of connexin-36 gap junctions in alcohol intoxication and consumption. *Synapse*. 2011; 65(8):695–707. DOI: 10.1002/syn.20885 [PubMed: 21638336]

- Stephens DN, Pistovcakova J, Worthing L, Atack JR, Dawson GR. Role of GABAA alpha5-containing receptors in ethanol reward: the effects of targeted gene deletion, and a selective inverse agonist. *Eur J Pharmacol.* 2005; 526(1-3):240–250. DOI: 10.1016/j.ejphar.2005.09.031 [PubMed: 16253225]
- Stewart A, Maity B, Andereg SP, Allamargot C, Yang J, Fisher RA. Regulator of G protein signaling 6 is a critical mediator of both reward-related behavioral and pathological responses to alcohol. *Proc Natl Acad Sci U S A.* 2015; 112(7):E786–795. DOI: 10.1073/pnas.1418795112 [PubMed: 25646431]
- Stork O, Kojima N, Stork S, Kume N, Obata K. Resistance to alcohol withdrawal-induced behaviour in Fyn transgenic mice and its reversal by ifenprodil. *Brain Res Mol Brain Res.* 2002; 105(1-2): 126–135. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/12399115>. [PubMed: 12399115]
- Szumlini KK, Ary AW, Lominac KD, Klugmann M, Kippin TE. Accumbens Homer2 overexpression facilitates alcohol-induced neuroplasticity in C57BL/6J mice. *Neuropsychopharmacology.* 2008; 33(6):1365–1378. DOI: 10.1038/sj.npp.1301473 [PubMed: 17568396]
- Szumlini KK, Lominac KD, Oleson EB, Walker JK, Mason A, Dehoff MH, et al. Kalivas PW. Homer2 is necessary for EtOH-induced neuroplasticity. *J Neurosci.* 2005; 25(30):7054–7061. DOI: 10.1523/JNEUROSCI.1529-05.2005 [PubMed: 16049182]
- Szumlini KK, Toda S, Middaugh LD, Worley PF, Kalivas PW. Evidence for a relationship between Group 1 mGluR hypofunction and increased cocaine and ethanol sensitivity in Homer2 null mutant mice. *Ann N Y Acad Sci.* 2003; 1003:468–471. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/14684491>. [PubMed: 14684491]
- Tammimaki A, Forsberg MM, Karayiorgou M, Gogos JA, Mannisto PT. Increase in free choice oral ethanol self-administration in catechol-o-methyltransferase gene-disrupted male mice. *Basic Clin Pharmacol Toxicol.* 2008; 103(4):297–304. DOI: 10.1111/j.1742-7843.2008.00267.x [PubMed: 18684228]
- Tanaka K, Kunishige-Yamamoto A, Hashimoto H, Shintani N, Hayata A, Baba A. Increased ethanol preference and serotonin 1A receptor-dependent attenuation of ethanol-induced hypothermia in PACAP-deficient mice. *Biochem Biophys Res Commun.* 2010; 391(1):773–777. DOI: 10.1016/j.bbrc.2009.11.136 [PubMed: 19944672]
- Thanos PK, Dimitrakakis ES, Rice O, Gifford A, Volkow ND. Ethanol self-administration and ethanol conditioned place preference are reduced in mice lacking cannabinoid CB1 receptors. *Behav Brain Res.* 2005; 164(2):206–213. DOI: 10.1016/j.bbr.2005.06.021 [PubMed: 16140402]
- Thanos PK, Rivera SN, Weaver K, Grandy DK, Rubinstein M, Umegaki H, et al. Volkow ND. Dopamine D2R DNA transfer in dopamine D2 receptor-deficient mice: effects on ethanol drinking. *Life Sci.* 2005; 77(2):130–139. DOI: 10.1016/j.lfs.2004.10.061 [PubMed: 15862598]
- Thiele TE, Koh MT, Pedrazzini T. Voluntary alcohol consumption is controlled via the neuropeptide Y Y1 receptor. *J Neurosci.* 2002; 22(3):RC208. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11826154>. [PubMed: 11826154]
- Thiele TE, Marsh DJ, Ste Marie L, Bernstein IL, Palmiter RD. Ethanol consumption and resistance are inversely related to neuropeptide Y levels. *Nature.* 1998; 396(6709):366–369. DOI: 10.1038/24614 [PubMed: 9845072]
- Thiele TE, Miura GI, Marsh DJ, Bernstein IL, Palmiter RD. Neurobiological responses to ethanol in mutant mice lacking neuropeptide Y or the Y5 receptor. *Pharmacol Biochem Behav.* 2000; 67(4): 683–691. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/11166058>. [PubMed: 11166058]
- Thiele TE, Navarro M. “Drinking in the dark” (DID) procedures: a model of binge-like ethanol drinking in non-dependent mice. *Alcohol.* 2014; 48(3):235–241. DOI: 10.1016/j.alcohol.2013.08.005 [PubMed: 24275142]
- Thiele TE, Naveilhan P, Ernfor P. Assessment of ethanol consumption and water drinking by NPY Y(2) receptor knockout mice. *Peptides.* 2004; 25(6):975–983. DOI: 10.1016/j.peptides.2004.03.009 [PubMed: 15203244]
- Thiele TE, Willis B, Stadler J, Reynolds JG, Bernstein IL, McKnight GS. High ethanol consumption and low sensitivity to ethanol-induced sedation in protein kinase A-mutant mice. *J Neurosci.* 2000; 20(10):RC75. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10783399>. [PubMed: 10783399]

- Thorsell A, Schank JR, Singley E, Hunt SP, Heilig M. Neurokinin-1 receptors (NK1R:s), alcohol consumption, and alcohol reward in mice. *Psychopharmacology (Berl)*. 2010; 209(1):103–111. DOI: 10.1007/s00213-010-1775-1 [PubMed: 20112009]
- Trudell JR, Messing RO, Mayfield J, Harris RA. Alcohol dependence: molecular and behavioral evidence. *Trends Pharmacol Sci*. 2014; 35(7):317–323. DOI: 10.1016/j.tips.2014.04.009 [PubMed: 24865944]
- van den Brink W, Sorensen P, Torup L, Mann K, Gual A. Group, Sense Study. Long-term efficacy, tolerability and safety of nalmefene as-needed in patients with alcohol dependence: A 1-year, randomised controlled study. *J Psychopharmacol*. 2014; 28(8):733–744. DOI: 10.1177/0269881114527362 [PubMed: 24671340]
- van Huijstee AN, Mansvelder HD. Glutamatergic synaptic plasticity in the mesocorticolimbic system in addiction. *Front Cell Neurosci*. 2014; 8466doi: 10.3389/fncel.2014.00466
- van Rijn RM, Whistler JL. The delta(1) opioid receptor is a heterodimer that opposes the actions of the delta(2) receptor on alcohol intake. *Biol Psychiatry*. 2009; 66(8):777–784. DOI: 10.1016/j.biopsych.2009.05.019 [PubMed: 19576572]
- Vanderlinden LA, Saba LM, Bennett B, Hoffman PL, Tabakoff B. Influence of sex on genetic regulation of drinking in the dark alcohol consumption. *Mamm Genome*. 2015; 26(1-2):43–56. DOI: 10.1007/s00335-014-9553-8 [PubMed: 25559016]
- Vanhanen J, Nuutinen S, Lintunen M, Maki T, Ramo J, Karlstedt K, Panula P. Histamine is required for H(3) receptor-mediated alcohol reward inhibition, but not for alcohol consumption or stimulation. *Br J Pharmacol*. 2013; 170(1):177–187. DOI: 10.1111/bph.12170 [PubMed: 23489295]
- Vengeliene V, Bilbao A, Spanagel R. The alcohol deprivation effect model for studying relapse behavior: a comparison between rats and mice. *Alcohol*. 2014; 48(3):313–320. DOI: 10.1016/j.alcohol.2014.03.002 [PubMed: 24811155]
- Vinod KY, Sanguino E, Yalamanchili R, Manzanares J, Hungund BL. Manipulation of fatty acid amide hydrolase functional activity alters sensitivity and dependence to ethanol. *J Neurochem*. 2008; 104(1):233–243. DOI: 10.1111/j.1471-4159.2007.04956.x [PubMed: 17944864]
- Vinod KY, Yalamanchili R, Thanos PK, Vadasz C, Cooper TB, Volkow ND, Hungund BL. Genetic and pharmacological manipulations of the CB(1) receptor alter ethanol preference and dependence in ethanol preferring and nonpreferring mice. *Synapse*. 2008; 62(8):574–581. DOI: 10.1002/syn.20533 [PubMed: 18509854]
- Walker AW, Smith CM, Chua BE, Krstew EV, Zhang C, Gundlach AL, Lawrence AJ. Relaxin-3 receptor (RXFP3) signalling mediates stress-related alcohol preference in mice. *PLoS One*. 2015; 10(4):e0122504.doi: 10.1371/journal.pone.0122504 [PubMed: 25849482]
- Wallace MJ, Newton PM, Oyasu M, McMahon T, Chou WH, Connolly J, Messing RO. Acute functional tolerance to ethanol mediated by protein kinase Cepsilon. *Neuropsychopharmacology*. 2007; 32(1):127–136. DOI: 10.1038/sj.npp.1301059 [PubMed: 16541084]
- Weinshenker D, Rust NC, Miller NS, Palmiter RD. Ethanol-associated behaviors of mice lacking norepinephrine. *J Neurosci*. 2000; 20(9):3157–3164. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/10777779>. [PubMed: 10777779]
- Werner C, Raivich G, Cowen M, Strekalova T, Sillaber I, Buters JT, et al. Hofmann F. Importance of NO/cGMP signalling via cGMP-dependent protein kinase II for controlling emotionality and neurobehavioural effects of alcohol. *Eur J Neurosci*. 2004; 20(12):3498–3506. DOI: 10.1111/j.1460-9568.2004.03793.x [PubMed: 15610182]
- Werner DF, Blednov YA, Ariwodola OJ, Silberman Y, Logan E, Berry RB, et al. Homanics GE. Knockin mice with ethanol-insensitive alpha1-containing gamma-aminobutyric acid type A receptors display selective alterations in behavioral responses to ethanol. *J Pharmacol Exp Ther*. 2006; 319(1):219–227. DOI: 10.1124/jpet.106.106161 [PubMed: 16785315]
- Wyatt LR, Finn DA, Khoja S, Yardley MM, Asatryan L, Alkana RL, Davies DL. Contribution of P2X4 receptors to ethanol intake in male C57BL/6 mice. *Neurochem Res*. 2014; 39(6):1127–1139. DOI: 10.1007/s11064-014-1271-9 [PubMed: 24671605]
- Yaka R, Tang KC, Camarini R, Janak PH, Ron D. Fyn kinase and NR2B-containing NMDA receptors regulate acute ethanol sensitivity but not ethanol intake or conditioned reward. *Alcohol Clin Exp*

Res. 2003; 27(11):1736–1742. DOI: 10.1097/01.ALC.0000095924.87729.D8 [PubMed: 14634488]

Zghoul T, Abarca C, Sanchis-Segura C, Albrecht U, Schumann G, Spanagel R. Ethanol self-administration and reinstatement of ethanol-seeking behavior in Per1(Brdm1) mutant mice. *Psychopharmacology (Berl)*. 2007; 190(1):13–19. DOI: 10.1007/s00213-006-0592-z [PubMed: 17051414]

Zhou Z, Karlsson C, Liang T, Xiong W, Kimura M, Tapocik JD, et al. Goldman D. Loss of metabotropic glutamate receptor 2 escalates alcohol consumption. *Proc Natl Acad Sci U S A*. 2013; 110(42):16963–16968. DOI: 10.1073/pnas.1309839110 [PubMed: 24082084]

Zindel LR, Kranzler HR. Pharmacotherapy of alcohol use disorders: seventy-five years of progress. *J Stud Alcohol Drugs Suppl*. 2014; 75 Suppl 17:79–88. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/24565314>. [PubMed: 24565314]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Alcohol consumption in GABA mutant mice

GeneKnockout/Knockin [#]	Background	Operant	2BC	DID	SHAC	References
γ -aminobutyric acid type A receptor (GABA _A R) subunit α -1 (<i>Gabra1</i>)	Not specified	↓ (30-60 min), males/females	↓ females	↓ (2 h) 1B, post-operant	—	(June et al., 2007)
<i>Gabra1</i> [#]	B6 × 129SvEv	—	—	—	—	(Blednov et al., 2003)
GABA _A R subunit α -2 (<i>Gabra2</i>)	B6 × 129SvEv	— males — females	—	—	—	(D. F. Werner et al., 2006)
GABA _A R subunit α -5 (<i>Gabra5</i>)	B6 × 129SvEv	↓ males — females	—	—	—	(Boehm, Ponomarev, et al., 2004)
	B6 × 129SvEv	— (1 h) females	—	—	—	(Boehm, Ponomarev, et al., 2004)
	B6 × 129SvEv	—	—	—	—	(Stephens et al., 2005)
GABA _A R subunit β -2 (<i>Gabrb2</i>)	B6 × 129SvEv	—	—	—	—	(Blednov et al., 2003)
GABA _A R subunit δ (<i>Gabrd</i>)	B6 × 129/Sv/SvJ	↓ males/females	↓ males/females	—	—	(Mihalek et al., 2001)
GABA _A R subunit ρ -1 (<i>Gabbr1</i>)	B6 × 129S4	↓ males — females — intermittent, males/females	↓ males — females — intermittent, males/females	— (2, 4 h), males/ females	—	(Blednov et al., 2014)
Sodium- and chloride-dependent GABA transporter 1 (<i>Slc6a1</i>)	B6	—	—	—	—	(Cai et al., 2006)
Glutamic acid decarboxylase 2 (<i>Gad2</i>)	B6 B6 × 129/SvJ N1 B6 × 129/SvJ N2	— — ↑	— — ↑	— (3 h) 2BC — (2, 4 h) 1B	— (30 min)	(Blednov et al., 2010)

[#] indicates gene knockin;

—, ↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in mutant vs. wild-type mice. Male mice were tested unless indicated otherwise. Ethanol intake in the two-bottle choice (2BC) tests was measured in 24-h sessions. Drinking session times for the other tests are indicated in parenthesis. DID, drinking in the dark; 1B, one bottle; SHAC, scheduled high alcohol concentration. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 2

Alcohol consumption in glutamate mutant mice

Gene Knockout/Overexpression *	Background	Operant	2BC	4BC	DID	SHAC	References
Glutamate receptor 1 (<i>Gria1</i>)	B6N	—	— ADE — post-stress				(Cowen, Schroff, et al., 2003)
Glutamate receptor 3 (<i>Gria3</i>)	B6N	—	— ↓ ADE, day 1				(Sanchis-Segura et al., 2006)
Glutamate receptor ionotropic, NMDA 2A (<i>Gria2a</i>)	B6	—	—				(Boyce-Rustay & Holmes, 2006)
Metabotropic glutamate receptor 2, mGluR2 (<i>Grm2</i>)	CD1	↑ intake	↑ intake				(Zhou et al., 2013)
mGluR4 (<i>Grm4</i>)	CD1 × 129/SvJ	—	— males/females				(Blednov, Walker, Osterndorf-Kahanek, et al., 2004)
mGluR5 (<i>Grm5</i>)	B6 × 129/SvJ	—	— males/females	↓ females	— (2 h) 1B, females ↓ (3 h) 2BC, females	— (30 min), females	(Blednov & Harris, 2008)
	B6		↓				(Bird et al., 2008)
Excitatory amino acid transporter 1, GLAST, EAAT1 (<i>Slc1a3</i>)	B6		↓ males/females				(Karlsson et al., 2012)
Homer protein homolog 2 (<i>Homer2</i>)	Not specified		↓				(Szumlinski et al., 2003)
	B6 × 129Xi/SvJ		↓ 12%, males/females				(Szumlinski et al., 2005)
	B6 × 129Xi/SvJ				— (2 h)		(Lum et al., 2014)
<i>Homer2b</i> *	B6	↑ (21 min), NAc					(Szumlinski et al., 2008)
Neuronal pentraxin-2, NARP (<i>Nptx2</i>)	129SV × B6		↓ intake, no escalation				(Ary et al., 2012)

* indicates gene overexpression;

—, ↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in mutant vs. wild-type mice. Males were tested unless otherwise indicated. Ethanol intake in the two- and four-bottle choice (2BC, 4BC) tests was measured in continuous 24-h sessions. Drinking session times for the other tests are indicated in parenthesis. DID, drinking in the dark; SHAC, scheduled high alcohol consumption; 1B, one bottle; ADE, alcohol deprivation effect. NAc, nucleus accumbens. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Alcohol consumption in dopamine mutant mice

Table 3

Gene Knockout/Overexpression *	Background	Operant	2BC	DID	SIP	References
D _{1A} dopamine receptor (<i>Drd1</i>)	B6	↓ limited/continuous access	↓			(El-Ghundi et al., 1998)
D ₂ dopamine receptor (<i>Drd2</i>)	B6	↓ (23 h)				(Risinger et al., 2000)
	B6	↓ males/females — ethanol-exposed				(Palmer et al., 2003)
	B6	↓				(Thanos, Rivera, et al., 2005)
	B6	↓ males/females				(Phillips et al., 1998)
	B6	↓ ↑ chronic stress				(Delis et al., 2013)
D ₂ long receptor	B6			↑ (4 h), males/ females		(Bulwa et al., 2011)
<i>Drd2</i> *	B6	↓ WT ↑ transient KO				(Thanos, Rivera, et al., 2005)
D ₃ dopamine receptor (<i>Drd3</i>)	B6	— (23 h)	—			(Boyce-Rustay & Risinger, 2003)
	B6	↓		↓ (4 h)		(Leggio et al., 2014)
D ₄ dopamine receptor (<i>Drd4</i>)	B6129/Ola × B6	—				(Falzone et al., 2002)
Sodium-dependent dopamine transporter, DAT (<i>Slc6a5</i>)	B6	— males ↓ females				(Savelieva, Caudle, Findlay, Caron, & Miller, 2002)
	B6				↓ (40 min), males/ females	(Mittleman et al., 2011)
	B6 × 129Sv	↑ 24%, males — females				(Hall, Sora, & Uhl, 2003)
Dopamine β-hydroxylase (<i>Dhdt</i>)	B6 × 129/SvEv	↓				(Weinschenker, Rust, Miller, & Palmiter, 2000)

* indicates gene overexpression;

—, ↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in knockout (KO) or mice overexpressing D2* vs. wild-type (WT)/control mice. Male mice were tested unless indicated otherwise. Deletion of the long form of D2 produces overexpression of the short relative to the long form. Ethanol intake in the two-bottle choice (2BC) tests was measured in 24-h sessions, unless otherwise indicated. Drinking session times for the other tests are indicated in parenthesis. DID, drinking in the dark; SIP, scheduled-induced polydipsia. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 4
Alcohol consumption in serotonin mutant mice

Gene Knockout/Knockin [#] /Overexpression [*]	Background	Operant	2BC	References
Tryptophan 5-hydroxylase 2 (<i>Tph2</i>) [#]	B6 × 129S6/SvEvTac		↑	(Sachs et al., 2014)
5-hydroxytryptamine receptor (5-HTR), 5-HTR _{1B} (<i>Htr1b</i>)	Not specified	↑ (23 h) 10% phase 1, males/females		(Risinger et al., 1999)
	129/Sv-Ter		↑ males/females	(Crabbe et al., 1996)
	129/Sv		— males	(Bouwknicht et al., 2000)
5-HTR _{3A} (<i>Htr3a</i>) [*]	B6DBA/2J		↓ —	(Metz et al., 2006)
5-HTR ₆ (<i>Htr6</i>)	B6		—	(Bonasera et al., 2006)
Sodium-dependent serotonin transporter (<i>Slc6a4</i>)	B6	↓ (90 min)		(Lamb & Daws, 2013)
	B6		↓	(Kelai et al., 2003)
	B6		↓ intake 7%, females	(Boyce-Rustay et al., 2006)

^{*} indicates gene overexpression;

—, ↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in mutant vs. wild-type mice. Male mice were tested unless indicated otherwise. Ethanol intake in the two-bottle choice (2BC) tests was measured in 24-h sessions. Drinking session times for the operant tests are indicated in parenthesis. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 5
Alcohol consumption in adenosine knockout mice

Gene Knockout	Background	Operant	2BC	References
Adenosine receptor A1 (<i>Adora1</i>)	B6 × 129	— (2 h)		(Houchi et al., 2013)
Adenosine receptor A2A (<i>Adora2a</i>)	CD1		↑ males/females	(Naassila et al., 2002)
	CD1B6		↑ males — males/females	(Houchi et al., 2008)
Equilibrative nucleoside transporter-1, ENT1 (<i>Slc29a1</i>)	B6 × 129X1/SvJ	↑		(J. Chen et al., 2010)
	B6 × 129X1/SvJ		↑	(Nam, Hinton, et al., 2013; Nam et al., 2011)
	B6 × 129X1/SvJ		↑	(Choi et al., 2004)
	B6 × 129X1/SvJ		↑ sex not specified	(M. R. Lee et al., 2013; Ruby et al., 2014)
	B6 × 129X1/SvJ		↑ females	(Ruby et al., 2011)

—, ↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in knockout *vs.* wild-type mice. Male mice were tested unless indicated otherwise. Ethanol intake in the operant and two-bottle choice (2BC) tests was measured in 30-min or 24-h sessions, respectively, unless otherwise indicated. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 6

Alcohol consumption in cannabinoid and opioid knockout mice

Gene Knockout	Background	Operant	2BC	DD	References
Cannabinoid receptor 1 (<i>Cnr1</i>)	B6 × 129/Ola		↓ (6 h)		(Poncellet et al., 2003)
	B6 × 129/Ola		↓ preference		(Lallemant & de Witte, 2005)
	B6		↓ males/females		(Hungund et al., 2003)
	CD1		↓ males/females		(Naassila et al., 2004)
	CD1			↓ (8 h)	(Thanos, Dimitrakakis, et al., 2005)
	B6 and DBA/2J		↓		(Vinod, Yalamanchili, et al., 2008)
	B6		— after 1 week		(Racz et al., 2003)
Cannabinoid receptor 2 (<i>Cnr2</i>)	CD1 and B6	↑ (1 h)	↑		(Ortega-Alvaro et al., 2015)
Fatty acid amide hydrolase (<i>Faah</i>)	B6 × 129/SvJ		↑ males/females		(Blednov et al., 2007)
	B6		↑ females — males		(Basavarajappa et al., 2006)
	B6		↑		(Vinod, Sanguino, et al., 2008)
μ-type opioid receptor (<i>Oprm1</i>)	B6 × 129/Sv	↓ nose-poke ↓ lever operant ↓ post-operant ↓ post-forced ethanol	↓ post-operant ↓ post-operant ↓ post-forced ethanol		(Roberts et al., 2000)
	B6 × 129/Sv		↓ females — males		(Hall et al., 2001)
	Not specified		↓ after several weeks		(Becker et al., 2002)
	B6			↓ (2 h) — (2h) ADE	(Contet et al., 2014)
	B6			— (4 h)	(van Rijn & Whistler, 2009)
δ-type opioid receptor (<i>Oprd1</i>)	B6			↑ (4 h)	(van Rijn & Whistler, 2009)
	B6Orl × 129/Sv	↑ post-2BC	— naive ↑ post-operant		(Roberts et al., 2001)
κ-type opioid receptor (<i>Oprk1</i>)	B6			↓ (4 h)	(van Rijn & Whistler, 2009)
	B6Orl		↓ males/females		(Kovacs et al., 2005)
β-endorphin (<i>Pomc</i>)	B6		↑ 7%, males/females — 10%, males/females		(Grisel et al., 1999)
	B6		— ↑ 2 days, ADE ↑ (2 h) days 1-10 ↑ day 2, ADE		(Grahame, Mosemiller, Low, & Froehlich, 2000)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Gene Knockout	Background	Operant	2BC	DID	References
	B6		↓ females — preference, males		(Racz et al., 2008)
Prodynorphin (<i>Pdyn</i>)	B6		↑ males/females		(Racz, Markert, Mauer, Stoffel-Wagner, & Zimmer, 2013)
	B6		—		(Sperling, Gomes, Sypek, Carey, & McLaughlin, 2010)
	B6		↑		(Femenia & Manzanares, 2012)
	B6 × 129/SvEv-Tac		↓ females — males		(Blednov, Walker, Martinez, & Harris, 2006)
Preproenkephalin (<i>Penk-l</i>)	B6		—		(Koenig & Olive, 2002)
	B6		— females ↑ 1-3 weeks, males		(Racz et al., 2008)

—, ↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in knockout *vs.* wild-type mice. Male mice were tested unless otherwise indicated. Ethanol intake in the operant and two-bottle choice (2BC) tests was measured in 30-min and 24-h sessions, respectively, unless otherwise indicated. Wild-type, but not knockout mice, showed a stress-induced increase in ethanol intake/preference (Racz et al., 2003; 2008; 2013; Sperling et al., 2010). Drinking in the dark (DID) session times are noted in parenthesis. ADE, alcohol deprivation effect. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 7
Alcohol consumption in immune-related receptor knockout mice

Gene Knockout	Background	2BC	DID	References
Cathepsin F (<i>Ctsf</i>)	B6 × 129/SvJ	↓males — females		(Blednov et al., 2012)
Cathepsin S (<i>Ctss</i>)	B6	↓males/females	— 2BC, males ↓ 1B, males ↓ 1B, 2BC, females	(Blednov et al., 2012)
C-C motif chemokine 2 (<i>Ccl2</i>)	B6	— males ↓ females		(Blednov et al., 2005)
C-C motif chemokine 3 (<i>Ccl3</i>)	B6	↓ males/females		(Blednov et al., 2005)
C-C chemokine receptor type 2 (<i>Ccr2</i>)	B6	↓ males/females		(Blednov et al., 2005)
C-C chemokine receptor type 5 (<i>Ccr5</i>)	B6	— males/females		(Blednov et al., 2005)
Apolipoprotein E (<i>ApoE</i>)	B6	— females		(Bechtholt, Smith, Raber, & Cunningham, 2004)
Monocyte differentiation antigen CD14 (<i>Cd14</i>)	B6	↓ males/females	— 1B, 2BC, males — 1B, females ↓ 2BC, females	(Blednov et al., 2012)
Interleukin 1 receptor antagonist (<i>Il1rn</i>)	B6 × 129/SvJ	↓ males/females	↓ 1B, 2BC, males ↓ 1B, 2BC, females	(Blednov et al., 2012)
Interleukin-6 (<i>Il6</i>)	B6	↓ males/females	— 1B, 2BC, males — 1B, 2BC, females	(Blednov et al., 2012)
β-2-microglobulin (<i>B2m</i>)	B6	↓ males/females	— 1B, 2BC, males — 1B, 2BC, females	(Blednov et al., 2012)
B2 bradykinin receptor (<i>Bdkrb2</i>)	B6	—		(Maul et al., 2005)
Toll-like receptor 4 (<i>Tlr4</i>)	B6	—		(Pascual et al., 2011)

—, ↓, ↑: no significant difference, decreased ethanol intake and/or preference, or increased ethanol intake and/or preference, respectively, in knockout vs. wild-type mice. Male mice were tested unless indicated otherwise. Ethanol intake in the two-bottle choice (2BC) tests was measured in 24-h sessions. Drinking in the dark (DID) session times were 2 and 4 h; 1B, one bottle. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 8

Alcohol consumption in ion channel mutant mice

Gene Knockout/Overexpression *	Background	2BC	CIE	DID	References
Transient receptor potential cation channel, subfamily M member 5 (<i>Trpm5</i>)	B6	↓			(Blednov et al., 2008)
Transient receptor potential cation channel subfamily V member 1 (<i>Trpv1</i>)	B6 × 129X1/SvJ	↑			(Blednov & Harris, 2009)
G protein-activated inward rectifier potassium channel 2, GIRK2 (<i>Kcnj6</i>)	B6 × 129Sv	— withbottle alternation; males/females ↑ without bottle alternation; females			(Blednov et al., 2001)
G protein-activated inward rectifier potassium channel 3, GIRK3 (<i>Kcnj9</i>)	B6	↑ (2 h) —			(Herman et al., 2015)
Voltage-dependent N-type calcium channel subunit α-1B (<i>Cacna1b</i>)	B6 × 129SvJae	↓			(Newton et al., 2004)
Calcium-activated potassium channel subunit β-1, BK β1 (<i>Kcnnb1</i>)	B6	— continuous, intermittent	↑ <i>a</i>	— (2 h)	(Kreifeldt et al., 2013)
Calcium-activated potassium channel subunit β-4, BK β4 (<i>Kcnnb4</i>)	B6	— continuous, intermittent	↓ <i>a</i>	— (2 h)	(Kreifeldt et al., 2013)
	B6			↑ (2 h) males/females	(Martin et al., 2008)
P2X purinoceptor 4 (<i>P2rx4</i>)	B6	↑ intake		↑ (4 h) 5%, 20% — (4 h) 10%	(Wyatt et al., 2014)
Glycine receptor subunit α-2 (<i>Glr2</i>)	B6	↓ — intermittent		— (2, 4 h) 1B	(Blednov, Benavidez, Black, Leiter, et al., 2015)
Glycine receptor subunit α-3 (<i>Glr3</i>)	B6	— ↑ intermittent		— (2, 4 h) 1B	(Blednov, Benavidez, Black, Leiter, et al., 2015)
Nicotinic acetylcholine receptor (nAChR) α4 subunit (<i>Chrna4</i>)	B6			↓ (2 h) 20% — (2 h) 2%	(Hendrickson et al., 2010)
nAChR α5 subunit (<i>Chrna5</i>)	B6			— (4 h)	(Santos et al., 2013)
nAChR α6 subunit (<i>Chrna6</i>)	B6	— males/females			(Kamens et al., 2012)
nAChR α7 subunit (<i>Chrna7</i>)	B6	↓ females — males			(Kamens et al., 2010)
nAChR β2 subunit (<i>Chrnβ2</i>)	B6	— males/females			(Kamens et al., 2010)
nAChR β3 subunit (<i>Chrnβ3</i>)	B6	— intermittent			(Dawson et al., 2013)
	B6	— males/females			(Kamens et al., 2012)
nAChR α5α3β4 *	B6SJLF1/J	↓ intake			(Gallego et al., 2012)

* indicates gene overexpression;

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

↓, ↑: no difference, decreased, or increased ethanol intake and/or preference, respectively, in mutant *vs.* wild-type mice. Male mice were tested unless otherwise indicated. Ethanol intake in the two-bottle (2BC) tests was measured in 24-h sessions unless otherwise indicated. Drinking session times for the other tests are indicated in parenthesis. CIE, chronic intermittent ethanol; DID, drinking in the dark; 1B, one bottle.

^aThese tests involved weekly limited access (2 h/day) 2BC drinking alternated with weekly CIE vapor (16 h/day) to create alcohol dependence/withdrawal. Deletion of BK β 4 attenuated, while deletion of BK β 1 accelerated, the escalation of ethanol drinking in dependent mice during withdrawal from CIE. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 9

Alcohol consumption in kinase mutant mice

Gene Knockout/Overexpression *	Background	Operant	2BC	DID	References
cAMP-dependent protein kinase type II- β regulatory subunit (<i>Prkar2b</i>)	B6 $\times$ 129/SvJ		$\uparrow$ males/females		(Thiele, Willis, et al., 2000)
	B6	genotype/sex interactions	$\uparrow$ males/females		(Ferraro, Sparta, Knapp, Breesse, & Thiele, 2006)
	B6		$\uparrow$ males/females		(Fee et al., 2004)
	B6 $\times$ 129/SvEv		$\uparrow$ males/females		(Fee et al., 2004)
Protein kinase C (PKC) ϵ type (<i>Prkce</i>)	B6 $\times$ 129/SvJae	$\downarrow$ ADE $\downarrow$ ADE			(Olive et al., 2000)
	B6 $\times$ 129/SvJae $\times$ FVB/N		$\downarrow$		(Choi et al., 2002)
	B6 $\times$ 129/SvJae		$\downarrow$ (23 h)		(Hodge et al., 1999)
	B6 $\times$ 129/SvJae		$\downarrow$ (16 h)		(Olive et al., 2005)
	B6 $\times$ 129S4	$\downarrow$ $\downarrow$ post-tolerance			(Wallace et al., 2007)
	B6 $\times$ 129/SvJae		$\downarrow$		(Besheer et al., 2006)
	B6 $\times$ 129S4/SvJae			$\downarrow$ (2 h) amygdala	(Lesscher et al., 2009)
PKC γ type (<i>Prkcg</i>)	B6 $\times$ 129/SvEvTac		$\uparrow$ males/females		(Bowers & Wehner, 2001)
PKC ζ type (<i>Prkcz</i>)	B6 $\times$ 129S6/SvEvTac		— continuous $\uparrow$ (4, 24 h) intermittent	— (4 h)	(A. M. Lee et al., 2014)
Tyrosine-protein kinase Fyn (<i>Fyn</i>)	B6 $\times$ 129SF2/J		$\downarrow$		(Boehm, Peden, Chang, Harris, & Blednov, 2003)
	129/SvJ		— (23 h)		(Yaka, Tang, Camarini, Janak, & Roni, 2003)
	B6 $\times$ CBA		— ADE — post-stress		(Coven, Schumann, Yagi, & Spanagel, 2003)
<i>Fyn</i> *	B6 $\times$ 129/Sv		$\downarrow$ males/females		(Boehm, Peden, et al., 2004)
	B6		$\downarrow$		(Stork, Kojima, Stork, Kume, & Obata, 2002)
cGMP-dependent protein kinase 2 (<i>Prkg2</i>)	B6N and 129/SvN		$\uparrow$		(C. Werner et al., 2004)
Serine/threonine-protein kinase, TAO2 (<i>Tao2</i>)	B6			$\uparrow$ (4 h)	(Kapfhamer et al., 2013)
ALK tyrosine kinase receptor (<i>Alk</i>)	B6			$\uparrow$ (4 h) intermittent	(Lasek et al., 2011)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

* indicates gene overexpression;
↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in mutant *vs.* wild-type mice. Male mice were tested unless indicated otherwise. Ethanol intake in the two-bottle choice (2BC) tests was measured in 24-h sessions, unless otherwise indicated. Session times for operant tests were 16 h, and drinking in the dark (DID) session times are indicated in parenthesis. ADE, alcohol deprivation effect. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 10

Alcohol consumption in enzyme knockout mice

Gene Knockout	Background	Operant	2BC	DID	References
Adenylate cyclase (AC) type 1 (<i>Adcy1</i>)	B6	—	—	—	(Maas et al., 2005)
AC type 5 (<i>Adcy5</i>)	B6	↑ males/females	—	—	(Kim, Kim, Baek, Lee, & Han, 2011)
AC type 8 (<i>Adcy8</i>)	B6	↓	—	—	(Maas et al., 2005)
<i>Adcy1/Adcy8</i> double knockout	B6	↓	—	—	(Maas et al., 2005)
Pituitary adenylate cyclase-activating polypeptide (<i>Adcyap1</i>)	Crlj;CD1	↑	—	—	(Tanaka et al., 2010)
Aldehyde dehydrogenase, mitochondrial (<i>Aldh2</i>)	B6	↓	—	—	(Isse et al., 2002)
Glutamyl aminopeptidase (Aminopeptidase A) (<i>Enpep</i>)	B6 × 129Sv/lex	↓ sex not specified	—	—	(Fernandez et al., 2006)
	B6 × 129Sv	—	— post-stress ^a	—	(Faber et al., 2006)
Catechol- <i>O</i> -methyltransferase (<i>Comt</i>)	B6	↑ males — females	—	—	(Tamminaki, Forsberg, Karayorgou, Gogos, & Mannisto, 2008)
Amine oxidase [flavin-containing] A (<i>Maoa</i>)	Tg8 and C3H/HeJ	— (2, 24 h)	—	—	(Popova, Vishnivetskaya, Ivanova, Skrinskaya, & Seif, 2000)
Nephrilysin, NEP (<i>Mme</i>)	B6N	—	↑ post-stress	—	(Maul et al., 2012)
	Not specified	↑	—	—	(Siems et al., 2000)
Nitric oxide synthase, brain (<i>Nos1</i>)	B6 × 129X1/SvJ	↑ 8-16%, sex not specified	—	—	(Spanagel et al., 2002)
Cytochrome P450 2E1 (<i>Cyp2e1</i>)	129S1/SV-Ter	↓ preference 4-8%, females	—	—	(Correa et al., 2009)
Protein phosphatase 1 regulatory subunit 1B, DARPP-32 (<i>Ppp1r1b</i>)	B6	↓ (23 h)	—	—	(Risinger, Freeman, Greengard, & Fienberg, 2001)
Histidine decarboxylase (<i>Hdc</i>)	129/SvB6	—	—	— (4 h), males/ females	(Vanhanen et al., 2013)
Tyrosine-protein phosphatase non-receptor type 5, STEP (<i>Ptpn5</i>)	B6	↑	—	—	(Legastelois, Darcq, Wegner, Lombroso, & Ron, 2015)
Ubiquitin carboxyl-terminal hydrolase 46 (<i>Usp46</i>)	B6	↓	—	—	(Imai, Kano, Nonoyama, & Ebihara, 2013)

—, ↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in knockout vs. wild-type mice. Male mice were tested unless otherwise indicated.

^a Social stress reduced alcohol consumption in both knockout and wild-type mice, but there was no genotype difference. Ethanol intake in the two-bottle choice (2BC) tests was measured in 24-h sessions, unless indicated otherwise. Drinking session times for the other tests are indicated in parenthesis. DID, drinking in the dark; 1B, one bottle. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.

Table 11

Alcohol consumption in neuropeptide/hormone mutant mice

Gene Knockout/Overexpression *	Background	Operant	2BC	DID	References
Neuropeptide Y, NPY (<i>Npy</i>)	129/SvEv	—	3-10%, ↑ 20%		(Thiele, Miura, Marsh, Bernstein, & Palminter, 2000)
<i>Npy</i> *	B6 × 129Sv	↑			
	B6 × 129Sv	↓			(Thiele, Marsh, Ste Marie, Bernstein, & Palminter, 1998)
NPY receptor type 1 (<i>Npy1r</i>)	B6	↑ 3-10%, males ↑ 10%, females			(Thiele, Koh, & Pedrazzini, 2002)
NPY receptor type 2 (<i>Npy2r</i>)	129/SvJ × Balb/cJ	↓			(Thiele, Naveilhan, & Ernfor, 2004)
	Balb/cJ	—			(Thiele et al., 2004)
NPY receptor Y5 (<i>Npy5r</i>)	129/SvEv	—			(Thiele, Miura, et al., 2000)
Agouti-related protein (<i>Agtrp</i>)	B6	↓ (2 h), males/ females	↓ (2 h), females	— (2 h), males/ females ↓ (4 h), males/ females	(Navarro, Cubero, Ko, & Thiele, 2009)
Angiotensinogen (<i>Agtr</i>)	Not specified	↓			(Maul et al., 2001)
<i>Agtr</i> *	Not specified	↑			(Maul et al., 2001)
Type-1A angiotensin II receptor, AT1A (<i>Agtr1a</i>)	Not specified	↓ sex not specified			(Maul et al., 2005)
<i>Agtr1a</i> *	Not specified	—			(Moore, Kristew, Kirchoff, Davison, & Lawrence, 2007)
Type-2 angiotensin II receptor, AT2 (<i>Agtr2</i>)	Not specified	— sex not specified			(Maul et al., 2005)
Cholecystokinin receptor type A, CCK-A (<i>Cckar</i>)	B6	— preference			(Miyasaka et al., 2005)
Gastrin/cholecystokinin type B receptor, CCK-B (<i>Cckbr</i>)	B6	—			(Miyasaka et al., 2005)
	B6	— males — preference, females			(Abramov et al., 2006)
Cocaine- and amphetamine-regulated transcript protein (<i>Cartpt</i>)	B6	↓, males/females			(Salinas, Nguyen, Ahmadi-Tehrani, & Morrisett, 2014)
Corticotropin-releasing factor (<i>Crtf</i>)	B6 × 129S	↑ (2, 23 h)			(Olive et al., 2003)
	B6			↓ (2, 4 h), males/ females	(Kaur, Li, Stenzel-Poore, & Ryabinin, 2012)
Corticotropin-releasing factor (CRF) receptor 1 (<i>Crtf1</i>)	B6			↓ (2, 4 h), males/ females	(Kaur et al., 2012)

Gene Knockout/Overexpression *	Background	Operant	2BC	DID	References
	B6 × 129SvJ	— (1 h) ↓ (1h) ADE			(Chu, Koob, Cole, Zorrilla, & Roberts, 2007)
	129/Ola × CD1		— ↑ post-stress		(Sillaber et al., 2002)
	B6, males/females		— 3-10%, ↓ 20% — (21 h) 10%, pre-/post-stress ↓ (21 h) 10%, chronic stress		(Pastor et al., 2011)
	B6			↓ intake (2, 4 h), males/females	(Giardino & Ryabinin, 2013)
	129 × 1/SvJ × CD1		— ↑ post-stress ↑ escalation post-dependence — ADE		(Molander et al., 2012)
<i>Chr1^{NestinCre}</i>	129S2/Sv × B6 × (B6 × SJL)		— ↓ post-stress ↓ escalation post-dependence — ADE		(Molander et al., 2012)
CRF receptor 2 (<i>Crhr2</i>)	B6			↓ (2 h) day 1, males/females	(Kaur et al., 2012)
	B6		— females	↑ (30 min), males/ females — (2 h), males/ females	(Sharpe et al., 2005)
<i>Crhr1/Crhr2</i> double knockout	B6, males/females		— (21 h) 10%, pre-/post-stress ↓ (21 h) 10%, chronic stress		(Pastor et al., 2011)
	B6			— (2, 4 h), males/ females	(Kaur et al., 2012)
Galanin peptides (<i>Gal</i>)	129Ola/Hsd		↓ 15%, females — males		(Karatayev et al., 2010)
<i>Gal</i> *	B6		↑ 15%, pre-/post-food deprivation, males; — females		(Karatayev, Baylan, & Leibowitz, 2009)
Neurokinin-1 receptor, Substance P receptor (<i>Tacr1</i>)	B6		↓		(Thorsell, Schank, Singley, Hunt, & Heilig, 2010)
Noiceptin receptor (<i>Oprl1</i>)	B6		↓ 20%, females		(Sakoori & Murphy, 2008)
Relaxin-3 receptor 1 (<i>Rxfp3</i>)	B6		— ↓ post-stress		(Walker et al., 2015)
Melanin-concentrating hormone receptor 1 (<i>Mchrl</i>)	B6		↑ males — females	— (1 h), males/ females	(Duncan et al., 2007)

Gene Knockout/Overexpression*	Background	Operant	2BC	DID	References
Neurotensin receptor type 1 (<i>Ntsr1</i>)	B6 × 129X1/SvJ		↑		(M. R. Lee et al., 2010)
Neurotensin receptor type 2 (<i>Ntsr2</i>)	B6 × 129X1/SvJ		↑		(M. R. Lee, Hinton, Unal, Richelson, & Choi, 2011)
Urocortin (<i>Ucn</i>)	B6			— (2, 4 h), males/ females	(Kaur et al., 2012)
	B6		↓ preference		(Giardino, Cocking, Kaur, Cunningham, & Ryabinin, 2011)
Vasopressin V1a receptor (<i>Avpr1a</i>)	B6Cr Slc × 129/Sv and B6Cr Slc		↑ males> females		(Sanbe et al., 2008)
	B6 × 129/Sv-CP, females		— homozygous mutant — post-stress		(Caldwell et al., 2006)
Vasopressin V1b receptor (<i>Avpr1b</i>)	B6 × 129/SvJ, females		— — post-stress		(Caldwell et al., 2006)
Adiponectin receptor protein 2 (<i>Adipor2</i>)	B6			— (2 h) ↓ (2 h) CIE	(Repunte-Canonigo et al., 2010)
Atrial natriuretic peptide receptor 1 (<i>Npr1</i>)	B6 × 129/SvJ		— ↑ post-stress		(Mutschler et al., 2010)
Ghrelin (<i>Ghr1</i>)	B6			— (90 min)	(Jerlhag, Landgren, Egecioglu, Dickson, & Engel, 2011)
	Not specified		↓		(Bahi et al., 2013)
Leptin (<i>Lep</i>)	B6-Lep ^{ob}		↓ males/females		(Blednov, Walker, & Harris, 2004)
Leptin receptor (<i>Lep^r</i>)	B6-m Lep rd /J		↓ males/females		(Blednov, Walker, & Harris, 2004)
Melanocortin receptor 3 (<i>Mcr3^r</i>)	B6		— males/females		(Navarro et al., 2005)
	B6			↓ (1 h) — (4 h), males/ females	(Olney, Sprow, Navarro, & Thiele, 2014)
Melanocortin receptor 4 (<i>Mcr4^r</i>)	B6		— (6, 24 h), males/females		(Navarro et al., 2011)

* indicates gene overexpression;

—, ↓, ↑: no significant difference, decreased, or increased intake and/or preference, respectively, in mutant *vs.* wild-type/control mice. Male mice were tested unless indicated otherwise. Ethanol intake in the two-bottle choice (2BC) tests was measured in 24-h sessions unless indicated otherwise. Drinking session times for operant and drinking in the dark (DID) tests are indicated in parenthesis. Stress and alcohol dependence tend to increase ethanol intake in both control and knockout mice, and the arrows indicate if the drinking in knockout mice was lower or higher compared to control under these conditions (Molander et al., 2012); however, the knockout mice in Pastor et al. (2011) showed reduced intake following chronic stress, whereas wild-type mice showed increased intake. ADE, alcohol deprivation effect; CIE, chronic intermittent ethanol vapor. Recommended mouse protein and gene (in *italics*) names are from Uniprot. B6 refers to C57BL/6J mice. Classical neuropeptides are listed followed by a putative neuropeptide and hormones and hormone receptors.

Alcohol consumption in other mutant mice

Table 12

Gene Knockout/Overexpression *	Background	Operant	2BC	DID	References
α -synuclein (<i>Sncal</i>)	B6		↑		(Lopez-Jimenez et al., 2013)
β -arrestin-1 (<i>Arrb1</i>)	B6		—		(Bjork et al., 2008)
β -arrestin-2 (<i>Arrb2</i>)	B6		↓ 9-15%		(Bjork et al., 2008)
	B6		↑		(H. Li, Tao, Ma, Liu, & Ma, 2013)
Gap junction delta-2 protein, Connexin-36 (<i>Gjd2</i>)	B6			↓ (2 h) 20%	(Steffensen et al., 2011)
G1/S-specific cyclin-D2 (<i>Ccnd2</i>)	129X1/SvJ × B6 × BALB/cAnNCrl		— 2%, 4% ↑ 8%, 16%		(Jaholkowski et al., 2011)
α -gustducin (<i>Gnat3</i>)	B6		↓		(Blednov et al., 2008)
Histamine H3 receptor (<i>Hrh3</i>)	B6		↓	↓ (2, 4 h)	(Nuutinen et al., 2011)
LIM domain only protein 3 (<i>Lmo3</i>)	B6		— males/females	↑ (2, 4 h), males/ females	(Savarese, Zou, Kharazia, Maiya, & Lasek, 2014)
Metallothionein-1/2 (<i>Mt1, Mt2</i> double knockout)	129S7/SvEvBrd and 129S1/SvImJ		↑ males/females		(Loney, Uddin, & Singh, 2006)
Period circadian protein homolog 1 (<i>Per1</i>)	<i>Per1^{B6/m1}</i> B6-Tyr ^{c-Brd} × 129S7		— drink-o-meter ↑ stress		(Dong et al., 2011)
	B6-Tyr ^{c-Brd} × 129SvEv ^{Brd}	—	— ADE		(Zghoul et al., 2007)
Period circadian protein homolog 2 (<i>Per2</i>)	<i>Per2^{B6/m1}</i> B6-Tyr ^{c-Brd} × 129SvEv ^{Brd}	↑	↑ 8-16% ↓ after acamprostate		(Spanagel et al., 2005)
	B6		↑ drink-o-meter		(A. J. Brager et al., 2011)
	B6		↑		(A. Brager et al., 2011)
Disks large homolog 4, PSD-95 (<i>Dlg4</i>)	B6		↓ males/females ↑ day 1, ADE		(Camp et al., 2011)
Protein fosB (<i>Fosb</i>)	129Sv × BALB/c		— males/females		(Korkosz et al., 2004)
Protransforming growth factor α (<i>Tgfa</i>) [*]	CD1		↑		(Hilakivi-Clarke & Goldberg, 1995)
Epidermal growth factor receptor kinase substrate 8 (<i>Eps8</i>)	B6		↑ males/females		(Offenhauser et al., 2006)
Ras-related protein Rab-3A (<i>Rab3a</i>)	B6		— (23 h)		(Kaphamer et al., 2008)
Ras-specific guanine nucleotide-releasing factor 2 (<i>Rasgrf2</i>)	B6		↓		(Stacey et al., 2012)
Regulator of G-protein signaling 6 (<i>Rgs6</i>)	B6 × 129/Sv		↓ males/females		(Stewart et al., 2015)
Taste receptor type 1 member 3 (<i>Tas1r3</i>)	B6		↓		(Blednov et al., 2008)

Gene Knockout/Overexpression *	Background	Operant	2BC	DID	References
Trace amine-associated receptor 1 (<i>Taar1</i>)	B6		↓ males/females		(Brasser, Norman, & Lemon, 2010)
Protein unc-79 homolog (<i>Unc79</i>)	B6 × 129S1/Sv		↑ females		(Lynch et al., 2013)
	B6 and B6 × DBA/2J		↑		(Specia et al., 2010)

* indicates gene overexpression;

–, ↓, ↑: no significant difference, decreased, or increased ethanol intake and/or preference, respectively, in mutant vs. wild-type mice. Male mice were tested unless indicated otherwise. Ethanol intake in the operant and two-bottle choice (2BC) tests was measured in 30-min and 24-h sessions, respectively, unless otherwise indicated. Drinking in the dark (DID) session times are indicated in parenthesis. ADE, alcohol deprivation effect. Recommended mouse protein and gene (in italics) names are from Uniprot. B6 refers to C57BL/6J mice.