Published in final edited form as:

J Biomol NMR. 2017 January ; 67(1): 23–34. doi:10.1007/s10858-016-0081-6.

Bacterial Production of Site Specific ^{13}C Labeled Phenylalanine and Methodology for High Level Incorporation into Bacterially Expressed Recombinant Proteins

Bhargavi Ramaraju, Hana McFeeters, Bernhard Vogler, and Robert L. McFeeters*

Department of Chemistry, University of Alabama in Huntsville, Huntsville, AL 35899

Abstract

Nuclear magnetic resonance spectroscopy studies of ever larger systems have benefited from many different forms of isotope labeling, in particular, site specific isotopic labeling. Site specific ^{13}C labeling of methyl groups has become an established means of probing systems not amenable to traditional methodology. However useful, methyl reporter sites can be limited in number and/or location. Therefore, new complementary site specific isotope labeling strategies are valuable. Aromatic amino acids make excellent probes since they are often found at important interaction interfaces and play significant structural roles. Aromatic side chains have many of the same advantages as methyl containing amino acids including distinct ^{13}C chemical shifts and multiple magnetically equivalent ^1H positions. Herein we report economical bacterial production and one-step purification of phenylalanine with ^{13}C incorporation at the $\text{C}\alpha$, $\text{C}\gamma$ and $\text{C}\epsilon$ positions, resulting in two isolated ^1H - ^{13}C spin systems. We also present methodology to maximize incorporation of phenylalanine into recombinantly overexpressed proteins in bacteria and demonstrate compatibility with ILV-methyl labeling. Inexpensive, site specific isotope labeled phenylalanine adds another dimension to biomolecular NMR, opening new avenues of study.

Keywords

Site Specific Isotope Labeling; Phenylalanine; Aromatic Amino Acids; NMR Spectroscopy; Isolated ^1H - ^{13}C Spin Systems; Stable Isotope Incorporation

INTRODUCTION

Large macromolecular systems and membrane proteins are at the forefront of structural biology. A number of site specific isotope labeling techniques have been developed to provide information on these systems. Popular ^{13}C methyl labeling of isoleucine, leucine, valine (Gardner et al. 1997; Rosen et al. 1996), alanine (McCaldon and Argos 1988; Tugarinov and Kay 2003), methionine (Fischer et al. 2007; Gelis et al. 2007), and threonine (Sinha et al. 2011) side chains all take advantage of magnetically equivalent hydrogens and fast methyl rotation that allow for observation in slow tumbling systems. While providing tremendous insight, ^{13}C methyl labeling can be limited by the number and location of sites

probed. To overcome these limitations and provide a more complete picture of the system, more site specific probes readily implemented with methyl labeling are needed.

Aromatic residues are excellent probes for reporting structure and structural changes. Aromatic residues are often found in protein “hot spots” (Ben-Shimon and Eisenstein; Bogan and Thorn 1998) and have significant structural roles (Butterfield et al. 2002; Churchill et al.; Duan et al. 2000; Madhusudan Makwana and Mahalakshmi 2015; Schneider 2009). Interactions with aromatics are also important for molecular recognition of carbohydrates (Chavez et al. 2005; Hudson et al. 2015). Additionally, identification of unambiguous aromatic side chain NOEs can be instrumental for accurate structure determination (McFeeters et al. 2007a; McFeeters et al. 2007b). Like methyl containing residues, phenylalanine and other aromatics are attractive targets for site specific labeling since they generally have side chains with multiple magnetically equivalent hydrogens and characteristic ^{13}C chemical shifts. In addition, distinct biosynthesis pathways make simultaneous labeling of both aromatic and methyl containing residues possible. Thus, aromatic-aromatic, aromatic-methyl, and methyl-methyl information can be acquired from one sample. Overall, site specific labeling of aromatic amino acids holds great potential to advance NMR studies of biological macromolecules.

Several methods for site specific labeling of phenylalanine have been previously reported. In early pioneering work, aromatic residues with alternating ^{13}C incorporation were generated from hydrolysates of proteins expressed in media containing a variety of ^{13}C containing carbon sources (LeMaster and Cronan 1982; LeMaster and Richards 1982). Site specific ^{13}C labeling of proteins was also generated using 1- and 2- ^{13}C acetate (Wand et al. 1995) or 1- and 2- ^{13}C -glucose (Lundstrom et al. 2007; Teilum et al. 2006). For biosynthesis of deuterated proteins with selective protonation of aromatic rings, shikimic acid was utilized as a precursor, with improved incorporation efficiency achieved using auxotrophic bacteria for recombinant protein expression (Rajesh et al. 2003). The first site specific labeling and generation of isolated ^1H - ^{13}C spin systems during recombinant protein expression was performed in bacteria grown in the presence of phenylalanine synthesized from commercially available site specifically labeled tyrosine (Wang et al. 1999). Improved ^1H - ^{13}C site specific labeling came from synthetically prepared stereo array isotope labeled (SAIL) aromatic amino acids (Zhang 2009) incorporated to proteins in a cell-free expression system. However, the cost of SAIL aromatic amino acids is prohibitive and in spite of considerable improvements, cell-free protein production is not always reliable and not necessarily scalable to the milligram quantities necessary for structural biology (Casteleijn et al. 2013). Alternatively, site specific labeling of phenylalanine has been achieved using synthetic ^{13}C ketoacid precursors, an excellent approach limited by their current commercial unavailability (Lichtenecker 2014; Lichtenecker et al. 2013).

Herein we report an inexpensive means of site specific isotopically labeled (SSIL) phenylalanine biosynthesis and general methodology that results in high level incorporation of phenylalanine into recombinantly expressed proteins in bacteria. Phenylalanine with isolated ^1H - ^{13}C spin systems at the C_α and C_ϵ positions, termed SSIL-Phe (Figure 1), was produced in bacteria grown in media containing 2- ^{13}C -glycerol as the sole carbon source. Production yield efficiency (defined as grams of phenylalanine produced per gram carbon

source) of approximately 30% was achieved, in line with other industrially oriented systems producing phenylalanine from glycerol (Thongchuang et al. 2012). To make scale-up and purification straightforward, the system was engineered to secrete phenylalanine into the growth media. In addition to production, methodology is also presented for high level phenylalanine incorporation into recombinantly expressed bacterial proteins that is compatible with ILV-methyl labeling. The current laboratory scale cost of producing SSIL-Phe for protein production is on the order of uniform ^{13}C labeling, making it economically feasible for more widespread utilization.

MATERIALS and METHODS

Production and Purification of SSIL-Phe

JB-161 *E. coli* with enhanced carbon flux to the aromatic biosynthesis pathways were transformed with a modified p2.274 plasmid (Barker and Frost 2001). For production of phenylalanine, the original *ubiC* gene (directing carbon flux to production of *p*-hydroxybenzoic acid) in p2.274 was replaced with genes in the phenylalanine biosynthesis pathway. These genes were cloned from *E. coli* K12 and placed under control of separate promoters. Transformed cells were grown to an OD_{600} of 0.8 at 37 °C in LB media supplemented with 30 µg/mL kanamycin and 25 µg/mL chloramphenicol. Cells were then pelleted by centrifugation and resuspended in an equal volume of M9 minimal media containing the appropriate carbon source and 15 µg/mL kanamycin, and 25 µg/mL chloramphenicol. Cultures were grown at 30 °C for a defined period of time, the media containing secreted phenylalanine clarified by centrifugation, and the pH adjusted to 2.0. The supernatant was then filtered through a 0.22 µm syringe filter and loaded onto a 10 × 250 mm PRP1-C18 column (Hamilton Corporation, Reno, NV, USA) at a constant flow rate of 3 mL/min in 40 mL batches. Sufficient separation was achieved without heating so the column was kept at room temperature (~22 °C). Phenylalanine was eluted by an acetonitrile gradient and detected by monitoring absorbance at 254 nm. Fractions containing phenylalanine were pooled and dried.

Phenylalanine NMR Spectroscopy

Dried phenylalanine was resuspended in D_2O and spectra were acquired at 25 °C on a 500 MHz Varian Inova spectrometer equipped with a room temperature probe. One dimensional ^1H spectra were acquired with an 8,000 Hz sweep width centered at 6 ppm. Typically 32k points were collected and 16 transients were averaged with a recycle delay of 3 seconds. One-dimensional ^{13}C spectra were acquired with a 26,400 Hz sweep width centered at 100 ppm. Generally, 65.5k points were collected. ^{13}C -HSQC spectra were acquired using the same ^1H parameters except the number of points was reduced to 2048. Typically 16 transients were averaged per point. ^{13}C parameters included a sweep width of 17,600 Hz centered at 70 ppm and 128 points (64 real + 64 imaginary) collected.

Mass Spectrometry

Phenylalanine was derivatized with *N*-(*tert*-butyldimethylsilyl)-*N*-methyl-trifluoroacetamide (MTBSTFA) for GC-MS analysis. A 200 µL sample of 500 µM phenylalanine in acetonitrile was combined with 200 µL of MTBSTFA also in acetonitrile and incubated for 4

hours at 55 °C. Following the derivatization, GC-MS analysis was conducted using an Agilent 6890 series gas chromatograph equipped with Agilent J&W HP-5MS capillary column directly connected to 5973 Network Mass Selective Detector. The column was composed of (5%-phenyl)-methylpolysiloxane with 0.25 µm film thickness and 30 m × 0.25 mm dimensions. Initial oven temperature was 100 °C which was held for 1 min and then raised to 280 °C at a rate of 35 °C/min. The temperature was further increased to 320 °C at a rate of 40 °C/min and held constant for 2.5 min. The injection volume was 1 µL, injector temperature 300 °C, and the split ratio 5:1. Helium was used as the carrier gas at a constant flow rate of 1 mL/min, initial pressure of 10.5 psi, and average velocity of 37 cm/sec. Other settings included 280 °C interface temperature, 230 °C ion source temperature, 150 °C quad temperature, and electron impact ionization at 70 eV. Mass spectra were recorded in the range of m/z 40–450 amu at a rate of 3.5 scans/min for a run time of 9.64 min. The solvent delay was 3.0 min.

Ubiquitin Expression

BL21(DE3) *E. coli* were transformed with pET28b carrying the ubiquitin gene (Wilson et al.). A single colony was used to inoculate 30 mL of M9 minimal media supplied with 30 µg/mL kanamycin and the culture was grown overnight at 37 °C. In the morning, the starter culture was spun down and cells resuspended in 1 Liter of M9 minimal media with 30 µg/mL kanamycin. The culture was grown at 37 °C to an OD₆₀₀ of 0.6. The cells were pelleted by centrifugation and resuspended in optimized M9+20 media (McFeeters 2002) (see Table 1, Results) containing 140 mg/L of glyphosate. Cells were then allowed to grow to an OD₆₀₀ of 0.8 where expression of ubiquitin was induced with IPTG added to a final concentration of 1 mM. Four hours later, cells were harvested and the cell pellet stored at -80 °C until purification. For purification, frozen cell pellets were resuspended in 50 mM Tris-HCl, pH 8.0 and supplemented with 5 mM MgCl₂, DNase at 0.8 units/mL, and 10µg/mL lysozyme. The mixture was left to incubate for 30 minutes at room temperature, followed by a 30 minute heat cut at 65 °C. Insoluble material was separated by centrifugation and ubiquitin was purified by HPLC. The supernatant was applied to a 4.6 × 250 mm C4 (Bio-Rad Laboratories, Hercules, CA, USA) column. At a flow rate of 1.0 mL/min, ubiquitin was eluted with a gradient from 0 to 90% acetonitrile in 0.05% TFA over the course of an hour. Ubiquitin containing fractions were pooled, dialyzed extensively against an NMR buffer of 20 mM NaOAc, 20 µM NaN₃, pH 5.0, and concentrated by ultrafiltration.

E. coli Pth1 Expression

W3110 *E. coli* cells were transformed with a pKQV4 plasmid bearing the *E. coli* peptidyl-tRNA hydrolase (Pth1) gene. From a single colony, cells were originally grown in M9 minimal media to an OD₆₀₀ of 0.4. Cells were then pelleted and resuspended in the same volume of modified M9+20 (see Table 1, Results) containing 140 mg/L glyphosate. At an OD₆₀₀ of 0.6, the temperature was dropped to 25°C and Pth1 expression was initiated with IPTG added to a final concentration of 1 mM. After 16 hours the cells were harvested by centrifugation and stored at -80 °C until purification. Pth1 was purified as previously described (Hughes et al. 2012; McFeeters et al. 2012; Taylor-Creel et al. 2014).

Protein NMR Spectroscopy

Protein NMR data were collected at 25 °C on a 500 MHz Varian Inova spectrometer equipped with a room temperature probe. For ^{15}N -HSQC data, the ^1H sweep width was 6,000 Hz centered on water (4.77 ppm). For the ^{15}N dimension, the sweep width was 1,600 Hz with the carrier at 118 ppm. Sixteen transients were collected for each of the 192 points (96 real + 96 imaginary). The recycle delay was 1.5 seconds. For non-constant time ^{13}C -HSQC data, ^1H parameters were the same as for ^{15}N -HSQC. Carbon parameters included a sweep width of 17,500 Hz with carrier centered at 70 ppm for the full carbon spectra and a sweep width of 4,400 Hz centered at 127 ppm for aromatic spectra. Resonance assignments for ubiquitin were obtained from the BMRB ID 5387 (Babu et al. 2001). Resonance assignments for *E. coli* Pth1 were adapted from BMRB entry 17404 (Giorgi et al. 2011) for the buffer conditions used.

For measurement of ^{13}C T_1 and $T_{1\rho}$, ^{13}C sweep widths were set to 1382 Hz with the carbon carrier set to 132.5 ppm. The recycle delay was 4 seconds and 16 transients were collected for 32 complex points. Relaxation delays for T_1 were 0*, 0.025, 0.05, 0.1*, 0.15, 0.2, 0.25, 0.3*, 0.45, 0.6 seconds and for $T_{1\rho}$ 0.01, 0.015, 0.02*, 0.03, 0.045*, 0.06, 0.08, 0.1 seconds (* indicates time points measured in triplicate). Data were processed with Gaussian window functions and peak volumes used for determination of relaxation parameters.

RESULTS AND DISCUSSION

Bacterial Production of Phenylalanine

Industrial production of amino acids has frequently been performed in *C. glutamicum*. However, extensive alterations are required for direct use of glycerol as a carbon source (Rittmann et al. 2008). Therefore, we adopted an *E. coli* system previously engineered to produce aromatic *p*-hydroxybenzoic acid (Barker and Frost 2001) as the starting point for production of phenylalanine. Elevated secretase activity was retained, resulting in secretion of phenylalanine into growth media, greatly facilitating purification. Modifications directing carbon flux to phenylalanine production consisted of replacing the original *ubiC* gene in the p2.274 plasmid with *E. coli* phenylalanine biosynthesis pathway genes. Focusing on the bottlenecks from both the common pathway leading to chorismate, the branched pathway from chorismate to phenylalanine (Ikeda 2006), and combinations known to elevate tyrosine production (Lutke-Eversloh and Stephanopoulos 2008), the effects of overexpressing *aroC* (chorismate synthase), *aroK* (shikimic acid kinase), *tyrB* (tyrosine aminotransferase, which also functions as a phenylalanine aminotransferase), and *ydiB* (shikimate dehydrogenase) on phenylalanine production were determined, Figure 2. It was found that overexpressing combinations of these genes produced more phenylalanine even when minimal changes were observed for individually overexpressed genes, in agreement with previous studies of tyrosine production in *E. coli* (Lutke-Eversloh and Stephanopoulos 2008). Also similar to tyrosine, the combination of *tyrB/ydiB* showed considerable increase in production of phenylalanine. However, the combination of *aroK/tyrB* produced the most phenylalanine under all conditions tested and was utilized for subsequent studies.

Preferred Carbon Source to Produce SSIL-Phe

Glucose, glycerol, and acetate were tested as carbon sources that could potentially be used for site specific ^{13}C isotope incorporation into phenylalanine. Glucose has been extensively utilized for industrial production of aromatic amino acids including phenylalanine (Baez-Viveros et al. 2004; Ikeda and Katsumata 1992; Sprenger 2007; Wang et al. 2011; Zhou et al. 2010). However, current commercial availability is limited to a single ^{13}C labeled site (i.e. 2- ^{13}C -glucose, not 2,5- ^{13}C -glucose), leading to diminished ^{13}C incorporation in phenylalanine since only half of the pyruvate from glycolysis would contain ^{13}C . While both glycerol and acetate are commercially available with individual ^{13}C sites labeled, glycerol was most efficiently converted to phenylalanine. Maximum production yield efficiency (i.e. grams of phenylalanine obtained per 1 gram of carbon source) was 32% for glycerol, 27% for glucose and 10% for acetate albeit under slightly different conditions for each. While the starting concentration of glycerol and glucose provided was 1 g/L, the concentration of acetate used was 5 g/L. The time required to reach maximum phenylalanine production was different for each carbon source, fastest for glucose (generally 1.5 days), intermediate for glycerol (3 days), and slowest for acetate (4.5 days) when comparing the same starting amount of carbon source present in the media. Slower production may be the result of slightly elevated media viscosity when using glycerol as well as the altered metabolic demands associated with using acetate (Oh et al. 2002). Regardless of conditions, glycerol led to the most efficient phenylalanine production and was used as the carbon source for subsequent studies.

Optimization of SSIL-Phe Production

In order to maximize efficiency, the impacts of growth temperature and starting glycerol concentration on phenylalanine production were determined. The starting amount of glycerol had the greatest impact on phenylalanine production. Most notably, more efficient production of phenylalanine was achieved at lower glycerol concentrations with maximal production of 150, 315, 385 and 750 mg of phenylalanine per liter from starting concentrations of 0.5, 1, 2, and 8 grams of glycerol, respectively. This translated to production yield efficiencies of 30, 31.5, 19.3, and 9.4%. At concentrations below 1 g/L, however, ^{13}C incorporation markedly decreased indicating the presence of an unlabeled “carbon reservoir” from the initial growth and accumulation of cell density in natural abundance media. Efficiency gains from an initial growth in isotopic labeled glycerol media to purge the unlabeled reservoir did not outweigh the additional cost. Unlike the concentration of glycerol, growth temperature had a more modest effect on phenylalanine production yield efficiency with maximized production at 30 °C.

For all conditions tested, no significant change in phenylalanine production was observed for increased aeration using baffled flasks or reduced culture volume to flask ratio. Thus, it appears that batch-fed fermentation would not significantly benefit this application with the exception of potentially reaching maximum production faster (Thongchuang et al. 2012). Using MOPs buffered media was reported to significantly increase bacterial production of tyrosine (Santos and Stephanopoulos 2008), but had no effect on phenylalanine production in this system. Regardless of growth temperature or glycerol concentration, maximum phenylalanine production was achieved at 72 $\pm$ 6 hours, or approximately 3 days.

Increasing the amount of nitrogen present, in the form of ammonium chloride, had no effect on phenylalanine production. Thus optimal phenylalanine production conditions were M9 minimal media with 1 g/L of glycerol (substituting for glucose) grown for 3 days at 30 °C which consistently produced over 300 mg of phenylalanine per liter after purification.

Characterization of SSIL-Phe Isotope Incorporation

The identity of phenylalanine as a purified product of the bacterial synthesis and isotopic incorporation was determined by NMR spectroscopy and mass spectrometry (Figure 3). From the one dimensional carbon spectrum, high levels of ^{13}C incorporation were observed at the $\text{C}\alpha$, $\text{C}\gamma$, and $\text{C}\epsilon$ positions of phenylalanine. Specifically for the ^1H - ^{13}C sites of interest, ^{13}C incorporation was calculated to be ~95% for $\text{C}\alpha$ and averaged ~70% for $\text{C}\epsilon$ from integrated peak intensities in the ^1H spectra. Whereas $\text{H}\alpha$ is clearly resolved, readily allowing integration of ^{13}C versus ^{12}C coupled peaks, only the downfield ^{13}C coupled resonance of $\text{H}\epsilon$ could be reliably quantified. This level of ^{13}C isotope incorporation is in agreement with previous characterization of aromatic amino acids produced in bacteria utilizing glycerol as the sole carbon source (LeMaster and Cronan 1982). ^{15}N incorporation was controlled by growth conditions with >95% incorporation readily achieved using ^{15}N ammonium chloride as the sole nitrogen source during SSIL-Phe production. Mindful of the potential for vastly different relaxation rate, in particular for aromatic resonances, ^{13}C incorporation was also characterized using GC-MS. MTBSTFA derivatives with SSIL-Phe had the same retention time as commercial phenylalanine and fragmentation patterns for natural abundance phenylalanine matched expected values from the NIST/EPA/NIH reference library. The increase in size observed for fragmentation patterns of SSIL-Phe agreed with the isotope incorporation observed via NMR. Best fits of simulated ^{13}C incorporation to the isotope distribution of individual fragment peaks closely matched the NMR derived values.

SSIL-Phe in Bacterially Expressed Recombinant Proteins

Incorporation of SSIL-Phe into bacterially expressed recombinant proteins was shown in 8.5 kDa ubiquitin (76 residues, 2 phenylalanine and 1 tyrosine residues). To show applicability of the method to other systems, recombinant incorporation was also demonstrated in another protein of different size and aromatic content *E. coli* Pth1 (20 kDa, 194 residues, 11 phenylalanine, 2 tyrosine, and 2 tryptophan residues). Since it was previously reported and we also observed that negligible differences are found in the degree of ^{13}C and ^{15}N isotope scrambling for phenylalanine (Krishnarajuna et al. 2011), readouts of reverse labeled ^{15}N -HSQC spectra (natural abundance phenylalanine in a uniform ^{15}N background) were initially used to monitor phenylalanine incorporation into recombinantly expressed ubiquitin and Pth1.

Direct Amino Acid Supplementation of M9 Minimal Media

As a first attempt to incorporate phenylalanine into recombinant proteins, natural abundance phenylalanine was added directly to uniform ^{15}N M9 minimal media. For the expression of both ubiquitin and *E. coli* Pth1, adding 200 mg/L of phenylalanine led to approximately 40% incorporation and the amount of expressed protein was unaffected. Though the level of phenylalanine incorporation increased with increasing concentration of phenylalanine, even

at 2 g/L, the highest concentration tested, phenylalanine incorporation only reached ~50%. Moreover, considerable isotope scrambling was observed when using such high phenylalanine concentration, primarily to tyrosine and tryptophan and to a lesser extent Asx/Glx residues. Supplementing media with 200 mg/L of these amino acids, in addition to phenylalanine, suppressed scrambling but did not improve phenylalanine incorporation.

To prevent biosynthesis of phenylalanine during recombinant protein production and thereby improve SSIL-Phe incorporation, glyphosate was utilized. Glyphosate is an inhibitor of 5-enolpyruvyl-shikimate-3-phosphate production, one of the final intermediates in the common aromatic amino acid biosynthetic pathway. The concentration of glyphosate, duration of aromatic suppression, and amount of SSIL-Phe added versus level of recombinant incorporation were optimized. For both proteins tested, maximal SSIL-Phe incorporation was achieved by addition of 140 mg/liter glyphosate at an OD₆₀₀ of 0.6 along with 400 mg/L of phenylalanine and 200 mg/L of tyrosine and tryptophan. Following this procedure, the highest level of phenylalanine incorporation still only reached 70%.

Improved SSIL-Phe Incorporation with Modified M9

To further improve phenylalanine incorporation into bacterially expressed recombinant proteins, a strategy using modified M9 minimal media was developed. In this case, isotope labeled phenylalanine was used, while the rest of the media reagents were unlabeled. Transformed cells were grown in unlabeled M9 minimal media to generate the initial cell density. At an OD₆₀₀ of 0.6, cells were pelleted and resuspended in an equal volume of modified M9, termed M9+20 (Table 1), containing 140 mg/L glyphosate. At an OD₆₀₀ of 0.8, roughly 45 minutes later when grown at 37 °C, recombinant protein expression was induced as usual. Cells were harvested after the same amount of time used for expression in regular M9 minimal media.

Originally, all 20 amino acids were added at a concentration of 200 mg/L with only phenylalanine isotopically labeled, a method previously established for site specific labeling of other amino acids (McFeeters 2002). While the amino acids are generally more concentrated than reported in other selective isotope labeling studies (Shortle 1994; Tong et al. 2008), this mixture maintained recombinant protein expression levels and resulted in high level SSIL-Phe incorporation for all conditions tested. However, isotopic scrambling from phenylalanine to tyrosine, aspartate, and glutamate was observed. Therefore, in line with previous observations, the concentration of these amino acids was increased to 400 mg/L. Isotopic scrambling was reduced to <5% with no reduction in phenylalanine incorporation. Using this approach, the protein yield remained unchanged compared to regular M9 for both ubiquitin (Figure 4) and *E. coli* Pth1. Phenylalanine incorporation was >95% in both cases.

Phenylalanine ¹He-¹³Ce Dynamics in Ubiquitin

Utilization of SSIL-Phe for ¹³Ce dynamics characterization in ubiquitin was evaluated. Relaxation parameters in the non-deuterated background phenylalanine are moderately affected by relaxation of adjacent hydrogens, in this case ¹Hδ and ¹Hη. From equations governing relaxation (Kleckner and Foster 2011), the relative contribution from lack of perdeuteration is not completely negligible, summing to ~3% that of the bonded ¹He

(calculated using distances of 1.09 Å for the $^1\text{H}\epsilon$ - $^{13}\text{C}\epsilon$ bond and 2.20 Å for both the $^1\text{H}\delta$ - and $^1\text{H}\eta$ - $^{13}\text{C}\epsilon$ distances). Even without background deuteration, relaxation parameters from aromatic side chains can provide meaningful insight as demonstrated using 1- ^{13}C glucose labeling (Teilum et al. 2006) which revealed an expanded network of dynamically coupled residues in eglin c (Boyer and Lee 2008). Similarly, though partially deuterated, labeling with 4- ^{13}C erythrose-4 phosphate and deuterated pyruvate exposed a wide range of motions in the sub-nanosecond time range for aromatic side chains of calcium saturated calmodulin (Kasinath et al. 2013). SSIL-Phe labeling has the advantage over these labeling methods of generating isolated ^1H - ^{13}C spin systems at multiple positions in the aromatic side chain (i.e. at both $\text{C}\epsilon$ positions), not just one. The cost is on par with single ^{13}C labeled glucose, but considerably less than using 4- ^{13}C erythrose even before adding the cost of deuterated pyruvate. To demonstrate the utility, ^{13}C relaxation parameters were measured for SSIL-Phe labeled ubiquitin (Figure 5). The faster transverse relaxation of Phe45 fits with recent characterization of supra tauC motions in ubiquitin (Lange et al. 2008) in which a pincer motion is hypothesized. Moreover, Phe45 is part of the conformationally dynamic binding face of ubiquitin while Phe4 is not.

ILV-Methyl Labeling Compatibility

To further extend utility of SSIL-Phe, we explored the possibility of simultaneous SSIL-Phe and ILV-methyl labeling. The simplest approach, substituting the appropriate α -ketoacid precursors for isoleucine, leucine, and valine in the M9+20 at concentration used in unmodified M9, proved successful. High level methyl labeling was achieved with no change observed in SSIL-Phe labeling (Figure 6). Thus simultaneous aromatic-ILV-methyl labeling is readily achievable.

CONCLUSIONS

Reported here is the efficient production of SSIL-Phe in which ^{13}C has been incorporated at the $\text{C}\alpha$, $\text{C}\gamma$, and $\text{C}\epsilon$ positions. Being inexpensive to produce and having isolated ^1H - ^{13}C spin systems makes this reagent widely applicable for probing phenylalanine positions in a variety of settings. Also reported is an undemanding method for high level incorporation of SSIL-Phe into bacterially expressed recombinant proteins. This provides a means for facile phenylalanine labeling for existing BL21 expression systems. Adding to utility, compatibility with ILV-methyl labeling is demonstrated. Overall, SSIL-Phe is readily accessible and is a first step towards inexpensive implementation in systems of increasing size.

Compared to existing site specific phenylalanine labeling strategies, the major benefit of the SSIL-Phe system is reduced cost. Enough phenylalanine can be produced from just over a gram of starting glycerol to label a liter of media for recombinant protein production. Other benefits, discussed below, include alternate isotope incorporation and adaptation to other molecules of interest. Compared to the erythrose/pyruvate method for dynamics (Kasinath et al. 2013), the SSIL-Phe system allows control of which amino acids are labeled. Also [4- ^{13}C] erythrose leads to labeling of one $\text{C}\delta$ position, not ^{13}C incorporation at both (at least in part) $\text{C}\epsilon$ positions. Compared to aromatic ketoacids, SSIL-Phe can be directly compatible

with *in vitro* labeling systems and existing SILAC protocols. The major disadvantages of the SSIL-Phe system are the lack of deuteration at ^{12}C positions and current incorporation methodology is not, short of using auxotrophic strains, readily amenable to uniform $^{13}\text{C}/^{15}\text{N}$ labeling or perdeuteration.

With the aim of widespread use in mind, the SSIL system presented offers many possibilities of further development and innovation. Phenylalanine produced by bacterial biosynthesis is one enzymatic step from the corresponding ketoacid precursor. Thus, it may be possible to inexpensively produce the ketoacid precursor from the full amino acid, in particular to utilize the previously reported straightforward media supplementation for high level incorporation (Lichtenecker et al. 2013) which is of particular importance for uniform $^{13}\text{C}/^{15}\text{N}$ labeling or perdeuteration. Produced in bacteria, alternative isotope labeling of SSIL-Phe is readily possible. Options certainly include the use of different isotopically labeled carbon source (i.e. 1- ^{13}C -glycerol) and “background” deuteration of protonated ^{12}C positions (currently underway). Background deuteration requires a partially deuterated precursor which is not yet commercially available with appropriate ^1H - ^{13}C labeling, however several straight forward strategies for appropriate deuteration of available precursors are being developed. Additional improvements may also be possible in the incorporation of aromatic amino acids into recombinant proteins, including reducing the amount of SSIL aromatic amino acids necessary for high level incorporation by engineering bacterial strains (elevating aromatic uptake via increased transporter/pump expression or biosynthesis knockout (Miyanoiri et al. 2016)) or extending SSIL-Phe incorporation to proteins expressed in *Pichia pastoris*.

Overall, the bacterial based production system for SSIL-Phe was designed to be “modular.” It readily accommodates other isotope labeling options as well as production of other biomolecules. Having the biosynthetic genes on a plasmid makes addition or substitution of other genes of interest straight forward. This opens the door for production of other amino acids, metabolites, natural products, or biomolecules of interest by replacing the phenylalanine biosynthetic genes with corresponding genes for production of the compound of interest. Different ^{13}C isotope distributions in the final product are achieved by simply changing the isotope distribution or carbon source. We focused on ^1H - ^{13}C labeling of the side chain epsilon position of phenylalanine. While eliminating carbon-carbon coupling, thereby the need for constant time NMR experiments, this isotopic labeling pattern also significantly simplifies the aromatic carbon spectra by reducing the number of ^1H - ^{13}C resonances per phenylalanine residue to one. Therefore, unambiguous resonance assignment of phenylalanine is possible even in systems too large for traditional assignment methods by site directed mutagenesis. Moreover, the ^{13}C chemical shift is significantly different between phenylalanine and tyrosine, allowing for better separation in future simultaneous site specific labeling of phenylalanine and tyrosine. Implementation in highly deuterated systems is still limited, but deuteration of background positions in SSIL produced molecules is possible as is efficient conversion of amino acids to corresponding ketoacids for straight forward incorporation using traditional M9 media, both areas of active investigation.

Acknowledgments

Research reported in this publication was supported by the NIGMS of the National Institute of Health under award number 1R15GM093912. The authors thank Mary Hames for cloning and molecular biology support.

References

- Babu CR, Flynn PF, Wand AJ. Validation of protein structure from preparations of encapsulated proteins dissolved in low viscosity fluids. *J Am Chem Soc.* 2001 Mar 21; 123(11):2691–2. [PubMed: 11456950]
- Baez-Viveros JL, Osuna J, Hernandez-Chavez G, Soberon X, Bolivar F, Gosset G. Metabolic engineering and protein directed evolution increase the yield of L-phenylalanine synthesized from glucose in *Escherichia coli*. *Biotechnol Bioeng.* 2004; 87:516–524. DOI: 10.1002/bit.20159 [PubMed: 15286989]
- Barker JL, Frost JW. Microbial synthesis of p-hydroxybenzoic acid from glucose. *Biotechnol Bioeng.* 2001; 76:376–390. DOI: 10.1002/bit.10160 [PubMed: 11745165]
- Ben-Shimon A, Eisenstein M. Computational mapping of anchoring spots on protein surfaces. *J Mol Biol.* 2010; 402:259–277. S0022-2836(10)00774-6. DOI: 10.1016/j.jmb.2010.07.021 [PubMed: 20643147]
- Bogan AA, Thorn KS. Anatomy of hot spots in protein interfaces. *J Mol Biol.* 1998; 280:1–9. DOI: 10.1006/jmbi.1998.1843 [PubMed: 9653027]
- Boyer JA, Lee AL. Monitoring aromatic picosecond to nanosecond dynamics in proteins via ¹³C relaxation: expanding perturbation mapping of the rigidifying core mutation, V54A, in *eglin c*. *Biochemistry.* 2008; 47:4876–4886. DOI: 10.1021/bi702330t [PubMed: 18393447]
- Butterfield SM, Patel PR, Waters ML. Contribution of Aromatic Interactions to α -Helix Stability. *J Am Chem Soc.* 2002; 124:9751–9755. DOI: 10.1021/ja026668q [PubMed: 12175233]
- Casteleijn MG, Urtti A, Sarkhel S. Expression without boundaries: Cell-free protein synthesis in pharmaceutical research. *International Journal of Pharmaceutics.* 2013; 440:39–47. <http://dx.doi.org/10.1016/j.ijpharm.2012.04.005>. [PubMed: 22521712]
- Chavez MI, et al. On the importance of carbohydrate-aromatic interactions for the molecular recognition of oligosaccharides by proteins: NMR studies of the structure and binding affinity of AcAMP2-like peptides with non-natural naphthyl and fluoroaromatic residues. *Chemistry.* 2005; 11:7060–7074. DOI: 10.1002/chem.200500367 [PubMed: 16220560]
- Churchill CD, Rutledge LR, Wetmore SD. Effects of the biological backbone on stacking interactions at DNA-protein interfaces: the interplay between the backbone... π and π ... π components. *Phys Chem Chem Phys.* 12:14515–14526. DOI: 10.1039/c0cp00550a
- Duan G, Smith VH, Weaver DF. Characterization of Aromatic–Amide(Side-Chain) Interactions in Proteins through Systematic *ab Initio* Calculations and Data Mining Analyses. *The Journal of Physical Chemistry A.* 2000; 104:4521–4532. DOI: 10.1021/jp993381f
- Fischer M, Kloiber K, Hausler J, Ledolter K, Konrat R, Schmid W. Synthesis of a ¹³C-methyl-group-labeled methionine precursor as a useful tool for simplifying protein structural analysis by NMR spectroscopy. *Chembiochem.* 2007; 8:610–612. DOI: 10.1002/cbic.200600551 [PubMed: 17328009]
- Gardner KH, Rosen MK, Kay LE. Global folds of highly deuterated, methyl-protonated proteins by multidimensional. *NMR Biochemistry.* 1997; 36:1389–1401. DOI: 10.1021/bi9624806bi9624806 [PubMed: 9063887]
- Gelis I, et al. Structural basis for signal-sequence recognition by the translocase motor SecA as determined by NMR. *Cell.* 2007; 131:756–769. DOI: 10.1016/j.cell.2007.09.039 [PubMed: 18022369]
- Giorgi L, et al. NMR-based substrate analog docking to *Escherichia coli* peptidyl-tRNA hydrolase. *J Mol Biol.* 2011; 412:619–633. S0022-2836(11)00671-1. DOI: 10.1016/j.jmb.2011.06.025 [PubMed: 21718701]

- Hudson KL, Bartlett GJ, Diehl RC, Agirre J, Gallagher T, Kiessling LL, Woolfson DN. Carbohydrate-Aromatic Interactions in Proteins. *J Am Chem Soc.* 2015; 137:15152–15160. DOI: 10.1021/jacs.5b08424 [PubMed: 26561965]
- Hughes RC, McFeeters H, Coates L, McFeeters RL. Recombinant production, crystallization and X-ray crystallographic structure determination of the peptidyl-tRNA hydrolase of *Pseudomonas aeruginosa*. *Acta Crystallogr Sect F Struct Biol Cryst Commun.* 2012; 68:1472–1476. DOI: 10.1107/S1744309112045770
- Ikedo M. Towards bacterial strains overproducing L-tryptophan and other aromatics by metabolic engineering. *Appl Microbiol Biotechnol.* 2006; 69:615–626. DOI: 10.1007/s00253-005-0252-y [PubMed: 16374633]
- Ikedo M, Katsumata R. Metabolic Engineering To Produce Tyrosine or Phenylalanine in a Tryptophan-Producing *Corynebacterium glutamicum*. *Strain Appl Environ Microbiol.* 1992; 58:781–785. [PubMed: 16348670]
- Kasinath V, Valentine KG, Wand AJ. A ¹³C labeling strategy reveals a range of aromatic side chain motion in calmodulin. *J Am Chem Soc.* 2013; 135:9560–9563. DOI: 10.1021/ja4001129 [PubMed: 23767407]
- Kleckner IR, Foster MP. An introduction to NMR-based approaches for measuring protein dynamics. *Biochim Biophys Acta.* 2011; 1814:942–968. DOI: 10.1016/j.bbapap.2010.10.012 [PubMed: 21059410]
- Krishnarajuna B, Jaipuria G, Thakur A, D'Silva P, Atreya HS. Amino acid selective unlabeled for sequence specific resonance assignments in proteins. *J Biomol NMR.* 2011; 49:39–51. DOI: 10.1007/s10858-010-9459-z [PubMed: 21153044]
- Lange OF, et al. Recognition dynamics up to microseconds revealed from an RDC-derived ubiquitin ensemble in solution. *Science.* 2008; 320:1471–1475. DOI: 10.1126/science.1157092 [PubMed: 18556554]
- LeMaster DM, Cronan JE Jr. Biosynthetic production of ¹³C-labeled amino acids with site-specific enrichment. *J Biol Chem.* 1982; 257:1224–1230. [PubMed: 7035446]
- LeMaster DM, Richards FM. Preparative-scale isolation of isotopically labeled amino acids. *Anal Biochem.* 1982; 122:238–247. [PubMed: 7051891]
- Lichtenecker RJ. Synthesis of aromatic (¹³C)/(²H)-α-ketoacid precursors to be used in selective phenylalanine and tyrosine protein labelling. *Org Biomol Chem.* 2014; 12:7551–7560. DOI: 10.1039/c4ob01129e [PubMed: 25136818]
- Lichtenecker RJ, Weinhaupl K, Schmid W, Konrat R. α-Ketoacids as precursors for phenylalanine and tyrosine labelling in cell-based protein overexpression. *J Biomol NMR.* 2013; 57:327–331. DOI: 10.1007/s10858-013-9796-9 [PubMed: 24264768]
- Lundstrom P, et al. Fractional ¹³C enrichment of isolated carbons using [1-¹³C]- or [2-¹³C]-glucose facilitates the accurate measurement of dynamics at backbone Cα and side-chain methyl positions in proteins. *J Biomol NMR.* 2007; 38:199–212. DOI: 10.1007/s10858-007-9158-6 [PubMed: 17554498]
- Lutke-Eversloh T, Stephanopoulos G. Combinatorial pathway analysis for improved L-tyrosine production in *Escherichia coli*: identification of enzymatic bottlenecks by systematic gene overexpression. *Metab Eng.* 2008; 10:69–77. S1096-7176(07)00071-7. DOI: 10.1016/j.ymben.2007.12.001 [PubMed: 18243023]
- Maniatis, T., Sambrook, J. *Molecular Cloning: A Laboratory Manual*. 1. Cold Spring Harbor Laboratory; 1982.
- Madhusudan Makwana K, Mahalakshmi R. Implications of aromatic-aromatic interactions: From protein structures to peptide models. *Protein Science.* 2015; 24:1920–1933. DOI: 10.1002/pro.2814 [PubMed: 26402741]
- McCaldon P, Argos P. Oligopeptide biases in protein sequences and their use in predicting protein coding regions in nucleotide sequences. *Proteins.* 1988; 4:99–122. DOI: 10.1002/prot.340040204 [PubMed: 3227018]
- McFeeters H, Gilbert MJ, Thompson RM, Setzer WN, Cruz-Vera LR, McFeeters RL. Inhibition of essential bacterial peptidyl-tRNA hydrolase activity by tropical plant extracts. *Natural Product Communications.* 2012; 7:1107–1110. [PubMed: 22978241]

- McFeeters, RL. Backbone Chemical Shift Assignments and Dynamics Characteriation of an Extracellular Ligand-Binding Domain form an Iontoroic Glutamate Receptor. Cornell University; 2002.
- McFeeters RL, Altieri AS, Cherry S, Tropea JE, Waugh DS, Byrd RA. The high-precision solution structure of Yersinia modulating protein YmoA provides insight into interaction with H-NS. *Biochemistry*. 2007a; 46:13975–13982. DOI: 10.1021/bi701210j [PubMed: 18001134]
- McFeeters RL, et al. The novel fold of scytovirin reveals a new twist for antiviral entry inhibitors. *J Mol Biol*. 2007b; 369:451–461. DOI: 10.1016/j.jmb.2007.03.030 [PubMed: 17434526]
- Miyanoi Y, Ishida Y, Takeda M, Terauchi T, Inouye M, Kainosho M. Highly efficient residue-selective labeling with isotope-labeled Ile, Leu, and Val using a new auxotrophic *E. coli* strain. *J Biomol NMR*. 2016; 65:109–119. DOI: 10.1007/s10858-016-0042-0 [PubMed: 27272978]
- Oh MK, Rohlin L, Kao KC, Liao JC. Global expression profiling of acetate-grown *Escherichia coli*. *J Biol Chem*. 2002; 277:13175–13183. DOI: 10.1074/jbc.M110809200M110809200 [PubMed: 11815613]
- Rajesh S, Nietlispach D, Nakayama H, Takio K, Laue ED, Shibata T, Ito Y. A novel method for the biosynthesis of deuterated proteins with selective protonation at the aromatic rings of Phe, Tyr and Trp. *J Biomol NMR*. 2003; 27:81–86. 5139765. [PubMed: 12878843]
- Rittmann D, Lindner SN, Wendisch VF. Engineering of a glycerol utilization pathway for amino acid production by *Corynebacterium glutamicum*. *Appl Environ Microbiol*. 2008; 74:6216–6222. AEM.00963-08. DOI: 10.1128/AEM.00963-08 [PubMed: 18757581]
- Rosen MK, Gardner KH, Willis RC, Parris WE, Pawson T, Kay LE. Selective methyl group protonation of perdeuterated proteins. *J Mol Biol*. 1996; 263:627–636. DOI: 10.1006/jmbi.1996.0603 [PubMed: 8947563]
- Santos CN, Stephanopoulos G. Melanin-based high-throughput screen for L-tyrosine production in *Escherichia coli*. *Appl Environ Microbiol*. 2008; 74:1190–1197. AEM.02448-07. DOI: 10.1128/AEM.02448-07 [PubMed: 18156325]
- Schneider HJ. Binding mechanisms in supramolecular complexes. *Angew Chem Int Ed Engl*. 2009; 48:3924–3977. DOI: 10.1002/anie.200802947 [PubMed: 19415701]
- Shortle D. Assignment of amino acid type in 1H-15N correlation spectra by labeling with 14N-amino acids. *Journal of magnetic resonance Series B*. 1994; 105:88–90. [PubMed: 7921674]
- Sinha K, Jen-Jacobson L, Rule GS. Specific labeling of threonine methyl groups for NMR studies of protein-nucleic acid complexes. *Biochemistry*. 2011; 50:10189–10191. DOI: 10.1021/bi201496d [PubMed: 22039762]
- Sprenger GA. From scratch to value: engineering *Escherichia coli* wild type cells to the production of L-phenylalanine and other fine chemicals derived from chorismate. *Appl Microbiol Biotechnol*. 2007; 75:739–749. DOI: 10.1007/s00253-007-0931-y [PubMed: 17435995]
- Taylor-Creel K, Hames MC, Holloway WB, McFeeters H, McFeeters RL. Expression, purification, and solubility optimization of peptidyl-tRNA hydrolase 1 from *Bacillus cereus*. *Protein Expr Purif*. 2014; 95:259–264. S1046-5928(14)00009-6. DOI: 10.1016/j.pep.2014.01.007 [PubMed: 24480186]
- Teilum K, Brath U, Lundstrom P, Akke M. Biosynthetic 13C labeling of aromatic side chains in proteins for NMR relaxation measurements. *J Am Chem Soc*. 2006; 128:2506–2507. DOI: 10.1021/ja055660o [PubMed: 16492013]
- Thongchuang M, Pongsawasdi P, Chisti Y, Packdibamrung K. Design of a recombinant *Escherichia coli* for producing L-phenylalanine from glycerol. *World J Microbiol Biotechnol*. 2012; 28:2937–2943. DOI: 10.1007/s11274-012-1104-4 [PubMed: 22806734]
- Tong KI, Yamamoto M, Tanaka T. A simple method for amino acid selective isotope labeling of recombinant proteins in *E. coli*. *J Biomol NMR*. 2008; 42:59–67. DOI: 10.1007/s10858-008-9264-0 [PubMed: 18762866]
- Tugarinov V, Kay LE. Ile, Leu, and Val Methyl Assignments of the 723-Residue Malate Synthase G Using a New Labeling Strategy and Novel NMR Methods. *J Am Chem Soc*. 2003; 125:13868–13878. DOI: 10.1021/ja030345s [PubMed: 14599227]
- Wand AJ, Bieber RJ, Urbauer JL, McEvoy RP, Gan Z. Carbon relaxation in randomly fractionally 13C-enriched proteins. *J Magn Reson B*. 1995; 108:173–175. [PubMed: 7648015]

- Wang H, Janowick DA, Schkeryantz JM, Liu X, Fesik SW. A Method for Assigning Phenylalanines in Proteins. *Journal of the American Chemical Society*. 1999; 121:1611–1612. DOI: 10.1021/ja983897x
- Wang J, Liu L, Zhou H, Li J, Du G, Chen J. Comparative study of L-phenylalanine production in the growing and stationary phases during high cell density cultivation of an auxotrophic *Escherichia coli*. *Biotechnology and Bioprocess Engineering*. 2011; 16:916–922. DOI: 10.1007/s12257-011-0135-2
- Wilson RC, Edmondson SP, Platt JW, Helms K, Twigg PD. The E2-25K ubiquitin-associated (UBA) domain aids in polyubiquitin chain synthesis and linkage specificity. *Biochem Biophys Res Commun*. 405:662–666. S0006-291X(11)00132-X. DOI: 10.1016/j.bbrc.2011.01.089
- Zhang G, Neubert TA. Use of stable isotope labeling by amino acids in cell culture (SILAC) for phosphotyrosine protein identification and quantitation. *Methods in Molecular Biology*. 2009; 527:79–92. [PubMed: 19241007]
- Zhou H, Liao X, Liu L, Wang T, Du G, Chen J. Enhanced L-phenylalanine production by recombinant *Escherichia coli* BR-42 (pAP-B03) resistant to bacteriophage BP-1 via a two-stage feeding approach. *J Ind Microbiol Biotechnol*. 2010; 38:1219–1227. DOI: 10.1007/s10295-010-0900-9 [PubMed: 21104105]

Fig. 1.
Phenylalanine Biosynthesis and SSIL-Phenylalanine from 2-¹³C Glycerol

Fig. 2. Effect of Aromatic Amino Acid Biosynthesis Genes on Phenylalanine Production
Representative of all conditions tested, the relative production of phenylalanine is shown for the indicated genes. Standard deviations for triplicate measurements were less than $\pm 5\%$. To the right is a vector map of modified p2.274 showing where phenylalanine biosynthesis genes were inserted.

Fig. 3.
 ^{13}C Incorporation into SSIL-Phe

A) The non-constant time ^{13}C -HSQC spectrum of SSIL-Phe shows the existence of isolated ^1H - ^{13}C spin systems at the $\text{C}\alpha$ and $\text{C}\epsilon$ carbons with (red) and without (black) ^{13}C decoupling. The ^{13}C spectrum (left) and ^1H spectrum without ^{13}C decoupling (above) are shown. All phenylalanine peaks are labeled in the 1D spectra. **B)** GC-MS of derivatized natural abundance phenylalanine (left) and ^{13}C -SSIL-Phe (right). The mass of the fragment is indicated above.

Fig. 4. Non-constant Time ^{13}C -HSQC Spectra of Ubiquitin
Spectra of SSIL-Phe and u- ^{13}C Ubiquitin are shown for **A)** all carbons and **B)** aromatic carbons.

Fig. 5. ^{13}C Relaxation of Phenylalanines in Ubiquitin

R_1 (a, c) and $R_{1\rho}$ (b, d) were measured for ^1H - ^{13}C SSIL-Phe labeled ubiquitin. Data are shown for Phe4 (top) and Phe45 (bottom).

Fig. 6.
Non-constant time ^{13}C -HSQC Spectra of SSIL/ILV-methyl Ubiquitin

Table 1

Optimized M9+20 Media for SSIL-Phe Incorporation into Recombinantly Expressed Proteins in Bacteria.

Component	g/L
Sodium Phosphate Dibasic	6.7
Potassium Phosphate Monobasic	3
Sodium Chloride	0.5
Nucleotide Triphosphates (A,U,C,G)	0.04
Phenylalanine (isotope labeled)	0.4
Aspartate, Glutamate, Tyrosine, Tryptophan	0.4
Remaining 15 Amino Acids	0.2
Divalent ions, vitamins, and trace elements according to (T. Maniatis 1982)	