Published in final edited form as:

Mitochondrion. 2015 November ; 25: 87–97. doi:10.1016/j.mito.2015.10.004.

IL-15R α deficiency in skeletal muscle alters respiratory function and the proteome of mitochondrial subpopulations independent of changes to the mitochondrial genome

Grant C. O'Connell¹, Cody Nichols¹, Ge Guo¹, Tara L. Croston¹, Dharendra Thapa¹, John M. Hollander^{1,2}, and Emidio E. Pistilli^{1,2,3,*}

¹Division of Exercise Physiology, West Virginia University, Morgantown, WV.

²Center for Cardiovascular and Respiratory Sciences, West Virginia University, Morgantown, WV.

³Mary Babb Randolph Cancer Center, West Virginia University, Morgantown, WV.

Abstract

Interleukin-15 receptor alpha knockout (IL15R α KO) mice exhibit a greater skeletal muscle mitochondrial density with an altered mitochondrial morphology. However, the mechanism and functional impact of these changes have not been determined. In this study, we characterized the functional, proteomic, and genomic alterations in mitochondrial subpopulations isolated from the skeletal muscles of IL15R α KO mice and B6129 background control mice. State 3 respiration was greater in interfibrillar mitochondria and whole muscle ATP levels were greater in IL15R α KO mice supporting the increases in respiration rate. However, the state 3/state 4 ratio was lower, suggesting some degree of respiratory uncoupling. Proteomic analyses identified several markers independently in mitochondrial subpopulations that are associated with these functional alterations. Next Generation Sequencing of mtDNA revealed a high degree of similarity between the mitochondrial genomes of IL15R α KO mice and controls in terms of copy number, consensus coding and the presence of minor alleles, suggesting that the functional and proteomic alterations we observed occur independent of alterations to the mitochondrial genome. These data provide additional evidence to implicate IL-15R α as a regulator of skeletal muscle phenotypes through effects on the mitochondrion, and suggest these effects are driven by alterations to the mitochondrial proteome.

Keywords

myokine; mitochondrial subpopulations; mitochondrial genome; NGS; IL-15; IL-15R α

1. Introduction

Interleukin-15 (IL-15) is a ubiquitously expressed cytokine which has been primarily characterized as a critical T-cell survival factor within the context of the immune system (6,

*Corresponding Author: Emidio E. Pistilli, PhD, Division of Exercise Physiology, Department of Human Performance and Applied Exercise Science, Center for Cardiovascular and Respiratory Sciences, West Virginia University, Morgantown, WV 26506-9227, Phone (304) 293-0291, Fax (304) 293-7105, epistilli2@hsc.wvu.edu.

19, 55). IL-15 is subject to multifaceted regulation by interleukin-15 receptor alpha (IL-15R α), an IL-15 binding protein which not only modulates secretion of IL-15, but also the affinity of IL-15 for its receptor complex of interleukin-2 receptor beta (IL-2R β) and the common cytokine receptor gamma chain (γ c) (5, 17, 18, 34, 44, 47). Despite widespread tissue expression of IL-15, its receptor complex, and IL-15R α , the functions of IL-15 in non-immune tissues have only recently begun to be recognized. Evidence is accumulating which suggests that IL-15 plays a role as a myokine associated with oxidative skeletal muscle phenotypes (39, 41-43).

In the absence of the regulatory action of IL-15R α , global tissue IL-15R α knockout (IL15R α KO) mice display greater circulating levels of IL-15 (20, 43). Phenotypically, these mice display an increased capacity for exercise and *ex vivo* studies of muscle function have shown that fast muscles from IL15R α KO mice are resistant to fatigue (20, 39). Furthermore, there is evidence of a shift towards an oxidative muscle phenotype driven by alterations to the mitochondrial network. Muscles from the IL15R α KO mouse display a greater mitochondrial density, an observation supported by greater transcriptional activity of genes associated with mitochondrial biogenesis such as peroxisome proliferator-activated receptor gamma coactivator 1 alpha (PGC-1 α) and peroxisome proliferator-activated receptor delta (PPAR δ) (39, 40). Interestingly, a separate transgenic mouse model bred to over-express a highly secretable form of IL-15 within skeletal muscle also displays greater circulating levels of IL-15, increases in exercise capacity and similar molecular markers indicative of an oxidative shift (42). Collectively, these data demonstrate an association between increased circulating IL-15, mitochondrial biogenesis and oxidative skeletal muscle phenotypes in two strains of transgenic mice, implicating IL-15 and IL-15R α as targets for altering the mitochondrial network in skeletal muscle.

In addition to a greater mitochondrial density, high magnification electron micrographs of muscle tissue from IL15R α KO mice reveal an atypical mitochondrial structure. Specifically, mitochondria from the skeletal muscles of IL15R α KO mice present a swollen phenotype with sparse cristae. In support of this visual observation, mitochondrial internal complexity, a flow cytometric measure of cristae structure, is lower compared to mitochondria isolated from control muscles (40). It remains unclear whether these structural alterations which are observed with IL-15R α deficiency impact mitochondrial function. Mitochondrial swelling has been shown to occur in response to long duration endurance exercise (21, 32); it is possible that the mitochondria from muscles of IL15R α KO mice are fully functional, and that the structural changes in mitochondria are the result of the increased propensity these mice show for physical activity (20, 39). Alternatively, it is possible that the structural mitochondrial phenotype observed in the IL15R α KO mouse is indicative of pathology, as mitochondria in several mitochondrial disorders, many of which are driven by mtDNA mutations or deletion, appear qualitatively similar (3, 9, 16, 31, 52). Increases in mitochondrial biogenesis are seen in several of these disorders as a compensatory mechanism to rescue cellular ATP levels and prevent a bioenergetics crisis (57).

Therefore, the purpose of this project was to determine the functional status of mitochondria isolated from the skeletal muscles of IL15R α KO mice, as well as analyze the proteomic and genomic profiles associated with these mitochondria. These experiments were performed in

mitochondrial subpopulations extracted from directly under the sarcolemma (i.e. subsarcolemmal mitochondria; SSM) and from between the muscle myofibrils (i.e. interfibrillar mitochondria; IFM). Based on the phenotypic changes observed at the whole muscle level in the IL15R α KO mouse (39), as well as the alterations in mitochondrial density and morphology (40), we hypothesized that the respiratory capacity of the IFM mitochondrial subpopulation isolated from muscle of IL15R α KO mice would be greater compared to the IFM isolated from muscles of control mice, and that proteomic and genomic analyses would highlight markers in support of this respiratory phenotype. Polarography experiments revealed differences in several respiratory parameters measured in mitochondrial subpopulations isolated from muscles of IL15R α KO mice and controls, and multi-dimensional protein identification suggested that these changes are associated with altered expression of mitochondrial-associated proteins. However, analysis of mtDNA content and Next Generation Sequencing (NGS) of mtDNA did not reveal any genomic differences when comparing mitochondrial subpopulations from IL15R α KO mice to control mice, suggesting that the functional and proteomic alterations we observed occurred independent of changes to the mitochondrial genome. Collectively, these data provide additional evidence to implicate IL-15 and IL-15R α as regulators of skeletal muscle phenotypes through effects on the mitochondrial proteome.

2. Methods

2.1. Experimental Animals

IL15R α KO mice (stock no. 003723) and B6129SF2/J background control mice (stock no. 101045) were obtained from Jackson Laboratories (Bar Harbor, Maine) and housed at 22°C under a standard 12 hour light/dark cycle. The IL15R α KO is a global tissue KO mouse that was originally generated to determine the effects of IL15R α within the immune system (29). Standard rodent chow and water were provided *ad libitum*. Experimental procedures were performed using mice between 9 and 11 weeks of age, to maintain consistency with our previously published data (39, 40). All experimental procedures were submitted to and approved by the West Virginia University Institutional Animal Care and Use Committee.

2.2. Isolation of Mitochondrial Subpopulations

Mitochondrial subpopulations were isolated from fresh skeletal muscle using methods of Palmer et al. (36), with modifications (4, 11). Mice were anesthetized using 4% isoflurane, quadriceps muscles were dissected and immediately placed in ice cold Chappel-Perry buffer (100 mM KCl, 50 mM MOPS, 5 mM MgSO₄, 1 mM EGTA, 1 mM ATP, pH 7.4). Muscles were homogenized on ice using a Potter-Elvehjem PTFE pestle. Homogenates were spun at 800g for 10 min at 4°C; the resultant SSM-containing supernatant was collected and the remaining pellet was resuspended in KME buffer (100mM KCl, 50 mM MOPS, 0.5 mM EGTA). Trypsin digestion was then used to release IFM from the collected myofibrils by adding porcine trypsin at 1 mg·ml⁻¹ and incubating at 37°C with agitation for 5 minutes. Following addition of protease inhibitors, myofibril suspensions were spun at 800g for 10 min at 4°C to pellet debris and the IFM containing supernatant was collected. Supernatants were then spun at 8,000g at 4°C for 10 min to pellet mitochondria. Final SSM and IFM

mitochondrial pellets were then resuspended in an appropriate buffer for functional, proteomic or genomic experiments.

2.3. Analysis of Mitochondrial Respiration

For polarographic analysis, SSM and IFM originating from quadriceps muscles were resuspended in KME and mitochondrial protein concentrations were determined using the BCA assay (BioRad). A total of 50 μ g of mitochondrial protein was added to a 1 mL polarography chamber (Qubit Systems OX1LP) containing mitochondrial respiration buffer (80 mM KCL, 50 mM MOPS, 1 mM EGTA, 5 mM KH_2PO_4 , and 1 $\text{mg}\cdot\text{mL}^{-1}$ BSA). Glutamate and malate were then added at a final concentration of 5 mM and state 3 respiration was initiated by the addition of ADP at a final concentration of 250 nM. Oxygen concentration was plotted using LoggerPro data acquisition software (Vernier) and state 3 and state 4 respiration rates, respiratory control ratios, and ADP to oxygen ratios were calculated based on the acquired traces. Due to the necessity to use freshly isolated mitochondria, and the amount of time required to complete a single polarography run, mitochondria from one knockout and one control mouse were assessed each day for four consecutive days.

2.4. Whole muscle ATP determination

Whole quadriceps muscles were homogenized in RIPA buffer (ThermoFisher Scientific; Waltham, MA). Bioluminescent ATP content was analyzed in muscle homogenates from B6129 control and IL15R α KO mice utilizing an ATP Determination Kit, per manufacturer's instructions (ThermoFisher Scientific). Briefly, cellular ATP was determined on a luminescence plate reader and normalized to citrate synthase activity. Spectrophotometric analysis of citrate synthase enzymatic activity was determined in muscle homogenates utilizing a Citrate Synthase Assay Kit (ScienCell Research Laboratories; Carlsbad, CA) per manufacturer's instructions.

2.5. iTRAQ Labeling of Mitochondrial Subpopulations

Proteomic analyses were performed as previously described (4, 11). Briefly, pooled SSM from control mice (n = 5), pooled SSM from IL15R α KO mice (n = 5), pooled IFM from control mice (n = 5), and pooled IFM from IL15R α KO mice (n = 5) were lysed and precipitated overnight in acetone at -20°C and pellets were resuspended in 20 μL of 0.5 M triethylammonium bicarbonate (TEAB; pH 8.5). Protein contents were determined using a 2-D Quant Kit (Amersham, Piscataway, NJ) and 100 μg of each pooled sample was then denatured with 0.1% SDS and reduced with 5mM tris-(2-carboxyethyl) phosphine (TCEP). After incubation for 1 hour at 60°C , cysteines were blocked with 10 mM methyl methane thiosulfonate (MMTS) in isopropanol, and the samples were incubated at room temperature for 10 minutes. Subsequently, 10 μL of sequencing grade trypsin (Applied Biosystems, Foster City, CA) was added in a trypsin/protein ratio of 1:20, and the samples were incubated at 37°C overnight. Digested samples were labeled with the iTRAQ^{TM} reagents following the protocol provided by the vendor (Applied Biosystems, Foster City, CA).

After digestion and iTRAQ labeling, the ultra-complex protein digests were combined to create a 400 μg pooled protein digest sample that contained equal fractions of each of the

four labeled samples for subsequent Multi-dimensional Protein Identification Technology (MudPIT) analysis (28). After lyophilization, the digest mixture was reconstituted in strong cation exchange (SCX) loading buffer (5mM ammonium formate in 20% acetonitrile; pH 3.0) to be fractionated with SCX SpinTips (Protea Biosciences, Morgantown, WV) per the manufacturer's protocol. Briefly, the sample solution was loaded centrifugally onto the SCX SpinTip. The non-adsorbing fraction that passed through the SCX SpinTip was collected. Eight different elution solutions were used to fractionate the peptides (20, 60, 100, 150, 200, 250, 400, and 500 mM ammonium formate in 10% acetonitrile) in a step-wise manner, for a total of nine sample fractions. The collected fractions were lyophilized and reconstituted in a 0.1 M acetic acid solution, and then lyophilized to dryness. The fractions were then submitted for LC-MALDI TOF/TOF mass spectral analysis for protein identification, characterization, and differential expression analysis.

2.6. Mass Spectrometry (MS) Analyses of iTRAQ Labeled Mitochondrial Subpopulations

The LC-MALDI mass spectrometry system utilized was an ABI Tempo LC MALDI spotter with Tempo LC MALDI v.2.00.09 data acquisition and processing software. Lyophilized SCX sample fractions were reconstituted in LC aqueous run buffer (0.1% trifluoroacetic acid, 2% acetonitrile) and was injected onto a Zorbax C₁₈ chromatographic column, 150 × 0.3 mm (Agilent Technologies, Wilmington, DE). The peptides were eluted from the column using an acetonitrile/trifluoroacetic acid gradient (2-72% acetonitrile in 35 minutes) and spotted directly onto a MALDI plate in 6 second spot fractions. The MALDI spots were analyzed using an ABI 4800 MALDI TOF/TOF analyzer operated with 4000 Series Explorer software. The MS acquisition was in positive ion reflector mode with 400 laser shots per spectrum performed. The 15 strongest precursors per spot were chosen for MS/MS and the MALDI spot was interrogated until at least 4 peaks in the MS/MS, spectra achieved a S/N = 70.

The resulting MS/MS spectra were analyzed using ABI Protein ProteinPilot software 2.0 (Applied Biosystems, Foster City, CA). The spectral data were searched against the mouse protein database (NCBI nr.fasta database customized to select for all mouse proteins) for identification of the peptides and corresponding proteins. In ProteinPilot, the sample type was selected as iTRAQ 4Plex for retrieval of the isotopic tag information from the mass spectra. After database correlation analysis, the proteins were grouped, scored, and normalized against one to four isotope correction factors. The Pro Group algorithm of ProteinPilot generated a ProtScore that is a cumulative score from each of the peptides used by the algorithm in the protein identification. Protein scores (ProtScore) above 2.0, 1.0, and 0.47 expressed the percent confidence levels of >99, >90, and >66%, respectively. Each peptide match showed the iTRAQ isotopic labels and identified the differential protein expression level compared against the other iTRAQ labeled samples for relative quantification.

2.7. mtDNA Content Analysis

SSM and IFM derived from quadriceps muscles were resuspended in PBS and mitochondrial protein concentrations were assessed using the BCA assay. DNA was isolated from 100ug of mitochondrial protein using a commercially available blood and tissue DNA extraction kit

(Qiagen) with modifications. Briefly, mitochondria were incubated in 200 μ L of lysis buffer with proteinase K at 37°C for 3 hours with agitation. Following incubation, DNA was collected on silica micro-spin columns per manufacture instructions. DNA was then eluted from spin columns using 25 μ L of clean H₂O pre-warmed to 37°C. Real-time qPCR was then used to quantify copies of the *mt-Co2* mitochondrial gene in the eluent. For each sample, 1 μ L of eluent was added to a 20 μ L PCR reaction containing TaqMan primers for *mt-Co2* (Invitrogen). All samples were assayed in triplicate. Threshold cycle (CT) values were obtained, and delta-delta CT ($\Delta\Delta$ CT) values were calculated.

This technique was used as a surrogate marker for the mtDNA content of the isolated mitochondrial subpopulations within skeletal muscle samples. Because a linear relationship exists between the DNA content of starting material and DNA yield obtained using the DNA extraction kit, and DNA was isolated from a normalized amount of mitochondrial protein per sample, differences in CT values between samples should be indicative of differences in the amount of mtDNA per unit of mitochondrial protein. This unique approach to assessing mtDNA content in terms of mitochondrial protein was experimentally validated by assaying test mitochondrial lysates of varying mtDNA content created by combining DNase-treated mtDNA-null mitochondrial lysate with untreated mtDNA-containing mitochondrial lysate at different ratios. A strong linear relationship ($r^2=0.98$) was observed between CT value and test lysate mtDNA content (**Supplementary Figure 1**).

2.8. Next Generation Sequencing of mtDNA

Mitochondrial subpopulations derived from pooled tibialis anterior and extensor digitorum longus muscles were resuspended in PBS and mtDNA was isolated using the same method employed for assessment of mitochondrial DNA content. Real-time qPCR targeting both the mitochondrial *mt-Co2* gene and the nuclear *RN18s* gene was used to verify mtDNA enrichment and quantify levels of nuclear DNA contamination through comparison of the fold difference between *mt-Co2* and *RN18s*. For library construction, isolated mtDNA was quantified using a Qubit Fluorometer (Life Technologies). For each sample, 1 ng of DNA was used to construct a dual-indexed library via the Nextera XT DNA sample prep kit (Illumina). The subsequent libraries were quantified using qPCR and pooled in equimolar ratios. Multiplexed libraries were then sequenced using the Illumina MiSeq next generation sequencing platform; a total of 600 μ L of multiplexed library was added to the flow cell (MiSeq Reagent Kit V2, Illumina) at a final concentration of 8 pM and 500 cycles of sequencing were performed under standard run conditions. Sample reads were then de-multiplexed and aligned to the reference mitochondrial genome using Bowtie2 under default parameters. To account for nuclear DNA contamination, reads were also mapped to the nuclear genome and matching reads were subtracted from mitochondrial genome assemblies. Positional depth of assembled mitochondrial genomes was determined using the Samtools *mpileup* function. To determine whether loss of IL-15Ra resulted in site-specific mutations, VCFtools and custom perl scripts were used to screen for potential SNPs. For SNP analysis, SNPs were defined as a non-reference matching allele representing 5% or greater of the total number of reads at any given position given that the read depth at said position was greater than 10. Called SNPs were compared across experimental groups to determine if they constituted a group-specific phenomenon. In order to determine whether

loss of IL-15R α resulted in changes in total mutagenic stress, minor allele frequency (MAF) was quantified in a site-specific, gene-specific, and genome-wide manner. Positional MAF was quantified by counting the reads at each position that disagreed with the reference genome and normalizing to total reads on positional basis. MAF of individual mitochondrial genes was calculated as the ratio of non-reference matching reads to total reads within the gene of interest. Total genomic MAF was calculated as the ratio of non-reference matching reads to total number of reads across the genome.

2.9. Statistics

Mitochondrial respiratory data and mtDNA content data were analyzed using the GraphPad Prism 5.0 statistical software package. Mitochondrial respiratory data were analyzed using matched-paired t-tests, to account for day-to-day variability. MtDNA content data were analyzed using standard student t-tests. Proteomic and genomic data were analyzed using student t-tests or one-way ANOVA where appropriate. For one-way ANOVA, either Bonferonni or Tukey post hoc testing was performed based on the number of post-hoc comparisons. Data are presented as mean $\pm$ SE unless otherwise indicated, and the null hypothesis was rejected when $p < 0.05$.

3. Results

3.1. Analysis of mitochondrial respiration

Figure 1A displays representative polarography generated by IFM isolated from quadriceps muscles of IL15R α KO and B6129 control mice. IFM obtained from IL15R α KO mice displayed significantly greater state 3 respiration rates (**Figure 1B**) and consumed more total O₂ during state 3 respiration compared to controls (**Figure 1C**). No statistical difference was observed in the IFM state 4 respiration rate between strains (**Figure 1D**). The state 3 to state 4 respiration ratio (i.e. respiratory control ratio; RCR) was lower in IFM isolated from muscles of IL15R α KO mice as compared to controls, providing evidence of some degree of respiratory uncoupling (**Figure 1E**). **Figure 1F** displays representative polarography generated by SSM isolated from quadriceps muscles of IL15R α KO and B6129 control mice. No statistical differences were observed between strains in terms of state 3 respiration rate (**Figure 1G**), total state 3 O₂ consumption (**Figure 1H**), state 4 respiration rate (**Figure 1I**), or RCR (**Figure 1J**). However, the pattern of response of these parameters was consistent with what was observed in the IFM population. Collectively, these data demonstrate greater mitochondrial oxygen consumption with IL-15R α deficiency, although this increase in respiration is, at least partly, uncoupled from ATP production. In order to determine the effects of these functional alterations on ATP production, we assessed the ATP content of quadriceps muscles. Quadriceps muscles from IL15R α KO mice displayed greater ATP content than muscle obtained from controls, supporting the observed increases in respiration rate and indicating that the uncoupled respiratory phenotype does not significantly impair ATP production (**Figure 2**).

3.2. Proteomic alterations in mitochondrial subpopulations

Proteomic analysis revealed several proteins that were either up-regulated or down-regulated in the SSM and IFM of IL15R α KO mice relative to B6129 control mice (**Figure 3**). These

protein alterations were distinct between the two subpopulations, suggesting intrinsic differences between the SSM and IFM or that the subpopulations were exposed to different physiological conditions within the muscle. In these muscles, a greater number of altered proteins were observed in the SSM compared to the IFM.

3.2.1. Interfibrillar Mitochondria (IFM)—Proteins that were altered in the IFM are shown in **Figure 3A**. Enzymes involved in ketone body metabolism as well as beta oxidation of fatty acids were up-regulated in IFM from muscles of the IL15R α KO mouse compared to muscles from control mice. These included: isocitrate dehydrogenase NADP; acetyl-CoA acetyltransferase; methylmalonate semialdehyde dehydrogenase; and hydroxyacyl-CoA dehydrogenase. Mitochondrial proteins that are associated with mitochondrial respiratory capacity, mitochondrial biogenesis, and overall muscle oxidative capacity were up-regulated in the IFM from muscles of the IL15R α KO mouse compared to muscles from control mice. These included: GRP75 and O-acetyl ADP ribose acetylase. Apolipoprotein O, a protein known to associate with mitochondria to increase mitochondrial respiration and uncoupling and is associated with increased PGC-1 α expression, was also up-regulated in the IFM. Two proteins associated with ketone body metabolism and fatty acid oxidation were down-regulated in the IFM from muscles of IL15R α KO mice compared to muscles from control mice. These proteins were: isocitrate dehydrogenase NAD subunit gamma 1 and leucine rich PPR motif containing protein.

3.2.2. Subsarcolemmal Mitochondria (SSM)—Proteins that were altered in the SSM are shown in **Figure 3B**. Trap1 protein, the protein most up-regulated in the SSM, is a member of the Hsp90 family of chaperone proteins and has been suggested to act as a metabolic switch between mitochondrial respiration and aerobic glycolysis. There were also a number of proteins associated with fatty acid oxidation, ketone body metabolism and oxidative metabolism that were up-regulated in the SSM. These included: leucine rich PPR motif containing protein; mitochondrial pyruvate carrier 2; methylmalonate semialdehyde dehydrogenase; isocitrate dehydrogenase NADP; propionyl-CoA carboxylase beta chain; and CDGSH iron-sulfur domain containing protein 1. Proteins associated with ETC complexes of the mitochondria that were down-regulated in the SSM included: NADH dehydrogenase 1 alpha subcomplex subunit 6; sulfurtransferase; and cytochrome c oxidase subunit 6A2. NADH dehydrogenase 1 alpha subcomplex subunit 13, also known as GRIM19, is a protein involved in cross-talk between mitochondria and the innate immune system and was also down-regulated.

3.3. mtDNA Content Analysis

Quantitative PCR revealed no statistical differences in the number of copies of *mt-Co2* recovered from mitochondria between IL15R α KO mice and controls in terms of both SSM (**Figure 4A**) and IFM (**Figure 4B**), suggesting that loss of IL-15R α does not negatively impact mitochondrial DNA content. Interestingly, isolated IFM displayed greater levels of *mt-Co2* in comparison to isolated SSM (**Figure 4C**).

3.4. Next Generation Sequencing of mtDNA

Quantitative PCR analysis of mitochondrial *mt-Co2* and nuclear *m18s* in DNA isolated from our mitochondrial preparation indicated that our isolation procedure yielded DNA product enriched for mitochondrial DNA with low levels of genomic DNA contamination; DNA isolated from SSM lysates displayed a mean *m18s* to *mt-Co2* copy ratio of $1:560 \pm 98$ while DNA isolated from IFM lysates displayed *m18s* to *mt-Co2* copy ratio of $1:6876 \pm 2739$ (Mean $\pm$ SD). Sequencing generated 2×251 bp reads with median quality scores of greater than 30 across all read positions, indicating high read accuracy (**Figure 5A**). Positional depth analysis indicated strong mitochondrial genome coverage, with most groups exhibiting 200x coverage or greater across a majority of positions (**Figure 5B,C**). SNP analysis and hierarchical clustering based on genomic similarity revealed no SNPs constituting minor or major alleles which were inherent to any of the four experimental groups (**Figure 6A**). Taken as a whole, this suggests that loss of IL-15R α does not result in any site-specific pathologic mutations to the mitochondrial genome. In addition, it also suggests that the mtDNA of the IFM and the SSM are sequentially identical in terms of consensus coding. However, a comparison of minor alleles revealed a low degree of homogeneity between IFM and SSM samples. Specifically, $39 \pm 3.4\%$ of total positions carrying a minor allele failed to exhibit intra-animal homogeneity between the DNA of the IFM and SSM. This low degree of minor allele homogeneity between IFM and SSM DNA is evident when samples are hierarchically clustered by genomic similarity; IFM and SSM samples from the same animal were often relatively dissimilar in terms of the total pool of samples (**Figure 6A**). Another interesting observation was that there was a highly uneven distribution in terms of the substituted bases observed at polymorphic positions across all samples. A disproportionate number of polymorphic positions ($\approx 40\%$) contained a substituted adenine, while significantly less polymorphic positions carried substitutions of the other three nucleotides, with less than 10% of polymorphic positions carrying a guanine substitution. This pattern of substitution was similarly observed across both the SSM (**Figure 6B**) and the IFM (**Figure 6C**).

Analysis of total genomic MAF revealed no differences between IL15R α .KO mice and B6129 controls in either mtDNA originating from the SSM (**Figure 6D**) or the IFM (**Figure 6E**), suggesting that loss of IL-15R α does not result in changes in mutagenic stress. Furthermore, no differences were observed in terms of total genomic MAF between IFM and SSM samples, suggesting no differences in overall mutagenic stress between the two subpopulations (**Figure 6F**). Interestingly, analysis of positional MAF revealed four clusters of high variability which were consistently observed across the SSM (**Figure 7A, B**) and IFM (**Figure 7C, D**) of samples from IL15R α .KO and B6129 control mice, potentially indicating areas of the genome which are more susceptible to polymorphism. The first of these clusters fully spanned the *mt-Tf* gene which encodes for the mitochondrial phenylalanine transfer RNA. The second of these clusters was observed within the *mt-Rnr2* gene which encodes for mitochondrial 16s ribosomal RNA. The third of these high MAF clusters was observed within the *mt-Nd1* gene, which encodes for subunit one of NADH dehydrogenase of the electron transport chain. The remaining cluster spanned the *mt-Tw*, *mt-Ta*, *mt-Tn*, *mt-Tc*, and *mt-Ty* genes, which encode for the mitochondrial transfer RNAs for tryptophan, alanine, asparagine, cysteine, and tyrosine, respectively. Analyzing the MAF

of the individual mitochondrial genes allowed for statistical comparison of MAF on a gene by gene basis. Confirming the observations made in our positional analysis of MAF, the *mt-Tf*, *mt-Rnr2*, *mt-Nd1*, *mt-Tw*, *mt-Ta*, *mt-Tn*, *mt-Tc*, and *mt-Ty* genes all displayed elevated MAFs when analyzed statistically. **Figures 8A and B** display the mean MAFs calculated for every gene of the mitochondrial genome in terms of both SSM and IFM. The detailed results of the statistical analysis can be found in **Supplemental Table 1**. Interestingly, protein coding genes displayed relatively low MAFs with the exception of *mt-Nd1*, which displayed a significantly higher frequency of minor alleles in comparison to others protein coding genes, a phenomena which was observed in both the SSM (**Figure 8C**) and the IFM (**Figure 8D**).

4. Discussion

This study represents an extension of our prior observations on the unique exercise and skeletal muscle phenotype of the IL15R α KO mouse. Loss of IL-15R α in the global tissue IL15R α KO mouse is associated with substantial increases in skeletal muscle mitochondrial mass which are believed to facilitate a fatigue resistant muscle phenotype and enhanced exercise capacity (20, 39). Despite the fact that skeletal muscle appears to be oxidatively enhanced with IL-15R α deficiency, muscle mitochondria from these mice appear structurally altered (40), displaying a phenotype commonly observed in mitochondrial pathologies (3, 9, 16, 31, 52). These structural observations suggested to us the possibility that the increases in mitochondrial mass observed in the IL15R α KO mouse were a result of compensatory mitochondrial biogenesis in response to an IL-15R α -related mitochondrial impairment. Based on the phenotype changes observed at the whole muscle level in the IL15R α KO mouse (39), as well as the alterations in mitochondrial density and morphology (40), we hypothesized that the respiratory capacity of the IFM mitochondrial subpopulation isolated from muscle of IL15R α KO mice would be greater compared to the IFM isolated from muscles of control mice, and that proteomic and genomic analyses would highlight affected markers in support of this respiratory phenotype. Our polarographic analysis of mitochondrial respiration demonstrated greater State 3 respiration in isolated mitochondria from muscles of IL15R α KO mice, with evidence of some degree of mitochondrial uncoupling. Proteomic analysis of these mitochondrial subpopulations highlight alterations in nuclear encoded proteins that explain, in part, the functional data we obtained in the isolated mitochondria. However, NGS analyses and mtDNA content analysis suggested that these changes occurred independent of alterations to the mitochondrial genome.

Polarographic analysis of isolated mitochondria represent a method to quantify the maximum respiratory capacity of mitochondria when exposed to specific substrates. In our studies, we exposed isolated mitochondria to the substrates glutamate and malate, to drive the oxidative phosphorylation of ADP to ATP. Alterations in the respiratory capacity of SSM and IFM mitochondria from muscles of IL15R α KO mice were evident. We observed increased respiration rates in mitochondria obtained from skeletal muscles of IL15R α KO mice, along with greater whole muscle ATP levels. However, RER and total state 3 O₂ consumption values suggested that this increase in respiration rate was accompanied by some degree of respiratory uncoupling. Our data are highly consistent with previous reports of increased body temperature, elevated basal metabolic rates, and leaner body composition

in the IL15R α KO mouse, providing an organelle-level sub-cellular mechanistic explanation for these previous observations (20). Furthermore, our observations of respiratory uncoupling may also explain the increases in mitochondrial mass which have been previously reported in the IL15R α KO mouse (40), as respiratory uncoupling has been shown to promote mitochondrial biogenesis (7).

Our proteomic analysis was consistent with the respiratory phenotype we observed, and further supports the total body phenotype which has been previously reported (20). Numerous nuclear-encoded and mitochondrial-associated proteins involved in oxidative metabolism and fatty acid oxidation, ketone body metabolism and electron transport chain (ETC) complexes were altered independently in the two mitochondrial subpopulations, while none of the 13 mitochondrial-encoded proteins were altered. This phenomenon has been observed previously in other models analyzing isolated mitochondria (4), although this may also reflect the proteomic methodology utilized in the study and not necessarily a lack of alteration in this subset of proteins. In the IFM subpopulation, there were 8 proteins up-regulated with only 2 proteins down-regulated. In contrast, in the SSM subpopulation, there were 10 proteins up-regulated and 10 proteins down-regulated. Previous studies have demonstrated distinct proteomic alterations in the IFM and SSM subpopulations isolated from the heart in response to pathologic stimuli including heart failure, ischemia/reperfusion injury, diabetes and aging (4, 11-13, 15, 22, 46). These distinct alterations in the spatially separated mitochondrial subpopulations may reflect the biochemical differences between the IFM and SSM as well as the hypothesized functional roles. Specifically, ATP generated from IFM is thought to contribute directly to skeletal muscle contraction while ATP generated from SSM is thought to be utilized for active transport of metabolites across the sarcolemma (45, 49). While the proteomic signature of IFM from muscles of IL15R α KO mice support the exercise and muscle functional phenotype we previously observed (39), there were a greater number of altered proteins in the SSM subpopulation. Since both membrane-bound and soluble isoforms of IL-15R α exist (14), these data suggest that the membrane-bound isoform of IL-15R α may contribute to SSM function based on the spatial location of this subpopulation. This hypothesis warrants further evaluation.

In terms of specific protein alterations, Apolipoprotein O was significantly up-regulated in the IFM isolated from muscles of IL15R α KO mice compared to control. Apolipoprotein O localizes to the mitochondria and exists as a member of the multi-protein complex known as the mitochondrial inner membrane organizing system (MitOS) (23). Studies have demonstrated that apolipoprotein O is over-expressed in the hearts of diabetic patients (27) as well as in hearts from animals after 9 weeks of high-fat feeding (37). More recently, it has been shown that apolipoprotein O promotes mitochondrial uncoupling and is associated with greater expression of PPAR δ and PGC1 α in cardiomyocytes (54). This is consistent with the respiratory uncoupling we observed in the current study, as well as the increases in PPAR δ and PGC-1 α which we previously reported in the muscles from IL15R α KO mice (39). A second protein up-regulated in our proteomic analysis was GRP75, a member of the heat shock protein 70 family. GRP75 expression is associated with greater mitochondrial activity and biogenesis, and therefore, greater amounts of muscle activity (35, 50, 58). This is consistent with prior observations by us and others of increased spontaneous cage activity in the IL15R α KO mouse (20, 39). In addition, GRP75 expression is associated with the rate of

protein import through the inner mitochondrial membrane of the mitochondria. Studies have shown that up-regulation and down-regulation of GRP75 results in greater or lesser protein import, respectively (4, 10). Therefore, greater protein expression of GRP75 in the IFM isolated from muscles of IL15R α KO mice would suggest a greater rate of import of nuclear-encoded proteins into the mitochondria. Trap1 protein, a member of the heat shock protein 90 family, was the most up-regulated protein in the SSM and has been suggested to act as a metabolic switch between mitochondrial respiration and aerobic glycolysis (59). GRIM19 is a nuclear-encoded mitochondrial protein that associates with complex I of the electron transport chain and has been identified as a link between mitochondrial respiratory capacity and the innate immune system (8). Specifically, knockout of GRIM19 in mice is associated with a greater susceptibility to bacterial infection and lower pro-inflammatory cytokine production by macrophages (8). Recently, mutations of GRIM19 was associated with early onset hypotonia, dyskinesia and decrease in complex I enzyme activities in muscle biopsies (2). The fact that GRIM19 protein is altered in the mitochondria isolated from muscles of IL15R α KO mice provides additional evidence for a link between mitochondrial respiration and the immune system, although it is not clear as to why GRIM19 protein was greater in IFM and lower in SSM in the current study. Collectively, these data suggest that the exercise and muscle phenotypes reported in the IL15R α KO mouse are facilitated by the up-regulation of proteins which promote mitochondrial biogenesis, although at the expense of a greater degree of mitochondrial respiratory uncoupling.

Whether or not the respiratory and proteomic alterations we observed here, along with the structural alterations we have previously reported, are pathologic is not fully clear. However, our analysis provided little evidence beyond the observation of uncoupled respiration to suggest a pathological mechanism, and the higher rates of uncoupling we observed did not appear to negatively impact ATP production. It is possible that the changes observed in functional respiratory parameters are simply a result of normal adaptations to the increases in physical activity displayed by the IL15R α mouse (20, 39), as physical activity can induce similar respiratory changes to those which we observed in terms of respiration rates and respiratory uncoupling via upregulation of OXPHOS and uncoupling proteins, respectively (25, 26, 48, 53). In addition, it has recently been demonstrated that the IL15R α KO mouse preferentially uses fat as a fuel source (30). A preferential use of fats has been shown to stimulate greater expression of uncoupling proteins in skeletal muscle (38, 56), which provides further support to explain our polarographic data. Our proteomic analysis did not reveal any deficiencies or differential regulation of proteins involved in mitochondrial organization and structure which would suggest a pathological mechanism for the phenotype we previously observed (40). Furthermore, our genomic analysis revealed no evidence of defects to the mitochondrial genome which would be indicative of pathology. Thus, these data, along with previous reports of a fully functional muscle phenotype (39), suggest that the gains in mitochondrial mass observed in the muscles of the global IL15R α KO mouse are most likely not due to compensatory mitochondrial biogenesis in response to a pathological alteration in the mitochondrial network.

In addition to providing insights as to the effects of IL-15R α deficiency on mitochondrial function and proteomic composition, this project produced novel data regarding the mouse mitochondrial genome in terms of skeletal muscle. A limited number of studies have

analyzed the mouse mitochondrial genome using NGS, none of which addressed skeletal muscle tissue, with the most extensive of these studies analyzing mtDNA from the liver (1). We observed mutational frequencies which were generally consistent with what was reported in liver mtDNA in terms of genome-wide mutational frequency. We observed four areas of high mutational frequency in all of our experimental groups, potentially indicating sites which are more prone to mutagenic stress. Similar mutational hotspots were reported in the liver; however, they were differentially located throughout the genome (1). This discrepancy may suggest that mtDNA mutational hotspots may vary across tissues, and that post-developmental mutations may be differentially acquired in different tissue types. One of these hotspots we observed in this study fully spanned the *ND-1* gene, which codes for NADH dehydrogenase I. This protein is one of a number of proteins that comprise Complex I of the ETC, which is responsible for the first step in the electron transport process. Mutations within the *ND-1* gene are associated with numerous mitochondrial pathologies and Complex I deficiency is the most common ETC defect in human mitochondrial disorders (24, 51). The large mutational frequency in this area of the mitochondrial genome supports this observation, and suggests a greater likelihood for altered function of Complex I in response to pathological stimuli, compared to other complexes within the ETC. Conversely, it has been shown that ETC proteins, such as matrix arm subunits of Complex I are actually decreased in younger, longer living mice, and this is associated with improved Complex I assembly and greater State 3 oxygen consumption rates (33). These data suggest that alterations in protein content of ETC complexes is not necessarily associated with a reduced efficiency of ATP production. Our data revealed lower mutational frequencies in genes encoding mitochondrial proteins than in genes encoding transfer RNAs; this was also reported in the liver, although not to the same degree (1). This may suggest that protein coding genes are somehow protected from mutagenic stress to a greater degree than the rest of the mitochondrial genome. We observed a highly uneven distribution in terms of the substituted bases observed at polymorphic positions across all samples, potentially suggesting that some aspect of mtDNA synthesis results in a predisposition for the production of polymorphic alleles carrying specific substitutions.

To the authors' knowledge, this study also constitutes the first independent sequencing of mtDNA originating from the mitochondrial subpopulations of skeletal muscle. No significant differences were observed between IFM and SSM samples between the global IL15RαKO mouse and the B6129 control mouse in terms of genome-wide mutational frequency or location of highly variable regions, suggesting that despite the fact that these two subpopulations exist in two distinctly different environments, it is possible that they are exposed to similar environmental stressors and exhibit a similar susceptibility to mutation. Interestingly, we observed a low degree of homogeneity between the IFM and the SSM of individual mice, in terms of polymorphic minor alleles, providing evidence that mtDNA mutations originating in one population remain isolated, potentially due to the spatial distinction between the two populations. We also observed significantly higher *mt-Co2* levels in isolated IFM as opposed to SSM, irrespective of the mouse genotype, supporting the spatial distinction between the two mitochondrial subpopulations.

The purpose of this project was to determine the functional status of mitochondria isolated from the skeletal muscles of IL15RαKO mice, as well as analyze the proteomic and

genomic alterations that are associated with these mitochondria, in an effort to extend our prior observations on the unique exercise and skeletal muscle phenotype in these mice. Collectively, our data demonstrate greater mitochondrial respiration and a greater degree of respiratory uncoupling driven by alterations to the mitochondrial proteome in the absence of IL-15R α . These functional and proteomic changes do not appear to negatively impact ATP production, and loss of IL-15R α does not appear to alter the mitochondrial genome in a manner which would be indicative of a mitochondrial pathology. Therefore, the increases in mitochondrial density we have previously reported and the functional changes we observed in respiratory parameters here may be related to the greater exercise capacity observed in these mice (20, 39) and/or their preferential use of fats as a fuel source (30), and are not likely the result of compensatory mechanisms in response to a pathological impairment in mitochondrial network. However, as the IL15R α KO mouse is a global tissue knockout (29), and IL-15R α is widely expressed in a number of tissues (18), we cannot discount the effects cross-talk from non-muscle tissues on these alterations in skeletal muscle mitochondria. In addition to these insights regarding the implications of IL-15R α deficiency on the mitochondrion, we also provided a novel assessment of the mouse mitochondrial genome in skeletal muscle. Collectively, these data provide additional support for a role of IL-15 and IL-15R α in altering skeletal muscle phenotypes through its effects on the mitochondrion.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by the Genomics Discovery Grant at West Virginia University (P30 GM103503; E. Pistilli), the West Virginia University Research Funding Developmental Grant (E. Pistilli), and by the National Institutes of Health from the National Institutes of Diabetes and Digestive and Kidney Diseases [DP2DK083095; J.Hollander). The authors would like to acknowledge Drs. B. Teng and J. Mustafa for assistance with the real-time qPCR experiments performed in this manuscript, as well as the WVU Genomics Core Facility for assistance with acquisition and analysis of NGS data.

References

1. Ameer A, Stewart JB, Freyer C, Hagstrom E, Ingman M, Larsson NG, Gyllenstein U. Ultra-deep sequencing of mouse mitochondrial DNA: mutational patterns and their origins. *PLoS Genetics*. 2011; 7:e1002028. [PubMed: 21455489]
2. Angebault C, Charif M, Guegen N, Piro-Megy C, Mousson de Camaret B, Procaccio V, Guichet PO, Hebrard M, Manes G, Leboucq N, Rivier F, Hamel CP, Lenaers G, Roubertie A. Mutation in NDUFA13/GRIM19 leads to early onset hypotonia, dyskinesia and sensorial deficiencies, and mitochondrial complex I instability. *Human Molecular Genetics*. 2015; 24:3948–3955. [PubMed: 25901006]
3. Arbustini E, Diegoli M, Fasani R, Grasso M, Morbini P, Banchieri N, Bellini O, Dal Bello B, Pilotto A, Magrini G, Campana C, Fortina P, Gavazzi A, Narula J, Vigano M. Mitochondrial DNA mutations and mitochondrial abnormalities in dilated cardiomyopathy. *The American Journal of Pathology*. 1998; 153:1501–1510. [PubMed: 9811342]
4. Baseler WA, Dabkowski ER, Williamson CL, Croston TL, Thapa D, Powell MJ, Razunguzwa TT, Hollander JM. Proteomic alterations of distinct mitochondrial subpopulations in the type 1 diabetic heart: contribution of protein import dysfunction. *Am J Physiol Regul Integr Comp Physiol*. 2011; 300:R186–R200. [PubMed: 21048079]

5. Bergamaschi C, Jalah R, Kulkarni V, Rosati M, Zhang GM, Alicea C, Zolotukhin AS, Felber BK, Pavlakis GN. Secretion and biological activity of short signal peptide IL-15 is chaperoned by IL-15 receptor alpha in vivo. *J Immunol.* 2009; 183:3064–3072. [PubMed: 19696432]
6. Burton JD, Bamford RN, Peters C, Grant AJ, Kurys G, Goldman CK, Brennan J, Roessler E, Waldmann TA. A lymphokine, provisionally designated interleukin T and produced by a human adult T-cell leukemia line, stimulates T-cell proliferation and the induction of lymphokine-activated killer cells. *Proceedings of the National Academy of Sciences of the United States of America.* 1994; 91:4935–4939. [PubMed: 8197160]
7. Cerqueira FM, Laurindo FR, Kowaltowski AJ. Mild mitochondrial uncoupling and calorie restriction increase fasting eNOS, akt and mitochondrial biogenesis. *PloS One.* 2011; 6:e18433. [PubMed: 21483800]
8. Chen Y, Lu H, Liu Q, Huang G, Lim CP, Zhang L, Hao A, Cao X. Function of GRIM-19, a mitochondrial respiratory chain complex I protein, in innate immunity. *The Journal of Biological Chemistry.* 2012; 287:27227–27235. [PubMed: 22665480]
9. Chinnery PF, Turnbull DM. Mitochondrial DNA and disease. *Lancet (London, England).* 1999; 354(Suppl 1):S117–21.
10. Craig EE, Hood DA. Influence of aging on protein import into cardiac mitochondria. *The American Journal of Physiology.* 1997; 272:H2983–2988. [PubMed: 9227577]
11. Dabkowski ER, Baseler WA, Williamson CL, Powell M, Razunguzwa TT, Frisbee JC, Hollander JM. Mitochondrial dysfunction in the type 2 diabetic heart is associated with alterations in spatially distinct mitochondrial proteomes. *Am J Physiol Heart Circ Physiol.* 2010; 299:H529–540. [PubMed: 20543078]
12. Dabkowski ER, Williamson CL, Bukowski VC, Chapman RS, Leonard SS, Peer CJ, Callery PS, Hollander JM. Diabetic cardiomyopathy-associated dysfunction in spatially distinct mitochondrial subpopulations. *Am J Physiol Heart Circ Physiol.* 2009; 296:H359–369. [PubMed: 19060128]
13. Dabkowski ER, Williamson CL, Hollander JM. Mitochondria-specific transgenic overexpression of phospholipid hydroperoxide glutathione peroxidase (GPx4) attenuates ischemia/reperfusion-associated cardiac dysfunction. *Free radical biology & medicine.* 2008; 45:855–865. [PubMed: 18638546]
14. Dubois S, Magrangeas F, Lehours P, Raher S, Bernard J, Boisteau O, Leroy S, Minvielle S, Godard A, Jacques Y. Natural splicing of exon 2 of human interleukin-15 receptor alpha-chain mRNA results in a shortened form with a distinct pattern of expression. *The Journal of Biological Chemistry.* 1999; 274:26978–26984. [PubMed: 10480910]
15. Fannin SW, Lesnefsky EJ, Slabe TJ, Hassan MO, Hoppel CL. Aging selectively decreases oxidative capacity in rat heart interfibrillar mitochondria. *Archives of Biochemistry and Biophysics.* 1999; 372:399–407. [PubMed: 10600182]
16. Gilkerson RW, Margineantu DH, Capaldi RA, Selker JM. Mitochondrial DNA depletion causes morphological changes in the mitochondrial reticulum of cultured human cells. *FEBS Letters.* 2000; 474:1–4. [PubMed: 10828440]
17. Giri JG, Ahdieh M, Eisenman J, Shanebeck K, Grabstein K, Kumaki S, Namen A, Park LS, Cosman D, Anderson D. Utilization of the beta and gamma chains of the IL-2 receptor by the novel cytokine IL-15. *The EMBO Journal.* 1994; 13:2822–2830. [PubMed: 8026467]
18. Giri JG, Kumaki S, Ahdieh M, Friend DJ, Loomis A, Shanebeck K, DuBose R, Cosman D, Park LS, Anderson DM. Identification and cloning of a novel IL-15 binding protein that is structurally related to the alpha chain of the IL-2 receptor. *The EMBO Journal.* 1995; 14:3654–3663. [PubMed: 7641685]
19. Grabstein KH, Eisenman J, Shanebeck K, Rauch C, Srinivasan S, Fung V, Beers C, Richardson J, Schoenborn MA, Ahdieh M, et al. Cloning of a T cell growth factor that interacts with the beta chain of the interleukin-2 receptor. *Science (New York, NY).* 1994; 264:965–968.
20. He Y, Wu X, Khan RS, Kastin AJ, Cornelissen-Guillaume GG, Hsueh H, Robert B, Halberg F, Pan W. IL-15 receptor deletion results in circadian changes of locomotor and metabolic activity. *J Mol Neurosci.* 2010; 41:315–321. [PubMed: 20012227]

21. Hikida RS, Staron RS, Hagerman FC, Sherman WM, Costill DL. Muscle fiber necrosis associated with human marathon runners. *Journal of the Neurological Sciences*. 1983; 59:185–203. [PubMed: 6854349]
22. Hoppel CL, Tandler B, Parland W, Turkaly JS, Albers LD. Hamster cardiomyopathy. A defect in oxidative phosphorylation in the cardiac interfibrillar mitochondria. *The Journal of Biological Chemistry*. 1982; 257:1540–1548. [PubMed: 6460026]
23. Hoppins S, Collins SR, Cassidy-Stone A, Hummel E, Devay RM, Lackner LL, Westermann B, Schuldiner M, Weissman JS, Nunnari J. A mitochondrial-focused genetic interaction map reveals a scaffold-like complex required for inner membrane organization in mitochondria. *The Journal of Cell Biology*. 2011; 195:323–340. [PubMed: 21987634]
24. Kirby DM, Crawford M, Cleary MA, Dahl HH, Dennett X, Thorburn DR. Respiratory chain complex I deficiency: an underdiagnosed energy generation disorder. *Neurology*. 1999; 52:1255–1264. [PubMed: 10214753]
25. Korzeniewski B. Regulation of oxidative phosphorylation in different muscles and various experimental conditions. *The Biochemical Journal*. 2003; 375:799–804. [PubMed: 12901719]
26. Korzeniewski B, Zoladz JA. Training-induced adaptation of oxidative phosphorylation in skeletal muscles. *The Biochemical Journal*. 2003; 374:37–40. [PubMed: 12741955]
27. Lamant M, Smih F, Harmancey R, Philip-Couderc P, Pathak A, Roncalli J, Galinier M, Collet X, Massabuau P, Senard JM, Rouet P. ApoO, a novel apolipoprotein, is an original glycoprotein up-regulated by diabetes in human heart. *The Journal of Biological Chemistry*. 2006; 281:36289–36302. [PubMed: 16956892]
28. Lin D. Multidimensional Protein Identification Technology as an Effective Tool for Proteomics. *American Genomic/Proteomic Technology*. 2001; 1:38–46.
29. Lodolce JP, Boone DL, Chai S, Swain RE, Dassopoulos T, Trettin S, Ma A. IL-15 receptor maintains lymphoid homeostasis by supporting lymphocyte homing and proliferation. *Immunity*. 1998; 9:669–676. [PubMed: 9846488]
30. Loro E, Seifert EL, Moffat C, Romero F, Mishra MK, Sun Z, Krajacic P, Anokye-Danso F, Summer RS, Ahima RS, Khurana TS. Interleukin-15 Receptor alpha (IL15Ra) is a determinant of muscle fuel utilization and its loss protects against obesity. *American Journal of Physiology*. 2015:ajpregu 00505 02014.
31. Mandel H, Hartman C, Berkowitz D, Elpeleg ON, Manov I, Iancu TC. The hepatic mitochondrial DNA depletion syndrome: ultrastructural changes in liver biopsies. *Hepatology (Baltimore, Md)*. 2001; 34:776–784.
32. McCutcheon LJ, Byrd SK, Hodgson DR. Ultrastructural changes in skeletal muscle after fatiguing exercise. *J Appl Physiol* (1985). 1992; 72:1111–1117. [PubMed: 1568966]
33. Miwa S, Jow H, Baty K, Johnson A, Czapiewski R, Saretzki G, Treumann A, von Zglinicki T. Low abundance of the matrix arm of complex I in mitochondria predicts longevity in mice. *Nature Communications*. 2014; 5:3837.
34. Nishimura H, Fujimoto A, Tamura N, Yajima T, Wajjwalku W, Yoshikai Y. A novel autoregulatory mechanism for transcriptional activation of the IL-15 gene by a nonsecretable isoform of IL-15 generated by alternative splicing. *Faseb J*. 2005; 19:19–28. [PubMed: 15629891]
35. Ornatsky OI, Connor MK, Hood DA. Expression of stress proteins and mitochondrial chaperonins in chronically stimulated skeletal muscle. *The Biochemical Journal*. 1995; 311(Pt 1):119–123. [PubMed: 7575442]
36. Palmer JW, Tandler B, Hoppel CL. Biochemical properties of subsarcolemmal and interfibrillar mitochondria isolated from rat cardiac muscle. *The Journal of Biological Chemistry*. 1977; 252:8731–8739. [PubMed: 925018]
37. Philip-Couderc P, Smih F, Pelat M, Vidal C, Verwaerde P, Pathak A, Buys S, Galinier M, Senard JM, Rouet P. Cardiac transcriptome analysis in obesity-related hypertension. *Hypertension*. 2003; 41:414–421. [PubMed: 12623937]
38. Pilegaard H, Keller C, Steensberg A, Helge JW, Pedersen BK, Saltin B, Neufer PD. Influence of pre-exercise muscle glycogen content on exercise-induced transcriptional regulation of metabolic genes. *The Journal of Physiology*. 2002; 541:261–271. [PubMed: 12015434]

39. Pistilli EE, Bogdanovich S, Garton F, Yang N, Gulbin JP, Conner JD, Anderson BG, Quinn LS, North K, Ahima RS, Khurana TS. Loss of IL-15 receptor alpha alters the endurance, fatigability, and metabolic characteristics of mouse fast skeletal muscles. *The Journal of Clinical Investigation*. 2011; 121:3120–3132. [PubMed: 21765213]
40. Pistilli EE, Guo G, Stauber WT. IL-15Ralpha deficiency leads to mitochondrial and myofiber differences in fast mouse muscles. *Cytokine*. 2013; 61:41–45. [PubMed: 23116661]
41. Pistilli EE, Quinn LS. From Anabolic to Oxidative: Reconsidering the Roles of IL-15 and IL-15Ralpha in Skeletal Muscle. *Exercise and Sport Sciences Reviews*. 2013; 41:100–106. [PubMed: 23072822]
42. Quinn LS, Anderson BG, Conner JD, Wolden-Hanson T. IL-15 overexpression promotes endurance, oxidative energy metabolism, and muscle PPARdelta, SIRT1, PGC-1alpha, and PGC-1beta expression in male mice. *Endocrinology*. 2013; 154:232–245. [PubMed: 23161867]
43. Quinn LS, Anderson BG, Conner JD, Wolden-Hanson T, Marcell TJ. IL-15 is required for postexercise induction of the pro-oxidative mediators PPARdelta and SIRT1 in male mice. *Endocrinology*. 2014; 155:143–155. [PubMed: 24169546]
44. Ring AM, Lin JX, Feng D, Mitra S, Rickert M, Bowman GR, Pande VS, Li P, Moraga I, Spolski R, Ozkan E, Leonard WJ, Garcia KC. Mechanistic and structural insight into the functional dichotomy between IL-2 and IL-15. *Nature Immunology*. 2012; 13:1187–1195. [PubMed: 23104097]
45. Riva A, Tandler B, Loffredo F, Vazquez E, Hoppel C. Structural differences in two biochemically defined populations of cardiac mitochondria. *Am J Physiol Heart Circ Physiol*. 2005; 289:H868–872. [PubMed: 15821034]
46. Rosca MG, Vazquez EJ, Kerner J, Parland W, Chandler MP, Stanley W, Sabbah HN, Hoppel CL. Cardiac mitochondria in heart failure: decrease in respirasomes and oxidative phosphorylation. *Cardiovascular Research*. 2008; 80:30–39. [PubMed: 18710878]
47. Rubinstein MP, Kovar M, Purton JF, Cho JH, Boyman O, Surh CD, Sprent J. Converting IL-15 to a superagonist by binding to soluble IL-15R{alpha}. *Proceedings of the National Academy of Sciences of the United States of America*. 2006; 103:9166–9171. [PubMed: 16757567]
48. Schrauwen P, Hesselink M. Uncoupling protein 3 and physical activity: the role of uncoupling protein 3 in energy metabolism revisited. *The Proceedings of the Nutrition Society*. 2003; 62:635–643. [PubMed: 14692599]
49. Shimada T, Horita K, Murakami M, Ogura R. Morphological studies of different mitochondrial populations in monkey myocardial cells. *Cell and Tissue Research*. 1984; 238:577–582. [PubMed: 6525619]
50. Takahashi M, Chesley A, Freyssenet D, Hood DA. Contractile activity-induced adaptations in the mitochondrial protein import system. *The American Journal of Physiology*. 1998; 274:C1380–1387. [PubMed: 9612226]
51. Triepels RH, Van Den Heuvel LP, Trijbels JM, Smeitink JA. Respiratory chain complex I deficiency. *American Journal of Medical Genetics*. 2001; 106:37–45. [PubMed: 11579423]
52. Trimmer PA, Swerdlow RH, Parks JK, Keeney P, Bennett JP Jr. Miller SW, Davis RE, Parker WD Jr. Abnormal mitochondrial morphology in sporadic Parkinson's and Alzheimer's disease cybrid cell lines. *Experimental Neurology*. 2000; 162:37–50. [PubMed: 10716887]
53. Tsuboyama-Kasaoka N, Tsunoda N, Maruyama K, Takahashi M, Kim H, Ikemoto S, Ezaki O. Up-regulation of uncoupling protein 3 (UCP3) mRNA by exercise training and down-regulation of UCP3 by denervation in skeletal muscles. *Biochemical and Biophysical Research Communications*. 1998; 247:498–503. [PubMed: 9642158]
54. Turkieh A, Caubere C, Barutaut M, Desmoulin F, Harmancey R, Galinier M, Berry M, Dambrin C, Polidori C, Casteilla L, Koukoui F, Rouet P, Smih F. Apolipoprotein O is mitochondrial and promotes lipotoxicity in heart. *The Journal of Clinical Investigation*. 2014; 124:2277–2286. [PubMed: 24743151]
55. Waldmann TA, Tagaya Y. The multifaceted regulation of interleukin-15 expression and the role of this cytokine in NK cell differentiation and host response to intracellular pathogens. *Annual Review of Immunology*. 1999; 17:19–49.

56. Weigle DS, Selfridge LE, Schwartz MW, Seeley RJ, Cummings DE, Havel PJ, Kuijper JL, BeltrandelRio H. Elevated free fatty acids induce uncoupling protein 3 expression in muscle: a potential explanation for the effect of fasting. *Diabetes*. 1998; 47:298–302. [PubMed: 9519732]
57. Wenz T. PGC-1alpha activation as a therapeutic approach in mitochondrial disease. *IUBMB Life*. 2009; 61:1051–1062. [PubMed: 19859975]
58. Williamson CL, Dabkowski ER, Dillmann WH, Hollander JM. Mitochondria protection from hypoxia/reoxygenation injury with mitochondria heat shock protein 70 overexpression. *Am J Physiol Heart Circ Physiol*. 2008; 294:H249–256. [PubMed: 17982016]
59. Yoshida S, Tsutsumi S, Muhlebach G, Sourbier C, Lee MJ, Lee S, Vartholomaïou E, Tatokoro M, Beebe K, Miyajima N, Mohny RP, Chen Y, Hasumi H, Xu W, Fukushima H, Nakamura K, Koga F, Kihara K, Trepel J, Picard D, Neckers L. Molecular chaperone TRAP1 regulates a metabolic switch between mitochondrial respiration and aerobic glycolysis. *Proceedings of the National Academy of Sciences of the United States of America*. 2013; 110:E1604–1612. [PubMed: 23564345]

Figure 1. Effects of IL-15R α loss on mitochondrial respiration

(A) Representative polarography traces generated by IFM isolated from the skeletal muscle of IL15R α KO mice and B6129 controls. (B) State 3 respiration rate in the IFM subpopulation. (C) Total state 3 oxygen consumption in the IFM subpopulation. (D) State 4 respiration rate in the IFM subpopulation. (E) Respiratory control ratio values in the IFM subpopulation. (F) Representative polarography traces generated by SSM isolated from the skeletal muscle of IL15R α KO mice and B6129 controls. (G) State 3 respiration rate in the SSM subpopulation. (H) Total state 3 oxygen consumption in the SSM subpopulation. (I) State 4 respiration rate in the SSM subpopulation. (J) Respiratory control ratio in the SSM subpopulation. *, $p < 0.05$.

Figure 2. Total muscle ATP

Total ATP content of the quadriceps muscle normalized to citrate synthase activity. *, $p < 0.5$.

Figure 3. Proteomic alterations in isolated mitochondrial subpopulations

Multi-dimensional protein identification, using iTRAQ, was used to identify differentially regulated proteins in isolated mitochondrial subpopulations. (A) Differentially regulated proteins in isolated IFM from muscles of IL15R α KO mice relative to B6129 control mice. (B) Differentially regulated proteins in isolated SSM from muscles of IL15R α KO mice relative to B6129 control mice.

Figure 4. Effects of IL-15R α loss on the mtDNA content of isolated mitochondria
 (A) *mt-Co2* levels in the IFM subpopulations from IL15R α KO and B6129 mice. (B) *mt-Co2* levels in the SSM subpopulations from IL15R α KO and B6129 mice. (C) *mt-Co2* levels in the SSM and IFM subpopulations. *, $p < 0.05$.

Figure 5. NGS sample quality and read depth

(A) Median quality scores in samples across all read positions. (B) Positional depth analysis in all experimental groups across the mitochondrial genome. (C) Positional depth analysis of IFM and SSM subpopulations across the mitochondrial genome.

Figure 6. Next Generation Sequencing of sub-population mtDNA

(A) Hierarchical clustering of samples based on mtDNA similarity. (B) Percentage of substituted bases at polymorphic sites in SSM mtDNA. (C) Percentage of substituted bases at polymorphic sites in IFM mtDNA. (D) Genome-wide minor allele frequency between SSM mtDNA from IL15RaKO and B6129 mice. (E) Genome-wide minor allele frequency between IFM mtDNA from IL15RaKO and B6129 mice. (F) Genome-wide minor allele frequency between SSM and IFM mtDNA. *, $p < 0.05$.

Figure 7. Genomic distribution of highly variable regions of skeletal muscle mtDNA
 (A) Analysis of positional minor allele frequency in SSM mtDNA from muscles of B6129 control mice. (B) Analysis of positional minor allele frequency in SSM mtDNA from muscles of IL15RaKO mice. (C) Analysis of positional minor allele frequency in IFM mtDNA from muscles of B6129 control mice. (D) Analysis of positional minor allele frequency in IFM mtDNA from muscles of IL15RaKO mice.

Figure 8. Gene specific minor allele frequencies in the mtDNA across all genotypes
(A) Gene specific minor allele frequency of SSM. (B) Gene specific minor allele frequency of IFM. (C) Gene specific minor allele frequency of the protein coding genes in SSM mtDNA. (D) Gene specific minor allele frequency of the protein coding genes in IFM mtDNA. *, $p < 0.05$. Supplementary Figure 1. mtDNA content validation experiment. Linear relationship between the percent of mtDNA content and the CT values obtained using qPCR.