Published in final edited form as:

Bio Protoc. 2016 November 20; 6(22): . doi:10.21769/BioProtoc.2009.

Assay to evaluate BAL Fluid regulation of Fibroblast α -SMA Expression

Jennifer L. Larson-Casey¹ and A. Brent Carter^{1,2,*}

¹Department of Medicine, Division of Pulmonary, Allergy, and Critical Care Medicine, University of Alabama at Birmingham, Birmingham, AL, USA 35294

²Birmingham Veterans Administration Medical Center, Birmingham, AL, USA 35294

Abstract

Because transforming growth factor- β (TGF- β 1) induces differentiation of fibroblasts to myofibroblasts, we developed a protocol to evaluate alveolar macrophage-derived TGF- β 1 regulation of lung fibroblast differentiation (Larson-Casey *et al.*, 2016). The protocol allows evaluating the ability of mouse bronchoalveolar lavage (BAL) fluid to alter fibroblast differentiation. Fibroblast differentiation was measured by the expression of α -smooth muscle actin (α -SMA).

Background—Alveolar macrophages play an integral role in pulmonary fibrosis development by increasing the expression of TGF- β 1 (He *et al.*, 2011). Our prior data demonstrate that alveolar macrophages are a critical source of TGF- β 1 as mice harboring a conditional deletion of TGF- β 1 in macrophages were protected from pulmonary fibrosis (Larson-Casey *et al.*, 2016). The expression of α -SMA is a defining feature of myofibroblasts, and TGF- β 1 is a well-characterized pro-fibrotic mediator that induces transformation of fibroblasts to myofibroblasts both *in vitro* (Desmoulière *et al.*, 1993) and *in vivo* (Sime *et al.*, 1997). Prior studies exposed fibroblasts to recombinant TGF- β 1 to show its effect on differentiation and function (Horowitz *et al.*, 2007). Here we have developed a protocol for determining the ability of mouse BAL fluid to alter the differentiation of human lung fibroblasts to myofibroblasts, the cells that produce extracellular matrix proteins.

Materials and Reagents

1. 6-well Cell Culture Plates (Corning, Costar[®], catalog number: [3516](#))
2. RPMI 1640 Medium, No Phenol Red (Gibco, catalog number: [11835030](#))
3. FBS (Gibco, catalog number: [26140095](#))
4. DPBS (Gibco, catalog number: [14190144](#))
5. Penicillin-Streptomycin 10,000 U/ml (Gibco, catalog number: 15140122)
6. Amphotericin B (Gibco, catalog number: 15290018)
7. Normal human fibroblasts (IMR-90) (ATCC, catalog number: CCL-186)

*For correspondence: bcarter1@uab.edu.

8. α -SMA Antibody (American Research Products, catalog number: 03-61001)
9. β -actin Antibody (Sigma, catalog number: A5441)
10. Fibroblast culture media (see Recipes)
11. Lysis Buffer (see Recipes)

Equipment

1. Cell culture incubator, 37°C, 5% CO₂: 95% air atmosphere (Thermo Scientific, Forma Direct Heat CO₂ Incubator)

Procedure

1. Seed fibroblasts at a density of 1×10^6 per well of a 6-well cell culture plate in a total volume of 1 ml. Allow cells to adhere to cell culture plate (2–6 h).
2. Harvest bronchoalveolar lavage (BAL) fluid from mice (Han and Ziegler, 2013). Mice can be treated with bleomycin (1.75U/kg) or saline as a negative control. Determine the protein concentration of bronchoalveolar lavage (BAL) fluid. Using RPMI 1640 media, normalize all BAL samples to the same protein concentration in a total volume of 1 ml. Typically, 0.5–1 ml of BAL fluid is used and RPMI is used to normalize the protein concentration to a final total volume of 1 ml. Alternatively, this assay can be performed *in vitro* using conditioned media from macrophage cell lines or bone marrow derived macrophages. Harvest the conditioned media from macrophages post-transfection or post-treatment, using 1 million cells per 1 ml of media.
3. Remove media from fibroblasts and rinse with 1.5 ml room temperature $1 \times$ PBS, add normalized BAL fluid to fibroblasts.
 - a. Incubate fibroblasts with BAL fluid for 24 h in a cell culture incubator, 37°C, 5% CO₂: 95% air atmosphere.
 - b. Harvest fibroblasts by lysing cells in lysis buffer (see recipes)
4. Determine α -SMA expression by immunoblot analysis. The expression of α -SMA protein was determined by Western blotting using 20 μ g of total cellular protein. After blocking in 5% milk, the membranes were probed with mouse α -SMA primary antibody using a 1:3000 dilution in Tris buffered saline containing 0.1% Tween 20, the membranes were incubated with the anti-mouse secondary antibody (1:2000) in antibody dilution buffer for 1 h.

Data analysis

Results can be shown as a representative immunoblot (Figure 1); also see Larson-Casey *et al.*, 2016.

Recipes

1. Fibroblast culture media
DMEM
10% heat-inactivated FBS
100 U/ml penicillin- streptomycin
1.25 µg/ml amphotericin B (fungizone)
2. Lysis Buffer
1% NP-40
0.15 M NaCl
0.05 M Tris pH 7.4
Protease and phosphatase inhibitor tablets

Acknowledgments

This work was supported by 2R01ES015981 & VA merit review BX001135.

References

1. He C, Murthy S, McCormick ML, Spitz DR, Ryan AJ, Carter AB. Mitochondrial Cu, Zn-superoxide dismutase mediates pulmonary fibrosis by augmenting H₂O₂ generation. *J Biol Chem.* 2011; 286(17):15597–15607. [PubMed: 21393238]
2. Larson-Casey JL, Deshane JS, Ryan AJ, Thannickal VJ, Carter AB. Macrophage Akt1 Kinase-Mediated Mitophagy Modulates Apoptosis Resistance and Pulmonary Fibrosis. *Immunity.* 2016; 44(3):582–596. [PubMed: 26921108]
3. Desmoulière A, Geinoz A, Gabbiani F, Gabbiani G. Transforming growth factor-beta 1 induces alpha-smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. *J Cell Biol.* 1993; 122(1):103–11. [PubMed: 8314838]
4. Sime PJ, Xing Z, Graham FL, Csaky KG, Gauldie J. Adenovector-mediated gene transfer of active transforming growth factor-beta1 induces prolonged severe fibrosis in rat lung. *J Clin Invest.* 1997; 100(4):768–76. [PubMed: 9259574]
5. Horowitz JC, Rogers DS, Sharma V, White ES, Cui Z, Thannickal VJ. Combinatorial activation of FAK and AKT by transforming growth factor-β1 confers an anoikis-resistant phenotype to myofibroblasts. *Cell Signal.* 2007; 19(4):761–771. [PubMed: 17113264]
6. Han H, Ziegler SF. Bronchoalveolar Lavage and Lung Tissue Digestion. *Bio-protocol.* 2013; 3(16):e859. <http://www.bio-protocol.org/e859>. [PubMed: 27390755]

Figure 1.

Immunoblot analysis of α -SMA in IMR-90 fibroblasts cultured in BAL fluid from saline or bleomycin (Bleo) exposed WT mice