Published in final edited form as:

Curr Opin Behav Sci. 2017 April ; 14: 78–85. doi:10.1016/j.cobeha.2016.12.010.

From attachment to independence: Stress hormone control of ecologically relevant emergence of infants' responses to threat

Adrienne Santiago^{1,2,3,*}, Chiye Aoki³, and Regina M Sullivan^{1,2,3}

¹Emotional Brain Institute, Nathan Kline Institute, New York University Langone Medical Center, New York, NY 10003

²Child and Adolescent Psychiatry, New York University Langone Medical Center, New York, NY 10003

³Center for Neural Science, New York University, New York, NY 10003

Abstract

Young infant rat pups learn to approach cues associated with pain rather than learning amygdala-dependent fear. This approach response is considered caregiver-seeking and ecologically relevant within the context of attachment. With maturation, increases in the stress hormone corticosterone permit amygdala-dependent fear, which is crucial for survival during independent living. During the developmental transition from attachment to fear learning, maternal presence suppresses corticosterone elevation to block amygdala-dependent fear learning and re-engage the attachment circuitry. Early life trauma disrupts this developmental sequence by triggering a precocious increase of corticosterone, which permits amygdala-dependent threat responses. In this review, we explore the importance of the stress hormone corticosterone in infants' transition from complete dependence on the caregiver to independence, with consideration for environmental influences on threat response ontogeny and mechanistic importance of social buffering of the stress response.

Graphical Abstract

*Address correspondences to: Adrienne Santiago (ans525@nyu.edu).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

stress; corticosterone; amygdala; social buffering; development

Introduction

Pioneering research on infant-caregiver dyads, which began in the mid 1900's, highlights stress as a major mediator of infant caregiving quality/experiences and is critical in programming neurobehavioral development [1,2]. Importantly, it was the combined insights from clinical and basic scientists, with diverse research areas across many species, that linked disturbed maternal care/separation and compromised threat response functioning [1,3,4]. This concept of stress as a mediator between infant experiences and programming of neurobehavioral function is still prevalent today and has been documented in many diverse species [5–7].

The stress hormone corticosterone is typically thought of as damaging to infant behavioral and neural development [8,9]. However, it is now clear that corticosterone is also critically important for normal brain development and normal infant neurobehavioral functioning [10,11]. Here we review the effects of corticosterone within the context of normal development using infant rat fear/threat learning and expression as a framework. The behavioral neurobiology of threat response is relatively well-defined, especially in the rodent and is a useful template to explore how corticosterone modulates the neurodevelopment of fear/threat learning and expression. While differences exist between humans and rodents in the stage of neurodevelopment at birth [12], we focus here on the experience-dependent learning and how it shapes threat response behaviors that are phylogenetically conserved among species and provide an important bridge for translating rodent and nonhuman primate findings to humans' [6,13,14].

In adults of many diverse species, threat presentation will prompt a defensive threat response specific to the environment and threat intensity: in rats and humans, these responses can range from hiding, freezing, fleeing or attack. As we explore the development of threat response and its modulation by corticosterone, it is important to consider the ecological context of this behavior as the infant transitions from complete dependence to independence. Altricial animals, such as the rat and human, require extensive caregiving to survive. A reciprocal bond between the infant and caregiver, termed attachment, must be learned by both to maintain this close contact. Infants of most altricial species are physically incapable of defending themselves from predators. Accordingly, an infant of altricial species will typically seek the caregiver for protection, and only later, begin to attack a predator or freeze. This review will discuss the neurobiology that supports this ecologically and age-appropriate change in fear/threat response and how this developmental switch is controlled by corticosterone to produce adaptive behaviors during early development.

Ontogeny of innate fear expression: Modulation by corticosterone

In altricial species, very young infants confined to the nest depend entirely on their caregiver for protection. The expression of fear changes during maturation in ways that are appropriate

to the developmental stage and ecological niche of the animal: the infant rat pup does not freeze to predator odor until it begins to crawl out of the nest [15]. For a rat, these brief excursions begin around postnatal day (PN) 10 [15], at which point the amygdala becomes functionally integrated to support innate species-specific defensive responses to predator odor, such as freezing [16–19]. Initially, it was thought that this reflected maturation of the amygdala at PN10; this, however, turned out not to be the case. While amygdala development is protracted and continues through adolescence, major nuclei of the amygdala become visible and support plasticity days before the amygdala begins to support threat response behavior [20,21], provided sufficient levels of corticosterone are present in the amygdala [22,23]. The importance of maturation of the hypothalamic-pituitary-adrenal (HPA) axis and increased corticosterone levels in shaping the ontogeny of threat response was uncovered by Takahashi, who found that an increase in corticosterone in younger animals enables freezing to predator odor [24]. Adrenalectomy at PN10 prevents development of freezing behavior, which can be reinstated by delivering exogenous CORT (Figure 1) [18].

Thus, in order to understand the impact of corticosterone on the developing brain, it is important to first consider the ontogeny of the HPA axis, which undergoes considerable changes in most altricial species. The period of reduced stress-induced corticosterone release prior to the age of PN10 has been termed the ‘stress hyporesponsive period’ (SHRP). Gradual increase in basal corticosterone levels over the course of this period reaches a critical threshold at PN10 to terminate the SHRP and permit the amygdala to become active with exposure to predator odor [25]. The SHRP is observed at multiple levels of the HPA-axis, including blunted pituitary adrenocorticotrophic hormone (ACTH) secretion, decreased sensitivity to corticotropin-releasing hormone (CRH) and an adrenal gland hyporesponsive to circulating ACTH levels [26]. Thus, the stress hormone corticosterone plays two roles in defining the neurodevelopment of threat response: 1) gradual increases in endogenous corticosterone reach a critical threshold and 2) an acute threat (shock or predator odor) will produce an immediate increase in stimulus-evoked corticosterone. Together, this change in the stress system permits the switch to trigger a specific threat response.

Research from our lab and others expanded on Takahashi’s to show that either ontogenetic or experimentally manipulated changes in corticosterone level control whether the amygdala is activated by predator odor, as measured by c-Fos expression. Fear is expressed if the amygdala is functionally activated by corticosterone, while decreasing corticosterone results in suppressed amygdala activity and blocked fear expression [17,27]. Importantly, corticosterone acts as a switch with the power to activate the amygdala to permit fear expression. While circadian regulation of corticosterone causes fluctuations in baseline corticosterone levels throughout the day, importantly, this fluctuation in corticosterone does not appear to have a detectable impact on the control switch that induces amygdala activation and threat response, since threat responses occur regardless of the time of day.

Ontogeny of learned fear

In addition to innate, naturally occurring threats in the environment, animals also learn to tag stimuli in the environment with threat value, which provides a necessary substrate for

behavioral plasticity. Learning about threat is phylogenetically conserved and is supported by a relatively simplistic circuit that appears homologous across mammalian species. Threat learning is a rapid, robust form of classical conditioning, where a neutral conditioned stimulus (CS; e.g. tone or odor) is paired with an aversive unconditioned stimulus (US; e.g. electric shock). After temporal pairing of the CS-US, an associative link between CS and US causes the neutral CS to take on threat value [28,29]. Once a CS takes on the threat value, the animal will show defensive/fear responses to presentations of the CS, even in the absence of the US. These threat responses are typically similar in form to responses to predator odor.

As we explore the development of learned fear responses, we will highlight the role of corticosterone in supporting ecologically adaptive behaviors as the infant transitions from complete dependence on the caregiver to independence [15,16,30,31]. The ability to tag stimuli with threat value is absent during the SHRP until PN10 [15,30]. As is the case for innate fear expression discussed above, the ontogenetic delay is due to the failure of the amygdala to become functionally active in response to threat [16]. Indeed, in PN10 pups and older, pairing odor with moderate shock (0.5 mA to the tail or foot) will activate the amygdala and support fear/threat learning [16]. A causal link between fear/threat learning and the amygdala is revealed by experiments in which pharmacological suppression of the amygdala in older pups (PN 10) will block odor-shock threat/fear learning, although this manipulation has no effect on pups still too young to engage in amygdala-dependent fear learning [32]. Likewise, tetanic stimulation of the basolateral amygdala does not induce long-term potentiation in pups younger than PN10, in contrast to older pups [33]. Together, these studies suggest that activation of the amygdala is causal to support associative fear learning in pups older than PN10, thus representing the functional emergence of the amygdala in threat response at that time point.

As discussed previously, basal corticosterone levels increase over the course of the SHRP of postnatal development. By PN10, basal corticosterone increases sufficiently to reach a threshold for engaging learning plasticity within the amygdala during odor-shock conditioning [34]. Bilateral amygdala administration of corticosterone in PN8 pups will prematurely induce amygdala plasticity in response to odor-shock conditioning and support fear/threat learning [23]. However, in pups younger than PN6, the amygdala is too immature to support odor-shock conditioning, even with corticosterone injection into the amygdala. Conversely, intra-amygdala blockade of corticosterone receptors in older pups (PN10–15) is sufficient to suppress amygdala activation and abolish odor-shock fear/threat conditioning [23] (Figure 1). Thus, between the ages of PN6 and PN15, corticosterone is necessary and sufficient to support both amygdala plasticity and subsequent fear/threat learning.

During the SHRP of early development (PN<10), pups paradoxically prefer to approach a conditioned stimulus (e.g., neutral odor, such as peppermint) that has been paired with a painful stimulus [31,35–37]. Indeed, a shock-paired odor becomes powerful enough to support social interactions with the mother, including nipple attachment, which otherwise requires maternal odor. Neither failure of the pup to detect pain or varied threshold for pain explain this pain-induced odor preference [38,39]. Importantly, odor-shock conditioning in pups during the SHRP does not functionally activate the amygdala threat response circuitry. Instead, this paradoxical preference is under control of the unique neurobiological circuitry

for attachment learning: the olfactory bulb, anterior piriform cortex, and hyperfunctioning noradrenergic locus coeruleus (Figure 2) [40–42]. During the SHRP, when corticosterone levels are low, increases in norepinephrine (NE) are integral for forming pup-caregiver attachment [40]. Noradrenergic cell bodies located in the locus coeruleus send axons to innervate the mitral-tufted cells of the olfactory bulb with NE [43,44]. In young pups, this produces learned attachment-related odor preferences and learning-induced plasticity in the bulb and piriform cortex [45].

Ecological significance of the late emergence of threat learning

Why would it be advantageous for pups to have a delayed onset for fear/threat learning? For species that require parental care for survival, infants must learn rapid and robust attachment to their caregiver, regardless of the quality of care that they receive [46]. As stated above, abused pups paradoxically approach cues associated with maltreatment [47,48]. Importantly, odor-shock conditioning in pups during the SHRP of early development (PN<10) does not functionally activate the adult-like fear/threat circuit, but rather engages the age-dependent learning circuit critical for survival in young pups – the attachment neural system used by pups to learn about and attach to the caregiver. Typical maternal behaviors, such as pup retrieval or entering/leaving the nest, are at times associated with brief painful stimuli as the dam occasionally handles pups roughly or tramples them [46,48]. Pup survival depends on strong, learned attachment to the caregiver, and any associative learning such as odor-pain associations with the caregiver, could pose a threat to this bond. Thus, delayed onset of amygdala dependent CS-US learning may have evolved to prevent pups from avoiding the caregiver, regardless of the quality of care.

In summary, the stress hormone corticosterone has an adaptive role in defining pup's immediate neurobehavioral response. The gradual increase in endogenous corticosterone over early neurodevelopment together with the stimulus-evoked increase in corticosterone control the development of pup threat response behavior, thus ensuring adaptive and ecologically appropriate response to threat. This illustrates the important role of this stress hormone in producing adaptive behaviors [10].

Maternal social buffering of corticosterone response

Next, we will describe a system controlled by corticosterone that enables pups to rapidly and repeatedly switch back and forth between the immature attachment behavior of younger pups in the nest and the more mature freezing behavior of the older pups (Figure 3). Understanding the immediate effects of corticosterone on developing pups requires an understanding of how maternal presence modifies pups' corticosterone levels [5]. This is termed 'social buffering' and is observed in many species and across development [49–51]. Social buffering has robust beneficial effects by reducing pain and anxiety while enhancing healing in both humans and rats [5,52–56].

Previous work has shown that maternal presence is sufficient to block pain-induced increases in corticosterone [57]. Due to the critical role of corticosterone in turning on and off fear conditioning in pups, we explored whether naturalistic control of corticosterone by

social buffering could control rapid switching between fear/threat learning and attachment learning in pups. Indeed, our lab has found that mother's presence controls pups' shock-induced preference vs. aversion learning. As illustrated in Figure 3 we have termed this brief developmental period the Transitional Sensitive Period, which extends from PN10 through PN15 in typically developing pups [22]. Specifically, maternal absence during odor-shock conditioning in pups at PN10–15 functionally activates the amygdala and induces a strong aversion to the conditioned odor. However, if this same conditioning takes place in the presence of the mother, amygdala plasticity is suppressed, blocking threat learning and reactivating the locus coeruleus-dependent neural circuitry supporting attachment learning [22,32]. The mechanism for maternal suppression of fear learning is through social buffering as determined by microdialysis of the hypothalamic paraventricular nucleus (PVN): that is, maternal presence reduces pups' basal corticosterone levels via reduction of NE release into the pup's PVN, resulting in blockade of HPA activation [58]. This link between PVN NE and amygdala-dependent threat behaviors has been shown to be causal: blockade of PVN NE prevents maternal social buffering of HPA activation. Furthermore, maternal social buffering of the HPA axis can be overridden by a microinjection of NE receptor agonists into the PVN, which increases corticosterone in the mother's presence and permits the amygdala-dependent behavioral response to threat. More recently, these findings have been paralleled in humans and non-human primates [5,7,59].

However, social buffering is compromised in children reared with an abusive caregiver [6]. This has been modeled in non-human primates reared with a naturally abusive caregiver [7]. Rodent research from our lab has modeled abusive caregiving using the Scarcity Model of low resources, where the mother is provided with insufficient nest-building material. This manipulation produces abnormal maternal behaviors, including increases in rough handling or trampling of pups as well as reduced nurturing behavior, even though pups gain weight normally [47]. Across species, these models have confirmed that early life trauma from the caregiver degrades the mother's ability to socially buffer both corticosterone levels and threat response behavior, suggesting loss of some beneficial effects of social buffering [5,6,60]. Furthermore, pups reared with a stressed mother exhibit enhanced levels of corticosterone, arising both from corticosterone delivered through the mother's milk and endogenously generated corticosterone release by the pup [61]. These increases in corticosterone induced by early-life abuse prematurely end the SHRP, resulting in a precocious termination of the sensitive period for attachment learning [62] (Figure 4). Thus early life trauma alters the neurobehavioral trajectory of threat response development.

Social transmission of fear

However, complete suppression of fear in offspring is likely not adaptive in all situations and indeed, the mother is capable of overriding this social buffering suppression of fear. The mother is also capable of increasing pups' corticosterone and this highlights a critical role for social transmission of fear from the mother to infants [63,64]. Specifically, if the caregiver expresses fear of an odor in the presence of the pups, the pups will learn to fear that odor. The mechanism for this learning appears to be the mother's release of an alarm pheromone, which increases basal corticosterone level in pups and permits amygdala plasticity for fear learning of the odor. Thus, although the caregiver's presence at default

buffers stress and fear, the presence of the frightened caregiver induces fear in pups. Similar social learning occurs throughout development and has been well documented in adults [65,66].

Conclusion

We reviewed the role of the stress hormone corticosterone with an emphasis on new directions in our understanding of how this stress hormone has immediate impact on pup adaptive behavior. Furthermore, while the negative effects of stress are typically emphasized, it is also clear that normal increases in corticosterone over development can produce adaptive behaviors in pups. Here we gave examples of how corticosterone in the pup is controlled by the mother for adaptive expression of fear: socially buffering pups' corticosterone to decrease fear but also increasing pups' corticosterone via caregiver's alarm pheromone to increase fear. The caregiver's control of pup behavior and learning enables the caregiver to regulate pups' expression of adaptive behaviors, ultimately benefiting pups' survival.

Acknowledgments

This work was supported by the National Institutes of Health grants MH105846, EY13079 (CA), DC009910, MH091451, HD083217 (RMS). AS was supported by NIH training grants 2T32MH096331 and T32NS086750.

References

1. Bowlby, J. Attachment. Attachment and Loss: Vol. I. Loss. New York Basic Books; 1969.
2. Ainsworth, MDS. The development of infant-mother attachment. In: Ricciuti, BCH., editor. Review of child developmental research. Vol. 3. University of Chicago Press; 1973. p. 1-94.
3. Harlow, HF., Harlow, MK. The affectional systems. In: Schrier HFH, A., Stollnitz, F., editors. Behavior of nonhuman primates. Vol. 2. Academic Press; 1965. p. 287-344.
4. Levine S. Maternal and environmental influences on the adrenocortical response to stress in weanling rats. Science. 1967; 156:258–260. [PubMed: 6021047]
5. Hostinar CE, Sullivan RM, Gunnar MR. Psychobiological mechanisms underlying the social buffering of the hypothalamic-pituitary-adrenocortical axis: a review of animal models and human studies across development. Psychol Bull. 2014; 140:256–282. [PubMed: 23607429] This review provides a translational approach to understanding the effects of social buffering on the HPA axis, while identifying major gaps in both the human and rodent literature.
6. Gunnar MR. The effects of early life stress on neurobehavioral development in children and adolescents: Mediation by the HPA axis. Psychoneuroendocrinology. 2015; 61:3.
7. Sanchez MM, McCormack KM, Howell BR. Social buffering of stress responses in nonhuman primates: Maternal regulation of the development of emotional regulatory brain circuits. Soc Neurosci. 2015; 10:512–526. [PubMed: 26324227] This review of primate literature highlights the long-term and short-term effects of infant maltreatment on the mother's ability to socially buffer the infant.
8. McEwen BS. Understanding the potency of stressful early life experiences on brain and body function. Metabolism. 2008; 57(Suppl 2):S11–S15. [PubMed: 18803958]
9. Lupien SJ, McEwen BS, Gunnar MR, Heim C. Effects of stress throughout the lifespan on the brain, behaviour and cognition. Nat Rev Neurosci. 2009; 10:434–445. [PubMed: 19401723]
10. McEwen BS. Protective and damaging effects of stress mediators. N Engl J Med. 1998; 338:171–179. [PubMed: 9428819]

11. Wilson MA, Grillo CA, Fadel JR, Reagan LP. Stress as a one-armed bandit: Differential effects of stress paradigms on the morphology, neurochemistry and behavior in the rodent amygdala. *Neurobiol Stress*. 2015; 1:195–208. [PubMed: 26844236]
12. Semple BD, Blomgren K, Gimlin K, Ferriero DM, Noble-Haeusslein LJ. Brain development in rodents and humans: Identifying benchmarks of maturation and vulnerability to injury across species. *Prog Neurobiol*. 2013; 106–107:1–16.
13. Fareri DS, Tottenham N. Effects of early life stress on amygdala and striatal development. *Dev Cogn Neurosci*. 2016; 19:233–247. [PubMed: 27174149]
14. Graham AM, Buss C, Rasmussen JM, Rudolph MD, Demeter DV, Gilmore JH, Styner M, Entringer S, Wadhwa PD, Fair DA. Implications of newborn amygdala connectivity for fear and cognitive development at 6-months-of-age. *Dev Cogn Neurosci*. 2016; 18:12–25. [PubMed: 26499255]
15. Bolles RC, Woods PJ. The ontogeny of behaviour in the albino rat. *Animal Learning and Behavior*. 1964:427–441.
16. Sullivan RM, Stackenwalt G, Nasr F, Lemon C, Wilson DA. Association of an odor with activation of olfactory bulb noradrenergic beta-receptors or locus coeruleus stimulation is sufficient to produce learned approach responses to that odor in neonatal rats. *Behavioral Neuroscience*. 2000; 114:957–962. [PubMed: 11085610]
17. Moriceau S, Roth TL, Okotoghaide T, Sullivan RM. Corticosterone controls the developmental emergence of fear and amygdala function to predator odors in infant rat pups. *Int J Dev Neurosci*. 2004; 22:415–422. [PubMed: 15380840]
18. Takahashi LK. Organizing action of corticosterone on the development of behavioral inhibition in the preweanling rat. *Brain Res Dev Brain Res*. 1994; 81:121–127. [PubMed: 7805277]
19. Wiedenmayer CP, Barr GA. Developmental changes in c-fos expression to an age-specific social stressor in infant rats. *Behav Brain Res*. 2001; 126:147–157. [PubMed: 11704260]
20. Berdel B, Morys J, Maciejewska B, Dziewiatkowski J. Volume and topographical changes of the basolateral complex during the development of the rat's amygdaloid body. *Folia Morphol (Warsz)*. 1997; 56:1–11. [PubMed: 9409074]
21. Saul ML, Helmreich DL, Rehman S, Fudge JL. Proliferating cells in the adolescent rat amygdala: Characterization and response to stress. *Neuroscience*. 2015; 311:105–117. [PubMed: 26476262]
22. Upton KJ, Sullivan RM. Defining age limits of the sensitive period for attachment learning in rat pups. *Dev Psychobiol*. 2010; 52:453–464. [PubMed: 20583142]
23. Moriceau S, Wilson DA, Levine S, Sullivan RM. Dual circuitry for odor-shock conditioning during infancy: corticosterone switches between fear and attraction via amygdala. *J Neurosci*. 2006; 26:6737–6748. [PubMed: 16793881]
24. Takahashi LK, Rubin WW. Corticosteroid induction of threat-induced behavioral inhibition in preweanling rats. *Behav Neurosci*. 1993; 107:860–866. [PubMed: 8280395]
25. Moriceau S, Rainecki C, Holman JD, Holman JG, Sullivan RM. Enduring neurobehavioral effects of early life trauma mediated through learning and corticosterone suppression. *Front Behav Neurosci*. 2009; 3:22. [PubMed: 19750195]
26. Dallman MF. Early life stress: nature and nurture. *Endocrinology*. 2014; 155:1569–1572. [PubMed: 24746330]
27. Wiedenmayer CP, Magarinos AM, McEwen BS, Barr G. Mother lowers glucocorticoid levels of preweaning rats after acute threat. *Ann N Y Acad Sci*. 2003; 1008:304–307. [PubMed: 14998902]
28. Fanselow MS, LeDoux JE. Why we think plasticity underlying Pavlovian fear conditioning occurs in the basolateral amygdala. *Neuron*. 1999; 23:229–232. [PubMed: 10399930]
29. Janak PH, Tye KM. From circuits to behaviour in the amygdala. *Nature*. 2015; 517:284–292. [PubMed: 25592533]
30. Haroutunian V, Campbell BA. Emergence of interoceptive and exteroceptive control of behavior in rats. *Science*. 1979; 205:927–929. [PubMed: 472715]
31. Camp LL, Rudy JW. Changes in the categorization of appetitive and aversive events during postnatal development of the rat. *Dev Psychobiol*. 1988; 21:25–42. [PubMed: 3338626]
32. Moriceau S, Sullivan RM. Maternal presence serves as a switch between learning fear and attraction in infancy. *Nat Neurosci*. 2006; 9:1004–1006. [PubMed: 16829957]

33. Thompson JV, Sullivan RM, Wilson DA. Developmental emergence of fear learning corresponds with changes in amygdala synaptic plasticity. *Brain Res.* 2008; 1200:58–65. [PubMed: 18295751]
34. Barr GA, Moriceau S, Shionoya K, Muzny K, Gao P, Wang S, Sullivan RM. Transitions in infant learning are modulated by dopamine in the amygdala. *Nat Neurosci.* 2009; 12:1367–1369. [PubMed: 19783994]
35. Sullivan RM, Landers M, Yeaman B, Wilson DA. Good memories of bad events in infancy. *Nature.* 2000; 407:38–39.
36. Sullivan RM, Brake SC, Hofer MA, Williams CL. Huddling and independent feeding of neonatal rats can be facilitated by a conditioned change in behavioral state. *Dev Psychobiol.* 1986; 19:625–635. [PubMed: 3803730]
37. Sullivan RM, Hofer MA, Brake SC. Olfactory-guided orientation in neonatal rats is enhanced by a conditioned change in behavioral state. *Dev Psychobiol.* 1986; 19:615–623. [PubMed: 3803729]
38. Barr GA. Ontogeny of nociception and antinociception. *NIDA Res Monogr.* 1995; 158:172–201. [PubMed: 8594484]
39. Moriceau S, Roth TL, Sullivan RM. Rodent model of infant attachment learning and stress. *Dev Psychobiol.* 2010; 52:651–660. [PubMed: 20730787]
40. Sullivan RM, Wilson DA. The role of norepinephrine in the expression of learned olfactory neurobehavioral responses in infant rats. *Psychobiology (Austin, Tex).* 1991; 19:308–312.
41. Sullivan RM, Wilson DA, Leon M. Norepinephrine and learning-induced plasticity in infant rat olfactory system. *J Neurosci.* 1989; 9:3998–4006. [PubMed: 2585063]
42. Sullivan RM, Zyzak DR, Skierkowski P, Wilson DA. The role of olfactory bulb norepinephrine in early olfactory learning. *Brain Res Dev Brain Res.* 1992; 70:279–282. [PubMed: 1477962]
43. McLean JH, Shipley MT. Postnatal development of the noradrenergic projection from locus coeruleus to the olfactory bulb in the rat. *J Comp Neurol.* 1991; 304:467–477. [PubMed: 2022760]
44. McLean JH, Shipley MT, Nickell WT, Aston-Jones G, Reyher CK. Chemoanatomical organization of the noradrenergic input from locus coeruleus to the olfactory bulb of the adult rat. *J Comp Neurol.* 1989; 285:339–349. [PubMed: 2547851]
45. Sullivan RM. Developing a sense of safety: the neurobiology of neonatal attachment. *Ann N Y Acad Sci.* 2003; 1008:122–131. [PubMed: 14998878]
46. Hofer, MA., Sullivan, RM. *Towards a neurobiology of attachment.* Cumberland, RI: MIT Press; 2001.
47. Rainecki C, Moriceau S, Sullivan RM. Developing a neurobehavioral animal model of infant attachment to an abusive caregiver. *Biol Psychiatry.* 2010; 67:1137–1145. [PubMed: 20163787]
48. Roth TL, Sullivan RM. Memory of early maltreatment: neonatal behavioral and neural correlates of maternal maltreatment within the context of classical conditioning. *Biol Psychiatry.* 2005; 57:823–831. [PubMed: 15820702]
49. Hennessy MB, Kaiser S, Sachser N. Social buffering of the stress response: diversity, mechanisms, and functions. *Front Neuroendocrinol.* 2009; 30:470–482. [PubMed: 19545584]
50. van Rooij SJ, Cross D, Stevens JS, Vance LA, Kim YJ, Bradley B, Tottenham N, Jovanovic T. Maternal buffering of fear-potentiated startle in children and adolescents with trauma exposure. *Soc Neurosci.* 2016:1–10.
51. Doom JR, Doyle CM, Gunnar MR. Social stress buffering by friends in childhood and adolescence: Effects on HPA and oxytocin activity. *Soc Neurosci.* 2016:1–14. This study addresses the ontogeny of stress-buffering in humans by peers, and correlates results to salivary cortisol levels
52. Vogt JL, Coe CL, Levine S. Behavioral and adrenocorticoid responsiveness of squirrel monkeys to a live snake: is flight necessarily stressful? *Behav Neural Biol.* 1981; 32:391–405. [PubMed: 7197154]
53. Hennessy MB. Presence of companion moderates arousal of monkeys with restricted social experience. *Physiol Behav.* 1984; 33:693–698. [PubMed: 6522489]
54. Kirschbaum C, Klauer T, Filipp SH, Hellhammer DH. Sex-specific effects of social support on cortisol and subjective responses to acute psychological stress. *Psychosom Med.* 1995; 57:23–31. [PubMed: 7732155]

55. Heinrichs M, Baumgartner T, Kirschbaum C, Ehlert U. Social support and oxytocin interact to suppress cortisol and subjective responses to psychosocial stress. *Biol Psychiatry*. 2003; 54:1389–1398. [PubMed: 14675803]
56. Taylor SE, Burklund LJ, Eisenberger NI, Lehman BJ, Hilmert CJ, Lieberman MD. Neural bases of moderation of cortisol stress responses by psychosocial resources. *J Pers Soc Psychol*. 2008; 95:197–211. [PubMed: 18605860]
57. Stanton ME, Levine S. Inhibition of infant glucocorticoid stress response: specific role of maternal cues. *Dev Psychobiol*. 1990; 23:411–426. [PubMed: 2253818]
58. Shionoya K, Moriceau S, Bradstock P, Sullivan RM. Maternal attenuation of hypothalamic paraventricular nucleus norepinephrine switches avoidance learning to preference learning in preweanling rat pups. *Horm Behav*. 2007; 52:391–400. [PubMed: 17675020]
59. Gee DG, Gabard-Durnam L, Telzer EH, Humphreys KL, Goff B, Shapiro M, Flannery J, Lumian DS, Fareri DS, Caldera C, et al. Maternal buffering of human amygdala-prefrontal circuitry during childhood but not during adolescence. *Psychol Sci*. 2014; 25:2067–2078. [PubMed: 25280904]
This study is the first to demonstrate that early life trauma is associated with developmentally accelerated connectivity between the amygdala and medial prefrontal cortex.
60. Sullivan RM, Perry RE. Mechanisms and functional implications of social buffering in infants: Lessons from animal models. *Soc Neurosci*. 2015; 10:500–511. [PubMed: 26324338]
61. Yeh KY. Corticosterone concentrations in the serum and milk of lactating rats: parallel changes after induced stress. *Endocrinology*. 1984; 115:1364–1370. [PubMed: 6541120]
62. Moriceau S, Shionoya K, Jakubs K, Sullivan RM. Early-life stress disrupts attachment learning: the role of amygdala corticosterone, locus ceruleus corticotropin releasing hormone, and olfactory bulb norepinephrine. *J Neurosci*. 2009; 29:15745–15755. [PubMed: 20016090]
63. Debiec J, Sullivan RM. Intergenerational transmission of emotional trauma through amygdala-dependent mother-to-infant transfer of specific fear. *Proc Natl Acad Sci U S A*. 2014; 111:12222–12227. [PubMed: 25071168] This study is the first to show that pairing of an odor with the odor released by a fearful mother is sufficient to induce amygdala-dependent threat learning in her pups.
64. Chang DJ, Debiec J. Neural correlates of the mother-to-infant social transmission of fear. *J Neurosci Res*. 2016; 94:526–534. [PubMed: 27091313]
65. Inagaki H, Kiyokawa Y, Tamogami S, Watanabe H, Takeuchi Y, Mori Y. Identification of a pheromone that increases anxiety in rats. *Proc Natl Acad Sci U S A*. 2014; 111:18751–18756. [PubMed: 25512532]
66. Kiyokawa Y. Social Odors: Alarm Pheromones and Social Buffering. *Curr Top Behav Neurosci*. 2015

Highlights

- Altricial infants' threat-induced caregiver approach is ecologically relevant.
- Infant approach to cues associated with threat is protection-seeking behavior
- Older pups switch protection seeking approach to amygdala-dependent defense
- Corticosterone controls neurobehavioral switch from protection seeking to defense
- Maternal presence lowers corticosterone to block defense and engage attachment

Figure 1. Corticosterone control of amygdala-dependent threat response and threat learning in development

During the stress hyporesponsive period (SHRP) that occurs prior to PN10, corticosterone (CORT) levels are low and the amygdala is unresponsive to predator odor as well as odor cue conditioning. This result is also observed in older pups that have received adrenalectomy, and is reversed by injecting exogenous CORT. In pups older than PN10, the amygdala is responsive to predator odor and supports fear conditioning. This is also observed in pups as young as PN6, if exogenous corticosterone is injected into the amygdala.

Figure 2. Corticosterone acts as a control switch between attachment circuitry and amygdala-dependent threat response

In pups younger than PN10, high levels of norepinephrine (NE) are secreted by the locus coeruleus (LC) into the olfactory bulb (OFB), thus activating learning-induced changes in the OFB and anterior piriform cortex (aPir) to support attachment to the caregiver. As corticosterone (CORT) levels increase at PN10, the OFB will respond to threats likely by co-activation of the posterior piriform cortex (pPir) and amygdala to produce an independent defensive response, such as freezing. Between PN10–15, maternal presence will socially buffer threats to the pup, thus reducing pup corticosterone, suppressing amygdala activity and eliminating the independent defensive response.

STRESS HORMONE CORT DETERMINES SWITCH IN ADAPTIVE BEHAVIORS

Transitional Sensitive Period in pups from 10-15 days old

Figure 3. Maternal control of corticosterone determines rapid and responsive switch in adaptive threat response behaviors during the Transitional Sensitive Period (PN10–15)

Pup corticosterone (CORT) levels in response to threat are under maternal control between PN10–PN15. In the presence of a mother that perceives no threat, pups corticosterone levels are low and attachment circuitry is active: norepinephrine (NE) is released by the locus coeruleus (LC) to produce the paradoxical behavior wherein pups approach a conditioned odor stimulus paired with shock. If the mother is fearful, she emits an alarm pheromone, which increases pup corticosterone and activates the amygdala dependent threat-response circuitry: the pup will then avoid conditioned odor stimulus paired with shock. This corticosterone induced amygdala dependent threat response is also observed when pups are alone.

Figure 4. Early life abuse increases pup corticosterone and induces premature closure of the sensitive period for attachment learning

In the Scarcity Model of early life abuse, mother rats are provided with insufficient nesting material for five days of early post-natal development (PN8–12). In response, the mother is more likely to scatter, trample or roughly transport her pups. Notably, pup corticosterone levels increase more rapidly in response to the maltreatment, which triggers early (~PN7) termination of the stress hyporesponsive period (SHRP), which is the sensitive period for attachment learning. Early onset of the Transitional Sensitive Period observed in the Scarcity Model is associated with reduced maternal buffering of pup corticosterone response to threat stimuli.