Published in final edited form as:

J Phys Chem B. 2015 July 16; 119(28): 8781–8792. doi:10.1021/acs.jpcb.5b05354.

The Effect of the Methylation and N-H Acidic Group on the Physicochemical Properties of Imidazolium-Based Ionic Liquids

Ana S.M.C. Rodrigues[†], Marisa A. A. Rocha[†], Hugo F. D. Almeida[§], Catarina M. S. S. Neves[§], José A. Lopes-da-Silva[‡], Mara G. Freire[§], João A. P. Coutinho[§], and Luís M. N. B. F. Santos^{†,*}

[†]Centro de Investigação em Química, Departamento de Química e Bioquímica, Faculdade de Ciências da Universidade do Porto, R. Campo Alegre 687, P-4169-007 Porto, Portugal

[§]CICECO - Aveiro Institute of Materials, Chemistry Department, University of Aveiro, Campus de Santiago, 3810-193 Aveiro, Portugal

[‡]QOPNA Unit, Departamento de Química, Universidade de Aveiro, 3810-193 Aveiro, Portugal

Abstract

This work presents and highlights the differentiation of the physicochemical properties of the [C₁Him][NTf₂], [C₂Him][NTf₂], [C₁²C₁Him][NTf₂], and [C₄²C₁³C₁im][NTf₂] that are related with the strong bulk interaction potential, which highlights the differentiation on the physicochemical arising from the presence of the acidic group (N-H) as well as the methylation in position 2 (C(2)) of the imidazolium ring. Densities, viscosities, refractive indices and surface tensions in a wide range of temperatures, as well as, isobaric heat capacities at 298.15 K, for this IL series are presented and discussed. It was found that the volumetric properties are barely affected by the geometric and structural isomerization, following a quite regular trend. A linear correlation between the glass transition temperature, *T*_g, and the alkyl chain size was found; however, ILs with the acidic N-H group present a significant higher *T*_g than the [C_{N-1}³C₁im][NTf₂] and [C_N³C_Nim][NTf₂] series. It was found that the most viscous ILs, ([C₁Him][NTf₂], [C₂Him][NTf₂] and [C₁²C₁Him][NTf₂]) have an acidic N-H group in the imidazolium ring in agreement with the observed increase of energy barrier of flow. The methylation in position 2, C(2), as well as, the N-H acidic group in the imidazolium ring, contribute to a significant variation in the cation-anion interactions and their dynamics, which is reflected in their charge distribution and polarizability leading to a significant differentiation of the refractive indices, surface tension and heat capacities. The observed differentiation of the physicochemical properties of the [C₁Him][NTf₂], [C₂Him][NTf₂], [C₁²C₁Him][NTf₂], and [C₄²C₁³C₁im][NTf₂] are an indication of the stronger bulk interaction potential, which highlights the effect that arises from the presence of the acidic group (N-H) as well as the methylation in position 2 of the imidazolium ring.

*Corresponding author: Luís M N B F Santos (lbsantos@fc.up.pt).

Keywords

Thermophysical properties; Thermal behavior; Isomerization; acidic hydrogen bond; methylation; short alkyl

1 Introduction

There is an increasing interest in Ionic Liquids (ILs) due to their unusual physical and transport properties which result from their peculiar type of cohesive interactions, charge distribution and nanostructuration. The molecular structure and supramolecular organization of an ionic liquid is complex, comprising polar and non-polar domains. This structural heterogeneity leads to nanostructuration in the bulk which was already observed both theoretically and experimentally.^{1–9} Among their unique properties, the high thermal and chemical stabilities, negligible vapour pressure at room temperature, high ionic conductivity, and improved solvation ability makes them good candidates in a wide variety of applications in the chemical industry, as well as, models for the academic understanding of the structural features and interionic interactions in the bulk phase. Most of the studies reported in the literature were focused on the effect of the alkyl chain length and the chemical nature of the ion pairs on the thermophysical properties. There are some work in literature that focus on the effect of the structural isomerism of ionic liquids and how this can affect their thermodynamic properties,^{10–15} however there is a lack of understanding of the effect of important features like the methylation and N-H acidic bond on physicochemical properties.

The variation in the substituent groups at the cation is found to have a drastic impact in the charge distribution and accessibility⁸. As a consequence, the anion-cation interaction potential can vary significantly. A case to point is the cation methylation in the position 2, C(2), of the imidazolium ring. This structural feature has been investigated by several groups, both experimentally and theoretically, in order to understand its effect on the thermophysical properties and transport properties.^{16–21} This structural modification has been shown to induce a change in the melting temperature, thermal stability, viscosity, surface tension, and heat capacities. The experimental evidence published by Bonhôte et al.¹⁶ showed that the methylation in C(2) of the imidazolium ring of NTf₂-based ionic liquids increases the viscosity by a factor of 2.6, relative to the hydrogenated cation based IL. They also found that the methylation in the C(4) and C(5) positions of the imidazolium have a minor effect¹⁶. Later on, Hunt¹⁷ used quantum chemical calculations to investigate the effect of the C(2) methylation in the viscosities of imidazolium-based ionic liquids. They found that the increase in the viscosities and melting points is explained by the decrease in melting entropy of the methylated cations, due to a reduction of the number of stable conformers (cation-anion interactions) leading to a decrease of the absolute liquid entropy. Additionally, the free rotation on the butyl chain is restricted by the steric bulk methyl, which limits the position of the anion around the cation. Noack et al.¹⁸ observed the great influence of the C(2) position on the electron density distribution of the molecular structure and on the macroscopic behavior of imidazolium-based ILs, using vibrational and NMR spectroscopy. According to Fumino et al.,^{19,20} the suppression of the hydrogen bonds formed in C(2) position upon methylation enhances the overall Coulomb interactions

between anion and cation. They found that the hydrogen bond formed in C(2)-H disrupted the ionic network by altering the charge symmetry of the ions and thus fluidize imidazolium ILs. These results were supported experimentally by IR spectroscopy where a red-shift and an intensity change of the bands were observed. A theoretical study by Izgorodina et al.²¹ revealed that there is a restriction in the anion movement around the methylated C(2) cation relative to the hydrogenated one with a potential energy barrier exceeding 40 kJ·mol⁻¹. Another structural feature that is scarcely addressed in the literature is the acidic hydrogen present in N-H of the imidazolium ring. The ILs comprising this acidic character are more likely to establish hydrogen bonds with the anions and with solvents, such as water.¹⁰

In this work, we explore the structural isomerization effect on the properties of the imidazolium NTf₂-based ILs. Heat capacity, density, viscosity, refractive index, surface tension and thermal behavior (glass transition, crystallization temperatures/profile, melting temperature, enthalpies and entropies of fusion) are here studied. The change of a group position (such as a methyl group), as well as, the presence of the acidic N-H group in the ionic liquid were used as an *in situ* probe to explore the molecular effects on the overall ionic liquid interaction potential, which will be reflected in their physicochemical properties. Table 1, presents the list and abbreviation of the studied ILs.

2 Experimental Section

2.1 Materials and purification

The twelve ILs samples were acquired from IOLITEC with the following state purity: [¹C₁Him][NTf₂] (1-methylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 353239-08-4), >98 %; [¹C₂Him][NTf₂] (1-ethylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 353239-10-8), >98 %; [¹C₁²C₁Him][NTf₂] (1,2-dimethylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 353239-12-0), 98%; [¹C₁³C₁im][NTf₂] (1,3-dimethylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 174899-81-1), 99%; [¹C₂³C₁im][NTf₂] (1-ethyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 174899-82-2), >99%; [¹C₃³C₁im][NTf₂] (1-methyl-3-propylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 216299-72-8), 99%; [¹C₂³C₂im][NTf₂] (1,3-diethylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 174899-88-8), >99%; [¹C₂³C₃im][NTf₂] (1-ethyl-3-propylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 347882-21-7), >99%; [¹C₄³C₁im][NTf₂] (1-butyl-3-methylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 174899-83-3), >99%; [¹C₅³C₁im][NTf₂] (1-methyl-3-pentylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: n.a., >99%; [¹C₃³C₃im][NTf₂] (1,3-dipropylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: n.a., >99%; [¹C₄²C₁³C₁im][NTf₂] (1-butyl-2,3-dimethylimidazolium bis(trifluoromethylsulfonyl)imide, CAS RN: 350493-08-2, 99%). The relative atomic masses used in this work were those recommended by the IUPAC Commission in 2007.²²

The commercial ILs samples were purified under vacuum (0.1 Pa) at moderate temperature (323 K) and constant stirring for 48 hours in order to remove traces of volatile impurities. The water mass fraction content was determined in a 151 Metrohm 831 Karl Fischer coulometer, using a Hydranal-152 Coulomat AG from Riedel-de Haën. The water content, in

all ILs, was below 100 ppm. This process was performed systematically, before any experimental measurement.

2.2 Thermal behavior

Temperatures and the standard molar enthalpies of fusion for the ILs were measured in a power compensation differential scanning calorimeter, PERKIN ELMER model Pyris Diamond DSC, using hermetically sealed aluminum crucibles with a constant flow of nitrogen ($50 \text{ mL}\cdot\text{min}^{-1}$). Samples of about 15 mg were used in each experiment. The temperature and heat flux scales of the power compensation DSC were calibrated by measuring the temperature and the enthalpy of fusion of reference materials,^{23,24} namely benzoic acid, 4-methoxybenzoic acid, triphenylene, naphthalene, anthracene, 1,3,5-triphenylbenzene, diphenylacetic acid, perylene, *o*-terphenyl and 9,10-diphenylanthracene, at different scanning rates (2, 5 and $10 \text{ K}\cdot\text{min}^{-1}$). Each IL sample was previously heated above the melting temperature for 30 minutes, followed by a quenching step consisting of a fast cooling ($\sim 50 \text{ K}\cdot\text{min}^{-1}$) until 173 K. This procedure avoids the crystallization on cooling and promotes the glass formation. After that, the ILs samples were heated ($5 \text{ K}\cdot\text{min}^{-1}$) to promote crystallization followed by cooling ($\sim 50 \text{ K}\cdot\text{min}^{-1}$) and heating cycles ($5 \text{ K}\cdot\text{min}^{-1}$) in the crystallization region, exceeding the glass transition and below the temperature of melting to assure complete crystallization before the melting. A final scan at $5 \text{ K}\cdot\text{min}^{-1}$ rate was performed to determine the temperatures and enthalpies of the solid-solid and isotropization phase transitions.

2.3 Densities and Viscosities

The density, ρ , and viscosity, η , for the $[\text{C}_1\text{Him}][\text{NTf}_2]$, $[\text{C}_2\text{Him}][\text{NTf}_2]$, $[\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$ and $[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ ILs were measured using an automated SVM 3000 Anton Paar rotational Stabinger viscometer–densimeter. The measurements were carried out at 0.1 MPa in the temperature range from (278.15 to 363.15) K. Only the $[\text{C}_1\text{Him}][\text{NTf}_2]$ and $[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ ILs, which are solids at room temperature, were measured in a narrower temperature range of (323.15 to 363.15) and (303.15 to 363.15) K, respectively. For each ionic liquid, at least two independent measurements were performed using the same experimental conditions and different samples. The apparatus was calibrated using the three standard calibration samples, APN7.5, APN26 and APN415 in the same experimental conditions of the ionic liquid measurements. The reproducibility of the viscosity and density measurements is, according to the manufacturer, 0.35 % and $\pm 0.5 \text{ kg}\cdot\text{m}^{-3}$, respectively from 288.15 to 378.15 K and the uncertainty of temperature is within $\pm 0.02 \text{ K}$. Further details regarding the equipment and method are available in the literature.^{25,26}

2.4 Heat Capacities

The heat capacities at $T = 298.15 \text{ K}$ of the $[\text{C}_1\text{Him}][\text{NTf}_2]$, $[\text{C}_2\text{Him}][\text{NTf}_2]$, $[\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$ and $[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ ILs were measured by a high-precision heat capacity drop calorimeter, described in the literature.^{27–30} The calorimeter was calibrated with water and sapphire ($\alpha\text{-Al}_2\text{O}_3$).²³ The calibration constant was found to be $\epsilon = (6.6040 \pm 0.0036) \text{ W}\cdot\text{V}^{-1}$. The accuracy of the apparatus for measurements of heat capacities of liquids and solids was evaluated before, using benzoic acid, hexafluorobenzene, *p*-terphenyl and $[\text{C}_6^3\text{C}_1\text{im}][\text{NTf}_2]$.²⁹ The ampoules were weighted in a Mettler Toledo AG245 dual range

analytical balance (sensitivity of 1×10^{-6} g and repeatability of 2×10^{-6} g) both empty and after filling with the ionic liquid. All the uncertainties are given as twice of the standard deviation of the mean, and include the calibration uncertainty. The buoyancy effect correction was considered both for the calibration and experiments of the ILs.

2.5 Refractive indices

The refractive indices for all the ILs presented in Table 1 were measured at the sodium D-line using a Bellingham model RFM340 refractometer ($\pm 3 \times 10^{-5}$ stated precision), as a function of temperature. The refractometer features a presser with a seal ring made of fluoropolymer Kalrez® which is closed over the sample on the prism preventing/decreasing the IL sample contact with water and atmospheric gases in particular, CO₂. The presser incorporates a micro flow cell, which is used to introduce the sample into the refractometer, without opening it to the atmosphere in order to avoid moisture and other gases contamination of the ILs samples. For the studied samples no time drift was detected along the measurements. The presser and the internal prism water jacket assembly is temperature controlled by an external bath through the presser hinge and integral channels in the presser arm. The temperature in the refractometer cell is controlled using an external thermostatic bath within a temperature fluctuation of ($\pm 5 \times 10^{-3}$) K, measured with a resolution better than 1×10^{-3} K and an uncertainty within ± 0.02 K. The apparatus was calibrated with degassed water (Millipore quality) and toluene (Spectranal, 99.9 %). Samples were directly introduced into the flow cell (prism assembly) using a syringe; the flow cell was kept closed after sample injection. For each ionic liquid at least two independent experiments were performed and in each experiment at least three measurements were taken at each temperature. The refractive indices were measured with respect to air and no corrections were applied.

2.6 Surface Tension

The surface tension of each ionic liquid sample was determined by the analysis of the shape of a pendant drop and measured using a Dataphysics (model OCA-20) contact angle system. Drop volumes of (9 ± 0.5) μ L were obtained using a Hamilton DS 500/GT syringe connected to a Teflon coated needle placed inside an aluminium air chamber able to maintain the temperature of interest within ± 0.1 K. The surface tension measurements were performed in the temperature range from (298.15 to 343.15) K, with the exception of [¹C₁Him][NTf₂] and [¹C₁²C₁Him][NTf₂], which was performed in the temperature range from (328.15 to 343.15) K and (303.15 to 343.15) K, respectively, due to their higher melting temperature. The temperature of the ionic liquid in each surface tension measurement was considered to be the same than as measured inside the aluminium chamber with a Pt100 within ± 0.1 K placed at a distance of approximately 20 mm from the liquid drop. After reaching a specific temperature inside the aluminium chamber, the measurements were carried out after 40 min to guarantee the thermal stabilization. Silica gel was kept inside the air chamber to maintain a dry environment.

For the surface tensions determination at each temperature, and for each ionic liquid, at least 5 drops were formed and analysed. For each drop, an average of 150 images was captured. The analysis of the drop shape was done with the software modules SCA 20 where the

gravitational acceleration ($g = 9.8018 \text{ m}\cdot\text{s}^{-2}$) and latitude ($\text{lat.} = 40^\circ$) were used according to the location of the assay. The surface tensions were calculated using the measured density data. Further details on the equipment and its validity to measure surface tensions of ILs were previously addressed.^{31–33}

3 Results and Discussion

3.1 Thermal behavior

The experimental results of the onset temperatures of glass, cold crystallization and (solid-solid and fusion) transitions are presented in Table 2 along with some available literature data for comparison. The enthalpy of transitions were obtained by numerical integration of the peak thus obtained. Estimation errors of $\pm 0.5 \text{ K}$ for the onset temperature and $\pm 0.7 \text{ kJ}\cdot\text{mol}^{-1}$ for enthalpies were assumed taking into account the combined uncertainty of the calibration and the ILs experiments. The values found in the literature concerning the phase behavior of the ionic liquids here studied are scarce and in poor agreement with each other.^{6,34–36} This discrepancy in the phase transitions comes from the fact that the process by which liquids are cooled to form a supercooled liquid and glasses affects the crystallization kinetics, crystal perfection /relaxation level and, as a consequence, affects the experimental results.^{37–39} The differences in the glass transition temperatures, T_g , determined in this work and those reported in literature do not exceed 5 K, which may result from the different cooling rates applied in the formation of the glass.³⁷ The structural symmetry of a supercooled liquid and the molecular weight affect the glass transition.

Figure 1, depicts the glass transitions and fusion temperatures of the studied ILs. As it was expected T_g increases with the molecular weight. The glass transition temperature of the $[\text{C}_{N-1}\text{C}_1\text{im}][\text{NTf}_2]$ series follows the order: $[\text{C}_1\text{C}_1\text{im}][\text{NTf}_2] < [\text{C}_2\text{C}_1\text{im}][\text{NTf}_2] < [\text{C}_3\text{C}_1\text{im}][\text{NTf}_2] < [\text{C}_4\text{C}_1\text{im}][\text{NTf}_2] < [\text{C}_5\text{C}_1\text{im}][\text{NTf}_2]$. The $[\text{C}_2\text{C}_3\text{im}][\text{NTf}_2]$ and $[\text{C}_3\text{C}_3\text{im}][\text{NTf}_2]$ have slightly lower T_g than their respective isomers. These results are consistent with the symmetric features. Liquids composed of asymmetric molecules form glasses much more readily than those consisting of symmetric ones.³⁸ Additionally, there is a correlation between the T_g and the viscosities measured in this work. The most viscous ILs are those with the highest T_g , which could be related with glass cohesive energy and their relation with their higher energy barrier to flow.

Among the short alkyl chain ILs with the acidic N-H group, only for $[\text{C}_1\text{Him}][\text{NTf}_2]$ it was not possible to obtain a glass state temperature due to the fast and easy crystallization mechanism even at fast cooling rate before the glass formation. The fast and easy crystallization mechanism could be related with small hindrance effect and high cohesive energy due to the acidic N-H group on the crystallization process. The $[\text{C}_4\text{C}_1\text{im}][\text{NTf}_2]$ isomer with a methylation in the position 2, C(2), presents a higher T_g than the $[\text{C}_{N-1}\text{C}_1\text{im}][\text{NTf}_2]$ series due to the hindrance effect of the $-\text{CH}_3$ that decreases the cation-anion dynamics as will be discussed below. By other hand, the significant increase in the T_g in the $[\text{C}_2\text{Him}][\text{NTf}_2]$ should be related to the increase of the cohesive energy due to the presence of the acidic N-H group. The empirical correlation between the T_g and T_m , $[T_g/T_m = 2/3]$, observed for polymers,⁴⁰ was also found here for the $[\text{C}_2\text{Him}][\text{NTf}_2]$ and $[\text{C}_2\text{C}_1\text{im}][\text{NTf}_2]$.

For the enthalpies and entropies of transitions, the results presented in this work show some deviations from the literature values.^{34,35} The entropy of transition was determined by the relation: $S_{trans.} = H_{trans.} / T_{trans.}$.

The discussion, comparison, and interpretation of the results with the literature data should be carried out with caution since the obtained temperatures and enthalpies of transitions are affected by the thermal history and the methodology applied to the phase behavior studies (e.g., temperature scan rate, quenching procedure, time of stabilization, impurities, crucible material, and sample size).

Figure 2, depicts the enthalpies and entropies of melting as a function of the total number of carbon atoms, N , in the alkyl side chains of the ILs. The initial decrease in the enthalpy of melting is reflected in the initial decrease of the melting temperatures along the total number of carbons of the cation. The enthalpic differentiation between the series $[^1C_{N-1}^3C_1im][NTf_2]$ and $[^1C_NHim][NTf_2]$ is, however, partially cancelled in the melting temperatures due to the entropic compensation. The higher temperature of melting of the $[^1C_1^2C_1Him][NTf_2]$ relative to the $[^1C_2Him][NTf_2]$ and $[^1C_1^3C_1im][NTf_2]$ isomers is ruled by the lower entropy of melting in agreement with the decrease of the anion-cation dynamics due to the methylation in the position 2, C(2), of the imidazolium ring. The melting temperature of the $[^1C_2^3C_2im][NTf_2]$ is similar to the $[^1C_1^3C_1im][NTf_2]$, besides their significant higher enthalpy of melting, reflecting the strong entropic compensation effect.

3.2 Densities

The experimental raw data for density for the $[^1C_1Him][NTf_2]$, $[^1C_2Him][NTf_2]$, $[^1C_2^3C_3im][NTf_2]$ and $[^1C_1^2C_1Him][NTf_2]$ ILs are presented in the Supporting Information, Table SI.2. The density data (ρ), in the studied temperature (T) range, was further correlated using a second order polynomial equation:

$$\ln(\rho/\text{kg} \cdot \text{m}^{-3}) = a + b \cdot T + c \cdot T^2 \quad (1)$$

where a , b , and c are the coefficients obtained from the least square fitting of equation (1), T is the temperature in K. The graphic representation of the logarithm of density as function the temperature is presented in Figure 3, together with literature data¹⁵ for $[^1C_{N-1}^3C_1im][NTf_2]$ (where $N = 3 - 6$) and $[^1C_{N/2}^3C_{N/2}im][NTf_2]$ (where $N = 2, 4, 6$).

The isobaric thermal expansion coefficient, α_p , which considers the volumetric changes with temperature, was derived using the equation (2):

$$\alpha_p = -\frac{1}{\rho} \left(\frac{\partial \rho}{\partial T} \right)_p = - \left(\frac{\partial \ln \rho}{\partial T} \right)_p = -[b + 2c \cdot (T/K)] \quad (2)$$

where, ρ is the density in $\text{kg} \cdot \text{m}^{-3}$, T is the temperature in K, p is the standard pressure (0.1 MPa), and b and c are the fitting coefficients of equation (1). The fitting parameters such as the molar volume and the thermal expansion coefficients, at $T = 323.15$ K and 0.1 MPa, for

all the studied ILs are listed in Table 3. Due to the fact that some of the ILs are solid at room temperature, the comparison of the data was done at $T = 323.15$ K.

The graphic representations of the density and molar volume and thermal expansion coefficients at 323.15 K and 0.1 MPa, against the total number of carbon atoms in the alkyl side chains of the cation, N , are presented in Figures 4 and 5, respectively. No significant differentiation in the isomers could be detected in the density and thermal expansion coefficients. Thus, it can be concluded that the volumetric properties are barely affected by the geometric and structural isomerization for these ILs, following a quite regular trend.

3.3 Viscosities

The experimental viscosity data for the $[^1\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$, and $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ ILs are presented in the Supporting Information, Table SI. 3. Figure 6 depicts the graphic representation of the $\ln(\eta/\text{mPa}\cdot\text{s})$ against the temperature, together with literature data for $[^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ (where $N = 3 - 6$) and $[^1\text{C}_{N/2}^3\text{C}_{N/2}\text{im}][\text{NTf}_2]$ (where $N = 2, 4, 6$) series.^{15,41}

The experimental viscosity data was correlated using the Vogel-Tammann-Fulcher (VTF) model described in equation (3):

$$\ln(\eta/\text{mPa}\cdot\text{s}) = \ln(A_\eta/\text{mPa}\cdot\text{s}) + \frac{B_\eta}{(T - C_\eta)} \quad (3)$$

Where, η is the viscosity in $\text{mPa}\cdot\text{s}$, T is the temperature in K, and A_η , B_η and C_η are the fitting parameters from the fitting of the experimental data. The energy barrier of the fluid to a shear stress was evaluated based on the viscosity dependence with the temperature using the following equation (4)

$$E = R \cdot \frac{\partial(\ln\eta)}{\partial(1/T)} = R \cdot \left(\frac{B_\eta}{\left(\frac{C_\eta^2}{T^2} - \frac{2\cdot C_\eta}{T} + 1\right)} \right) \quad (4)$$

The derived coefficients of the VTF equation (3), the viscosity and the derived energy barrier, E , at $T = 323.15$ K, for the studied ILs, are presented in Table 4.

The graphic representation of viscosity, η , at $T = 323.15$ K, as a function of the total number of carbon atoms in the alkyl chains in the cation, for the $[^1\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$ and $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ ILs and their comparison with the literature data^{15,41} for the $[^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ (where $N = 3 - 6$) and $[^1\text{C}_{N/2}^3\text{C}_{N/2}\text{im}][\text{NTf}_2]$ (where $N = 2, 4, 6$) series, are depicted in Figure 7(I). The energy barrier at $T = 323.15$ K, $E(T = 323.15 \text{ K})$, as a function of the total number of carbon atoms in the alkyl side chains of the cation, N , are presented in Figure 7 (II).

The $[^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ series is systematically more viscous than the $[^1\text{C}_{N/2}^3\text{C}_{N/2}\text{im}][\text{NTf}_2]$ series, as described in a previous work.¹⁵ A strong differentiation in the viscosities

of the isomers with short alkyl chains, such as $[^1\text{C}_2\text{Him}][\text{NTf}_2]$, $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ and $[^1\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$ was observed. The shorter alkyl substituted ILs present higher viscosities when compared with the long alkyl chain series. This trend of the viscosity along the total number of carbons of the cation is very similar to that observed in the energy barrier profile, as depicted in Figure 7, which is an indication that the viscosity trend is ruled by the differentiation in the cohesive energy. The most viscous ILs, $[^1\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2\text{Him}][\text{NTf}_2]$ and $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$, have an acidic N-H group in the imidazolium ring. The acidic hydrogen atom is able to establish hydrogen bonds with the oxygen atoms of the $[\text{NTf}_2]^-$ anion, leading to an increase of energy barrier of shear stress. The analysis of the IR spectra of the $[^1\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$, $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ and $[^1\text{C}_1\text{Him}][\text{NTf}_2]$ (data is presented as support information) in the range of 400- 4000 cm^{-1} do not show any significant differentiation that could be associated to a change in the sulfonyl vibrations, however the enhancement of the anion-cation interaction by increasing the number and the strength of H-bond abilities was previously proposed by Ludwig et al. 43 based on FIR (far IR) spectroscopy in agreement with our conclusion and remarks concerning the effect of the acid N-H group in the thermophysical properties.

The effect of methylation of the C(2) position in the viscosity was previously discussed by Hunt17, based on computational studies, considering the reduction of ion-pair configurational variation, which leads to an additional increase of the energy barrier in relation to other isomers, as observed in this work. Our results also indicate the differentiation (increase) in the viscosity by the methylation in C(2) in the $[^1\text{C}_4^2\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$ with the same magnitude that was observed in $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$, giving thus an additional support for the previous rationalization concerning the effect of the substitution in the position 2 of the imidazolium.

3.4 Heat Capacities

The molar, $C_{p,m}^o$ ($\text{J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$), specific, c_p^o ($\text{J} \cdot \text{K}^{-1} \cdot \text{g}^{-1}$), and volumetric, C_p^o/V ($\text{J} \cdot \text{K}^{-1} \cdot \text{cm}^{-3}$) heat capacities at $T = 298.15 \text{ K}$ and 0.1 MPa , of the $[^1\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$ and $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ ILs are presented in Table 5 together with the total number of drop experiments, N_{drop} , for each ionic liquid. Due to the high metastability of the supercooled liquid $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$, it was possible to measure the heat capacity in the liquid phase, which is also presented in Table 5.

The heat capacity data obtained for the $[^1\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$ and $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$, was compared with the data for the $[^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ (where $N = 3 - 5$) and $[^1\text{C}_{N/2}^3\text{C}_{N/2}\text{im}][\text{NTf}_2]$ (where $N = 2, 4, 6$) ILs available in the

literature.^{44,45} Figure 8 (I) shows the representation of the molar heat capacity data ($C_{p,m}^o$) against the total number of carbon atoms in the alkyl side chains of the cation, N , of the $[^1\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$ and $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ ILs, together with the data for the $[^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ (where $N = 3 - 5$)⁴⁴ and $[^1\text{C}_{N/2}^3\text{C}_{N/2}\text{im}][\text{NTf}_2]$ (where $N = 2, 4, 6$)⁴⁵ and (II) the deviation from the linear fitting of the molar heat capacities. Figure 9 presents the plots of the specific heat capacity (I) and the volumetric

heat capacities (II) against the total number of carbon atoms in the alkyl side chains of the cation, N , of the considered ILs.

From Figure 8 (II) it can be seen that the $N=2$ isomers, namely $[^1\text{C}_2\text{Him}][\text{NTf}_2]$ $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$, are outliers, with a significant positive deviation from the linear trend of the ILs series presented in this work. From the analysis of the volumetric heat capacities depicted in Figure 9 (II), the differentiation is more evident. The heat capacity *per* volume units is significantly higher for the $[^1\text{C}_2\text{Him}][\text{NTf}_2]$ and $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$. The presence of the acidic N-H contributes to a significant interaction potential profile which leads to a heat capacity increment of $\sim 12 \text{ J}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$. In agreement with the previous findings, the $[^1\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$ (symmetrical series) presents a slightly lower heat capacity than the asymmetric series.⁴⁵ However, the $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$ nicely fits the trend of the asymmetric series, as expected.

3.5 Refractive indices

The refractive indices fitting data for $[^1\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$, $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ (where $N=3-6$) and $[^1\text{C}_{N/2}^3\text{C}_{N/2}\text{im}][\text{NTf}_2]$ (where $N=2, 4, 6$), in the temperature range from 289 K to 342 K, are presented in supporting information, Table SI.4. The graphic representation of the refractive indices, as function of the temperature for the studied ILs, is depicted in Figure 10 (I). Table 6, lists the refractive indices of all studied ILs, at $T=298.15 \text{ K}$, together with available literature values, and the temperature derivative of the temperature dependence of the refractive index, $d n_D/dT$. The plots of the refractive indices, at $T=298.15 \text{ K}$, as a function of the total number of carbon atoms in the alkyl chains in the imidazolium cations, for the measured ILs, are shown in Figure 10 (II). The refractive indices obtained in this work are in good agreement with the available literature values, with relative deviations under 2%. 46–57

As shown in Figure 10 (II), the ILs with shorter alkyl chain length are clearly differentiated with respect to long-chain isomers. A regular increase trend in the refractive indices from $[^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ (where $N=3-6$) and $[^1\text{C}_{N/2}^3\text{C}_{N/2}\text{im}][\text{NTf}_2]$ (where $N=2, 4, 6$) was observed. Concerning the short alkyl chain ILs, the following order for the refractive indices was found: $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2] > [^1\text{C}_2\text{Him}][\text{NTf}_2] > [^1\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$. The methylation in C(2) position, as well as the N-H acidic group in the imidazolium ring contributes to a significant differentiation in the cation-anion interactions which is reflected in their charge distribution and polarizability compared with their respective isomers.

3.6 Surface Tensions

The surface thermodynamic properties, namely surface entropy and surface enthalpy, were estimated using the quasi-linear dependence of the surface tension with temperature.⁵⁹

The surface entropy, $S^\gamma(T)$, was evaluated according to equation (5):

$$S^\gamma(T) = - \left(\frac{d\gamma}{dT} \right)_T \quad (5)$$

Whereas the surface enthalpy, $H^\gamma(T)$, was determined according to equation (6):

$$H^\gamma(T) = \gamma - T \left(\frac{d\gamma}{dT} \right)_T \quad (6)$$

Where γ stands for the surface tension and T for the temperature.

The values of the surface tensions and the thermodynamic functions, at $T = 330\text{K}$, of all the bis[(trifluoromethyl)sulfonyl]imide based ILs derived from the temperature dependence of the surface tension, $\gamma = f(T)$, in combination with the associated deviation⁶⁰ are presented in Table 7. Figure 11, depicts the dependence of the surface tension with temperature among the ILs. The experimental raw data for the ILs studied is presented in the Supporting Information, Table SI.5. The surface tension, at $T = 330\text{ K}$, as a function of the total number of carbons is depicted in Figure 12. A strong differentiation in the surface tension between the $N = 2$ isomers was found to follow the trend: $[^1\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2] > [^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2] > [^1\text{C}_2\text{Him}][\text{NTf}_2]$.

The derived surface enthalpies and entropies of the ILs series are depicted in Figure 13. The higher values of surface tension of $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$ and $[^1\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$ are in agreement with their higher interaction potential in the bulk. It is interesting to notice that the present behavior reflects the low basicity of the bistriflamide in where the full aprotic character of this ionic liquid is preserved even with the acidic N-H group when compared with the recent results concerning the acetate derivatives.³² the remaining isomers follow a regular trend as observed previously,¹³ with an initial decrease of the surface tension until $N = 6$, critical alkyl chain length (CAL), reflecting the decrease of the polar interaction.

3.7 Final remarks

This work presents an extended study of the isomerization effect in the physicochemical properties of short chain length imidazolium $[\text{NTf}_2]^-$ ionic liquid series. A strong differentiation in the physicochemical properties arising from the presence of the acidic group, N-H, as well the methylation of the position 2, C(2), in the imidazolium ring was found and interpreted. The observed differentiation of $[^1\text{C}_1\text{Him}][\text{NTf}_2]$, the isomers $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$, $[^1\text{C}_2\text{Him}][\text{NTf}_2]$ and $[^1\text{C}_4^2\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$, when compared with the regular trend of the remaining IL member series, is in agreement with their higher interaction potential in the bulk. This increase in the interaction potential arises from two main effects, which contribute in the same direction to the differentiation: the acidic hydrogen N-H, that is able to form hydrogen bonds with the oxygen atoms of the $[\text{NTf}_2]^-$ anion; and the methylation in the C(2), which reduces the ion-pair configurational variation, leading to a more localized charge distribution and a significant decrease of the entropy of the liquid phase in agreement with the observed lower enthalpy of melting. The obtained results, and especially their comparative analysis, are in full agreement with the rationalization based in the higher interaction potential: higher heat capacities, higher viscosities, higher refractive indices, thermal behavior and the bulk structuration derived from the surface tension results.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Thanks are due to Fundação para a Ciência e Tecnologia (FCT), Lisbon, Portugal and to FEDER for financial support to Centro de Investigação em Química, University of Porto through the project Pest-C/QUI/UI0081/2013, and CICECO and QOPNA, University of Aveiro, through the projects UID/CTM/50011/2013 and PEst-C/QUI/UI0062/2013; FCOMP-01-0124-FEDER-037296, respectively, financed by national funds through the FCT/MEC and when applicable co-financed by FEDER under the PT2020 Partnership Agreement, and COST Action CM1206 - EXIL - Exchange on Ionic Liquids. The authors also thank FCT for the PhD and post-doctoral grants SFRH/BD/81261/2011, SFRH/BD/60513/2009, SFRH/BD/88369/2012 and SFRH/BD/70641/2010 from A.S.M.C. Rodrigues, M.A.A. Rocha, H.F.D. Almeida and C.M.S.S.N., respectively. M. G. Freire acknowledges the European Research Council (ERC) for the Starting Grant ERC-2013-StG-337753.

References

- Urahata SM, Ribeiro MCC. Structure of Ionic Liquids of 1-Alkyl-3-Methylimidazolium Cations: A Systematic Computer Simulation Study. *J Chem Phys.* 2004; 120:1855–1863. [PubMed: 15268318]
- Wang Y, Voth GA. Unique Spatial Heterogeneity in Ionic Liquids. *J Am Chem Soc.* 2005; 127:12192–12193. [PubMed: 16131175]
- Canongia Lopes JNA, Pádua AAH. Nanostructural Organization in Ionic Liquids. *J Phys Chem B.* 2006; 110:3330–3335. [PubMed: 16494347]
- Bhargava BL, Devane R, Klein ML, Balasubramanian S. Nanoscale Organization in Room Temperature Ionic Liquids: A Coarse Grained Molecular Dynamics Simulation Study. *Soft Matter.* 2007; 3:1395–1400.
- Triolo A, Russina O, Bleif H-J, Di Cola E. Nanoscale Segregation in Room Temperature Ionic Liquids. *J Phys Chem B.* 2007; 111:4641–4644. [PubMed: 17388414]
- Zheng W, Mohammed A, Hines LG, Xiao D, Martinez OJ, Bartsch RA, Simon SL, Russina O, Triolo A, Quitevis EL. Effect of Cation Symmetry on the Morphology and Physicochemical Properties of Imidazolium Ionic Liquids. *J Phys Chem B.* 2011; 115:6572–6584. [PubMed: 21534579]
- Rocha MAA, Lima CFRAC, Gomes LR, Schröder B, Coutinho JAP, Marrucho IM, Esperança JMSS, Rebelo LPN, Shimizu K, Lopes JNC, et al. High-Accuracy Vapor Pressure Data of the Extended [C(n)C1im][Ntf2] Ionic Liquid Series: Trend Changes and Structural Shifts. *J Phys Chem B.* 2011; 115:10919–10926. [PubMed: 21815684]
- Bernardes CES, Shimizu K, Lobo Ferreira AIMC, Santos LMNBF, Canongia Lopes JN. Structure and Aggregation in the 1,3-Dialkyl-Imidazolium Bis(trifluoromethylsulfonyl)imide Ionic Liquid Family: 2. From Single to Double Long Alkyl Side Chains. *J Phys Chem B.* 2014; 118:6885–6895. [PubMed: 24873822]
- Shimizu K, Bernardes CES, Canongia Lopes JN. Structure and Aggregation in the 1-Alkyl-3-Methylimidazolium Bis(trifluoromethylsulfonyl)imide Ionic Liquid Homologous Series. *J Phys Chem B.* 2014; 118:567–576. [PubMed: 24358956]
- Martins MAR, Neves CMSS, Kurnia KA, Santos LMNBF, Freire MG, Pinho SP, Coutinho JAP. Analysis of the Isomerism Effect on the Mutual Solubilities of Bis(trifluoromethylsulfonyl)imide-Based Ionic Liquids with Water. *Fluid Phase Equilib.* 2014; 381:28–35.
- Rocha MAA, Coutinho JAP, Santos LMNBF. Vapor Pressures of 1,3-Dialkylimidazolium Bis(trifluoromethylsulfonyl)imide Ionic Liquids with Long Alkyl Chains. *J Chem Phys.* 2014; 141:134502–134508. [PubMed: 25296816]
- Kurnia KA, Sintra TE, Neves CMSS, Shimizu K, Canongia Lopes JN, Gonçalves F, Ventura SPM, Freire MG, Santos LMNBF, Coutinho JAP. The Effect of the Cation Alkyl Chain

- Branching on Mutual Solubilities with Water and Toxicities. *Phys Chem Chem Phys*. 2014; 16:19952–19963. [PubMed: 25119425]
- (13). Almeida HFD, Freire MG, Fernandes AM, Lopes-da-Silva JA, Santos LMNBF, Coutinho JAP, Morgado P, Shimizu K, Filipe EJM, Lopes JNC, et al. Cation Alkyl Side Chain Length and Symmetry Effects on the Surface Tension of Ionic Liquids. *Langmuir*. 2014; 30:6408–6418. [PubMed: 24834955]
  - (14). Rocha MAA, Ribeiro FMS, Schröder B, Coutinho JAP, Santos LMNBF. Volatility Study of [C1C1im][NTf2] and [C2C3im][NTf2] Ionic Liquids. *J Chem Thermodyn*. 2014; 68:317–321.
  - (15). Rocha MAA, Neves CMSS, Freire MG, Russina O, Triolo A, Coutinho JAP, Santos LMNBF. Alkylimidazolium Based Ionic Liquids: Impact of Cation Symmetry on Their Nanoscale Structural Organization. *J Phys Chem B*. 2013; 117:10889–10897. [PubMed: 23915346]
  - (16). Bonhôte P, Dias A-P, Papageorgiou N, Kalyanasundaram K, Grätzel M. Hydrophobic, Highly Conductive Ambient-Temperature Molten Salts †. *Inorg Chem*. 1996; 35:1168–1178. [PubMed: 11666305]
  - (17). Hunt PA. Why Does a Reduction in Hydrogen Bonding Lead to an Increase in Viscosity for the 1-Butyl-2,3-Dimethyl-Imidazolium-Based Ionic Liquids? *J Phys Chem B*. 2007; 111:4844–4853. [PubMed: 17388550]
  - (18). Noack K, Schulz PS, Paape N, Kiefer J, Wasserscheid P, Leipertz A. The Role of the C2 Position in Interionic Interactions of Imidazolium Based Ionic Liquids: A Vibrational and NMR Spectroscopic Study. *Phys Chem Chem Phys*. 2010; 12:14153–14161. [PubMed: 20871893]
  - (19). Fumino K, Wulf A, Ludwig R. Strong, Localized, and Directional Hydrogen Bonds Fluidize Ionic Liquids. *Angew Chem Int Ed Engl*. 2008; 47:8731–8734. [PubMed: 18846527]
  - (20). Fumino K, Wulf A, Ludwig R. The Cation-Anion Interaction in Ionic Liquids Probed by Far-Infrared Spectroscopy. *Angew Chem Int Ed Engl*. 2008; 47:3830–3834. [PubMed: 18412214]
  - (21). Izgorodina EI, Maganti R, Armel V, Dean PM, Pringle JM, Seddon KR, MacFarlane DR. Understanding the Effect of the C2 Proton in Promoting Low Viscosities and High Conductivities in Imidazolium-Based Ionic Liquids: Part I. Weakly Coordinating Anions. *J Phys Chem B*. 2011; 115:14688–14697. [PubMed: 22023104]
  - (22). Wieser ME, Berglund M. Atomic Weights of the Elements 2007 (IUPAC Technical Report). *Pure Appl Chem*. 2009; 81:2131–2156.
  - (23). Sabbah RC, (France), Xu-wu A, (China), Chickos JS, (USA), Leitão MLP, (Portugal), Roux MV, (Spain), Torres LA, (México), Sabbah R, Xu-wu A, Chickos JS, Leitão MLP, et al. Reference Materials for Calorimetry and Differential Thermal Analysis. *Thermochim Acta*. 1999; 331:93–204.
  - (24). Roux MV, Temprado M, Chickos JS, Nagano Y. Critically Evaluated Thermochemical Properties of Polycyclic Aromatic Hydrocarbons. *J Phys Chem Ref Data*. 2008; 37:1855–1996.
  - (25). Paredes X, Fandiño O, Comuñas MJP, Pensado AS, Fernández J. Study of the Effects of Pressure on the Viscosity and Density of Diisodecyl Phthalate. *J Chem Thermodyn*. 2009; 41:1007–1015.
  - (26). Carvalho PJ, Regueira T, Santos LMNBF, Fernandez J, Coutinho JAP. Effect of Water on the Viscosities and Densities of 1-Butyl-3-Methylimidazolium Dicyanamide and 1-Butyl-3-Methylimidazolium Tricyanomethane at Atmospheric Pressure†. *J Chem Eng Data*. 2009; 55:645–652.
  - (27). Konicek IWJ, Suurkuusk J. A Precise Drop Heat Capacity Calorimeter for Small Samples. *Chem Scr*. 1971; 1:217–220.
  - (28). Suurkuusk J, Wadsö I. Design and Testing of an Improved Precise Drop Calorimeter for the Measurement of the Heat Capacity of Small Samples. *J Chem Thermodyn*. 1974; 6:667–679.
  - (29). Santos LMNBF, Rocha MAAA, Rodrigues ASMCMC, Štejfa V, Fulem M, Bastos M. Reassembling and Testing of a High-Precision Heat Capacity Drop Calorimeter. Heat Capacity of Some Polyphenyls at T=298.15K. *J Chem Thermodyn*. 2011; 43:1818–1823.
  - (30). Bernardes CES, Santos LMNBF, da Piedade MEM. A New Calorimetric System to Measure Heat Capacities of Solids by the Drop Method. *Meas Sci Technol*. 2006; 17:1405–1408.
  - (31). Almeida HFD, Lopes-Da-Silva JA, Freire MG, Coutinho JAP. Surface Tension and Refractive Index of Pure and Water-Saturated Tetradecyltriethylphosphonium-Based Ionic Liquids. *J Chem Thermodyn*. 2013; 57:372–379.

- (32). Almeida HFD, Passos H, Lopes-Da-Silva JA, Fernandes AM, Freire MG, Coutinho JAP. Thermophysical Properties of Five Acetate-Based Ionic Liquids. *J Chem Eng Data*. 2012; 57:3005–3013.
- (33). Almeida HFD, Teles ARR, Lopes-Da-Silva JA, Freire MG, Coutinho JAP. Influence of the Anion on the Surface Tension of 1-Ethyl-3-Methylimidazolium-Based Ionic Liquids. *J Chem Thermodyn*. 2012; 54:49–54.
- (34). Tokuda H, Hayamizu K, Ishii K, Susan MABH, Watanabe M. Physicochemical Properties and Structures of Room Temperature Ionic Liquids. 2. Variation of Alkyl Chain Length in Imidazolium Cation. *J Phys Chem B*. 2005; 109:6103–6110. [PubMed: 16851672]
- (35). Doma ska U, R kawek A, Marciniak A. Solubility of 1-Alkyl-3-Ethylimidazolium-Based Ionic Liquids in Water and 1-Octanol. *J Chem Eng Data*. 2008; 53:1126–1132.
- (36). Blokhin AV, Paulechka YU, Strechan AA, Kabo GJ. Physicochemical Properties, Structure, and Conformations of 1-Butyl-3-Methylimidazolium Bis(trifluoromethanesulfonyl)imide [C4mim]NTf2 Ionic Liquid. *J Phys Chem B*. 2008; 112:4357–4364. [PubMed: 18341327]
- (37). Ediger MD, Angell CA, Nagel SR. Supercooled Liquids and Glasses. *J Phys Chem*. 1996; 100:13200–13212.
- (38). Turnbull D. Under What Conditions Can a Glass Be Formed? *Contemp Phys*. 1969; 10:473–488.
- (39). Debenedetti PG, Stillinger FH. Supercooled Liquids and the Glass Transition. *Nature*. 2001; 410:259–267. [PubMed: 11258381]
- (40). Nicholson, JW. *The Chemistry of Polymers*. RSC Publishing, The Royal Society of Chemistry; 2006.
- (41). Tariq M, Carvalho PJ, Coutinho JAP, Marrucho IM, Lopes JNC, Rebelo LPN. Viscosity of (C2–C14) 1-Alkyl-3-Methylimidazolium Bis(trifluoromethylsulfonyl)amide Ionic Liquids in an Extended Temperature Range. *Fluid Phase Equilib*. 2011; 301:22–32.
- (42). Gaciño FM, Regueira T, Lugo L, Comuñas MJP, Fernández J. Influence of Molecular Structure on Densities and Viscosities of Several Ionic Liquids. *J Chem Eng Data*. 2011; 56:4984–4999.
- (43). Wulf A, Fumino K, Ludwig R. Spectroscopic Evidence for an Enhanced Anion-Cation Interaction from Hydrogen Bonding in Pure Imidazolium Ionic Liquids. *Angew Chem Int Ed Engl*. 2010; 49:449–453. [PubMed: 20029865]
- (44). Rocha MAA, Bastos M, Coutinho JAP, Santos LMNBF. Heat Capacities at 298.15 K of the Extended [CnC1im][Ntf2] Ionic Liquid Series. *J Chem Thermodyn*. 2012; 53:140–143.
- (45). Rocha MAA, Coutinho JAP, Santos LMNBF. Evidence of Nanostructuration from the Heat Capacities of the 1,3-Dialkylimidazolium Bis(trifluoromethylsulfonyl)imide Ionic Liquid Series. *J Chem Phys*. 2013; 139:104502–104507. [PubMed: 24050354]
- (46). Seki S, Tsuzuki S, Hayamizu K, Umebayashi Y, Serizawa N, Takei K, Miyashiro H. Comprehensive Refractive Index Property for Room-Temperature Ionic Liquids. *J Chem Eng Data*. 2012; 57:2211–2216.
- (47). Miran Beigi AA, Abdouss M, Yousefi M, Pourmortazavi SM, Vahid A. Investigation on Physical and Electrochemical Properties of Three Imidazolium Based Ionic Liquids (1-Hexyl-3-Methylimidazolium Tetrafluoroborate, 1-Ethyl-3-Methylimidazolium Bis(trifluoromethylsulfonyl) Imide and 1-Butyl-3-Methylimidazolium Methylsulfate). *J Mol Liq*. 2013; 177:361–368.
- (48). Tariq M, Forte PAS, Gomes MFC, Lopes JNC, Rebelo LPN. Densities and Refractive Indices of Imidazolium- and Phosphonium-Based Ionic Liquids: Effect of Temperature, Alkyl Chain Length, and Anion. *J Chem Thermodyn*. 2009; 41:790–798.
- (49). Seoane RG, Corderí S, Gómez E, Calvar N, González EJ, Macedo EA, Domínguez A, Domínguez Á. Temperature Dependence and Structural Influence on the Thermophysical Properties of Eleven Commercial Ionic Liquids. *Ind Eng Chem Res*. 2012; 51:2492–2504.
- (50). Seoane RG, González EJ, González B. 1-Alkyl-3-Methylimidazolium Bis(trifluoromethylsulfonyl)imide Ionic Liquids as Solvents in the Separation of Azeotropic Mixtures. *J Chem Thermodyn*. 2012; 53:152–157.
- (51). Corderí S, González B. Ethanol Extraction from Its Azeotropic Mixture with Hexane Employing Different Ionic Liquids as Solvents. *J Chem Thermodyn*. 2012; 55:138–143.

- (52). Lago S, Rodríguez H, Soto A, Arce A. Deterpenation of Citrus Essential Oil by Liquid–Liquid Extraction with 1-Alkyl-3-Methylimidazolium Bis(trifluoromethylsulfonyl)amide Ionic Liquids. *J Chem Eng Data*. 2011; 56:1273–1281.
- (53). Gómez E, Calvar N, Macedo EA, Domínguez Á. Effect of the Temperature on the Physical Properties of Pure 1-Propyl 3-Methylimidazolium Bis(trifluoromethylsulfonyl)imide and Characterization of Its Binary Mixtures with Alcohols. *J Chem Thermodyn*. 2012; 45:9–15.
- (54). Andreatta AE, Arce A, Rodil E, Soto A. Physico-Chemical Properties of Binary and Ternary Mixtures of Ethyl Acetate + Ethanol + 1-Butyl-3-Methyl-Imidazolium Bis(trifluoromethylsulfonyl)imide at 298.15 K and Atmospheric Pressure. *J Solution Chem*. 2010; 39:371–383.
- (55). Andreatta AE, Francisco M, Rodil E, Soto A, Arce A. Isobaric vapour–liquid equilibria and physical properties for isopropyl acetate + isopropanol + 1-butyl-3-methyl-imidazolium bis(trifluoromethylsulfonyl)imide mixtures. *Fluid Phase Equilib*. 2011; 300:162–171.
- (56). Jin H, O'Hare B, Dong J, Arzhantsev S, Baker GA, Wishart JF, Benesi AJ, Maroncelli M. Physical Properties of Ionic Liquids Consisting of the 1-Butyl-3-Methylimidazolium Cation with Various Anions and the Bis(trifluoromethylsulfonyl)imide Anion with Various Cations. *J Phys Chem B*. 2008; 112:81–92. [PubMed: 18069817]
- (57). Deetlefs M, Seddon KR, Shara M. Predicting Physical Properties of Ionic Liquids. *Phys Chem Chem Phys*. 2006; 8:642–649. [PubMed: 16482306]
- (58). Seki S, Tsuzuki S, Hayamizu K, Umebayashi Y, Serizawa N, Takei K, Miyashiro H. Comprehensive Refractive Index Property for Room-Temperature Ionic Liquids. *J Chem Eng Data*. 2012; 57:2211–2216.
- (59). Adamson, AW., Gast, AP. *Physical Chemistry of Surfaces*. John Wiley; New York: 1997.
- (60). Miller, JC., Miller, JN. *Statistical for Analytical Chemistry*. Hall, PTR Prentice; Chichester, NY: 1993.
- (61). Carvalho PJ, Freire MG, Marrucho IM, Queimada AJ, Coutinho JAP. Surface Tensions for the 1-Alkyl-3-Methylimidazolium Bis(trifluoromethylsulfonyl)imide Ionic Liquids. *J Chem Eng Data*. 2008; 53:1346–1350.

Figure 1.

Graphic representation of the glass transitions temperatures, T_g /K, **(I)**, and melting temperatures, T_m /K, **(II)** as a function of the total number of carbon atoms in the alkyl side chains of the ILs: $\blacktriangle$ - $[\text{C}_2\text{Him}][\text{NTf}_2]$; $\blacklozenge$ - $[\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$; $\diamond$ - $[\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$; $\star$ - $[\text{C}_2^3\text{C}_1\text{im}][\text{NTf}_2]$; $\circ$ - $[\text{C}_3^3\text{C}_1\text{im}][\text{NTf}_2]$; $\ast$ - $[\text{C}_4^3\text{C}_1\text{im}][\text{NTf}_2]$; $\triangleright$ - $[\text{C}_5^3\text{C}_1\text{im}][\text{NTf}_2]$; $\blacklozenge$ - $[\text{C}_3^3\text{C}_3\text{im}][\text{NTf}_2]$; $\blacktriangle$ - $[\text{C}_4^2\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$; $\blacksquare$ - $[\text{C}_1\text{Him}][\text{NTf}_2]$; $\bullet$ - $[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$; $\square$ - $[\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$. The dash and dots view-lines highlight the $[\text{C}_N\text{Him}][\text{NTf}_2]$ and $[\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ series which have no physical meaning.

Figure 2.

Graphic representation of the enthalpies (I) and entropies (II), as a function of the total number of carbon atoms in the alkyl side chains of the ILs, N : ■ - $[\text{}^1\text{C}_1\text{Him}][\text{NTf}_2]$; ▲ - $[\text{}^1\text{C}_2\text{Him}][\text{NTf}_2]$; ● - $[\text{}^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$; ★ - $[\text{}^1\text{C}_2^3\text{C}_1\text{im}][\text{NTf}_2]$; ◇ - $[\text{}^1\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$; □ - $[\text{}^1\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$. The dash and dots view-lines highlight the $[\text{}^1\text{C}_N\text{Him}][\text{NTf}_2]$ and $[\text{}^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ series which have no physical meaning.

Figure 3.

Logarithm of density as a function of temperature. This work: ■ - $[^1\text{C}_1\text{Him}][\text{NTf}_2]$; ▲ - $[^1\text{C}_2\text{Him}][\text{NTf}_2]$; ● - $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$; ◆ - $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$; Literature15: ★ - $[^1\text{C}_2^3\text{C}_1\text{im}][\text{NTf}_2]$; ○ - $[^1\text{C}_3^3\text{C}_1\text{im}][\text{NTf}_2]$; * - $[^1\text{C}_4^3\text{C}_1\text{im}][\text{NTf}_2]$; ▷ - $[^1\text{C}_5^3\text{C}_1\text{im}][\text{NTf}_2]$; ◇ - $[^1\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$; □ - $[^1\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$; ● - $[^1\text{C}_3^3\text{C}_3\text{im}][\text{NTf}_2]$.

Figure 4.

Density(I) and molar volume(II), at 323.15 K and 0.1MPa, as a function of the total number of carbon atoms in the alkyl side chains of the cation, N . This work: ■ - $[\text{C}_1\text{Him}][\text{NTf}_2]$; ▲ - $[\text{C}_2\text{Him}][\text{NTf}_2]$; ● - $[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$; ◆ - $[\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$; Literature: ★ - $[\text{C}_2^3\text{C}_1\text{im}][\text{NTf}_2]$; ○ - $[\text{C}_3^3\text{C}_1\text{im}][\text{NTf}_2]$; * - $[\text{C}_4^3\text{C}_1\text{im}][\text{NTf}_2]$; ▷ - $[\text{C}_5^3\text{C}_1\text{im}][\text{NTf}_2]$; ◇ - $[\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$; □ - $[\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$; ⬠ - $[\text{C}_3^3\text{C}_3\text{im}][\text{NTf}_2]$ .15

Figure 5.

Thermal expansion coefficient, α_p , at 323.15 K and 0.1 MPa, as a function of the total number of carbon atoms in the alkyl side chains of the cation, N . This work: ■ - [$^1\text{C}_1\text{Him}$][NTf₂]; ▲ - [$^1\text{C}_2\text{Him}$][NTf₂]; ● - [$^1\text{C}_1^2\text{C}_1\text{Him}$][NTf₂]; ◆ - [$^1\text{C}_2^3\text{C}_3\text{im}$][NTf₂]; Literature: ★ - [$^1\text{C}_2^3\text{C}_1\text{im}$][NTf₂]; ○ - [$^1\text{C}_3^3\text{C}_1\text{im}$][NTf₂]; * - [$^1\text{C}_4^3\text{C}_1\text{im}$][NTf₂]; ▷ - [$^1\text{C}_5^3\text{C}_1\text{im}$][NTf₂]; ◇ - [$^1\text{C}_1^3\text{C}_1\text{im}$][NTf₂]; □ - [$^1\text{C}_2^3\text{C}_2\text{im}$][NTf₂]; ⬠ - [$^1\text{C}_3^3\text{C}_3\text{im}$][NTf₂]. 15

Figure 6.

Logarithm of viscosity as a function of temperature for studied ILs. The solid lines represents the Vogel-Tammann-Fulcher fitting from equation (3) This work: ■ - [¹C₁Him][NTf₂]; ▲ - [¹C₂Him][NTf₂]; ● - [¹C₁²C₁Him][NTf₂]; ◆ - [¹C₂³C₃im][NTf₂]; Literature: ★ - [¹C₂³C₁im][NTf₂]; ○ - [¹C₃³C₁im][NTf₂]; * - [¹C₄³C₁im][NTf₂]; ▷ - [¹C₅³C₁im][NTf₂]; ◇ - [¹C₁³C₁im][NTf₂]; □ - [¹C₂³C₂im][NTf₂]; ◆ - [¹C₃³C₃im][NTf₂].15,41

Figure 7.

Viscosity (η /mPa.s) at $T = 323.15$ K and 0.1 MPa, (I) and energy barrier (E / kJ.mol $^{-1}$) at 323.15 K, (II) for ILs under study as function of the total number of carbons, N . This work:

■ - [^1C_1Him][NTf $_2$]; ▲ - [^1C_2Him][NTf $_2$]; ● - [$^1C_1^2C_1Him$][NTf $_2$]; ◆ - [$^1C_2^3C_3im$][NTf $_2$]; Literature: ★ - [$^1C_2^3C_1im$][NTf $_2$]; ○ - [$^1C_3^3C_1im$][NTf $_2$]; * - [$^1C_4^3C_1im$][NTf $_2$]; ▷ - [$^1C_5^3C_1im$][NTf $_2$]; ◇ - [$^1C_1^3C_1im$][NTf $_2$]; □ - [$^1C_2^3C_2im$][NTf $_2$]; ● - [$^1C_3^3C_3im$][NTf $_2$]; 15,41 ◀ - [$^1C_4^2C_1^3C_1im$][NTf $_2$]. The dashed line highlights the $[^1C_{N-1}^3C_1im][NTf_2]$ series which has no physical meaning.

Figure 8.

Molar heat capacities, at 298.15 K, as a function of the total number of carbon atoms in the alkyl side chains of the cation, N (I): ■ - $[\text{C}_1\text{Him}][\text{NTf}_2]$ (cr); ▲ - $[\text{C}_2\text{Him}][\text{NTf}_2]$; ● - $[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$; ◆ - $[\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$; Literature: ★ - $[\text{C}_2^3\text{C}_1\text{im}][\text{NTf}_2]$; ○ - $[\text{C}_3^3\text{C}_1\text{im}][\text{NTf}_2]$; * - $[\text{C}_4^3\text{C}_1\text{im}][\text{NTf}_2]$; ▷ - $[\text{C}_5^3\text{C}_1\text{im}][\text{NTf}_2]$; ◇ - $[\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$; □ - $[\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$; ◆ - $[\text{C}_3^3\text{C}_3\text{im}][\text{NTf}_2]$. 45 Deviation from the $C_{p,m}^o$ (linear fitting) of the ILs presented in (I) as a function of total number of carbons in the alkyl side chains of the cation (II). The dash view-line highlight the $[\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ series which has no physical meaning.

Figure 9.

Specific heat capacities **(I)** and the volumic heat capacities **(II)**, at 298.15 K, as a function of the total number of carbon atoms in the alkyl side chains of the cation, N . ■ - $[\text{C}_1\text{Him}][\text{NTf}_2]$ (cr); ▲ - $[\text{C}_2\text{Him}][\text{NTf}_2]$; ● - $[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$, ◆ - $[\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$; Literature: ★ - $[\text{C}_2^3\text{C}_1\text{im}][\text{NTf}_2]$; ○ - $[\text{C}_3^3\text{C}_1\text{im}][\text{NTf}_2]$; * - $[\text{C}_4^3\text{C}_1\text{im}][\text{NTf}_2]$; ▷ - $[\text{C}_5^3\text{C}_1\text{im}][\text{NTf}_2]$;44 ◇ - $[\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$; □ - $[\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$; ◆ - $[\text{C}_3^3\text{C}_3\text{im}][\text{NTf}_2]$ .45 The dash view-line highlight the $[\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ series which has no physical meaning.

Figure 10.

Refractive indices as a function of temperature for studied ILs (I): Refractive indices, n_D ($T = 298.15$ K) as function of the total number of carbon atoms in the alkyl side chains of the cation, N (II). ■ - $[^1C_1Him][NTf_2]$; ▲ - $[^1C_2Him][NTf_2]$; ● - $[^1C_1^2C_1Him][NTf_2]$; ◆ - $[^1C_2^3C_1im][NTf_2]$; ★ - $[^1C_2^3C_1im][NTf_2]$; ○ - $[^1C_3^3C_1im][NTf_2]$; * - $[^1C_4^3C_1im][NTf_2]$; ◇ - $[^1C_1^3C_1im][NTf_2]$; □ - $[^1C_2^3C_2im][NTf_2]$; ● - $[^1C_3^3C_3im][NTf_2]$. The dash view-line highlight the $[^1C_{N-1}^3C_1im][NTf_2]$ series which has no physical meaning.

Figure 11.

Surface tension values for the ILs as function of temperature: ■ - $[^1C_1Him][NTf_2]$; ▲ - $[^1C_2Him][NTf_2]$; ● - $[^1C_1^2C_1Him][NTf_2]$; ◆ - $[^1C_2^3C_3im][NTf_2]$; △ - $[^1C_4^2C_1^3C_1im][NTf_2]$; Literature data: ★ - $[^1C_2^3C_1im][NTf_2]$; ○ - $[^1C_3^3C_1im][NTf_2]$; ✱ - $[^1C_4^3C_1im][NTf_2]$; ▷ - $[^1C_5^3C_1im][NTf_2]61$; ◇ - $[^1C_1^3C_1im][NTf_2]$; □ - $[^1C_2^3C_2im][NTf_2]$; ◀ - $[^1C_3^3C_3im][NTf_2]13$.

Figure 12.

Surface tension dependence, at 330 K, as a function of the total number of carbons in the aliphatic chains, N , ■ - $[^1\text{C}_1\text{Him}][\text{NTf}_2]$; ▲ - $[^1\text{C}_2\text{Him}][\text{NTf}_2]$; ● - $[^1\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$; ◆ - $[^1\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$; ▲ - $[^1\text{C}_4^2\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$. Literature data: ◇ - $[^1\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$; □ - $[^1\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$; ● - $[^1\text{C}_3^3\text{C}_3\text{im}][\text{NTf}_2]$ 13; ★ - $[^1\text{C}_2^3\text{C}_1\text{im}][\text{NTf}_2]$; ○ - $[^1\text{C}_3^3\text{C}_1\text{im}][\text{NTf}_2]$; * - $[^1\text{C}_4^3\text{C}_1\text{im}][\text{NTf}_2]$; ▷ - $[^1\text{C}_5^3\text{C}_1\text{im}][\text{NTf}_2]$ 61; The dash view-line highlight the $[^1\text{C}_{N-1}^3\text{C}_1\text{im}][\text{NTf}_2]$ series which has no physical meaning.

Figure 13.

Surface enthalpies (I) and entropies (II) as a function of the total number of carbon atoms N ,

■ - [¹C₁Him][NTf₂]; ▲ - [¹C₂Him][NTf₂]; ● - [¹C₁²C₁Him][NTf₂]; ◆ - [¹C₂³C₃im][NTf₂]; ▲ - [¹C₄²C₁³C₁im][NTf₂]. Literature data: ◇ - [¹C₁³C₁im][NTf₂]; □ - [¹C₂³C₂im][NTf₂]; ● - [¹C₃³C₃im][NTf₂]; ★ - [¹C₂³C₁im][NTf₂]; ○ - [¹C₃³C₁im][NTf₂]; * - [¹C₄³C₁im][NTf₂]; ▽ - [¹C₅³C₁im][NTf₂];. The dash view-line highlight the [¹C _{$N-1$} ³C₁im][NTf₂] series which has no physical meaning.

Table 1

Schematic structures and abbreviations of the ionic liquids under study.

Imidazolium-Based Cations			Abbreviations
R ₁ : methyl	R ₂ : hydrogen	R ₃ : hydrogen	[¹ C ₁ Him][NTf ₂]
R ₁ : ethyl	R ₂ : hydrogen	R ₃ : hydrogen	[¹ C ₂ Him][NTf ₂]
R ₁ :methyl	R ₂ :methyl	R ₃ :hydrogen	[¹ C ₁ ² C ₁ Him][NTf ₂]
R ₁ :methyl	R ₂ :hydrogen	R ₃ :methyl	[¹ C ₁ ³ C ₁ im][NTf ₂]
R ₁ :ethyl	R ₂ :hydrogen	R ₃ :methyl	[¹ C ₂ ³ C ₁ im][NTf ₂]
R ₁ :propyl	R ₂ :hydrogen	R ₃ :methyl	[¹ C ₃ ³ C ₁ im][NTf ₂]
R ₁ :ethyl	R ₂ :hydrogen	R ₃ :ethyl	[¹ C ₂ ³ C ₂ im][NTf ₂]
R ₁ :ethyl	R ₂ :hydrogen	R ₃ :propyl	[¹ C ₂ ³ C ₃ im][NTf ₂]
R ₁ :butyl	R ₂ :hydrogen	R ₃ :methyl	[¹ C ₄ ³ C ₁ im][NTf ₂]
R ₁ :pentyl	R ₂ :hydrogen	R ₃ :methyl	[¹ C ₅ ³ C ₁ im][NTf ₂]
R ₁ :propyl	R ₂ :hydrogen	R ₃ :propyl	[¹ C ₃ ³ C ₃ im][NTf ₂]
R ₁ :butyl	R ₂ :methyl	R ₃ :methyl	[¹ C ₄ ² C ₁ ³ C ₁ im][NTf ₂]

Table 2

Experimental glass, T_g , cold crystallization, T_{cc} , solid-solid, T_{ss} , melting, T_m , temperatures, enthalpies, $H(T)$, and entropies of transitions, $S(T)$ for the studied ILs.

Ionic Liquid	T / K	$H(T) / \text{kJ}\cdot\text{mol}^{-1}$	$S(T) / \text{J}\cdot\text{K}^{-1}\cdot\text{mol}^{-1}$
$[\text{C}_1\text{Him}][\text{NTf}_2]$	$321.5 \pm 0.5 (T_m)$	24.1 ± 0.7	75.0 ± 2.2
$[\text{C}_2\text{Him}][\text{NTf}_2]$	$184.2 \pm 0.5 (T_g)$		
	$206.6 \pm 0.5 (T_{cc})$		
	$263.5 \pm 0.5 (T_{ss})$	1.8 ± 0.7	6.7 ± 2.6
	$275.5 \pm 0.5 (T_m)$	14.2 ± 0.7	51.5 ± 2.5
$[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$	$311.0 \pm 0.5 (T_m)$	19.2 ± 0.7	61.7 ± 2.2
$[\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$	$176.4 \pm 0.5 (T_g)$		
	$293.2 \pm 0.5 (T_m)$	22.0 ± 0.7	75.0 ± 2.4
	$[299 (T_m)]_{34}$	$[24.5]_{34}$	$[81.7]_{34}$
$[\text{C}_2^3\text{C}_1\text{im}][\text{NTf}_2]$	$178.3 \pm 0.5 (T_g)$		
	$[186 (T_g)]_{34}$		
	$206.8 \pm 0.5 (T_{cc})$		
	$260.0 \pm 0.5 (T_m)$	21.3 ± 0.7	81.9 ± 2.7
	$[291 (T_m)]_{34}$	$[24.8 H(T_m)]_{34}$	$[97.1 S(T_m)]_{34}$
$[\text{C}_3^3\text{C}_1\text{im}][\text{NTf}_2]$	$179.6 \pm 0.5 (T_g)$	-	-
	$[184.0(T_g)]_6$		
$[\text{C}_4^3\text{C}_1\text{im}][\text{NTf}_2]$	$182.1 \pm 0.5(T_g)$	-	-
	$[186 (T_g)]_{34}$		
	$[181.5 \pm 0.1(T_g)]_{36}$		
$[\text{C}_2^3\text{C}_2\text{im}][\text{NTf}_2]$	$286.3 \pm 0.5 (T_m)$	29.1 ± 0.7	101.6 ± 2.4
	$[262.6 \pm 0.1(T_m)]_{35}$	$[20.4 \pm 0.3]_{35}$	
	$[285.5 (T_m)]_6$		
$[\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$	$179.5 \pm 0.5(T_g)$	-	-
$[\text{C}_5^3\text{C}_1\text{im}][\text{NTf}_2]$	$183.8 \pm 0.5(T_g)$	-	-
	$[186.5(T_g)]_6$		
$[\text{C}_3^3\text{C}_3\text{im}][\text{NTf}_2]$	$180.9 \pm 0.5(T_g)$	-	-
$[\text{C}_4^2\text{C}_1^3\text{C}_1\text{im}][\text{NTf}_2]$	$191.5 \pm 0.5(T_g)$	-	-

Table 3

List of the fitted parameters (equation (1)), density, molar volume, and the thermal expansion coefficients, α_p , at 323.15 K and 0.1 MPa for the studied ILs.

Ionic Liquid	a	$10^4 \times b / \text{K}^{-1}$	$10^7 \times c / \text{K}^{-2}$	$\rho / (\text{kg} \cdot \text{m}^{-3})$	$V_m / (\text{cm}^3 \cdot \text{mol}^{-1})$	$10^3 \times \alpha_p / \text{K}^{-1}$
$T=323.15 \text{ K}$						
$[\text{C}_1\text{Him}][\text{NTf}_2]$	7.6198 ± 0.0036	-8.42 ± 0.21	2.90 ± 0.30	1600.6	227.0	0.654 ± 0.029
$[\text{C}_2\text{Him}][\text{NTf}_2]$	7.5678 ± 0.0010	-7.59 ± 0.07	1.43 ± 0.10	1537.0	245.5	0.666 ± 0.009
$[\text{C}_2^3\text{C}_3\text{im}][\text{NTf}_2]$	7.4844 ± 0.0014	-7.62 ± 0.09	1.35 ± 0.14	1411.4	297.1	0.675 ± 0.013
$[\text{C}_1^2\text{C}_1\text{Him}][\text{NTf}_2]$	7.5632 ± 0.0032	-7.07 ± 0.19	0.79 ± 0.29	1545.2	244.2	0.656 ± 0.027

Table 4

Fitting parameters of VTF equation for the viscosity data of the studied ILs, viscosity and the derived energy barrier at $T = 323.15$ K.

Ionic Liquid	$A_\eta / (\text{mPa}\cdot\text{s})$	B_η / K	C_η / K	$\eta / (\text{mPa}\cdot\text{s})$	$E / (\text{kJ}\cdot\text{mol}^{-1})$
($T=323.15$ K)					
$[\text{}^1\text{C}_1\text{Him}][\text{NTf}_2]$	0.177 ± 0.005	833 ± 9	163.0 ± 1.0	32.31	28.22 ± 0.76
$[\text{}^1\text{C}_2\text{Him}][\text{NTf}_2]$	0.214 ± 0.004	724 ± 5	168.5 ± 0.5	23.15	26.30 ± 0.36
$[\text{}^1\text{C}_2\text{}^3\text{C}_3\text{im}][\text{NTf}_2]$	0.150 ± 0.003	789 ± 7	157.1 ± 0.6	17.43	24.85 ± 0.42
$[\text{}^1\text{C}_1\text{}^2\text{C}_1\text{Him}][\text{NTf}_2]$	0.234 ± 0.005	678 ± 6	188.2 ± 0.6	35.66	32.34 ± 0.78

Table 5

Number of drop experiments, N_{drop} , the molar heat capacity, $C_{p,m}^o$ ($\text{J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$), specific heat capacities, c_p^o ($\text{J} \cdot \text{K}^{-1} \cdot \text{g}^{-1}$), and volumic heat capacities, C_p^o/V ($\text{J} \cdot \text{K}^{-1} \cdot \text{cm}^{-3}$) at 298.15 K.

Ionic Liquid	N_{drop}	$C_{p,m}^o$ ($\text{J} \cdot \text{K}^{-1} \cdot \text{mol}^{-1}$),	C_p^o ($\text{J} \cdot \text{K}^{-1} \cdot \text{g}^{-1}$),	C_p^o/V ($\text{J} \cdot \text{K}^{-1} \cdot \text{cm}^{-3}$)
[¹ C ₁ Him][NTf ₂] (cr)	12	441.64 ± 0.33	1.2158 ± 0.0009	1.9785 ± 0.0015
[¹ C ₂ Him][NTf ₂] (l)	33	484.08 ± 0.44	1.2831 ± 0.0012	2.0055 ± 0.0019
[¹ C ₂ ³ C ₃ im][NTf ₂] (l)	19	565.08 ± 0.57	1.3475 ± 0.0014	1.9342 ± 0.0020
[¹ C ₁ ² C ₁ Him][NTf ₂] (l)	23	483.08 ± 0.38	1.2804 ± 0.0010	2.0113 ± 0.0016

Table 6

Experimental refractive indices at the sodium D-line, n_D^a , for the studied ILs as a function of temperature T at 0.1 MPa. Available literature data at $T = 298.15$ K.

Ionic liquid	n_D (298.15 K)	$10^4 \cdot (dn_D/dT) / K^{-1}$	n_D (298.15 K) Literature
[¹ C ₁ Him][NTf ₂]	1.42203	-2.93 ± 0.01	n.a.
[¹ C ₂ Him][NTf ₂]	1.42317	-3.012 ± 0.002	n.a.
[¹ C ₁ ² C ₁ Him][NTf ₂]	1.42490	-2.959 ± 0.003	n.a.
[¹ C ₁ ³ C ₁ im][NTf ₂]	1.42101	-2.97 ± 0.02	n.a.
[¹ C ₂ ³ C ₁ im][NTf ₂]	1.42319	-2.99 ± 0.01	1.4230(9)58 1.422047 1.4225148 1.4230749–51 1.4229852
[¹ C ₃ ³ C ₁ im][NTf ₂]	1.42534	-3.02 ± 0.01	1.4252553
[¹ C ₂ ³ C ₂ im][NTf ₂]	1.42512	-3.04 ± 0.02	n.a.
[¹ C ₂ ³ C ₃ im][NTf ₂]	1.42668	-3.08 ± 0.01	n.a.
[¹ C ₄ ³ C ₁ im][NTf ₂]	1.42705	-3.10 ± 0.01	1.4265348 1.4269254,55 1.42756 1.451 57 1.4267250,51
[¹ C ₃ ³ C ₃ im][NTf ₂]	1.42805	-3.10 ± 0.01	n.a.

^a in the temperature interval, n_D , at a specific temperature, T , can be estimated using the following equation: $n_D(T/K) = n_D(298.15 K) + dn_D/dT \cdot (T/K - 298.15 K)$.

Table 7

Values of the surface tension γ (mN·m⁻¹) at 330.0 K, and surface thermodynamic functions $S\gamma$ (J·K⁻¹·m⁻²) and $H\gamma$ (J·m⁻²).

Ionic liquid	γ (330 K)/ mN·m ⁻¹	$(S\gamma \pm \sigma^a) \times 10^5 /$ (J·K ⁻¹ ·m ⁻²)	$(H\gamma \pm \sigma^a) \times 10^2 /$ (J·m ⁻²)
[¹ C ₁ Him][NTf ₂]	36.2	5.1 ± 0.5	5.3 ± 0.2
[¹ C ₂ Him][NTf ₂]	33.1	5.5 ± 0.2	5.13 ± 0.08
[¹ C ₂ ³ C ₃ im][NTf ₂]	32.0	4.6 ± 0.1	4.71 ± 0.02
[¹ C ₁ ² C ₁ Him][NTf ₂]	36.7	6.2 ± 0.1	5.73 ± 0.04
[¹ C ₄ ² C ₁ ³ C ₁ im][NTf ₂]	32.7	5.0 ± 0.1	4.91 ± 0.03

^aStandard deviation.