Multilocus Phylogeny and Antifungal Susceptibility of *Aspergillus* Section *Circumdati* from Clinical Samples and Description of *A. pseudosclerotiorum* sp. nov.

J. P. Z. Siqueira,^{a,b} D. A. Sutton,^c J. Gené,^a D. García,^a N. Wiederhold,^c S. W. Peterson,^d J. Guarro^a

Unitat de Micologia, Facultat de Medicina i Ciències de la Salut, IISPV, Universitat Rovira i Virgili, Reus, Spain^a; Laboratório de Microbiologia, Faculdade de Medicina de São José do Rio Preto, São José do Rio Preto, Brazil^b; Fungus Testing Laboratory, University of Texas Health Science Center, San Antonio, Texas, USA^c; Mycotoxin Prevention and Applied Microbiology Research Unit, National Center for Agricultural Utilization Research, Peoria, Illinois, USA^d

ABSTRACT A multilocus phylogenetic study was carried out to assess species identity of a set of 34 clinical isolates from *Aspergillus* section *Circumdati* from the United States and to determine their *in vitro* antifungal susceptibility against eight antifungal drugs. The genetic markers used were the internal transcribed spacer (ITS) region, and fragments of the beta-tubulin (*BenA*), calmodulin (*CaM*), and RNA polymerase II second largest subunit (*RPB2*) genes. The drugs tested were amphotericin B, itraconazole, posaconazole, voriconazole, anidulafungin, caspofungin, micafungin, and terbinafine. The most common species sampled was *A. westerdijkiae* (29.4%), followed by a novel species, which was described here as *A. pseudosclerotiorum* (23.5%). Other species identified were *A. sclerotiorum* (17.6%), *A. ochraceus* (8.8%), *A. subramaninii* (8.8%), and *A. insulicola* and *A. ochraceopetaliformis*, with two isolates (5.9%) of each. The drugs that showed the most potent activity were caspofungin, micafungin, and terbinafine, while amphotericin B showed the least activity.

KEYWORDS *Aspergillus*, *Circumdati* section, clinical isolates, molecular identification, phenotypic identification

Section *Circumdati* includes aspergilli with biserial conidial heads in shades of yellow to ochre, with mostly globose vesicles, and sclerotia variable in shape and color (1–3). It contains 26 species (3), with *A. ochraceus* being the best known and described as an important producer of many extrolites, including the mycotoxin ochratoxin A (3–5). This metabolite has nephrotoxic, immunosuppressive, teratogenic, and carcinogenic properties (6, 7) and is commonly found in coffee, rice, beverages, and other contaminated foodstuffs (3, 8). Several species in this section have been involved in different types of infections, such as onychomycosis, caused by *A. insulicola*, *A. melleus*, *A. ochraceopetaliformis*, *A. persii*, *A. sclerotiorum*, and *A. westerdijkiae* (9–14); otomycosis, caused by *A. sclerotiorum* (15); skin infection, caused by *A. westerdijkiae* (12); and pulmonary aspergillosis and osteomyelitis, caused by *A. ochraceus* (16, 17). Moreover, *A. ochraceus*, *A. sclerotiorum*, and *A. westerdijkiae* have been repetitively isolated from clinical specimens of immunocompromised patients, although in such cases their pathogenic role is uncertain (18–22).

There are few data on the *in vitro* antifungal susceptibility of species within section *Circumdati*. The azoles, especially itraconazole, appear to have good activity against *A.*

Received 30 September 2016 Returned for modification 22 November 2016 Accepted 30 December 2016

Accepted manuscript posted online 4 January 2017

Citation Siqueira JPZ, Sutton DA, Gené J, García D, Wiederhold N, Peterson SW, Guarro J. 2017. Multilocus phylogeny and antifungal susceptibility of *Aspergillus* section *Circumdati* from clinical samples and description of *A. pseudosclerotiorum* sp. nov. *J Clin Microbiol* 55:947–958. <https://doi.org/10.1128/JCM.02012-16>.

Editor David W. Warnock, University of Manchester

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to J. Gené, josepa.gene@urv.cat.

ochraceus and *A. sclerotiorum* (18, 23). In contrast, amphotericin B shows limited activity against species in this section (18, 23, 24), particularly against *A. westerdijkiae* (25).

Identification of *Aspergillus* species, traditionally based on morphological and physiological aspects (2), has changed recently with the use of DNA sequencing and multilocus analyses (26). Therefore, to assess the diversity of clinically relevant species within this section, a set of isolates with features characteristic of *Circumdati* section were identified molecularly. These clinical isolates were recovered between 2003 and 2015 in a U.S. reference laboratory. Moreover, the antifungal susceptibility of the most frequent species was determined against eight antifungal drugs.

RESULTS

Single-gene analyses of sequences revealed similar topologies for all of them, especially for the terminal branches. The internal transcribed spacer (ITS) marker was the least informative, being unable to discriminate between closely related species. However, the most basal clades could still be discerned in the analysis of this region, providing useful data in the concatenated tree. A limitation of the concatenated analysis that included all of the species in the *Circumdati* section was the lack of RNA polymerase II second largest subunit (*RPB2*) sequences for the ex-type strains of *A. affinis*, *A. occultus*, *A. pulvericola*, *A. salwaensis*, *A. sesamicola*, and *A. westlandensis*. However, analyses of the other three markers, i.e., ITS, beta-tubulin (*BenA*), and calmodulin (*CaM*), unequivocally demonstrated that none of the strains studied here corresponded to any of the above-mentioned species.

The final concatenated sequence alignment, with 58 strains and the 4 sequenced markers, consisted of 2,451 bp (ITS, 482 bp; *BenA*, 470 bp; *CaM*, 481 bp; *RPB2*, 1,018 bp), of which 941 sites were variable (ITS, 85; *BenA*, 250; *CaM*, 231; *RPB2*, 375) and 686 parsimony informative (ITS, 57; *BenA*, 182; *CaM*, 159; *RPB2*, 288). Topology trees inferred by the two phylogenetic methods were basically the same, with only minor differences in the support values of the internal nodes. The ML phylogenetic tree and the bootstrap and posterior probability values (Fig. 1) show that 26 of the strains included in this study clustered with the ex-type strains of six species from section *Circumdati*, i.e., *A. westerdijkiae* ($n = 10$; 29.4%), *A. sclerotiorum* ($n = 6$; 17.6%), *A. ochraceus* ($n = 3$; 8.8%), *A. subramanianii* ($n = 3$; 8.6%), *A. insulicola* ($n = 2$; 5.7%), and *A. ochraceopetaliformis* ($n = 2$; 5.9%). Interestingly, a group of eight isolates (25.7%) formed a well-supported clade together with sequences of two unidentified *Aspergillus* strains (NRRL 35028 and NRRL 35056). This clade represents an undescribed species, proposed here as *Aspergillus pseudosclerotiorum*.

The isolates examined here showed typical morphology of section *Circumdati* and matched those of the respective species. We found, however, that identification to the species level based only on phenotypic characteristics is difficult, but combining some of the phenotypic characteristics can make this feasible (Table 1). Among the species identified here, *A. westerdijkiae* and *A. ochraceus* were the only ones with finely roughened conidia; these two species could be distinguished from each other by the lack of or only slight growth at 37°C (0 to 9 mm) for *A. westerdijkiae*, while *A. ochraceus* reached 23 to 26 mm in diameter in 7 days at the same temperature. The other species identified here had smooth-walled conidia. In addition, *A. insulicola* was the only species that did not produce sclerotia but did produce a reddish-brown soluble pigment on Czapek yeast autolysate agar (CYA); *A. subramanianii* showed good growth at 37°C (39 to 46 mm in 7 days); the colonies of *A. ochraceopetaliformis* had dense white mycelial areas and poor sporulation after 7 days; and *A. sclerotiorum* produced yellow (3A7) to brownish-orange (6C3) colonies, which reached 56 to 58 mm in diameter in 7 days on CYA, with white sclerotia, abundant sporulation, and profuse growth at 37°C (32 to 36 mm). *Aspergillus pseudosclerotiorum* shares similar morphological features with *A. sclerotiorum* but with a slightly lower growth rate at 25°C (45 to 55 diameter in 7 days) and at 37°C (22 to 38 mm), smaller metulae (3 to 9 by 2.5 to 6 μm , compared with 7 to 15 by 4 to 7 μm in *A. sclerotiorum*), and its sclerotia become yellow to orange yellow with age.

FIG 1 Maximum likelihood tree obtained from analysis of combined ITS, *BenA*, *CaM*, and *RPB2* data set. Branch lengths are proportional to phylogenetic distance. Bootstrap support values/Bayesian posterior probability scores over 70/0.95 are indicated on the nodes. Fully supported branches (100/1) and ex-type strains are shown in boldface. UTHSCSA, University of Texas Health Science Center (San Antonio, Texas, USA).

In vitro susceptibility testing showed that the drugs with the most potent activity against all of the isolates tested were caspofungin (CFG), micafungin (MFG), and terbinafine (TBF), while amphotericin B (AMB) showed the lowest activity. The azoles (itraconazole [ITC], posaconazole [PSC], and voriconazole [VRC]), showed good activity

TABLE 1 Key morphological features of *Aspergillus* section *Circumdati* species identified in this study

Species	Sclerotium	Metula dimensions (μm)	Conidial ornamentation	CYA colony diam (mm) in 7 days at:	
				25°C	37°C
<i>A. insulicola</i>	Absent	6.5–12 by 3–5	Smooth	46–49	14–15
<i>A. ochraceopetaliformis</i>	Present	9–18 by 3.5–6	Smooth	38–46	27–29
<i>A. ochraceus</i>	Present	7–14 by 3–6	Finely roughened	44–49	23–26
<i>A. pseudosclerotiorum</i>	Present	3–9 by 2.5–6	Smooth	45–55	22–38
<i>A. sclerotiorum</i>	Present	7–15 by 4–7	Smooth	56–58	32–36
<i>A. subramanianii</i>	Present	8.5–14 by 3.5–6.5	Smooth	52–53	39–46
<i>A. westerdijkiae</i>	Present	8–18 by 4–7	Finely roughened	41–51	0–9

in general, with the exception of ITC against *A. sclerotiorum*. Interestingly, according to statistical analyses based on the Mann-Whitney test, the ITC MIC values showed significant differences between *A. sclerotiorum*, *A. ochraceus*, and *A. westerdijkiae* (GM of 11.31 $\mu\text{g}/\text{ml}$, 1.0 $\mu\text{g}/\text{ml}$, and 0.46 $\mu\text{g}/\text{ml}$, respectively; $P < 0.05$); however, differences were not significant between *A. sclerotiorum* and *A. pseudosclerotiorum* (0.89 $\mu\text{g}/\text{ml}$; $P = 0.06$) and *A. subramanianii* (4.0 $\mu\text{g}/\text{ml}$; $P = 0.43$). Regarding the new species, in general the drugs tested showed good activity against *A. pseudosclerotiorum*. Higher MIC values were observed only for AMB and VRC. Results of the *in vitro* susceptibility test are summarized in Table 2.

Taxonomy. *Aspergillus pseudosclerotiorum* J. P. Z. Siqueira, Deanna A. Sutton & Gené sp. nov. (Mycobank accession no. [MB818572](https://www.mycobank.org/MB818572)) (Fig. 2). Etymology: the name refers to the morphological similarity to *A. sclerotiorum*. Holotype: USA, Pennsylvania, isolated from

TABLE 2 Results of *in vitro* antifungal susceptibility test for 30 isolates of *Aspergillus* section *Circumdati*

Species (no. of isolates) and parameter ^a	MIC or MEC ($\mu\text{g}/\text{ml}$) for ^b :							
	AMB	AFG	CFG	MFG	ITC	PSC	VRC	TBF
<i>A. ochraceus</i> (3)								
GM	16.0	0.25	0.04	0.03	1.0	0.31	2.0	0.03
MIC range	16.0	0.12–0.5	0.03–0.06	0.03	1.0	0.25–0.5	2.0	0.03
Mode	16.0	0.5	0.03	0.03	1.0	0.25	2.0	0.03
<i>A. subramanianii</i> (3)								
GM	>16.0	0.10	0.03	0.03	4.0	0.79	4.0	0.03
MIC range	16.0–>16.0	0.03–0.25	0.03	0.03	4.0	0.5–1.0	4.0	0.03
Mode	>16.0	0.25	0.03	0.03	4.0	1.0	4.0	0.03
<i>A. sclerotiorum</i> (6)								
GM	4.76	0.03	0.04	0.03	11.31	1.0	3.36	0.03
MIC range	4.0–8.0	0.03	0.03–0.06	0.03	4.0–>16.0	1.0	2.0–4.0	0.03
Mode	4.0	0.03	0.03	0.03	>16.0	1.0	4.0	0.03
<i>A. pseudosclerotiorum</i> (8)								
GM	5.04	0.04	0.03	0.03	0.89	0.25	1.41	0.03
MIC range	2.0–>16	0.03–0.12	0.03–0.06	0.03	0.25–>16.0	0.12–0.5	1.0–2.0	0.03
Mode	4.0	0.03	0.03	0.03	0.5	0.25	2.0	0.03
<i>A. westerdijkiae</i> (10)								
GM	>16.0	0.14	0.03	0.03	0.46	0.29	1.08	0.03
MIC range	>16.0	0.03–1.0	0.03–0.06	0.03–0.06	0.12–1.0	0.12–0.5	1.0–2.0	0.03
Mode	>16.0	0.25	0.03	0.03	0.5	0.25	1.0	0.03
MIC ₉₀	>16.0	0.5	0.06	0.06	0.5	0.5	1.0	0.03
Total (30)								
GM	12.82	0.08	0.03	0.03	1.28	0.39	1.74	0.03
MIC range	2.0–>16.0	0.03–1.0	0.03–0.06	0.03–0.06	0.12–>16.0	0.12–1.0	1.0–4.0	0.03
Mode	>16.0	0.03	0.03	0.03	0.5	0.25	1.0	0.03
MIC ₉₀	>16.0	0.5	0.06	0.03	4.0	1.0	4.0	0.03

^aGM, geometric mean.

^bAMB, amphotericin B; AFG, anidulafungin; CFG, caspofungin; MFG, micafungin; ITC, itraconazole; PSC, posaconazole; VRC, voriconazole; TBF, terbinafine; MEC, minimum effective concentration for AFG, CFG, and MFG.

FIG 2 Morphological features of *Aspergillus pseudosclerotiorum* sp. nov. (UTHSCA DI 15-13 [a to n] and UTHSCSA DI16-383 [o]). (a, b, e, and f) Front and reverse of colonies on CYA and MEA, respectively, after 7 days at 25°C. (c, d, g, and h) Front of colonies on DG18, OA, YES, and CREA, respectively, after 7 days at 25°C. (i) Enlarged view of conidial heads on CYA after 7 days at 25°C. (j) Sclerotia on CYA after 14 days at 25°C. (k) Conidia. (l) Conidiophores and a sclerotium. (m) Detail of conidiophore stipe. (n and o) Details of conidial heads. Scale bars: 10 μ m (k, m, n, and o) and 100 μ m (l).

TABLE 3 List of *Aspergillus* section *Circumdati* species, their isolate information, sequences generated in this study, and those retrieved from GenBank^d

Species	Isolate no. ^a	Origin ^b	Yr	GenBank/EMBL accession no. ^c			
				ITS	BenA	CaM	RPB2
<i>A. affinis</i>	ATCC MYA-4773 ^T			GU721090	GU721092	GU721091	
<i>A. auricomus</i>	NRRL 391 ^T			EF661411	EF661320	EF661379	EF661301
<i>A. bridgeri</i>	NRRL 13000 ^T			EF661404	EF661335	EF661358	EF661290
<i>A. cretensis</i>	NRRL 35672 ^T			FJ491572	AY819977	FJ491534	EF661311
<i>A. elegans</i>	NRRL 4850 ^T			EF661414	EF661349	EF661390	EF661316
<i>A. fresenii</i>	NRRL 407 ^T			EF661409	EF661341	EF661382	EF661296
<i>A. insulicola</i>	NRRL 6138 ^T			EF661430	EF661353	EF661396	EF661286
	UTHSCSA DI16-374	Marine	2003	LT574681	LT574716	LT574751	LT574786
	UTHSCSA DI16-402	Marine	2009	LT574682	LT574717	LT574752	LT574787
<i>A. melleus</i>	NRRL 5103 ^T			EF661425	EF661326	EF661391	EF661309
<i>A. muricatus</i>	NRRL 35674 ^T			EF661434	EF661356	EF661377	EF661314
<i>A. neobridgeri</i>	NRRL 13078 ^T			EF661410	EF661345	EF661359	EF661298
<i>A. occultus</i>	CBS 137330 ^T			KJ775443	KJ775061	KJ775239	
<i>A. ochraceopetaliformis</i>	NRRL 4752 ^T			EF661429	EF661350	EF661388	EF661283
	UTHSCSA DI16-387	BAL	2006	LT574683	LT574718	LT574753	LT574788
	UTHSCSA DI16-392	Marine	2007	LT574684	LT574719	LT574754	LT574789
<i>A. ochraceus</i>	NRRL 398 ^T			EF661419	EF661322	EF661381	EF661302
	UTHSCSA DI15-10	BAL	2012	LT574686	LT574721	LT574756	LT574791
	UTHSCSA DI15-11	Heart valve	2013	LT574687	LT574722	LT574757	LT574792
	UTHSCSA DI16-384	Ear	2006	LT574685	LT574720	LT574755	LT574790
<i>A. ostianus</i>	NRRL 420 ^T			EF661421	EF661324	EF661385	EF661304
<i>A. pallidofulvus</i>	NRRL 4789 ^T			EF661423	EF661328	EF661389	EF661306
<i>A. persii</i>	NRRL 35669 ^T			FJ491580	AY819988	FJ491559	EF661295
<i>A. pseudoelegans</i>	CBS 112796 ^T			FJ491590	AY819962	FJ491552	EF661282
<i>A. pseudosclerotiorum</i>	NRRL 35028			EF661407	EF661343	EF661362	EF661293
	NRRL 35056			EF661405	EF661344	EF661364	EF661294
	UTHSCSA DI15-13^T	Lung biopsy	2014	LT574713	LT574748	LT574783	LT574818
	UTHSCSA DI15-14	BAL	2014	LT574714	LT574749	LT574784	LT574819
	UTHSCSA DI15-15	Lung tissue	2015	LT574715	LT574750	LT574785	LT574820
	UTHSCSA DI16-373	Sputum	2003	LT574707	LT574742	LT574777	LT574812
	UTHSCSA DI16-380	BAL	2006	LT574708	LT574743	LT574778	LT574813
	UTHSCSA DI16-383	BAL	2006	LT574709	LT574744	LT574779	LT574814
	UTHSCSA DI16-385	Sputum	2006	LT574710	LT574745	LT574780	LT574815
	UTHSCSA DI16-386	Lung mass	2006	LT574711	LT574746	LT574781	LT574816
<i>A. pulvericola</i>	CBS 137327 ^T			KJ775440	KJ775055	KJ775236	
<i>A. robustus</i>	NRRL 6362 ^T			EF661176	EU014101	EF661357	EF661033
<i>A. roseoglobulosus</i>	NRRL 4565 ^T			FJ491583	AY819984	FJ491555	EF661299
<i>A. salwaensis</i>	DTO 297B3 ^T			KJ775447	KJ775056	KJ775244	
<i>A. sclerotiorum</i>	NRRL 415 ^T			EF661400	EF661337	EF661384	EF661287
	UTHSCSA DI15-12	Sputum	2014	LT574693	LT574728	LT574763	LT574798
	UTHSCSA DI16-395	Sputum	2007	LT574688	LT574723	LT574758	LT574793
	UTHSCSA DI16-398	BAL	2008	LT574689	LT574724	LT574759	LT574794
	UTHSCSA DI16-404	Sputum	2009	LT574690	LT574725	LT574760	LT574795
	UTHSCSA DI16-399	BAL	2009	LT574691	LT574726	LT574761	LT574796
	UTHSCSA DI16-409	Eye	2014	LT574692	LT574727	LT574762	LT574797
<i>A. sesamicola</i>	CBS 137324 ^T			KJ775437	KJ775063	KJ775233	
<i>A. steinii</i>	NRRL 35675 ^T			EF661416	EF661347	EF661378	JN121428
<i>A. subramanianii</i>	NRRL 6161 ^T			EF661403	EF661339	EF661397	EF661289
	UTHSCSA DI16-378	Lung tissue	2005	LT574694	LT574729	LT574764	LT574799
	UTHSCSA DI16-389	Wound	2006	LT574695	LT574730	LT574765	LT574800
	UTHSCSA DI16-390	Foot	2006	LT574696	LT574731	LT574766	LT574801
<i>A. tanneri</i>	NRRL 62425 ^T			JN853798	JN896582	JN896583	JN896585

(Continued on next page)

TABLE 3 (Continued)

Species	Isolate no. ^a	Origin ^b	Yr	GenBank/EMBL accession no. ^c			
				ITS	BenA	CaM	RPB2
<i>A. westerdijkiae</i>	NRRL 3174 ^T			EF661427	EF661329	EF661360	EF661307
	UTHSCSA DI15-5	BAL	2014	LT574703	LT574738	LT574773	LT574808
	UTHSCSA DI15-6	Sputum	2014	LT574704	LT574739	LT574774	LT574809
	UTHSCSA DI15-7	Nail	2015	LT574705	LT574740	LT574775	LT574810
	UTHSCSA DI15-8	Marine	2011	LT574706	LT574741	LT574776	LT574811
	UTHSCSA DI16-376	Unknown	2004	LT574697	LT574732	LT574767	LT574802
	UTHSCSA DI16-377	Unknown	2004	LT574698	LT574733	LT574768	LT574803
	UTHSCSA DI16-379	BAL	2005	LT574699	LT574734	LT574769	LT574804
	UTHSCSA DI16-388	Lung mass	2006	LT574700	LT574735	LT574770	LT574805
	UTHSCSA DI16-391	Lung nodule	2007	LT574701	LT574736	LT574771	LT574806
	UTHSCSA DI16-393	Sputum	2007	LT574702	LT574737	LT574772	LT574807
<i>A. westlandensis</i>	CBS 137321 ^T			KJ775434	KJ775066	KJ775230	

^aATCC, American Type Culture Collection; CBS, CBS-KNAW Fungal Biodiversity Centre (Utrecht, the Netherlands); DTO, Applied and Industrial Mycology Department Collection (Utrecht, Netherlands); NRRL, Agriculture Research Service Culture Collection (Peoria, NY); UTHSCSA, University of Texas Health Science Center (San Antonio, TX). A superscript T indicates an ex-type strain.

^bBAL, bronchoalveolar lavage fluid specimens.

^cITS, internal transcribed spacer regions of the rDNA and 5.8S region; *BenA*, β -tubulin; *CaM*, calmodulin; *RPB2*, partial RNA polymerase II, second largest subunit.

^dSequences generated in this study are in boldface.

lung biopsy specimen (human), D. A. Sutton, 2014 (CBS H-22808; culture ex-types: UTHSCSA DI15-13, FMR 14449, CBS 141845).

Colonies on CYA at 7 days reached 45- to 55-mm diameter at 25°C; at 30°C exhibited optimum growth, reaching 55- to 64-mm diameter; at 37°C reached 22- to 38-mm diameter; and at 40°C showed restricted growth. Colonies on CYA were pale yellow (3A3) to reddish white (7A3) at the center, white toward the periphery, cottony to floccose, and usually granulose due to the presence of abundant sclerotia, margin fimbriate; reverse yellow (3A7) to greyish yellow (3B5); colorless exudates present in most isolates; little soluble pigment produced, yellow (3A6), or absent. On malt extract agar (MEA), colonies similar to CYA but with slower growth, reaching 34 to 42 mm at 7 days. On yeast extract sucrose agar (YES), colonies showed fastest growth, reaching 56 to 66 mm at 7 days, white, cottony to floccose, with abundant sclerotia; reverse yellow (3A6) to greyish yellow (4B5), sulcate; exudates abundant, colorless to yellowish white (3A2). On dichloran 18% glycerol agar (DG18), colonies reaching 28 to 34 mm at 7 days, with white to light orange (5A4) compact center, and white fluffy mycelium toward periphery; reverse yellowish white (3A2) to pale yellow (3A3); sporulation sparsely produced only with age; sclerotia absent. On oatmeal agar (OA), colonies reaching 24 to 27 mm at 7 days, yellowish white (3A2) to greyish yellow (4B4), sandy to dusty, with a more compact center, margin regular; reverse yellowish white (4A2) to greyish yellow (4B6). On creatine-sucrose agar (CREA), colonies reaching 22 to 28 mm at 7 days, white, dense at the center, sparse aerial mycelium toward the periphery; acid production absent. Micromorphology consisting of conidiophores with biserial and radiating conidial heads; stipes septate with rough walls, subhyaline to pale brown, 120 to 980 μ m long by 2.5 to 8 μ m wide; vesicles mainly globose, occasionally subglobose, 7- to 31- μ m diameter; metulae cylindrical, 3 to 9 by 2.5 to 6 μ m, usually covering 100% of vesicle, with exception of the strain UTHSCSA DI16-383, which covered 75% of vesicle; phialides ampulliform, 4.5 to 8 by 1.25 to 3 μ m; conidia globose, smooth-walled, 1.5- to 3- μ m diameter; sclerotia present (except in UTHSCSA DI16-380), 150- to 507- μ m diameter, white to light orange (5A4), becoming yellow (3A6) to orange yellow (4A6) with age.

DISCUSSION

In this study, we identified a total of six species in the section *Circumdati* from clinical samples, some of which contained a relatively large number of isolates. Although their role as etiologic agents in these cases is unknown, detection of 34 isolates

of this section over a period of 12 years in a single reference center, together with some reports on infections produced by members of this section in the same period (15, 17, 18, 22, 27), highlights the importance of these fungi in the clinical setting. The degree of morphological similarity among the species of the *Circumdati* section, as with other groups of *Aspergillus*, requires DNA sequencing analysis for a definitive identification.

As was mentioned above, the most common *Aspergillus* species in the set of isolates studied here was *A. westerdijkiae*, a species described in 2004 and known to produce ochratoxin (28). It is noteworthy that the *A. ochraceus* strain from which ochratoxin A was discovered was later reidentified as *A. westerdijkiae*. This means that some isolates reported as *A. ochraceus*, especially the ones identified before 2004, in fact may be *A. westerdijkiae* (29). Growth rates at 37°C can be a useful feature to differentiate between these species without sequencing (3). *Aspergillus westerdijkiae* is commonly found in environmental samples (30) and as a food (31) and indoor contaminant (31–35). In the clinical setting, *A. westerdijkiae* has been linked to superficial infections (12) and isolated from sputum of immunocompromised patients in Tunisia (19). In our case, this species was mainly identified from respiratory specimens but also from a nail and in a sample from a marine animal (Table 3).

It is worth noting that the second most frequently identified species in the present study was a novel one, *A. pseudosclerotiorum*. This species is closely related to *A. bridgeri*, *A. persii*, *A. salwaensis*, *A. sclerotiorum*, and *A. subramanianii*. While these species could not be discriminated from each other using the ITS-based fungal barcode, *A. pseudosclerotiorum* was noted to have unique sequences for the other three markers (*BenA*, *CaM*, and *RPB2*). Phenotypically, *A. pseudosclerotiorum* generally can be distinguished from the above-mentioned aspergilli by its growth rate on different media and temperatures, colony pigmentation, and degree of sporulation, as well as sclerotia and conidiophore features. *Aspergillus bridgeri* produces brown colonies (3, 36). *A. persii* grows faster on OA (35- to 38-mm diameter in 7 days) and DG18 (45- to 50-mm diameter in 7 days) (3). *Aspergillus salwaensis* produces a characteristic yellowish-orange soluble pigment and usually has conidiophores with vesicles flattened at the apex (3). *Aspergillus subramanianii* grows faster on CYA at 37°C (39- to 46-mm diameter in 7 days). *Aspergillus sclerotiorum* grows faster on CYA at 25°C (54- to 57-mm diameter in 7 days), and at 37°C (32- to 36-mm diameter at 37°C) it shows a higher level of sporulation and its sclerotia are white to cream colored. However, one of the eight isolates of *A. pseudosclerotiorum* (UTHSCSA DI16-380), which showed 99.6% similarity with the other isolates, produced atypical colonies (i.e., brownish and profusely sporulated). The size of metulae is also a diagnostic feature for *A. pseudosclerotiorum*, because they are smaller (3 to 9 by 2.5 to 6 μm) than those of the related species (6.5 to 10 by 3.5 to 5.5 μm in *A. bridgeri*, 9 to 17.5 by 4 to 7.5 μm in *A. persii*, 8 to 21 by 3.5 to 6 μm in *A. salwaensis*, 8 to 16 by 4.5 to 7 μm in *A. sclerotiorum*, and 9 to 14 by 4 to 6.5 μm in *A. subramanianii*) (3). Although all isolates of *A. pseudosclerotiorum* were from the human respiratory tract (i.e., BAL fluid samples, sputum, and lung tissue), further studies are needed to determine the pathogenic role of this new fungus.

The third most common species sampled was *A. sclerotiorum*, which has been reported to cause superficial infections, such as onychomycosis and otomycosis (10, 14, 15). Here, most of the isolates were also from the human respiratory tract. *Aspergillus sclerotiorum* is found worldwide, commonly isolated from soil, and reported as a species of biotechnological importance due to its ability to produce a wide range of compounds (37–39).

The best-known species in the section, *A. ochraceus*, was poorly represented in this study (8.8%). In contrast, it is commonly found on coffee, rice, dried fruits, and nuts (8, 40, 41) and is capable of producing different metabolites (42–44). It was reported previously in pulmonary infections based on morphological identifications (16, 20). More recently, it has been identified in a case of osteomyelitis (17) and has also been isolated from immunocompromised patients (18, 19). Carpagnano et al. often found *A. ochraceus* in exhaled breath condensate of lung cancer patients (27). In other mammals,

it was associated with a case of otomycosis in a dog (45). Here, the three isolates were from different clinical origins (i.e., BAL fluid, ear, and heart valve).

Of the three other species identified, *A. insulicola* and *A. ochraceopetaliformis* have been reported from cases of onychomycoses (9, 12), while *A. subramanianii* was recovered for the first time from clinical specimens. Concerning the latter species, it is noteworthy that two isolates (UTHSCSA DI16-378 and UTHSCSA DI16-389) formed a clade slightly separate from the other *A. subramanianii* isolates (Fig. 1); however, the genetic identity (99.3%) with the ex-type strain and phenotypic similarity confirm their identification as *A. subramanianii*. This species could be considered a potential agent of human infections because of its ability to grow at 37°C and the deep-tissue origin of the isolates (lung tissue and wound).

Data available on the *in vitro* susceptibility of section *Circumdati* aspergilli against antifungal drugs are limited to a few reports with a small number of isolates tested. Here, the three echinocandins and TBF exhibited potent activity against the fungi tested. Similar results were obtained in our previous study on *Aspergillus* section *Versicolores* (46). TBF also has been reported to be highly effective *in vitro* against clinically relevant *Aspergillus* species, such as *A. flavus*, *A. niger*, *A. nidulans*, or *A. terreus*, and even against numerous isolates of *A. fumigatus sensu stricto* (47–49). To our knowledge, however, there is no previous information available on the activity of TBF against section *Circumdati* species. Results observed for echinocandins, especially MFG and anidulafungin (AFG), could be expected since, in general, they have been reported to be effective *in vitro* on *Aspergillus* species (50, 51). With respect to *Circumdati* aspergilli, Arabatzis et al. (18) tested three echinocandins against two isolates of *A. ochraceus* and one of *A. sclerotiorum* and reported high MICs only for CFG. In contrast, Gheith et al. (21) tested CFG against one isolate of *A. ochraceus* and one of *A. westerdijkiae* and reported low MICs, which is similar to our findings. AMB showed the least activity against the isolates tested, especially for *A. ochraceus*, *A. subramanianii*, and *A. westerdijkiae*. High AMB MICs were also observed for species in section *Circumdati* (i.e., *A. melleus*, *A. ochraceus*, and *A. pallidofulvus*), recently identified from human clinical specimens in India, in contrast to the results obtained in the same study for most isolates of *A. fumigatus*, *A. flavus*, and *A. terreus*, which were susceptible to antifungals tested there (51). PSC was the azole with the most potent activity against the strains tested, which agrees with Alastruey-Izquierdo et al. (25), Gheith et al. (21), and Masih et al. (51); however, the study of Arabatzis et al. (18) showed higher MICs for PSC. Recently, Babamahmoodi et al. (17) reported a case of osteomyelitis by *A. ochraceus*, for which the strain showed azole MICs (PSC, 0.032 µg/ml; VRC and ITC, 1.0 µg/ml) similar to ours (Table 2), and the patient improved after 4 months of treatment with VRC.

In conclusion, taxonomic studies are very important to assess the distribution of fungal species and their identity in clinical settings. In our study of clinical isolates within section *Circumdati* from a reference collection in the United States, we not only identified *A. subramanianii* as being associated with human specimens for the first time but also described a new taxon, *A. pseudosclerotiorum*, as one of the most frequent species of the section in this set of isolates. However, data from more isolates are needed to determine more reliable MICs of the different antifungal drugs against the species of this section and to determine the pathogenic role of these fungi in human and animal infections.

MATERIALS AND METHODS

Fungal isolates. A total of 34 *Aspergillus* isolates received from the Fungus Testing Laboratory at the University of Texas Health Science Center (San Antonio, TX, USA) were investigated. Based on morphological features, the isolates were identified as belonging to section *Circumdati*. Most isolates studied were from human clinical specimens, mainly from the respiratory tract ($n = 22$; 64.7%), although other human clinical sources were noted as well ($n = 8$; 23.5%). In addition, four isolates were from marine animals (Table 3).

Morphological characterization. The isolates were characterized morphologically by following the criteria recommended by Samson et al. (1). Briefly, colony morphology and growth rates were determined after 7 days of incubation on CYA (Becton, Dickinson and Company, Sparks, MD, USA) at 25°C and 37°C and on MEA (Pronadisa, Madrid, Spain) at 25°C. After 10 to 14 days of incubation, microscopic

structures were examined and measured from MEA cultures in wet mounts with 60% lactic acid and a drop of 70% ethanol to wash out the excess conidia. A minimum of 20 of each structure was measured in order to cover all of the size ranges. Photographs were made using a Zeiss Axio Imager M1 light microscope (Zeiss, Oberkochen, Germany) with a mounted DeltaPix Infinity X digital camera using Nomarski differential interference contrast and phase-contrast optics.

DNA extraction, amplification, and sequencing. Total genomic DNA was extracted from MEA cultures after 7 days of incubation at 25°C using the FastDNA kit and the FastPrep instrument (MP Biomedicals, Irvine, CA, USA) according to the manufacturer's specifications. Four genetic markers were amplified, i.e., the ITS region of the rRNA, which comprises ITS1, the 5.8S gene, and ITS2 regions, and fragments of the *BenA*, *CaM*, and *RPB2* genes (1, 26). The primers used were ITS5 and ITS4 for the ITS region (52), Bt2a and Bt2b for *BenA* (53), Cmd5 and Cmd6 for *CaM* (54), and 5F and 7CR for *RPB2* (55). PCR products were sequenced in both directions, using the same primers, at Macrogen Europe (Macrogen Inc., Amsterdam, Netherlands). Sequences were assembled and edited using SeqMan v.7.0.0 (DNASTAR, Madison, WI, USA).

Molecular identification and phylogenetic analysis. Phylogenetic analyses were first performed individually for each gene. Since the topologies proved to be congruent with the incongruence length difference test (56), a concatenated analysis was performed. Sequences of the ex-type strains of all the species in section *Circumdati* were obtained from GenBank and added to the analyses. *Aspergillus tanneri* (section *Tanneri*) and *A. robustus* (section *Robusti*) were used as outgroups. In addition, GenBank sequences of two strains identified only as *Aspergillus* spp. (NRRL 35028 and NRRL 35026) were also added to the analyses because they formed a distinct lineage in section *Circumdati* (26). For multiple-sequence alignment, ClustalW was used together with MUSCLE in MEGA v.6 (57), followed by manual adjustments. The maximum likelihood (ML) analysis was conducted with MEGA v.6, as well as to estimate the best nucleotide substitution model. Support of the internal branches was assessed by the bootstrap method with 1,000 replications, where values of ≥ 70 were considered significant. Bayesian inference (BI) was performed using MrBayes v.3.1.2 (58). The evolutionary model that best fit each gene was assessed by MrModelTest (59). Markov chain Monte Carlo (MCMC) sampling was performed with two simultaneous runs for 1 million generations, with samples taken every 100 generations. The 50% majority rule consensus trees and posterior probability (pp) values were calculated after removing the first 25% of the resulting trees for burn-in. A pp value of ≥ 0.95 was considered significant.

Antifungal susceptibility testing. Isolates of the most frequent *Aspergillus* species identified here were tested against eight antifungal drugs using the methods in the CLSI M38-A2 reference standard (60). The antifungal agents, obtained as pure powders, were AMB (Sigma-Aldrich Quimica S.A., Madrid, Spain), ITC (Jansen Pharmaceuticals, Beerse, Belgium), PSC (Schering-Plough Research Institut, NJ, USA), VRC (Pfizer S.A., Madrid, Spain), AFG (Pfizer S.A., Madrid, Spain), CFG (Merk & Co., Inc., Rahway, USA), MFG (Astellas Pharma, Madrid, Spain), and TBF. The MIC was defined as the lowest drug concentration that produced 100% inhibition of visible fungal growth for AMB and the azoles (ITC, PSC, and VRC) and 80% for TBF. The minimum effective concentration (MEC) was determined for the echinocandins (AFG, CFG, and MFG) and was defined microscopically as the lowest concentration of drug that permitted growth of small, rounded, compact hyphal forms, as opposed to the long, unbranched hyphal clusters that were seen in the growth control. The quality control strain *Candida krusei* ATCC 6258 was used in each test, and the MIC values were according to CLSI guideline ranges. All tests were carried out in duplicate, on different days, to assess reproducibility. Statistical analyses were performed using Prism software for Windows v.6.0 (GraphPad Software, San Diego, CA).

Accession number(s). Newly generated sequences from this study were deposited in GenBank/EMBL databases under the accession numbers listed in Table 3 and in MycoBank under accession number MB818572.

ACKNOWLEDGMENTS

This study was supported by the Spanish Ministerio de Economía y Competitividad, grant CGL2013-43789-P, and by CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil), grant BEX 0623/14-8.

REFERENCES

- Samson RA, Visagie CM, Houbaken J, Hong SB, Hubka V, Klaassen CH, Perrone G, Seifert KA, Susca A, Tanney JB, Varga J, Kocsube S, Szigeti G, Yaguchi T, Frisvad JC. 2014. Phylogeny, identification and nomenclature of the genus *Aspergillus*. *Stud Mycol* 78:141–173. <https://doi.org/10.1016/j.simyco.2014.07.004>.
- Raper KB, Fennell DL. 1965. The genus *Aspergillus*. Robert E. Krieger Publishing Co., Inc., Huntington, NY.
- Visagie CM, Varga J, Houbaken J, Meijer M, Kocsube S, Yilmaz N, Fotedar R, Seifert KA, Frisvad JC, Samson RA. 2014. Ochratoxin production and taxonomy of the yellow aspergilli (*Aspergillus* section *Circumdati*). *Stud Mycol* 78:1–61. <https://doi.org/10.1016/j.simyco.2014.07.001>.
- Ciegler A. 1972. Bioproduction of ochratoxin A and penicillic acid by members of the *Aspergillus ochraceus* group. *Can J Microbiol* 18: 631–636. <https://doi.org/10.1139/m72-100>.
- Varga J, Kevei E, Rinyu E, Téren J, Kozakiewicz Z. 1996. Ochratoxin production by *Aspergillus* species. *Appl Environ Microbiol* 62:4461–4464.
- Lea T, Steien K, Stormer FC. 1989. Mechanism of ochratoxin A-induced immunosuppression. *Mycopathologia* 107:153–159. <https://doi.org/10.1007/BF00707553>.
- Pfohl-Leszkowicz A, Manderville RA. 2007. Ochratoxin A: an overview on toxicity and carcinogenicity in animals and humans. *Mol Nutr Food Res* 51:61–99. <https://doi.org/10.1002/mnfr.200600137>.
- Duarte SC, Pena A, Lino CM. 2010. A review on ochratoxin A occurrence and effects of processing of cereal and cereal derived food products. *Food Microbiol* 27:187–198. <https://doi.org/10.1016/j.fm.2009.11.016>.
- Brasch J, Varga J, Jensen JM, Egberts F, Tintelnot K. 2009. Nail infection by *Aspergillus ochraceopetaliformis*. *Med Mycol* 47:658–662. <https://doi.org/10.1080/13693780902803032>.

10. Amri M, Gorcii M, Essabbah N, Belhajali H, Letscher-Bru V, Zili J, Azaiez R, Babba H. 2010. *Aspergillus sclerotiorum*: à propos d'un cas d'onychomycose en Tunisie. *J Med Mycol* 20:128–132. <https://doi.org/10.1016/j.mycmed.2010.03.004>.
11. Zotti M, Machetti M, Perotti M, Barabino G, Persi A. 2010. A new species, *Aspergillus persii*, as an agent of onychomycosis. *Med Mycol* 48:656–660. <https://doi.org/10.3109/13693780903420641>.
12. Hubka V, Kubatova A, Mallatova N, Sedlacek P, Melichar J, Skorepova M, Mencl K, Lyskova P, Sramkova B, Chudickova M, Hamal P, Kolarik M. 2012. Rare and new etiological agents revealed among 178 clinical *Aspergillus* strains obtained from Czech patients and characterized by molecular sequencing. *Med Mycol* 50:601–610. <https://doi.org/10.3109/13693786.2012.667578>.
13. Zotti M, Agnoletti AF, Vizzini A, Cozzani E, Parodi A. 2015. Onychomycosis from *Aspergillus melleus*, a novel pathogen for humans. Fungal identification and *in vitro* drug susceptibility. *Exp Dermatol* 24:966–968.
14. García-Martos P, Guarro J, Gené J, Mira J, Linares M, Ortoneda M. 2005. Onychomycosis caused by *Aspergillus sclerotiorum*. *J Mycol Med* 11: 222–224.
15. Harima N, Inoue T, Kubota T, Okada O, Ansai S, Manabe M, Ichinoe M, Kasai T. 2004. A case of otomycosis caused by *Aspergillus sclerotiorum*. *J Dermatol* 31:949–950. <https://doi.org/10.1111/j.1346-8138.2004.tb00635.x>.
16. Novey HS, Wells ID. 1978. Allergic bronchopulmonary aspergillosis caused by *Aspergillus ochraceus*. *Am J Clin Pathol* 70:840–843. <https://doi.org/10.1093/ajcp/70.5.840>.
17. Babamahmoodi F, Shokohi T, Ahangarkani F, Nabili M, Afzalian Ashkezari E, Alinezhad S. 2015. Rare case of *Aspergillus ochraceus* osteomyelitis of calcaneus bone in a patient with diabetic foot ulcers. *Case Rep Med* 2015:509827.
18. Arabatzis M, Kambouris M, Kyprianou M, Chrysaki A, Foustoukou M, Kanellopoulou M, Kondyli L, Kouppari G, Koutsia-Karouzou C, Lebesi E, Pangalis A, Petinaki E, Stathi A, Trikkas-Graphakos E, Vartzioti E, Vogiatzi A, Vyzantiadis TA, Zerva L, Velegraki A. 2011. Polyphasic identification and susceptibility to seven antifungals of 102 *Aspergillus* isolates recovered from immunocompromised hosts in Greece. *Antimicrob Agents Chemother* 55:3025–3030. <https://doi.org/10.1128/AAC.01491-10>.
19. Gheith S, Saghruni F, Bannour W, Ben Youssef Y, Khelif A, Normand A-C, Ben Said M, Piarroux R, Njah M, Ranque S. 2014. Characteristics of invasive aspergillosis in neutropenic haematology patients (Sousse, Tunisia). *Mycopathologia* 177:281–289. <https://doi.org/10.1007/s11046-014-9742-8>.
20. Wierzbicka M, Podsiadlo B, Janczarski M. 1997. Inwazyjna aspergiloza płuc wywołana przez *Aspergillus ochraceus*. *Pneumonol Alergol Pol* 65: 254–260.
21. Gheith S, Saghruni F, Bannour W, Ben Youssef Y, Khelif A, Normand A-C, Piarroux R, Ben Said M, Njah M, Ranque S. 2014. *In vitro* susceptibility to amphotericin B, itraconazole, voriconazole, posaconazole and caspofungin of *Aspergillus* spp. isolated from patients with haematological malignancies in Tunisia. *Springerplus* 3:19–26. <https://doi.org/10.1186/2193-1801-3-19>.
22. Negri CE, Goncalves SS, Zafranski H, Bergamasco MD, Aquino VR, Castro PT, Colombo AL. 2014. Cryptic and rare *Aspergillus* species in Brazil: prevalence in clinical samples and *in vitro* susceptibility to triazoles. *J Clin Microbiol* 52:3633–3640. <https://doi.org/10.1128/JCM.01582-14>.
23. García-Martos P, García-Agudo L, Gutiérrez-Calzada J, Ruiz-Aragón J, Saldarreja A, Marín P. 2005. Actividad *in vitro* de anfotericina B, itraconazol y voriconazol frente a 20 especies de *Aspergillus* empleando el método de microdilución Sensititre. *Enferm Infecc Microbiol Clin* 23: 15–18. <https://doi.org/10.1157/13070403>.
24. Wildfeuer A, Seidl HP, Paule I, Haberleiter A. 1998. *In vitro* evaluation of voriconazole against clinical isolates of yeasts, moulds and dermatophytes in comparison with itraconazole, ketoconazole, amphotericin B and griseofulvin. *Mycoses* 41:309–319. <https://doi.org/10.1111/j.1439-0507.1998.tb00344.x>.
25. Alastruey-Izquierdo A, Mellado E, Cuenca-Estrella M. 2012. Current section and species complex concepts in *Aspergillus*: recommendations for routine daily practice. *Ann N Y Acad Sci* 1273:18–24. <https://doi.org/10.1111/j.1749-6632.2012.06822.x>.
26. Peterson SW. 2008. Phylogenetic analysis of *Aspergillus* species using DNA sequences from four loci. *Mycologia* 100:205–226. <https://doi.org/10.3852/mycologia.100.2.205>.
27. Carpaghano GE, Lacedonia D, Palladino GP, Logrieco G, Crisetti E, Susca A, Logrieco A, Foschino-Barbaro MP. 2014. *Aspergillus* spp. colonization in exhaled breath condensate of lung cancer patients from Puglia region of Italy. *BMC Pulm Med* 14:22. <https://doi.org/10.1186/1471-2466-14-22>.
28. Frisvad JC, Frank JM, Houbraken JAMP, Kuijpers AFA, Samson RA. 2004. New ochratoxin A producing species of *Aspergillus* section *Circumdati*. *Stud Mycol* 50:23–43.
29. Samson RA, Hong SB, Frisvad JC. 2006. Old and new concepts of species differentiation in *Aspergillus*. *Med Mycol* 44:S133–S148.
30. Sabino R, Verissimo C, Parada H, Brandao J, Viegas C, Carolino E, Clemons KV, Stevens DA. 2014. Molecular screening of 246 Portuguese *Aspergillus* isolates among different clinical and environmental sources. *Med Mycol* 52:519–529. <https://doi.org/10.1093/mmy/myu006>.
31. Gil-Serna J, Patino B, Cortes L, Gonzalez-Jaen MT, Vazquez C, Patiño B, Cortes L, Gonzalez-Jaen MT, Vazquez C. 2015. *Aspergillus steinii* and *Aspergillus westerdijkiae* as potential risk of OTA contamination in food products in warm climates. *Food Microbiol* 46:168–175. <https://doi.org/10.1016/j.fm.2014.07.013>.
32. Mikkola R, Andersson MA, Hautaniemi M, Salkinoja-Salonen MS. 2015. Toxic indole alkaloids avrainvillamide and stephacidin B produced by a biocide tolerant indoor mold *Aspergillus westerdijkiae*. *Toxicon* 99:58–67. <https://doi.org/10.1016/j.toxicon.2015.03.011>.
33. Bacha N, Atoui A, Mathieu F, Liboz T, Lebrihi A. 2009. *Aspergillus westerdijkiae* polyketide synthase gene “aoks1” is involved in the biosynthesis of ochratoxin A. *Fungal Genet Biol* 46:77–84. <https://doi.org/10.1016/j.fgb.2008.09.015>.
34. Marino A, Nostro A, Fiorentino C. 2009. Ochratoxin A production by *Aspergillus westerdijkiae* in orange fruit and juice. *Int J Food Microbiol* 132:185–189. <https://doi.org/10.1016/j.jfoodmicro.2009.03.026>.
35. Gil-Serna J, Patiño B, Cortés L, González-Jaén MT, Vázquez C. 2011. Mechanisms involved in reduction of ochratoxin A produced by *Aspergillus westerdijkiae* using *Debaryomyces hansenii* CYC 1244. *Int J Food Microbiol* 151:113–118. <https://doi.org/10.1016/j.jfoodmicro.2011.08.012>.
36. Christensen M. 1982. The *Aspergillus ochraceus* group: two new species from western soils and a synoptic key. *Mycologia* 74:210–225. <https://doi.org/10.2307/3792887>.
37. Whyte A, Gloer J. 1996. Sclerotamide: a new member of the paraherquamide class with potent antiseptic activity from the sclerotia of *Aspergillus sclerotiorum*. *J Nat Prod* 59:1093–1095. <https://doi.org/10.1021/np960607m>.
38. Whyte A, Joshi B. 2000. New cyclic peptide and bisindolyl benzenoid metabolites from the sclerotia of *Aspergillus sclerotiorum*. *J Nat Prod* 63:1006–1009. <https://doi.org/10.1021/np000103v>.
39. Zheng J, Zhu H, Hong K, Wang Y, Liu P. 2009. Novel cyclic hexapeptides from marine-derived fungus, *Aspergillus sclerotiorum* PT06-1. *Org Lett* 11:5262–5265. <https://doi.org/10.1021/ol902197z>.
40. Logrieco A, Bottalico A, Mulé G, Moretti A, Perrone G. 2003. Epidemiology of toxigenic fungi and their associated mycotoxins for some Mediterranean crops. *Eur J Plant Pathol* 109:645–667. <https://doi.org/10.1023/A:1026033021542>.
41. Taniwaki MH, Teixeira AA, Teixeira ARR, Copetti MV, Iamanaka BT. 2014. Ochratoxigenic fungi and ochratoxin A in defective coffee beans. *Food Res Int* 61:161–166. <https://doi.org/10.1016/j.foodres.2013.12.032>.
42. Trenk H, Butz M, Chu F. 1971. Production of ochratoxins in different cereal products by *Aspergillus ochraceus*. *Appl Microbiol* 21:1032–1035.
43. Steyn P, Holzapfel C. 1967. The synthesis of ochratoxins A and B: metabolites of *Aspergillus ochraceus* Wilh. *Tetrahedron* 23:4449–4461. [https://doi.org/10.1016/S0040-4020\(01\)88843-8](https://doi.org/10.1016/S0040-4020(01)88843-8).
44. Dai J, Carté B. 2001. Circumdatin G, a new alkaloid from the fungus *Aspergillus ochraceus*. *J Nat Prod* 64:125–126. <https://doi.org/10.1021/np000381u>.
45. Ghibaud G, Peano A. 2010. Chronic monolateral otomycosis in a dog caused by *Aspergillus ochraceus*. *Vet Dermatol* 21:522–526. <https://doi.org/10.1111/j.1365-3164.2010.00884.x>.
46. Siqueira JPZ, Sutton DA, García D, Gené J, Thomson P, Wiederhold N, Guarro J. 2016. Species diversity of *Aspergillus* section *Vesicicolores* in clinical samples and antifungal susceptibility. *Fungal Biol* 120: 1458–1467. <https://doi.org/10.1016/j.funbio.2016.02.006>.
47. Jessup CJ, Ryder NS, Ghannoum MA. 2000. An evaluation of the *in vitro* activity of terbinafine. *Med Mycol* 38:155–159. <https://doi.org/10.1080/mmy.38.2.155.159>.
48. Moore CB, Walls CM, Denning DW. 2001. *In vitro* activities of terbinafine against *Aspergillus* species in comparison with those of itraconazole and amphotericin B. *Antimicrob Agents Chemother* 45:1882–1885. <https://doi.org/10.1128/AAC.45.6.1882-1885.2001>.

49. Fernández MS, Rojas FD, Cattana ME, Sosa Mde L, Iovannitti CA, Lass-Flörl C, Giusiano GE. 2015. *In vitro* activities of amphotericin B, terbinafine, and azole drugs against clinical and environmental isolates of *Aspergillus terreus sensu stricto*. *Antimicrob Agents Chemother* 59: 3619–3622. <https://doi.org/10.1128/AAC.00045-15>.
50. Walsh TJ, Anaissie EJ, Denning DW, Herbrecht R, Kontoyiannis DP, Marr KA, Morrison VA, Segal BH, Steinbach WJ, Stevens DA, van Burik J-A, Wingard JR, Patterson TF. 2008. Treatment of aspergillosis: clinical practice guidelines of the Infectious Diseases Society of America. *Clin Infect Dis* 46:327–360. <https://doi.org/10.1086/525258>.
51. Masih A, Singh PK, Kathuria S, Agarwal K, Meis JF, Chowdhary A. 2016. Identification by molecular methods and matrix-assisted laser desorption ionization–time of flight mass spectrometry and antifungal susceptibility profiles of clinically significant rare *Aspergillus* species in a referral chest hospital in Delhi, India. *J Clin Microbiol* 54:2354–2364. <https://doi.org/10.1128/JCM.00962-16>.
52. White T, Bruns T, Lee S, Taylor J. 1990. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics, p 315–322. *In* Innis MA, Gelfand DH, Sninsky JJ, White TJ (ed), *PCR protocols: a guide to methods and applications*. Academic Press Inc., New York, New York.
53. Glass NL, Donaldson GC. 1995. Development of primer sets designed for use with the PCR to amplify conserved genes from filamentous ascomycetes. *Appl Environ Microbiol* 61:1323–1330.
54. Hong S-B, Go S-J, Shin H-D, Frisvad JC, Samson RA. 2005. Polyphasic taxonomy of *Aspergillus fumigatus* and related species. *Mycologia* 97: 1316–1329. <https://doi.org/10.3852/mycologia.97.6.1316>.
55. Liu YJ, Whelen S, Hall BD. 1999. Phylogenetic relationships among Ascomycetes: evidence from an RNA Polymerase II Subunit. *Mol Biol Evol* 16:1799–1808. <https://doi.org/10.1093/oxfordjournals.molbev.a026092>.
56. Farris J, Källersjö M, Kluge A, Bult C. 1994. Testing significance of incongruence. *Cladistics* 10:315–319. <https://doi.org/10.1111/j.1096-0031.1994.tb00181.x>.
57. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* 30: 2725–2729. <https://doi.org/10.1093/molbev/mst197>.
58. Ronquist F, Huelsenbeck JP. 2003. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19:1572–1574. <https://doi.org/10.1093/bioinformatics/btg180>.
59. Nylander JAA. 2004. MrModeltest v2. Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden.
60. Clinical and Laboratory Standards Institute. 2008. Reference method for broth dilution antifungals susceptibility testing of conidium-forming filamentous fungi: approved standard, 2nd ed. M38-A2. CLSI, Wayne, PA.